



Download this poster:



tinyurl.com/idbac2019

IDBac Website



chasemc.github.io/IDBac

Using **IDBac** to Investigate the Microbial and Natural Product Potential of Freshwater Sponges

Chase M. Clark, Antonio Hernandez, Jeongho Lee, Michael Mallowney, Maria S. Costa, Brian T. Murphy

Department of Pharmaceutical Sciences, Center for Biomolecular Sciences, College of Pharmacy,


University of Illinois at Chicago, Illinois 60612, United States

**PHARMACEUTICAL
SCIENCES
COLLEGE
OF PHARMACY**




CBS Center for Biomolecular Sciences
College of Pharmacy


Freshwater Sponge Microbiomes are an Understudied Resource



Since 2015 our lab and citizen scientist divers have collected 90 freshwater sponges across the Great Lakes. For this antibiotic-discovery study we chose two *Eunapius fragilis* var. *minuta* sponges collected one mile apart.




New IDBac Algorithms Automate the Creation of Small Highly-Diverse Microbial Libraries



[3] Costa MS, Clark CM, Omarsdottir S, Sanchez LM, Murphy BT. Minimizing Taxonomic and Natural Product Redundancy in Microbial Libraries using MALDI-TOF MS and the bioinformatics pipeline IDBac. *J Nat Prod*. In Press.

[4] Automation: Unpublished- Chase M Clark

Deprioritize Bioactive Isolates Producing Known Antibiotics




The red line represents the theoretical MS¹ spectrum of Rifamycin W. Grey bars represent experimental data. LC-MS/MS data acquired using UHPLC(C18)-QTOF.


Before scaling cultures, we screened bioactive colonies from agar for commonly-isolated antibiotics using UHPLC-QTOF and GNPS^[7]. In this case isolate 1043, while active, also produced a well-known antibiotic, rifamycin W, and was de-prioritized.

[7] Wang, Mingxun, et al. "Sharing and community curation of mass spectrometry data with Global Natural Products Social Molecular Networking." *Nature Biotechnology* 34.8 (2016): 828-837.


Every Distinguishable Colony of a Cultured Microbiome is Isolated



Sponge-associated bacteria are cultivated using diverse sample treatments and media.




Isolated 900 bacteria and inoculated onto a single media type.



Sample preparation and data collection for ~125 isolates x 3 replicates can be acquired in 4 hours.


[1] Clark CM, Costa MS, Conley E, Li E, Sanchez LM, Murphy BT. Using the Open-Source MALDI-TOF-MS IDBac Pipeline for Analysis of Microbial Protein and Specialized Metabolite Data. *J Vis Exp*. 2019;(147):e59219. doi:10.3791/59219



Tentatively Identify Prioritized Bacteria with IDBac Databases of Known Organisms


IDBac v1.0 or greater allows users to create shareable reference databases of known bacteria. Upcoming releases will include the ability to rapidly query spectra of unknowns against these databases.

We are working to make our database of 110 genera available soon (few replicates) and have begun digitizing a portion of the NRRL microbial library (high-quality, eight biological replicates).





[5] Unpublished- Chase M Clark;

Available: github.com/chasemc/maldiSearch



Remaining Prioritized Isolates Undergo Isolation and Elucidation

Another freshwater isolate, 112, exhibited bioactivity but dereplication has afforded no known antibiotics. Isolation and elucidation efforts are underway.




Above: A GNPS Feature-Based Molecular Network of isolate fractions and controls is used to quickly compare and prioritize LC-MS/MS features for further isolation and elucidation.^[8]

Left: Active and non-active C18-SPE fractions from isolate 112 are compared for differential occurrence/abundance of LC-MS/MS features.^[9]

[8] Nothias L-F, Nothias-Esposto M, da Silva R, et al. Bioactivity-Based Molecular Networking for the Discovery of Drug Leads in Natural Product Bioassay-Guided Fractionation. *J Nat Prod*. 2018;81(4):758-767. doi:10.1021/acs.jnatprod.7b00737


[9] Pluskal T, Castillo S, Villar-Briones A, Orešič M. MZmine 2: Modular framework for processing, visualizing, and analyzing mass spectrometry-based molecular profile data. *BMC Bioinformatics*. 2010;11(1):395. doi:10.1186/1471-2105-11-395

MALDI-TOF MS Data is Collected for Protein and Specialized Metabolite Mass Regions



Protein Data

Allows IDBac to create pseudo-phylogenetic groups based on Matrix Assisted Laser Desorption Time of Flight Mass Spectrometry (MALDI-TOF MS) spectra. These groupings are highly specific and correlate well to 16S rRNA gene groupings.




Specialized Metabolite Data

Once isolates are grouped using protein data, IDBac is used to create sub-groups of isolates with similar specialized metabolite MALDI-TOF MS profiles.

[2] Clark CM, Costa MS, Sanchez LM, Murphy BT. Coupling MALDI-TOF mass spectrometry protein and specialized metabolite analyses to rapidly discriminate bacterial function. *Proc Natl Acad Sci U S A*. 2018;115(19):4981-4986. doi:10.1073/pnas.1801247115



Further Prioritize the Highly-Diverse Microbial Library Using a Custom Agar-Based OSMAC Screen



Results of a screen of isolate 1043, which was found to inhibit *P. aeruginosa*. For more information on this assay, see poster P-016.


An agar-based OSMAC screen against the fluorescently-labelled pathogen *Pseudomonas aeruginosa* 10145GFP (among other ESKAPE pathogens) was used to further prioritize isolates for antibiotic discovery.

Some benefits:

- Screen doesn't require liquid culture.
- GFP result confirmed with a secondary MTT assay
- Differential activity in different growth conditions is easily exposed.
- 3D-printed 96-well plates are autoclavable and reusable.

[6] Unpublished

Future Directions



Antibiotic Discovery:

In addition to elucidating the bioactive compound(s) from isolate 112, we are preparing to screen all 120 prioritized sponge isolates against a panel of ESKAPE pathogens using our newly-developed agar-based OSMAC assay.

Freshwater Sponge Biogeography:

Utilizing existing, and not-yet-released IDBac analyses, we are working to understand the biogeographic distribution of Great Lakes freshwater sponges (map above), their associated bacteria, and bacterial specialized metabolites.

Software Links:

chasemc.github.io/IDBac

chasemc.github.io/mzEasy

github.com/chasemc/mzPlotter

github.com/chasemc/electricShine

Funding:



Grant CP-044R-17

Research reported in this presentation was supported by the National Institute Of General Medical Sciences of the National Institutes of Health under Award Number R01GM125943 and by the National Center for Complementary Health & Integrative Health of the National Institutes of Health under Award Number F31AT010419. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.