

Summer School in Quantitative Fisheries Stock Assessment

Day 4: An introduction to Statistical Catch-at-Age models

Cóilín Minto, Chato Osio, Alessandro Orio, Alessandro Ligas

July 20, 2017

Outline

Goal

Build a fully statistical model to estimate the parameters of the stock assessment


Cohort

Cohort progression

	Age →				
	$N_{a1,t1}$	$N_{a2,t1}$	$N_{a3,t1}$	$N_{a4,t1}$	$N_{a5,t1}$
	$N_{a1,t2}$	$N_{a2,t2}$	$N_{a3,t2}$	$N_{a4,t2}$	$N_{a5,t2}$
	$N_{a1,t3}$	$N_{a2,t3}$	$N_{a3,t3}$	$N_{a4,t3}$	$N_{a5,t3}$
	$N_{a1,t4}$	$N_{a2,t4}$	$N_{a3,t4}$	$N_{a4,t4}$	$N_{a5,t4}$
Time ↓	$N_{a1,t5}$	$N_{a2,t5}$	$N_{a3,t5}$	$N_{a4,t5}$	$N_{a5,t5}$


Cohort

Cohort progression



Cohort

Cohort progression



$$N_{a+1,t+1} = N_{a,t} e^{-(F_{a,t} + M_{a,t})}$$

Starting a cohort

But how do we start them?

	Age →				
Time ↓	$N_{a1,t1}$	$N_{a2,t1}$	$N_{a3,t1}$	$N_{a4,t1}$	$N_{a5,t1}$
	$N_{a1,t2}$	$N_{a2,t2}$	$N_{a3,t2}$	$N_{a4,t2}$	$N_{a5,t2}$
	$N_{a1,t3}$	$N_{a2,t3}$	$N_{a3,t3}$	$N_{a4,t3}$	$N_{a5,t3}$
	$N_{a1,t4}$	$N_{a2,t4}$	$N_{a3,t4}$	$N_{a4,t4}$	$N_{a5,t4}$
	$N_{a1,t5}$	$N_{a2,t5}$	$N_{a3,t5}$	$N_{a4,t5}$	$N_{a5,t5}$


Year one

We also need to start all ages in the first year

	Age →				
	$N_{a1,t1}$	$N_{a2,t1}$	$N_{a3,t1}$	$N_{a4,t1}$	$N_{a5,t1}$
Time ↓	$N_{a1,t2}$	$N_{a2,t2}$	$N_{a3,t2}$	$N_{a4,t2}$	$N_{a5,t2}$
	$N_{a1,t3}$	$N_{a2,t3}$	$N_{a3,t3}$	$N_{a4,t3}$	$N_{a5,t3}$
	$N_{a1,t4}$	$N_{a2,t4}$	$N_{a3,t4}$	$N_{a4,t4}$	$N_{a5,t4}$
	$N_{a1,t5}$	$N_{a2,t5}$	$N_{a3,t5}$	$N_{a4,t5}$	$N_{a5,t5}$

Year one

We also need to start all ages in the first year



How about F ?

	Age				
Time ↓	$F_{a1,t1}$	$F_{a2,t1}$	$F_{a3,t1}$	$F_{a4,t1}$	$F_{a5,t1}$
	$F_{a1,t2}$	$F_{a2,t2}$	$F_{a3,t2}$	$F_{a4,t2}$	$F_{a5,t2}$
	$F_{a1,t3}$	$F_{a2,t3}$	$F_{a3,t3}$	$F_{a4,t3}$	$F_{a5,t3}$
	$F_{a1,t4}$	$F_{a2,t4}$	$F_{a3,t4}$	$F_{a4,t4}$	$F_{a5,t4}$
	$F_{a1,t5}$	$F_{a2,t5}$	$F_{a3,t5}$	$F_{a4,t5}$	$F_{a5,t5}$

How about F ?

Age

	$F_{a1,t1}$	$F_{a2,t1}$	$F_{a3,t1}$	$F_{a4,t1}$	$F_{a5,t1}$
	$F_{a1,t2}$	$F_{a2,t2}$	$F_{a3,t2}$	$F_{a4,t2}$	$F_{a5,t2}$
	$F_{a1,t3}$	$F_{a2,t3}$	$F_{a3,t3}$	$F_{a4,t3}$	$F_{a5,t3}$
	$F_{a1,t4}$	$F_{a2,t4}$	$F_{a3,t4}$	$F_{a4,t4}$	$F_{a5,t4}$
	$F_{a1,t5}$	$F_{a2,t5}$	$F_{a3,t5}$	$F_{a4,t5}$	$F_{a5,t5}$

Time \longrightarrow

That's a lot of parameters!

How about F ?

Need to conserve degrees of freedom in the model but also need the \mathbf{F} matrix. One approach is a separable model

$$\mathbf{F} = \begin{pmatrix} s_{a1} & s_{a2} & s_{a3} & s_{a4} & s_{a5} \\ s_{a1} & s_{a2} & s_{a3} & s_{a4} & s_{a5} \\ s_{a1} & s_{a2} & s_{a3} & s_{a4} & s_{a5} \\ s_{a1} & s_{a2} & s_{a3} & s_{a4} & s_{a5} \\ s_{a1} & s_{a2} & s_{a3} & s_{a4} & s_{a5} \end{pmatrix} \times \begin{pmatrix} f_{t1} & f_{t1} & f_{t1} & f_{t1} & f_{t1} \\ f_{t2} & f_{t2} & f_{t2} & f_{t2} & f_{t2} \\ f_{t3} & f_{t3} & f_{t3} & f_{t3} & f_{t3} \\ f_{t4} & f_{t4} & f_{t4} & f_{t4} & f_{t4} \\ f_{t5} & f_{t5} & f_{t5} & f_{t5} & f_{t5} \end{pmatrix}$$

How about F ?

Need to conserve degrees of freedom in the model but also need the \mathbf{F} matrix. One approach is a separable model

$$\mathbf{F} = \begin{pmatrix} s_{a1}f_{t1} & s_{a2}f_{t1} & s_{a3}f_{t1} & s_{a4}f_{t1} & s_{a5}f_{t1} \\ s_{a1}f_{t2} & s_{a2}f_{t2} & s_{a3}f_{t2} & s_{a4}f_{t2} & s_{a5}f_{t2} \\ s_{a1}f_{t3} & s_{a2}f_{t3} & s_{a3}f_{t3} & s_{a4}f_{t3} & s_{a5}f_{t3} \\ s_{a1}f_{t4} & s_{a2}f_{t4} & s_{a3}f_{t4} & s_{a4}f_{t4} & s_{a5}f_{t4} \\ s_{a1}f_{t5} & s_{a2}f_{t5} & s_{a3}f_{t5} & s_{a4}f_{t5} & s_{a5}f_{t5} \end{pmatrix}$$

Separable model

So the parameters we have in the model are:

- $N_{1,t=1\dots T}$ number of recruits
- $N_{a=2\dots A,t=1}$ numbers in first year
- $s_{a=1\dots A}$ exploitation pattern (age effects)
- $f_{t=1\dots T}$ fishing mortality level (year effects)

Predictions

From this set of parameters we can predict $\hat{\mathbf{F}}$ and $\hat{\mathbf{N}}$ from which we can predict catch

$$\hat{C}_{a,t} = \frac{\hat{F}_{a,t}}{\hat{F}_{a,t} + M_{a,t}} \left(1 - e^{-(\hat{F}_{a,t} + M_{a,t})} \right) \hat{N}_{a,t}$$

Or, with the addition of catchability, predict a survey index

$$\hat{I}_{a,t} = q_{a,t} \hat{N}_{a,t}$$

Estimation

We observe $C_{a,t}$, $I_{a,t}$ so in principle we can estimate the parameters by maximising the likelihood of the parameters given the data

$$L(\mathbf{N}_{a=1}, \mathbf{N}_{t=1}, \mathbf{s}, \mathbf{f}, \mathbf{q}, \boldsymbol{\sigma} | \mathbf{C}, \mathbf{I}) = \prod_{a=1}^A \prod_{t=1}^T (\mathcal{N}(\ln(I_{a,t}) - \ln(\hat{I}_{a,t}), \sigma_I^2) \\ \times \mathcal{N}(\ln(C_{a,t}) - \ln(\hat{C}_{a,t}), \sigma_C^2))$$

We maximise the (log) likelihood to estimate the best fitting parameters of the model given the data!

intro to splines