

LECTURE 4: Counting

Discrete uniform law


- Assume Ω consists of n equally likely elements
- Assume A consists of k elements

Then :
$$P(A) = \frac{\text{number of elements of } A}{\text{number of elements of } \Omega} = \frac{k}{n}$$

- Basic counting principle

- Applications

| | |
|--------------|------------------------|
| permutations | number of subsets |
| combinations | binomial probabilities |
| partitions | |



Basic counting principle

4 shirts


3 ties

2 jackets

Number of possible attires?

- r stages
- n_i choices at stage i

Number of choices is: $n_1 \cdot n_2 \cdots n_r$




Basic counting principle examples

- Number of license plates with 2 letters followed by 3 digits:

$$26 \cdot 26 \cdot 10 \cdot 10 \cdot 10$$

- ... if repetition is prohibited: $26 \cdot 25 \cdot 10 \cdot 9 \cdot 8$

- Permutations:** Number of ways of ordering n elements:



$$n \cdot (n-1) \cdot (n-2) \cdot \dots \cdot 1 = n!$$

- Number of subsets of $\{1, \dots, n\}$:



$$2 \cdot 2 \cdot \dots \cdot 2 = 2^n$$

$$n=1 \quad \{1\} \quad 2^1=2$$
$$\{1\} \quad \emptyset$$

Example

- Find the probability that:
six rolls of a (six-sided) die all give different numbers.

$\swarrow A$

(Assume all outcomes equally likely.)

typical outcome

$$P(2, 3, 4, 3, 6, 2) = 1/6^6$$

" element of A :

$$(2, 3, 4, 1, 6, 5) = 6!$$

$$P(A) = \frac{\# \text{ in } A}{\# \text{ possible outcomes}} = \frac{6!}{6^6}.$$

Combinations


- Definition: $\binom{n}{k}$: number of k -element subsets of a given n -element set

$$= \frac{n!}{k!(n-k)!}$$



$$n = 0, 1, 2, \dots$$

- Two ways of constructing an **ordered** sequence of k **distinct** items: $k = 0, 1, \dots, n$
 - Choose the k items one at a time
 - Choose k items, then order them



$$\binom{n}{k} = \frac{n!}{k!(n-k)!}$$

$$\binom{n}{n} = 1 \quad \frac{n!}{n! \cdot 0!}$$

$0! = 1$ convention

$$\binom{n}{0} = \frac{n!}{0! \cdot n!} = 1 \quad \emptyset$$

$$\sum_{k=0}^n \binom{n}{k} = \binom{n}{0} + \binom{n}{1} + \dots + \binom{n}{n} = \# \text{ all subsets} = 2^n$$

Binomial coefficient $\binom{n}{k}$ \longrightarrow **Binomial probabilities**

- $n \geq 1$ independent coin tosses; $P(H) = p$


$$P(k \text{ heads}) = \binom{n}{k} p^k (1-p)^{n-k}$$

- $n=6$
 $P(HTTTHH) = p(1-p)(1-p)ppp = p^4(1-p)^2$

- $P(\text{particular sequence}) = p^{\# \text{ heads}} (1-p)^{\# \text{ tails}}$

- $P(\text{particular } k\text{-head sequence}) = p^k (1-p)^{n-k}$

$$P(k \text{ heads}) = p^k (1-p)^{n-k} \cdot (\# \text{ } k\text{-head sequences})$$



$$\binom{n}{k}$$

A coin tossing problem

- Given that there were 3 heads in 10 tosses, what is the probability that the first two tosses were heads?

- event A : the first 2 tosses were heads
- event B : 3 out of 10 tosses were heads

Assumptions:

- independence
- $P(H) = p$

$$P(k \text{ heads}) = \binom{n}{k} p^k (1-p)^{n-k}$$

- First solution:

$$P(A | B) = \frac{P(A \cap B)}{P(B)} = \frac{P(H_1, H_2 \text{ and one } H \text{ in tosses } 3, \dots, 10)}{P(B)}$$

$$= \frac{p^2 \cdot \binom{8}{1} p^1 \cdot (1-p)^7}{\binom{10}{3} p^3 (1-p)^7} = \frac{\binom{8}{1}}{\binom{10}{3}} = \frac{8}{\binom{10}{3}} \cdot$$

A coin tossing problem


- Given that there were 3 heads in 10 tosses, what is the probability that the first two tosses were heads?
 - event A : the first 2 tosses were heads
 - event B : 3 out of 10 tosses were heads

Assumptions:

- independence
- $P(H) = p$

$$P(k \text{ heads}) = \binom{n}{k} p^k (1-p)^{n-k}$$


- Second solution: Conditional probability law (on B) is uniform



Partitions

- $n \geq 1$ distinct items; $r \geq 1$ persons give n_i items to person i
 - here n_1, \dots, n_r are given nonnegative integers
 - with $n_1 + \dots + n_r = n$
- Ordering n items: $n!$
 - Deal n_i to each person i , and then order

$$C \ n_1! \ n_2! \ \dots \ n_r! = n!$$



$$r=2 \quad n_1=k \quad n_2=n-k$$

$$\text{number of partitions} = \frac{n!}{n_1! n_2! \dots n_r!} \quad (\text{multinomial coefficient})$$

Example: 52-card deck, dealt (fairly) to four players.
Find $P(\text{each player gets an ace})$

- Outcomes are: *partition equally likely*
 - number of outcomes: $\frac{52!}{13! 13! 13! 13!}$
- Constructing an outcome with one ace for each person:
 - distribute the aces $4 \cdot 3 \cdot 2 \cdot 1$
 - distribute the remaining 48 cards $\frac{48!}{12! 12! 12! 12!}$

- Answer:
$$\frac{4 \cdot 3 \cdot 2 \cdot \frac{48!}{12! 12! 12! 12!}}{\frac{52!}{13! 13! 13! 13!}}$$

Example: 52-card deck, dealt (fairly) to four players.
Find $P(\text{each player gets an ace})$

A smart solution

Stack the deck, aces on top

