

Behavioral and biochemical effects of ethanol withdrawal in zebrafish

Suianny Nayara da Silva Chaves¹, Gabriel Felício Rocha¹, Bruna Patrícia Dutra Costa¹, Witallo Etevaldo Araújo de Oliveira¹, Monica Gomes Lima², Diógenes Henrique de Siqueira Silva¹, Caio Maximino^{1,3*}

¹ Laboratório de Neurociências e Comportamento, Instituto de Estudos em Saúde e Biológicas, Universidade Federal do Sul e Sudeste do Pará

² Laboratório de Neurociências e Comportamento “Frederico Guilherme Graeff”, Campus VIII, Universidade do Estado do Pará

³ Programa de Pós-Graduação em Neurociências e Comportamento, Universidade Federal do Pará

*** Corresponding author:**

Caio Maximino

Laboratório de Neurociências e Comportamento – Instituto de Estudos em Saúde e Biológicas, Universidade Federal do Sul e Sudeste do Pará, Unidade III

Av. dos Ipês, S/N, s/ CEP, Bairro Cidade Jardim, Marabá/PA, Brazil

cmaximino@unifesspa.edu.br

Abstract: Chronic alcohol use induces adaptations and toxicity that can induce symptoms of anxiety, autonomic hyperarousal, and seizures when alcohol is removed (withdrawal syndrome). Zebrafish has recently gained wide attention as a behavioral model to study the neurobehavioral effects of acute and chronic alcohol use, including withdrawal. The literature, however, is very contradictory on findings regarding withdrawal effects, with some studies reporting increased anxiety, while others report no effect. A meta-analytic approach was taken to find the sources of this heterogeneity, and ethanol concentration during exposure and exposure duration were found to be the main sources of variation. A conceptual replication was also made using continuous exposure for 16 days in waterborne ethanol (0.5%) and assessing anxiety-like behavior in the light/dark test after 60 min withdrawal. Withdrawal was shown to reduce preference for darkness, consistent with decreased anxiety, but to increase risk assessment, consistent with increased anxiety. Animals were also subjected to the withdrawal protocol and injected with pilocarpine in a sub-convulsive dose to assess susceptibility to seizures. The protocol was sufficient to increase susceptibility to seizures in animals exposed to ethanol. Finally, withdrawal also decreased catalase activity in the brain, but not in the head kidney, suggesting that oxidative stress could be a mechanism associated with the behavioral effects of ethanol withdrawal.

Keywords: Anxiety; *Danio rerio*; Ethanol withdrawal; Oxidative stress; Seizures.

1. Introduction

Chronic alcohol (ethanol, EtOH) use produces adaptations and toxicity that can lead to tolerance and dependence, manifested as physical and mental distress when EtOH is removed (withdrawal); symptoms of ethanol withdrawal include symptoms of anxiety, insomnia, and autonomic hyperarousal (Krystal & Tabakoff, 2002). In more serious conditions, patients presenting this EtOH withdrawal syndrome can present perceptive changes, agitation, mental confusion, significant increases in autonomic arousal, and seizures (Gatch & Lal, 2001; Krystal & Tabakoff, 2002). The most serious condition involves *delirium tremens* and death by hyperthermia, cardiac arrhythmia, and complications from withdrawal-induced seizures (Longo & Schuckit, 2014). Since withdrawal symptoms are usually reduced after EtOH consumption, EtOH dependence can be maintained by negative reinforcement (Koob & Le Moal, 2008), and therefore investigating these motivational mechanisms could open new avenues for the treatment of EtOH consumption-related disorders.

In animal models, EtOH withdrawal changes the excitability of neurons located in brain regions associated with defensive behavior, anxiety, and fear (Bonassoli, Milani, & Oliveira, 2011; Chakravarty & Faingold, 1998; Long, Yang, Faingold, & Evans, 2007; L Yang, Long, Randall, & Faingold, 2003; Li Yang, Long, Evans, & Faingold, 2002; Li Yang, Long, & Faingold, 2001). Moreover, EtOH withdrawal also dysregulates the activity of the hypothalamus-pituitary-adrenal axis that modulates behavioral and endocrine responses to stress (Rasmussen, Boldt, Wilkinson, & Mitton, 2002). It makes sense, then, that the majority of animal models of EtOH withdrawal are focused on anxiety-like behavior, with consistent effects observed in rodent models such as the elevated plus-maze, light/dark box, social interaction test, and a drug discrimination assay using pentylentetrazole (Gatch & Lal, 2001).

Zebrafish (*Danio rerio*) species is increasingly being considered as useful model organisms for studying both behavioral genetics and behavioral neuroscience, neuropsychopharmacology, and neurotoxicology (Bonan & Norton, 2015; Kalueff et al., 2012; Norton & Bally-Cuif, 2010; Stewart

et al., 2015). The main advantages associated with this species are its low cost of acquisition and upkeep, ease handling, short lifespan, and readiness of reproduction in laboratory environments (Gerlai, 2014; Kalueff, Echevarria, & Stewart, 2014). The relatively high degree of genetic, neural, and endocrine homology with rodents and human beings is also cited as an advantage (Kokel & Peterson, 2008). Zebrafish is also a good model for studying anxiety and stress, with well-validated assays for novelty- and conflict-induced anxiety, social interaction, and antipredatory behavior (Gerlai, 2010; Maximino, Marques, et al., 2010; Oliveira, 2013). Importantly for EtOH withdrawal, seizure-like phenotypes were also characterized in the species (Hortopan, Dinday, & Baraban, 2010).

Zebrafish anxiety-like behavior has been used to demonstrate the effects of drug withdrawal, including cocaine (López-Patiño, Yu, Cabral, & Zhdanova, 2008; López-Patiño, Yu, Yamamoto, & Zhdanova, 2008), morphine (Cachat et al., 2010; Khor, Jamil, Adenan, & Shu-Chien, 2011; Wong et al., 2010), and EtOH (S. Tran, Facciol, & Gerlai, 2016). In that last case, the literature is inconsistent, with some studies (e.g., Tran et al., 2015) reporting significant effects of withdrawal on anxiety-like behavior, while others (e. g., Cachat et al., 2010) were unable to detect effects of EtOH withdrawal. Procedural differences, such as assay type, strain, concentration during exposure, or withdrawal duration, could be responsible for this difference.

In addition to these behavioral endpoints, neurochemical analyses were also reported in zebrafish models of EtOH withdrawal (Müller, Nunes, Silveira, Loro, & Rosemberg, 2017; Steven Tran et al., 2015a). After exposure to 0.5% EtOH for 22 days, increases in brain levels of dopamine, serotonin, and aspartate were observed with 60 min withdrawal (Gerlai, Chatterjee, Pereira, Sawashima, & Krishnannair, 2009). After exposure for 8 days to 1% EtOH (for 20 min per day), decreases in superoxide dismutase and catalase activity and consequent increases in oxidative stress (Müller et al., 2017). While a mechanistic explanation is still lacking, these results are congruent with the hyperexcitability and EtOH-induced neurotoxicity observed in other models (Krystal & Tabakoff, 2002; Zenki, Mussulini, Rico, de Oliveira, & Rosemberg, 2014).

The aim of the present work is to study the heterogeneity of effects of EtOH withdrawal on zebrafish anxiety-like behavior in the literature, by applying meta-analytical techniques. Moreover, the present work attempts a conceptual replication of findings on EtOH withdrawal by assessing the effects of a withdrawal protocol on behavior in the light/dark test. It also attempts to expand the breadth of endpoints used in withdrawal research by using a chemically-induced seizure model with sub-convulsive doses. Finally, the present work also attempted to replicate previous research on oxidative stress in EtOH withdrawal by analyzing the activity of the enzyme catalase.

2. Methods

2.1 Systematic review and metanalysis

The protocol for the meta-analysis was pre-registered in the CAMARADES-NC3Rs Preclinical Systematic Review & Meta-analysis Facility (SyRF) database (<https://drive.google.com/file/d/0B7Z0eAxKc8ApUjcyQjhwVnFjRFE/view?usp=sharing>). No modification from the pre-registered protocol was made. Article with the descriptors ‘ethanol withdrawal’ and ‘zebrafish’ were searched for in PubMed (<https://www.ncbi.nlm.nih.gov/pubmed>), using a search filter optimized to finding studies on animal experimentation on PubMed (Hooijmans, Tillema, Leenaars, & Ritskes-Hoitinga, 2010). Bibliographic data (including DOI, publication date, title, and abstract) from the studies identified in the systematic review were exported to a spreadsheet. Each article from the list was reviewed in four levels of detail (title, abstract, full text, and a detailed revision of the experimental design) in order to determine its eligibility to meta-analysis. Following Mohammad et al. (2016), studies should include (1) primary behavioral data obtained in tests for anxiety-like behavior in zebrafish (light/dark test, novel tank test, antipredator responses, shoaling responses); (2) reporting of appropriate controls; and (3) reporting of at least sample sizes and summary statistics (central tendency and dispersion measures) for control and withdrawal groups. Only control and withdrawal groups were considered in experiments that evaluated the effects of drugs or other interventions on behavior under withdrawal.

Possible confounds in relation to the role of development were reduced by excluding studies which were not performed on adult fish.

The following data were extracted from each included study: identification (DOI, authors, publication year); strain/phenotype; concentration of ethanol during exposure; duration of EtOH treatment; duration of withdrawal; behavioral test that was used; means and standard deviations, as well as a test statistics and degrees of freedom; and sample sizes (N) for each group. Data, which were represented graphically, were extracted from figures using PlotDigitizer (<http://plotdigitizer.sourceforge.net/>). When multiple dependent variables were reported, only the primary endpoint was used (time on white, time on bottom and distance from stimulus). All estimates were transformed to standardized mean differences (SMD), corrected for its positive bias (Hedges & Olkin, 1985), with unbiased estimates of sampling variances and confidence intervals at the 95% level, I^2 and τ^2 heterogeneity values, and p-values using a mixed-effects model, with concentration, exposure duration, and withdrawal duration used as moderators. Influential case diagnostics was made by inspecting plots for externally standardized residues, DFFITS values, Cook's distances, covariance ratios, estimates of τ^2 and test statistics for residual heterogeneity when each study is removed in turn, hat values, and weights for each study included in the analysis. Publication bias was assessed by inspection of a contour-enhanced funnel plot, with contours at the 90%, 95% and 99% confidence intervals. Moreover, funnel plot asymmetry was analyzed using a meta-regression test, with total samples size as predictor (Egger, Smith, Schneider, & Minder, 1997). Observed power for each study was calculated based on effect sizes, sample sizes and standard deviations, and fitted against SMDs by a generalized additive model with Gaussian curve family and identity link function. Finally, a sensitivity analysis was performed by adding study quality, assessed using SYRCLE's Risk of Bias (RoB) tool (Hooijmans et al., 2014), in the meta-regression model. The meta-analysis was made with the metafor package from R (Viechtbauer, 2010).

2.2. *Effects of EtOH withdrawal on anxiety-like behavior and seizure susceptibility in zebrafish*

2.1.1. Animals, housing, and baseline characteristics

Outbred populations were used due to their increased genetic variability, decreasing the effects of random genetic drift which could lead to the development of uniquely heritable traits (Parra, Adrian Jr, & Gerlai, 2009; Speedie & Gerlai, 2008). Thus, the animals used in the experiments are expected to better represent the natural populations in the wild. Twenty-four-adult zebrafish from the wildtype strain (longfin phenotype) were used in this experiment. Animals were bought from a commercial seller, and arrived in the laboratory with 3 months of age, approximately (standard length = 13.2 ± 1.4 mm), and were quarantined for two weeks; the experiment begun when animals had an approximate age of 4 months (standard length = 23.0 ± 3.2 mm). Animals were kept in mixed-sex tanks during acclimation, with an approximate ratio of 50 male: 50 female. The breeder was licensed for aquaculture under Ibama's (Instituto Brasileiro do Meio Ambiente e dos Recursos Naturais Renováveis) Resolution 95/1993. Animals were group-housed in 40 L tanks, with a maximum density of 25 fish per tank, for at least 2 weeks before experiments begun. Tanks were filled with non-chlorinated water at room temperature (28°C) and a pH of 7.0-8.0. Lighting was provided by fluorescent lamps in a cycle of 14-10 hours (LD), according to standards of care for zebrafish (Lawrence, 2007). Water quality parameters were as follow: pH 7.0-8.0; hardness 100-150 mg/L CaCO₃; dissolved oxygen 7.5-8.0 mg/L; ammonia and nitrite < 0.001 ppm. All manipulations minimized their potential suffering of animals, and followed Brazilian legislation (Conselho Nacional de Controle de Experimentação Animal - CONCEA, 2017). Animals were used for only one experiment and in a single behavioral test, to reduce interference from apparatus exposure.

2.2.2. Ethanol exposure and withdrawal

The EtOH exposure and withdrawal regimen was adapted from Gerlai et al. (2009). In brief, animals were exposed to increasing EtOH concentrations (0.125%-0.5%), dispersed on the tank

water, for 16 days, therefore decreasing mortality that is associated with prolonged exposure to EtOH in zebrafish. Concentrations doubled every four days, reaching a final concentration of 0.5% (v/v). Animals were kept in this final concentration for 4 days. Another group was exposed to the same manipulation, without EtOH exposure; animals were randomly allocated to treatments *via* generation of random numbers using the randomization tool in <http://www.randomization.com/>. Caretakers were blinded for treatment. After treatment, animals were transferred to a tank with system water for 60 min (withdrawal stage). All animals were included in the experiments, and no gross physical abnormalities were observed during the exposure period.

2.2.3. Light/dark test

After EtOH withdrawal, animals (10 animals in the control group, 14 in the withdrawal group) were individually tested in a tank (15 cm height X 10 cm width X 45 cm length) that was divided in half into a black compartment and a white compartment. The tank was made of matte acrylic. The apparatus was illuminated from above by two fluorescent 25 W lamps, which produced an average of 270 lumens above the tank (light levels were measured using a hand photometer). The tank contained sliding doors that defined a central compartment in which the animal was positioned for a 3-min acclimation. After this stage, the sliding doors were removed, allowing the animal to freely explore the apparatus for 15 min.

The order with which animals were exposed to the tank was randomized and balanced across treatments. Experimenters were blinded to treatment. Videos were manually transcribed by two observers, blinded to treatment, using X-Plo-Rat (<https://github.com/lanec-unifesspa/x-plo-rat>). The following variables were analyzed: time on the white compartment (s); transitions to white; duration of entries in the white compartment (total duration divided by the number of transitions); time freezing (s); number of erratic swimming events; time in thigmotaxis (s); number of risk assessment events. Operational definitions of these endpoints can be found in Table 1.

Data were analyzed via approximative two-sample Fisher-Pitman permutation tests with 10.000 Monte-Carlo re-samplings. The data analyst was blinded to treatment by using coding to reflect treatments in the resulting datasets; after analysis, data was unblinded. Data are presented using individual dot plots combined with summaries of mean and bootstrapped confidence intervals at 95% level. Standardized mean differences with unbiased variance estimates were calculated using the R package ‘metafor’. All analyses and graphs were made using R version 3.3.0 and packages ‘ggplot2’ and ‘coin’.

2.2.4. Pilocarpine-induced seizures

Immediately after the light/dark test, animals were injected with a dose of 150 mg/kg of pilocarpine. This dose has been shown to be insufficient to produce clonic and tonic-clonic seizures in adult animals (Pinto, 2015). Fifteen minutes after injection, animals were individually transferred to 1.5 L tanks, and filmed for 15 min to analyze the profile of seizures. Seizures were scored according to Mussulini et al. (2013), as in Table 2.

Score 4 is the minimum behavioral phenotype that can be considered epileptiform (Mussulini et al., 2013); therefore, the latency to reach Score 4 was considered as the main endpoint for seizure. Latencies were analyzed by fitting a Kaplan-Meier model to survival curves, using the R package ‘survival’.

2.3. Effects of EtOH withdrawal on catalase activity

A separate group of 12 animals were used in this experiment. Animals were subjected to the exposure and withdrawal regimen described in 2.2.1. Sixty minutes after withdrawal, animals were euthanized in cold water followed by spinal section, and their brains and head kidneys were dissected. Catalase activity in those organs was measured using the rate of disappearance of H₂O₂ spectrophotometrically, following the method described by Aebi (Aebi, 1984). Within-laboratory

validation yielded a linearity of $r^2 = 0.9849$, and intermediary repeatability of 0.2364 (IC95% [0.0042, 0.4686]; Horwitz ratio). Enzyme activity was corrected by protein levels, quantified by the Bradford method. Differences between groups were analyzed using Approximative Two-Sample Fisher-Pitman Permutation Tests.

2.4. Data availability

Datasets and scripts for all analyses are available from <https://github.com/lanec-unifesspa/etoh-withdrawal>.

3. Results

3.1. Metanalysis

To begin evaluating the effect of EtOH withdrawal on zebrafish anxiety-like behavior, we applied a mixed-effects meta-regression model on the results from the systematic review. Characteristics from the papers found in the systematic review can be found in Table 3; raw data and analysis scripts for this metanalysis can be found in our GitHub repository (<https://github.com/lanec-unifesspa/etoh-withdrawal/tree/master/metanalysis>). EtOH concentrations ranged from 0.25% to 3% v/v (median 1%); exposure durations ranged from 7 to 63 days (median 14 days), and withdrawal durations ranged from 2 to 1.512 h (median 48 h). In general, a high risk of bias was observed, since most studies did not report blinding or random allocation (<https://github.com/lanec-unifesspa/etoh-withdrawal/blob/master/metanalysis/etoh-withdrawal-metanalysis-rob.csv>). Behavioral test, strain/phenotype, EtOH concentration, exposure duration, and withdrawal duration were used as moderators. Results from this analysis are presented in the forest plot found in Fig. 1A. Residual heterogeneity was estimated as $\tau^2 = 0.1667$, significantly high ($QE_{[df = 11]} = 22.6338$, $p = 0.0199$), suggesting that although about 88% of the total heterogeneity can be explained by including the five moderators in the model, other factors might influence the effect. After applying a permutation test

with 1.000 replications, a suggestive effect of exposure duration ($p = 0.029$) and a significant effect EtOH concentration ($p = 0.001$) were found, but the other mediators did not affect withdrawal-like behavior (Figures 1B-1D). The contour-enhanced funnel plot (Figure 1E) suggest that most studies failed to reach statistical significance, with only one study falling in the 95% CI range, and one in the 99% CI range. Egger's test on this funnel plot did not suggest publication bias ($t_{[df = 10]} = 0.439$, $p = 0.67$). An analysis of influential observations suggests that the Mathur and Guo (2011) study on the effects on the NTT with the longer withdrawal duration and the Dewari et al. (2016) study produced most residual heterogeneity (Figure S1). Absolute SMDs were significantly explained by observed power (slope = 1.2819, $p = 0.0174$; Figure S2). Finally, sensitivity analysis suggested that study quality did not influence results, as including RoB scores in the model did not improve fit (model without RoB: AIC = 46.4816, BIC = 50.88=585; model with RoB: AIC = 47.2908, BIC = 50.9218).

3.2. *Effects of EtOH withdrawal on anxiety-like behavior in zebrafish*

Withdrawal increased time on white ($Z = 2.1207$, $p = 0.0261$; $SMD_{UB} = -0.995$; observed power = 0.6414; Figure 2A), without affecting entry duration ($Z = 0.92305$, $p = 0.4301$; $SMD_{UB} = -0.392$; observed power = 0.147; Figure 2B). Withdrawal did not increase erratic swimming ($Z = 1.9389$, $p = 0.0564$, $SMD_{UB} = 0.890$; observed power = 0.5467; Figure 2C), but it increased risk assessment ($Z = 1.9895$, $p = 0.0405$, $SMD_{UB} = 0.918$; observed power = 0.5724; Figure 2D). Freezing ($Z = 0.8082$, $p = 0.7003$, $SMD_{UB} = 0.286$; observed power = 0.098; Figure 2E) and thigmotaxis ($Z = 0.41291$, $p = 0.7024$, $SMD_{UB} = -0.2133$; observed power = 0.0718; Figure 2F) were unaffected. As for motor effects, transitions to white were not affected by withdrawal ($Z = 0.91765$, $p = 0.3763$, $SMD_{UB} = 0.39$; observed power = 0.1469; Figure 2G), but the number of squares crossed was higher in animals exposed to withdrawal ($Z = 2.5965$, $p = 0.0034$, $SMD_{UB} = 1.31$; observed power = 0.863; Figure 2H).

3.3. *Effects of EtOH withdrawal on seizure susceptibility*

Only one animal from the control group exhibited Score IV seizures after 150 mg/kg pilocarpine, replicating findings from Pinto (2015); therefore, this dose is indeed sub-convulsive in zebrafish. All animals from the withdrawal group exhibited Score IV seizures after 150 mg/kg pilocarpine; after applying a log-rank model for the differences in latencies to Score IV, a suggestive difference was seen in the withdrawal group ($\chi^2_{[df = 1]} = 8.8$, $p = 0.00308$; Figure 3). This data can be found in our GitHub repository (<https://github.com/lanec-unifesspa/etoh-withdrawal/tree/master/seizure>).

3.4. *Effects of EtOH withdrawal on catalase activity*

Catalase activity was reduced in the brain of animals exposed to the withdrawal regime ($Z = 2.0885$, $p = 0.0156$; Figure 4A), while no significant differences were found in the head kidney ($Z = 1.2547$, $p = 0.1863$; Figure 4B). These results are in line with what was observed by Müller et al. (2017), which observed decreased catalase activity in the brains of zebrafish in withdrawal. Data and scripts for this experiment are available at <https://github.com/lanec-unifesspa/etoh-withdrawal/tree/master/catalase>.

4. Discussion

The present work contributed with the use of zebrafish as a model organism in studying ethanol withdrawal by searching for broader patterns in the literature, providing a conceptual replication of some findings regarding anxiety-like behavior and oxidative stress, as well as expanding the breadth of behavioral domains for study. We found that the literature is inconsistent in what regards the effects of ethanol withdrawal. This heterogeneity is associated with the great procedure differences which are reported; the results of the metanalysis suggest that the main

driving factors are ethanol concentration during exposure and exposure duration, with lower concentrations and longer durations inducing more anxiety-like behavior. Moreover, we found that ethanol withdrawal (after exposing animals for 16 days to 0.5% ethanol and 1 h withdrawal) decreased scototaxis, but increased risk assessment, in the light/dark test, and decreased the threshold for chemically-induced seizure.

Zebrafish is increasingly being considered as a model organism in behavioral research (Bonan & Norton, 2015; Kalueff et al., 2012; Norton & Bally-Cuif, 2010; Stewart et al., 2015), with a great deal of studies on alcohol (S. Tran et al., 2016). Behavioral effects of drug withdrawal have been demonstrated with different drugs, including cocaine (López-Patiño, Yu, Cabral, et al., 2008; López-Patiño, Yu, Yamamoto, et al., 2008) and morphine (Cachat et al., 2010; Khor et al., 2011; Wong et al., 2010); in all cases, the behavioral endpoint assessed was anxiety-like behavior. In the same direction, ethanol withdrawal has been studied mainly with models for anxiety-like behavior (Table 3). In our conceptual replication, using the exposure method described in Gerlai et al. (2009), we showed that withdrawal reduces scototaxis. This is in line with the “daily-moderate” condition in Holcombe et al. (2013), which showed a decrease in scototaxis after using the same exposure profile we presented. As can be deprehended from the meta-analysis, the effect sizes calculated for scototaxis in Holcombe et al.’s (2013) experiment are similar in direction and magnitude. The Mathur and Guo (2011) scototaxis study used a higher concentration of EtOH (1%) but a shorter exposure period (8 days). While Pittman and Ichikawa (2013) used a similar exposure period (14 days), the concentration was much higher (3%); in both studies, EtOH withdrawal did not affect scototaxis.

While following only effects on scototaxis confirms findings reported in Holcombe et al. (2013), these findings contradict the overall results from the meta-analysis, which suggested that EtOH withdrawal increases anxiety-like behavior in zebrafish. If other variables are considered, however, the picture changes, with increases in risk assessment and number of transitions. The increase in risk assessment is consistent with increased anxiety-like behavior (Maximino et al.,

2014), but, since scototaxis was *decreased*, this conclusion is only provisional. The increase in transitions without an apparent increase in locomotion in the white compartment could be interpreted as psychomotor agitation, an important symptom of EtOH withdrawal.

In addition to this behavioral profile, EtOH withdrawal was also shown to decrease the threshold for chemically-induced seizures in zebrafish. Pilocarpine is a muscarinic agonist which induces seizures in rodents (Scorza et al., 2009), and has recently been shown to induce a similar profile in zebrafish (Pinto, 2015). The rationale of using a sub-convulsive dose is that, if EtOH withdrawal increases susceptibility to seizures, zebrafish should enter a seizure-like state with a dose which does not induce this state in control animals. We observed a much increased probability of entering Stage 4 seizures in EtOH withdrawal animals, and a shorter latency to this event, suggesting that the protocol presented here is able to model susceptibility to seizures, increasing the breadth of endpoints for studying EtOH withdrawal in zebrafish.

Finally, we also observed decreased catalase activity in the brain, but not in the head kidney of EtOH withdrawal animals. Catalase is a detoxifying enzyme that catalyzes the transformation of hydrogen peroxide, a free radical, into water and oxygen (Aebi, 1984). The inhibition of enzymatic activity observed here replicates what was observed by Müller et al. (2017) in the zebrafish brain; in that study, however, the exposure duration was shorter (8 days), the concentration of EtOH was higher (1%), and the pattern of exposure was different (animals were exposed for 20 min per day, instead of continuously, to ethanol). The lack of effect on the head kidney, where interrenal cells (the teleost functional equivalent of the mammalian adrenal cortex) lie, suggest a brain-specific effect, and also suggests that possible effects on cortisol (e.g., Cachat et al., 2010) are not due to effects in these cells, but upstream in the hypothalamus-hypophyseal-interrenal axis.

The present work advanced the use of zebrafish as a model in EtOH withdrawal research by: A) identifying sources of heterogeneity in the literature on EtOH withdrawal and anxiety-like behavior in the species; B) presenting a conceptual replication of withdrawal-induced anxiogenesis in the light/dark test; C) extending the behavioral phenotypes to include seizure susceptibility; and

D) replicating the effects on catalase activity, suggesting a mechanism of angiogenesis related to EtOH withdrawal in the species. Further work will characterize these mechanisms with care.

Acknowledgements

SNSC was the recipient of a CNPq/PIBIC studentship. GRF was the recipient of a CNPq/PIBITI studentship. WEO was the recipient of a CNPq/PIBIC-EM studentship to high school students.

References

- Aebi, H. (1984). Catalase in vitro. In *Methods in Enzymology* (Vol. 105, pp. 121–126).
[https://doi.org/10.1016/S0076-6879\(84\)05016-3](https://doi.org/10.1016/S0076-6879(84)05016-3)
- Benneh, C. K., Biney, R. P., Mante, P. K., Tandoh, A., Adongo, D. W., & Woode, E. (2017). *Maerua angolensis* stem bark extract reverses anxiety and related behaviours in zebrafish—Involvement of GABAergic and 5-HT systems. *Journal of Ethnopharmacology*, 207, 129–145.
<https://doi.org/10.1016/j.jep.2017.06.012>
- Bonan, C. D., & Norton, W. H. J. (2015). The utility of zebrafish as a model for behavioural genetics. *Current Opinion in Behavioral Sciences*, 2, 34–38.
<https://doi.org/10.1016/j.cobeha.2014.07.003>
- Bonassoli, V. T., Milani, H., & Oliveira, R. M. W. de. (2011). Ethanol withdrawal activates nitric oxide-producing neurons in anxiety-related brain areas. *Alcohol*, 45, 641–652.
<https://doi.org/10.1016/j.alcohol.2010.11.007>
- Cachat, J., Canavello, P., Elegante, M., Bartels, B., Hart, P., Bergner, C., & Kalueff, A. V. (2010). Modeling withdrawal syndrome in zebrafish. *Behavioural Brain Research*, 208, 371–376.
- Chakravarty, D. N., & Faingold, C. L. (1998). Comparison of neuronal response patterns in the external and central nuclei of inferior colliculus during ethanol administration and ethanol withdrawal. *Brain Research*, 783, 55–57.
- Conselho Nacional de Controle de Experimentação Animal - CONCEA. Diretriz brasileira para o cuidado e a utilização de animais para fins científicos e didáticos - DBCA. Anexo I. Peixes mantidos em instalações de instituições de ensino ou pesquisa científica, Pub. L. No. Resolução Normativa 30/2016 (2017). Brasil.
- Dewari, P. S., Ajani, F., Kushawah, G., Kumar, D. S., & Mishra, R. K. (2016). Reversible loss of reproductive fitness in zebrafish on chronic alcohol exposure. *Alcohol*, 50, 83–89.
<https://doi.org/10.1016/j.alcohol.2015.11.006>

- Egger, M., Smith, G. D., Schneider, M., & Minder, C. (1997). Bias in meta-analysis detected by a simple, graphical test. *British Medical Journal*, 315, 629–634.
- Gatch, M. B., & Lal, H. (2001). Animal models of the anxiogenic effects of ethanol withdrawal. *Drug Development Research*, 54, 95–115.
- Gerlai, R. (2010). Zebrafish antipredatory responses: A future for translational research? *Behavioural Brain Research*, 207, 223–231. <https://doi.org/10.1016/j.bbr.2009.10.008>
- Gerlai, R. (2014). Fish in behavior research: Unique tools with a great promise! *Journal of Neuroscience Methods*. <https://doi.org/10.1016/j.jneumeth.2014.04.015>
- Gerlai, R., Chatterjee, D., Pereira, T., Sawashima, T., & Krishnannair, R. (2009). Acute and chronic alcohol dose: Population differences in behavior and neurochemistry of zebrafish. *Genes, Brain, and Behavior*, 8, 586–599. <https://doi.org/10.1111/j.1601-183X.2009.00488.x>
- Hedges, L. V, & Olkin, I. (1985). *Statistical methods for meta-analysis*. San Diego: Academic Press.
- Holcombe, A., Howorko, A., Powell, R. A., Schalomon, M., & Hamilton, T. J. (2013). Reversed scototaxis during withdrawal after daily-moderate, but not weekly-binge, administration of ethanol in zebrafish. *PLoS ONE*, 8, e63319. <https://doi.org/10.1371/journal.pone.0063319>
- Hooijmans, C. R., Rovers, M. M., de Vries, R. B. M., Leenaars, M., Ritskes-Hoitinga, M., & Langendam, M. W. (2014). SYRCLE’s risk of bias tool for animal studies. *BMC Medical Research Methodology*, 14, 43.
- Hooijmans, C. R., Tillema, A., Leenaars, M., & Ritskes-Hoitinga, M. (2010). Enhancing search efficiency by means of a search filter for finding all studies on animal experimentation in PubMed. *Laboratory Animals*, 44, 170–175.
- Hortopan, G. A., Dinday, M. T., & Baraban, S. C. (2010). Zebrafish as a model for studying genetic aspects of epilepsy. *Disease Models & Mechanisms*, 3, 144–148. <https://doi.org/10.1242/dmm.002139>
- Kalueff, A. V, Echevarria, D. J., & Stewart, A. M. (2014). Gaining translational momentum: More zebrafish models for neuroscience research. *Progress in Neuro-Psychopharmacology & Biological Psychiatry*, 55, 1–6. <https://doi.org/10.1016/j.pnpbp.2014.01.022>
- Kalueff, A. V, Gebhardt, M., Stewart, A. M., Cachat, J. M., Brimmer, M., Chawla, J. S., ... Zebrafish Neuroscience Research Consortium. (2013). Towards a comprehensive catalog of zebrafish behavior 1.0 and beyond. *Zebrafish*, 10, 70–86. <https://doi.org/10.1089/zeb.2012.0861>
- Kalueff, A. V, Stewart, A. M., Kyzar, E. J., Cachat, J., Gebhardt, M., Landsman, S., ... Zebrafish Neuroscience Research Consortium. (2012). Time to recognize zebrafish “affective” behavior. *Behaviour*, 149, 1019–1036. <https://doi.org/10.1163/1568539X-00003030>
- Khor, B. S., Jamil, M. F., Adenan, M. I., & Shu-Chien, A. C. (2011). Mitragynine attenuates withdrawal syndrome in morphine-withdrawn zebrafish. *PLoS ONE*, 6, e28340. <https://doi.org/10.1371/journal.pone.0028340>

- Kokel, D., & Peterson, R. T. (2008). Chemobehavioural phenomics and behaviour-based psychiatric drug discovery in the zebrafish. *Briefings in Functional Genomics and Proteomics*, 7, 483–490.
- Koob, G. F., & Le Moal, M. (2008). Neurobiological mechanisms for opponent motivational processes in addiction. *Philosophical Transactions of the Royal Society Part B*, 363, 3113–3123. <https://doi.org/10.1098/rstb.2008.0094>
- Krystal, J. H., & Tabakoff, B. (2002). Ethanol abuse, dependence, and withdrawal: Neurobiology and clinical implications. In *Neuropsychopharmacology: The Fifth Generation of Progress* (pp. 1425–1443). Philadelphia: Lippincott, Williams & Wilkins.
- Lawrence, C. (2007). The husbandry of zebrafish (*Danio rerio*): A review. *Aquaculture*, 269, 1–20. <https://doi.org/10.1016/j.aquaculture.2007.04.077>
- Long, C., Yang, L., Faingold, C. L., & Evans, M. S. (2007). Excitatory amino acid receptor-mediated responses in periaqueductal gray neurons are increased during ethanol withdrawal. *Neuropharmacology*, 52, 802–811. <https://doi.org/10.1016/j.neuropharm.2006.09.019>
- Longo, D. L., & Schuckit, M. A. (2014). Recognition and management of withdrawal delirium (*delirium tremens*). *New England Journal of Medicine*, 371, 2109–2113. <https://doi.org/10.1056/NEJMr1407298>
- López-Patiño, M. A., Yu, L., Cabral, H., & Zhdanova, I. V. (2008). Anxiogenic effects of cocaine withdrawal in zebrafish. *Physiology & Behavior*, 93, 160–171. <https://doi.org/10.1016/j.physbeh.2007.08.013>
- López-Patiño, M. A., Yu, L., Yamamoto, B. K., & Zhdanova, I. V. (2008). Gender differences in zebrafish responses to cocaine withdrawal. *Physiology & Behavior*, 95, 36–47. <https://doi.org/10.1016/j.physbeh.2008.03.021>
- Mathur, P., & Guo, S. (2011). Differences of acute versus chronic ethanol exposure on anxiety-like behavioral responses in zebrafish. *Behavioural Brain Research*, 219, 234–239. <https://doi.org/10.1016/j.bbr.2011.01.019>
- Maximino, C., Brito, T. M. De, Dias, C. A. de M., Gouveia Jr., A., & Morato, S. (2010). Scototaxis as anxiety-like behavior in fish. *Nature Protocols*, 5(2), 209–216. <https://doi.org/10.1038/nprot.2009.225>
- Maximino, C., da Silva, A. W. B., Araújo, J., Lima, M. G., Miranda, V., Puty, B., ... Herculano, A. M. (2014). Fingerprinting of psychoactive drugs in zebrafish anxiety-like behaviors. *PLoS ONE*, 9, e103943. <https://doi.org/10.1371/journal.pone.0103943>
- Maximino, C., Marques, T., Brito, D., Waneza, A., Manoel, A., Morato, S., & Gouveia, A. (2010). Measuring anxiety in zebrafish: A critical review. *Behavioural Brain Research*, 214, 157–171. <https://doi.org/10.1016/j.bbr.2010.05.031>
- Mohammad, F., Ho, J., Lim, C. L., Woo, J. H., Poon, D. J. J., Lamba, B., & Claridge-chang, A. (2016). Concordance and incongruence in preclinical anxiety models: Systematic review and meta-analyses. *bioRxiv*. <https://doi.org/10.1101/020701>

- Müller, T. E., Nunes, S. Z., Silveira, A., Loro, V. L., & Rosemberg, D. B. (2017). Repeated ethanol exposure alters social behavior and oxidative stress parameters of zebrafish. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*.
<https://doi.org/10.1016/j.pnpbp.2017.05.026>
- Mussulini, B. H. M., Leite, C. E., Zenki, K. C., Moro, L., Baggio, S., Rico, E. P., ... de Oliveira, D. L. (2013). Seizures induced by pentylenetetrazole in the adult zebrafish: A detailed behavioral characterization. *PLOS ONE*, 8, e54515. <https://doi.org/10.1371/journal.pone.0054515>
- Norton, W., & Bally-Cuif, L. (2010). Adult zebrafish as a model organism for behavioural genetics. *BMC Neuroscience*, 11, 90. <https://doi.org/10.1186/1471-2202-11-90>
- Oliveira, R. F. (2013). Mind the fish: Zebrafish as a model in cognitive social neuroscience. *Frontiers in Neural Circuits*, 7, Article 131. <https://doi.org/10.3389/fncir.2013.00131>
- Parra, K. V, Adrian Jr, J. C., & Gerlai, R. (2009). The synthetic substance hypoxanthine 3-N-oxide elicits alarm reactions in zebrafish (*Danio rerio*). *Behavioural Brain Research*, 205, 336–341. <https://doi.org/10.1016/j.bbr.2009.06.037>
- Pinto, C. B. (2015). *Caracterização do perfil de crises epilépticas e dos efeitos comportamentais induzidos por pilocarpina em peixe-zebra adulto*. Universidade Federal do Rio Grande do Sul.
- Pittman, J., & Hylton, A. (2015). Behavioral, endocrine, and neuronal alterations in zebrafish (*Danio rerio*) following sub-chronic coadministration of fluoxetine and ketamine. *Pharmacology, Biochemistry & Behavior*, 139, 158–162. <https://doi.org/10.1016/j.pbb.2015.08.014>
- Pittman, J. T., & Ichikawa, K. M. (2013). iPhone® applications as versatile video tracking tools to analyze behavior in zebrafish (*Danio rerio*). *Pharmacology Biochemistry and Behavior*, 106, 137–142. <https://doi.org/10.1016/j.pbb.2013.03.013>
- Rasmussen, D. D., Boldt, B. M., Wilkinson, C. W., & Mitton, D. R. (2002). Chronic daily ethanol and withdrawal: 3. Forebrain pro-opiomelanocortin gene expression and implications for dependence, relapse, and deprivation effect. *Alcoholism: Clinical and Experimental Research*, 26, 535–546.
- Scorza, F., Arida, R., Naffah-Mazzacoratti, M. G., Scerni, D., Calderazzo, L., & Cavalheiro, E. (2009). The pilocarpine model of epilepsy: What have we learned? *Anais Da Academia Brasileira de Ciências*, 81, 345–365.
- Speedie, N., & Gerlai, R. (2008). Alarm substance induced behavioral responses in zebrafish (*Danio rerio*). *Behavioural Brain Research*, 188, 168–177. <https://doi.org/10.1016/j.bbr.2007.10.031>
- Stewart, A. M., Ullmann, J. F. P., Norton, W. H. J., Parker, M. O., Brennan, C. H., Gerlai, R., & Kalueff, A. V. (2015). Molecular psychiatry of zebrafish. *Molecular Psychiatry*, 20, 2–17. <https://doi.org/10.1038/mp.2014.128>
- Tran, S., Chatterjee, D., & Gerlai, R. (2015a). An integrative analysis of ethanol tolerance and withdrawal in zebrafish (*Danio rerio*). *Behavioural Brain Research*, 276, 161–170. <https://doi.org/10.1016/j.bbr.2014.02.034>

- Tran, S., Chatterjee, D., & Gerlai, R. (2015b). An integrative analysis of ethanol tolerance and withdrawal in zebrafish (*Danio rerio*). *Behavioural Brain Research*, 276, 161–170. <https://doi.org/10.1016/j.bbr.2014.02.034>
- Tran, S., Faccioli, A., & Gerlai, R. (2016). The zebrafish, a novel model organism for screening compounds affecting acute and chronic ethanol-induced effects. *International Review of Neurobiology*, 126, 467–484. <https://doi.org/10.1016/bs.irn.2016.02.016>
- Viechtbauer, W. (2010). Conducting meta-analyses in R with the metafor package. *Journal of Statistical Software*, 36, 1–48. Retrieved from <https://cran.r-project.org/web/packages/metafor/index.html>
- Wong, K., Elegante, M., Bartels, B., Elkhayat, S., Tien, D., Roy, S., ... Kalueff, A. V. (2010). Analyzing habituation responses to novelty in zebrafish (*Danio rerio*). *Behavioural Brain Research*, 208, 450–457. <https://doi.org/10.1016/j.bbr.2009.12.023>
- Yang, L., Long, C., Evans, M. S., & Faingold, C. (2002). Ethanol withdrawal results in aberrant membrane properties and synaptic responses in periaqueductal gray neurons associated with seizure susceptibility. *Brain Research*, 957, 99–108.
- Yang, L., Long, C., & Faingold, C. L. (2001). Neurons in the deep layers of superior colliculus are a requisite component of the neuronal network for seizures during ethanol withdrawal. *Brain Research*, 920, 134–141.
- Yang, L., Long, C., Randall, M. E., & Faingold, C. L. (2003). Neurons in the periaqueductal gray are critically involved in the neuronal network for audiogenic seizures during ethanol withdrawal. *Neuropharmacology*, 44, 275–281. [https://doi.org/10.1016/S0028-3908\(02\)00367-2](https://doi.org/10.1016/S0028-3908(02)00367-2)
- Zenki, K. C., Mussulini, B. H. M., Rico, E. P., de Oliveira, D. L., & Rosemberg, D. B. (2014). Effects of ethanol and acetaldehyde in zebrafish brain structures: An in vitro approach on glutamate uptake and on toxicity-related parameters. *Toxicology in Vitro*, 28, 822–828. <https://doi.org/10.1016/j.tiv.2014.03.008>

Tables and table legends

Table 1. Operational definitions for behavioral endpoints assessed in the light/dark test. Whenever available, codes used in the Zebrafish Behavioral Catalog (Kalueff et al., 2013) are provided.

Behavioral endpoint	Definition
Freezing	Time spent immobile, with the exception of opercular and eye movements, in seconds, observed in the white compartment (ZBC 1.68)
Erratic swimming	Sharp changes in direction or velocity of swimming, associated with repeated fast acceleration bouts, observed in the white compartment (ZBC 1.51)
Thigmotaxis	Percentage of time swimming in a distance of 2 cm or less from the white compartment's walls (ZBC 1.173)
Risk assessment	Fast (< 1 second) entries in the white compartment, followed by re-entry in the black compartment, or partial entries in the white compartment (i.e., the pectoral fin does not cross the midline)

Table 2. Behavioral phenotypes scored in the pilocarpine-induced seizure model. Scores were based on Mussulini et al. (2013).

Score	Behavioral phenotype
0	Short swim mainly in the bottom of the tank.
1	Increased swimming activity and high frequency of opercular movement.
2	Burst swimming, left and right movements, and erratic movements.
3	Circular movements.
4	Clonic seizure-like behavior (abnormal whole-body rhythmic muscular contraction).
5	Fall to the bottom of the tank, tonic seizure-like behavior (sinking to the bottom of the tank, loss of body posture, and principally by rigid extension of the body).
6	Death

Table 3. Studies included in the metanalysis. Studies were ordered by strain used (BSF = blue shortfin; WT = non-specified wild-type); behavioral assay (LDT = light/dark test; NTT = novel tank test); ethanol concentration during exposure ([EtOH]); exposure duration; and withdrawal duration.

Study	Strain	Assay	[EtOH]	Exposure duration	Withdrawal duration
Holcombe et al., 2013	BSF	LDT	0.2%	21 d	48 h
Tran et al., 2015	AB	NTT	0.5%	22 d	2 h
Mathur and Guo, 2011, study 1	AB	NTT	1%	8 d	48 h
Mathur and Guo, 2011, study 2	AB	LDT	1%	8 d	24 h
Mathur and Guo, 2011, study 3	AB	NTT	1%	8 d	144 h
Mathur and Guo, 2011, study 4	AB	LDT	1%	8 d	168 h
Pittman and Hylton, 2015	WT	NTT	3%	14 d	48 h
Cachat et al., 2010	BSF	NTT	0.3%	7 d	12 h
Gerlai et al., 2009, study 1	WT	Predator avoidance	0.25%	22 d	2 h
Gerlai et al., 2009, study 2	AB	Predator avoidance	0.25%	22 d	2 h
Gerlai et al., 2009, study 3	SF	Predator avoidance	0.25%	22 d	2 h
Gerlai et al., 2009, study 4	WT	Shoaling	0.25%	22 d	2 h
Gerlai et al., 2009, study 5	AB	Shoaling	0.25%	22 d	2 h
Gerlai et al., 2009, study 6	SF	Shoaling	0.25%	22 d	2 h
Pittman and Ichikawa, 2013, study 1	WT	NTT	3%	14 d	48 h
Pittman and Ichikawa, 2013, study 2	WT	LDT	3%	14 d	48 h
Müller et al., 2017	BSF	Shoaling	1%	8 d	24 h
Benneh et al., 2017, study 1	WT	NTT	0.5%	8 d	96 h
Benneh et al., 2017, study 2	WT	NTT	0.5%	8 d	192 h
Benneh et al., 2017, study 3	WT	LDT	0.5%	8 d	96 h
Benneh et al., 2017, study 4	WT	LDT	0.5%	8d	192 h
Dewari et al., 2016	WT	NTT	0.5%	63 d	1512 h

Figure legends

Figure 1 – (A) Forest plot showing the results of 16 studies examining the effect of ethanol withdrawal on zebrafish anxiety-like behavior. The figure shows the standardized mean difference (SMD) between control and withdrawal-exposed groups with corresponding 95% confidence intervals in the individual studies, based on a mixed-effects model. A negative standardized mean difference (SMD) corresponds to decreased anxiety-like behavior, while a positive SMD corresponds to increased anxiety-like behavior after ethanol withdrawal. Studies are ordered by total degrees of freedom. (B-D) Permutation distribution of the test statistic for the mediators: behavioral test (B), ethanol concentration during exposure (C), exposure duration, in days (D), withdrawal duration, in hours (E), and strain/phenotype (F). Distributions were based on a permutation test with 1,000 replications. The blue contour represents kernel density estimates of the permutation distributions; the red curve represents the standard normal density; the full line represents the null hypothesis of no difference; and the dashed line represents the observed values of test statistics. (G) Contour-enhanced funnel plot of meta-analysis. Estimated standardized mean differences were plotted against precision (1/standard error) were A negative estimate corresponds to decreased anxiety-like behavior, while a positive estimate corresponds to increased anxiety-like behavior after ethanol withdrawal. The unshaded region corresponds to p-values greater than 0.1, the gray-shaded region to p-values between 0.1 and 0.05, the dark gray-shaded region corresponds to p-values between 0.05 and 0.01, and the region outside of the funnel corresponds to p-values below 0.01.

Figure 2 – Effects of EtOH withdrawal (increasing concentration of up to 0.5% for 16 days, followed by 1 h withdrawal) on behavior in the light/dark test. (A) Scototaxis; (B) Erratic swimming; (C) Risk assessment; (D) Freezing; (E) Thigmotaxis; (F) Transitions to white; (G) Locomotion on white. Red bars represent mean, and red error bars represent nonparametric bootstrapped confidence intervals for the mean at the 95% level. To facilitate visualization, data

points were jittered; therefore, their absolute position does not reflect the actual value, and values can appear to be below 0.

Figure S2 – Relationship between observed power and modulus SMD values. Curve fitting was made using a linear model.


A

Author(s) and Year


SMD [95% CI]

Holcombe et al., 2013
 Tran et al., 2015
 Mathur et al., 2011.1
 Mathur et al., 2011.2
 Pittman et al., 2015
 Cachat et al., 2010
 Gerlai et al., 2009.1
 Gerlai et al., 2009.2
 Gerlai et al., 2009.3
 Gerlai et al., 2009.4
 Gerlai et al., 2009.5
 Pittman & Ichikawa, 2013.1
 Pittman & Ichikawa, 2013.2
 Müller et al., 2017
 Mathur et al., 2011.3
 Mathur et al., 2011.4
 Benneh et al., 2017.1
 Benneh et al., 2017.2
 Benneh et al., 2017.3
 Benneh et al., 2017.4
 Dewari et al., 2016


-1.08 [-1.85, -0.32]
 -0.08 [-0.82, 0.66]
 -0.30 [-1.28, 0.69]
 -0.02 [-1.00, 0.96]
 1.17 [0.22, 2.11]
 0.56 [-0.32, 1.45]
 -2.30 [-3.27, -1.33]
 -0.21 [-0.95, 0.53]
 0.37 [-0.38, 1.12]
 -1.33 [-2.14, -0.51]
 -0.06 [-0.80, 0.68]
 0.66 [-0.24, 1.56]
 0.53 [-0.36, 1.42]
 1.17 [0.11, 2.23]
 3.54 [2.03, 5.06]
 -0.01 [-0.99, 0.97]
 -1.09 [-1.75, -0.43]
 -0.60 [-1.23, 0.03]
 -2.72 [-3.75, -1.70]
 -2.15 [-3.09, -1.22]
 0.00 [-0.75, 0.75]




B




C




D




E











F




G






Cumulative risk of seizure



A

CAT activity (U per mg protein)

**B**

CAT activity (U per mg protein)

