Where is Neptune?

Evolution on graphs and vestibular schwannoma

Chay Paterson


University of Manchester

September 2022

Warning

There will be very few equations in this talk!

This network...



corresponds to this stochastic process:...

$$\emptyset \to X, \quad \text{rate } \alpha$$
 $X \to X + Y, \quad \text{rate } \mu X$
 $Y \to Y + Y, \quad \text{rate } \beta Y$
 $Y \to \emptyset, \quad \text{rate } \delta Y$

and approximately this linear system...


$$\frac{d}{dt} \begin{pmatrix} E[X] \\ E[Y] \end{pmatrix} = \begin{bmatrix} 0 & 0 \\ \mu & \beta - \delta \end{bmatrix} \cdot \begin{pmatrix} E[X] \\ E[Y] \end{pmatrix} + \begin{pmatrix} \alpha \\ 0 \end{pmatrix}$$

Most of our models are linear, high-dimensional and sparse¹

¹C. Paterson, I. Bozic, H. Clevers, PNAS 2020; 117(34): 20681-20688

Multi-stage models

P. Armitage and R. Doll²




¹P. Armitage and R. Doll, British Journal of Cancer 1954; 8: 1–12

²note that P(t) = 1 - S(t), other authors (e.g. Knudson)

Network models

- 1. Study **specific genes** and mechanisms of interest
- 2. Fix parameters from sequences and experiments
- 3. Distinguish different orders of events


Predict copy number alterations (etc.)




This gets us the incidence of specific karyotypes

Vestibular schwannoma

3-event model



- Fitness suspiciously low, $s \approx 0.005/\text{yr}$
- Suggests nearly-neutral 3-hit model





¹R. Woods et al. Genetic Epidemiology (2003)24: 265-272


Our model for sporadic VS

- ► Include *NF2*, *SMARCB1* and (simplified) linkage
- ► Add hypothetical oncogene *GFX*

Risk of each subtype looks like


$$P(\square) \propto \frac{t^3}{3!}$$

$$P(\square) \approx N_{WT} \mu_{GFX} r_{LOH} \frac{1}{2} \mu_{NF2} \frac{t^3}{3!} \times 6$$



¹C Paterson, I Bozic, MJ Smith, X Hoad, DGR Evans, https://doi.org/10.1101/2021.10.03.457528

Our model for sporadic VS



- ightharpoonup + \square = have LOH on 22q
- ► frequency of LOH = $f_{LOH} = (\square + \square)/(\square + \square + \square)$
- $ightharpoonup = SMARCB1^{-/-}$
- ▶ frequency of SMARCB1^{-/-}
 = f_{SMARCB1}


$$= \square/(\square + \square + \square)$$

Can use these to fix parameters!

Our model for sporadic VS


+ 🔲 + 🛄

+ 🔲



 $^{^{1}\}text{C Paterson, I Bozic, MJ Smith, X Hoad, DGR Evans, } \text{https://doi.org/} 10.1101/2021.10.03.457528}$

New parameter estimates



0 1.2 1e-9

0.00

0.0

0.2 0.4 0.6

и

0.0

0.0 0.5 1.0


1e-7

 $p_{LOH}(22q)$

Malignant transformation in vestibular schwannoma

Very rare, very bad

- ightharpoonup Risk pprox 0.1% of VS cases
- ▶ 5-year survival $\approx 12 20\%$






²AK Demetriades et al. Skull Base (2010)20:381–387.

Malignant schwannoma: two models

Timing and identity of drivers

Oncogene activation:

TSX inactivation:




Malignant VS is extremely rare!

 $^{^{1}\}mathsf{C}\ \mathsf{Paterson}, \ \mathsf{I}\ \mathsf{Bozic}, \ \mathsf{MJ}\ \mathsf{Smith}, \ \mathsf{X}\ \mathsf{Hoad}, \ \mathsf{DGR}\ \mathsf{Evans}, \ \mathsf{https:}//\mathsf{doi.org}/10.1101/2021.10.03.457528$

Malignant schwannoma: first model


Oncogene activation




- ▶ Oncogene activation ⇒ high risk
- ▶ But it's a rare outcome
- So it's probably not caused by oncogene activation

Malignant schwannoma: second model

Tumour suppressor *TSX* inactivation






- ightharpoonup TSX inactivation \implies low risk
- \triangleright Can also estimate n_{TSX} that's consistent with incidence

Who is *TSX*?

Parameter estimates for n_{TSX}




Probably multiple (10?) distinct tumour suppressors

i.e. not (just) TP53: $n_{TP53} = 73$

Malignant schwannoma

Why do we care about TSX anyway?





3 dose-dependent effects:

- ▶ DSB induction $p_{DSB}(D)$
- ▶ DSB misrepair $\varepsilon(D)$
- ▶ cell survival S(D)

Malignant schwannoma

Why do we care about TSX anyway?





Main outputs

- 1. Better estimates of event rates in Schwann cells
- 2. Can constrain timing of "TSX" (resp. for malignancy)
- 3. Can constrain size of GFX and TSX
- 4. Radiotherapy probably OK (w. caveats + huge error bars)

To do list

Next gen sequencing...

- ▶ Sporadic VS to constrain $f_{SMARCB1}$: n > 300
- ► CNA/NGS in MPNST (rare!): n > 30

but also...

- ▶ Better models!!! CNA/VAF models, machine learning, optimal clustering, $\mu_{C>T}$ etc.
- SEER data too?
- ► Multiple genes *GFX* and *TSX*?
- ► Haploinsufficiency, selection?

Lots to do!

I need collaborators, send me data

Acknowledgements + collaborators

for their in kind support





The University of Washington

In order of appearance...

- Ivana Božić
- Hans Clevers
- Gareth Evans
- Xanthe Hoad
- ► Miriam Smith