

Evolution on graphs and the transition to cancer




Chay Paterson

University of Manchester

June 25, 2024


Introduction

Differential equations




Introduction

Driver mutations



Multi-stage models

P. Armitage and R. Doll¹²



¹P. Armitage and R. Doll, British Journal of Cancer 1954; 8: 1–12

²P. Armitage and R. Doll, British Journal of Cancer 1957; 11(2): 161–169

Multi-stage models

A.G. Knudson¹²

Mutation and Retinoblastoma 823





FIG. 1. Semilogarithmic plot of fraction of cases of retinoblastoma not yet diagnosed (S) vs. age in months (t). The one-hit curve was calculated from $\log S = -t/30$, the two-hit curve from $\log S = -4 \times 10^{-6} t^2$.

¹AG. Knudson, PNAS 68.4 (1971): 820-823.


²F. Michor, Y. Iwasa and MA. Nowak, Nature Reviews Cancer 2004; 4: 197-205 doi:10.1038/nrc1295

Graphs

This network...



corresponds to this
stochastic process:...



How do these studies work?


What are the relevant observables in this type of longitudinal study & data analysis?

Track a cohort that initially contains N patients in the study:

- ▶ Age-specific incidence $I(a)$: rate at which new cases are recorded in the cohort with ages between a and $a + da$
- ▶ Survival function $S(a)$: probability to survive to age a without being diagnosed.

$$I(a) = -N \frac{dS}{da} = -N S(a) \frac{d \ln S}{da} \quad (1)$$

So what?



- ▶ The survival curve $S(a)$ determines the incidence curve $I(a)$
- ▶ The model determines the survival curve: the probability not to end up at one of the end nodes of the graph. We want to know what model+parameters best agree with data from studies.
- ▶ To fit (or “train”) the model to longitudinal data, we need to compute $S(a)$!

This is the central mathematical problem in cancer epidemiology.
How do we compute S ?

Multi-stage clonal expansion models

2-3 rate limiting steps¹²³



Problem: how to compute $S(t)$ for a given model?

Different methods: Fast:

- ▶ Armitage + Doll's approximation¹
- ▶ Moolgavkar + Venzon's quadrature²

Very slow:

- ▶ Gillespie algorithm + sampling³



↓?



¹P. Armitage and R. Doll, British Journal of Cancer 1957; 11(2): 161-169

²S. Moolgavkar and G. Luebeck, JNCI 1992; 84(8): 610-618

³C. Paterson, I. Bozic, H. Clevers, PNAS 2020; 117(34): 20681-20688

(supp. material)

Armitage and Doll's approximation



- ▶ Assume all the probabilities are small: $1 - S \ll 1$
- ▶ Then the relevant probability $S(a)$ is expressible in terms of expected values/population means, which implies


$$S(a) \propto a^k (e^{sa} - 1) \quad (2)$$

with constants k and s .

- ▶ Don't use correlations, variances, or higher moments in stem cell populations – just ignore these.


Models on graphs

1. Most methods for computing $S(a)$ do not consider graphs with multiple end nodes
2. To study **specific genes** and mechanisms of interest (SNVs, LOH, CNA, etc.), we need to evaluate $S(a)$ for a model defined on a graph (right)
3. What methods do we actually have?



This gets us the incidence of *specific karyotypes*


Colorectal adenocarcinoma model



- ▶ Can use A-D type approximation, or stochastic simulations

but:

- ▶ Mean-field breaks down at old ages / large probabilities
- ▶ Stochastic simulations are *extremely slow*



Alternative approach: Kolmogorov forward equations

Define a more general generating function Ψ :

$$\Psi(t, \vec{q}) = \mathbb{E}\left[\prod_j q_j^{N_j}\right] \quad (3)$$

and derive Kolmogorov forward equations instead. Then we can numerically integrate these, and get survival curves $S_i(a)$ for different types of cancer i . E.G. tumours with clonal LOH, or no clonal LOH.


$$S_i = \Psi(t, q_j = 1, \dots, q_i = 0) \quad (4)$$

People knew about this approach for a long time (since 1970s) but it was never considered as useful as backward equations.

¹SH Moolgavkar and DJ Venzon, Math. biosc. 1979; 47(1): 55-77

²DW Quinn, Risk Analysis 1989; 9(3): 407-13

Kolmogorov forward equations as wave equations



Briefly: the Kolmogorov forward equations in $P(N_0, \dots)$

$$\begin{aligned} \frac{dP(N_0, N_1, \dots)}{dt} = & \sum_{\text{vertices}} \alpha(N_j - 1)P(\dots, N_j - 1, \dots) \\ & - \alpha N_j P(\dots, N_j, \dots) \\ & + \beta \dots + \mu \dots + \delta \dots \end{aligned}$$

get transformed...

Kolmogorov forward equations as wave equations

They transform with $P \rightarrow \Psi$:

$$\frac{\partial \Psi}{\partial t} = \sum_{\text{vertices}} \alpha(q_j - 1) q_j \frac{\partial \Psi}{\partial q_j} + \cdots = \mathcal{H} \Psi \quad (5)$$

where \mathcal{H} is a hyperbolic differential operator.

Because this is a hyperbolic wave equation, we can solve for future values of Ψ if we have initial values, by evolving them along *characteristics*.

Kolmogorov forward equations


Using the big generating function Ψ , find the corresponding wave equation:

$$\frac{d\Psi}{dt} = \mathcal{H}\Psi \quad (6)$$

This can be solved using the method of characteristics, so we can instead integrate

$$\frac{d\vec{\gamma}}{dt} = \vec{X} \quad (7)$$

numerically, using an appropriate time stepper.



The vector field \vec{X} and a characteristic $\vec{\gamma}$

Random sampling

Error analysis

To compare methods, ask under what conditions the errors are similar. Stochastic algorithms have

$$\epsilon \sim N^{-1/2} \quad (8)$$

and will thus need

$$N \sim \mathcal{O}(\epsilon^{-2}) \quad (9)$$

runs, so overall runtime $T \propto N$.

Method of characteristics

Error analysis

Why wasn't this method ever used? Wave equation+characteristic methods were studied before, but used Euler integration, which has error

$$\epsilon \sim \Delta t \quad (10)$$

and required two passes, so it ran in

$$T \sim \Delta t^{-2} \sim \mathcal{O}(\epsilon^{-2}) \quad (11)$$

this is asymptotically just as bad as random sampling!

¹SH Moolgavkar and DJ Venzon, Math. biosc. 1979; 47(1): 55-77

²DW Quinn, Risk Analysis 1989; 9(3): 407-13

Fast forward method

Error analysis

But what happens if we only need one pass, and replace Euler integration with a Runge-Kutta scheme?

$$\epsilon \sim \Delta t^2 \tag{12}$$


and runs in

$$T \sim \Delta t^{-1} \sim \mathcal{O}(\epsilon^{-1/2}) \tag{13}$$

so new runtime $\sim \mathcal{O}(\text{old runtime}^{1/4})$.

Amazing!

Fast forward method




Random sampling vs fast forward method:


Monte Carlo: $\approx 5000s$ Fast forward: 4ms

Fast forward method

Error analysis





How do they compare?




Fast forward method

Efficiently computing $S_k(a)$ means we can evaluate likelihood functions directly, just sampling them.....





Fast forward method

Parameter inference



Fast forward method

Parameter inference



Thank you!

What is my message?

- ▶ Don't study continua – study PROBABILITIES!
- ▶ Age structure is important & informative, genes are discrete

Where next?

- ▶ Run these analyses on real studies!
- ▶ Combine genomic and age data TOGETHER
- ▶ Ongoing: sequencing schwannoma tumours in NF2 and oesophageal cancer in Barrett's cases

Acknowledgements



funded this research



The University of Manchester



Manchester University
NHS Foundation Trust



The University of Washington

All my collaborators...

- ▶ Ivana Božić
- ▶ Gareth Evans
- ▶ Miaomiao Gao
- ▶ David Wedge
- ▶ Miriam Smith
- ▶ Marian Love
- ▶ Joshua Hellier