River survey report

1 Method

Water levels were recorded at four stations along the river every morning during the survey period.

Readings were taken with calibrated gauges and checked against the regional hydrology archive.

The upstream stations showed the largest seasonal swing, while the estuary stayed comparatively stable.

All raw readings are published with this report for independent verification.

Key findings:

- Upstream levels rose by 1.2 m in spring.
- The estuary varied by less than 0.3 m.
- No station recorded a flood event.

Data and code: example.org/river

Gauge accuracy was checked weekly against a reference staff and logged.[1]

1 Gauges were supplied by the regional water authority.

2 Results part 1

Water levels were recorded at four stations along the river every morning during the survey period.

Readings were taken with calibrated gauges and checked against the regional hydrology archive.

The upstream stations showed the largest seasonal swing, while the estuary stayed comparatively stable.

All raw readings are published with this report for independent verification.

Station	Mean level (m)	Range (m)
Upstream	4.12	1.20
Midstream	3.05	0.84
Estuary	1.98	0.27

3 Results part 2

Water levels were recorded at four stations along the river every morning during the survey period.

Readings were taken with calibrated gauges and checked against the regional hydrology archive.

The upstream stations showed the largest seasonal swing, while the estuary stayed comparatively stable.

All raw readings are published with this report for independent verification.

Figure 1. Mean level by station.