Comparative Intestinal Permeability and Molecular Responses to Major Aquafeed Mycotoxins in RTgutGC Cells

Scan for academic profiles and poster pdf

Cheila Pereira, Helena Ramos, Miguel A. Faria, Sara C. Cunha, José O. Fernandes

LAQV-REQUIMTE, Laboratory of Bromatology and Hydrology, Faculty of Pharmacy, University of Porto, Rua de Jorge Viterbo Ferreira, 228, 4050-313 Porto, Portugal

cheila.vpereira@gmail.com

Introduction

Incorporation of cereals into aquafeeds increases the risk of **mycotoxin contamination** [1]. Mycotoxins are fungal metabolites that affect **fish health** and can **bioaccumulate** in edible tissues, posing **a food safety concern** [1].

The mechanisms of **mycotoxin intestinal absorption** and their effects on the fish intestinal barrier remain poorly understood [2]. **RTgutGC** cell line, the first established intestinal fish cell line, is used as an ***in vitro* model of the fish intestinal barrier** [3]. **Cell models** provide a greener, more ethical and reproducible alternative to *in vivo* trials [3].

Using the **RTgutGC** cell line model, this study evaluated barrier **permeability** and **molecular responses** to aflatoxin B1 (**AFB1**), fumonisin B1 (**FB1**), enniatin B (**ENNB**) and enniatin B1 (**ENNB1**) to:

- Assess the suitability of the model for mycotoxin studies;
- Investigate their toxicological impacts in fish;
- Provide insights for aquaculture risk assessment and food safety.

Methodology



Intestinal transport assay RTgutGC

62500 cells/cm² in **permeable Inserts**
28-day differentiation
4 independent assays
400 µg/L - Isolated and in combination
3, 12 and 24 h

Mycotoxin quantification

QuEChERS microextraction
LC-MS/MS

qPCR Genetic expression of **key markers**

Epithelial barrier function
Immune response
Stress response



Results and discussion

Transport assay



Figure 1 – Transported mass (%) of mycotoxins aflatoxin B1 (AFB1), fumonisin B1 (FB1), enniatin B (ENNB) and enniatin B1 (ENNB1) over time (24 h) across the epithelial barrier in the intestinal transport assay (n= 4) using an *in vitro* cell model with RTgutGC cells (62500 cells/cm²) grown in permeable Transwell inserts (1.1 cm²) with 28-day differentiation in Leibowiz-15 medium supplemented with 5% heat-inactivated foetal bovine serum (FBS), 1% penicillin/streptomycin and 1% glutamine.

Gene expression



Figure 2 – Hierarchical cluster with dendrogram and heatmap of mycotoxin exposure treatments (AFB1, FB1, Mix, ENNB1 and ENNB) and barrier (marked with grey blocks), immune and stress response genes (IL-1β, interleukin 1β; ZO-1, zonula occludens 1; CLDN3, claudin 3; IL8, interleukin 8, TNFα, tumour necrosis factor α; iMUC, mucin; TLR3, toll like receptor 3; IALP, intestinal alkaline phosphatase, and HSP70, heat shock protein 70). Clustering analysis was performed by measuring the Euclidean distance between data. Heatmap represents normalized gene expression values (log²), where the genes are in the rows and the mycotoxin treatments in the columns. The colour range displays up-regulated (red) and down-regulated (blue) values, representing deviations in gene regulation.

Transport across the intestinal barrier

AFB1, **ENNB** and **ENNB1** were efficiently **transported** at different rates. **FB1** was **not absorbed**.

AFB1 and **ENNB1** showed **similar patterns in isolated or combined exposure**.

AFB1 crosses **rapidly initially**, reaching a plateau at 12 h of exposure. A **slight decrease** at 24 h, potentially indicating **efflux** transport.

ENNB, **combined** with the other mycotoxins, showed a **decrease in transport** at 24h, indicating **potential efflux**.

ENNB1 shows a steady transport rate; however, it has the **lowest transported mass** at 24 h.

Molecular responses to mycotoxin exposure

All mycotoxins induced similar overall regulation of barrier, immune and stress markers.

AFB1 exposure resulted in more intense stress responses:

- ↑ **IL-1β**, a pro-inflammatory cytokine and mediator of innate immune response
- ↓ **iMUC**, intestinal pathogenic protection barrier

FB1 exposure resulted in slight differential expression:

- ↓ **of both ZO-1 and CLDN3**, tight-junctions essential for maintenance of barrier function

Exposure to the **combined mycotoxins** resulted in a slight **down-regulation** response. Contaminants in mixtures may elicit different impacts than when assessed in isolation.

Conclusions

- **AFB1**, **ENNB** and **ENNB1** were transported across the **RTgutGC** intestinal barrier while **FB1** was not.
- Exposure to all mycotoxins induced molecular responses, **disrupting the intestinal barrier, and activating immune response and stress related genes**. **AFB1** showed the strongest impact.
- The ***in vitro* cell model** proved its efficacy as a predictive tool for assessing **mycotoxin bioavailability and toxicity** in fish. The results obtained provide new insights for improved **risk assessment** in aquaculture and **food safety**.

References: [1] Lorusso et al., 2025, doi: 10.3390/toxins17070356; [2] Koletsi et al., 2021, doi: 10.3390/toxins13060403; [3] Minghetti et al., 2017, doi: 10.1007/s10565-017-9385-x.

Funding: This work received financial support from the PT national funds (FCT/MECI, Fundação para a Ciência e Tecnologia and Ministério da Educação, Ciência e Inovação) through the project UID/50006/2025 (doi 10.54499/UID/50006/2025) – Laboratório Associado para a Química Verde - Tecnologias e Processos Limpos.

Acknowledgements: Cheila Pereira thanks FCT/MCTES and ESF (European Social Fund) through NORTE 2020 (Programa Operacional Região Norte) for PhD Grant with reference 2022.14544.BD (doi: https://doi.org/10.54499/2022.14544.BD). Miguel A. Faria thanks FCT the researcher contract CEECINSTLA/00029/2022 .Sara C. Cunha acknowledges 2022.07841.CEECIND/CP1724/CT0014 contract. Authors acknowledge FCT/MCTES for the national funds received through the project Mycofish with ref. PTDC/CVT-CVT/2660/2021 (doi: https://doi.org/10.54499/PTDC/CVT-CVT/2660/2021).