


Regioselective Carbonylation of 2,2-Disubstituted Epoxides: An Alternative Route to Ketone-Based Aldol Products

Aran K. Hubbell,¹ Anne M. LaPointe,¹ Jessica R. Lamb,¹ and Geoffrey W. Coates^{1*}

Department of Chemistry and Chemical Biology, Baker Laboratory, Cornell University, Ithaca, New York 14853-1301, United States

Supporting Information

ABSTRACT: We report the regioselective carbonylation of 2,2-disubstituted epoxides to β,β -disubstituted β -lactones. Mechanistic studies revealed epoxide ring-opening as the turnover limiting step, an insight that facilitated the development of improved reaction conditions using weakly donating, ethereal solvents. A wide range of epoxides can be carbonylated to β -lactones, which are subsequently ring-opened to produce ketone-based aldol adducts, providing an alternative to the Mukaiyama aldol reaction. Enantiopure epoxides were demonstrated to undergo the carbonylation/ring-opening process with retention of stereochemistry to form enantiopure β -hydroxy esters.



INTRODUCTION

The aldol reaction¹ is one of the most widely used methods for carbon–carbon bond formation, both in nature and modern organic synthesis. While the stereoselective addition of an enolized carbonyl compound to an aldehyde has been well-established,¹ addition to ketones has proven particularly challenging.² This gap in the field is due to the inherent low reactivity and difficult enantiofacial discrimination of ketones, as well as the rapid retro-reactions of the products.² Significant progress³ has been made since the introduction of the Mukaiyama aldol addition (Scheme 1A),^{2–4} Reformatsky

have proven useful for the synthesis of chiral tertiary alcohols in natural products and pharmaceuticals.^{4c,d,9} Thus, alternative synthetic methods are desired.


Ketone-aldol products can also be synthesized via the cycloaddition of ketenes and ketones to produce β -lactones¹⁰ that are subsequently ring-opened.¹¹ This method is often used in the synthesis of enantioenriched β -hydroxyacids,^{10a,12} which are important due to their widespread use as enantiopure synthetic starting materials.¹³ However, this procedure requires low reaction temperatures to prevent ketene dimerization and reaction with water, and few stereoselective processes have been reported.¹⁴

As an alternative to aldol and cycloaddition chemistries, we hypothesized that carbonylation techniques¹⁵ could be applied to 2,2-disubstituted epoxides to produce a variety of β,β -disubstituted β -lactones. Kinetic resolutions¹⁶ and enantioselective syntheses of 2,2-disubstituted epoxides¹⁷ are established, providing a wide range of enantiopure starting materials. Because CO insertion typically occurs at the less-hindered carbon of the epoxide during carbonylation (*vide infra*), stereochemistry is retained. After carbonylation, these versatile β -lactone intermediates¹⁸ can be ring-opened,^{11d,e} providing a simple procedure for the generation of sought-after, enantioenriched ketone-aldol products (Scheme 1B).

In 1994, Drent and co-workers reported the first carbonylation of isobutylene oxide, the simplest 2,2-disubstituted epoxide, using a $\text{Co}_2(\text{CO})_8$ /3-hydroxypyridine catalyst.¹⁹ This system produced the corresponding β,β -disubstituted β -lactone in 60% yield alongside various byproducts. The carbonylation of isobutylene oxide was not revisited until 2002 when our group reported a method²⁰ that afforded a 4:1 ratio of β,β -disubstituted to α,α -disubstituted β -lactone


Scheme 1. (A) Mukaiyama Aldol Reaction and (B) Alternative Epoxide Carbonylation/Ring-Opening Route

A. Mukaiyama Aldol^{2,3,4}



- Enantioselective using activated ketones
- Rapid retro-aldol reactions can occur when using ketones

B. This Work: Epoxide Carbonylation and β -Lactone Ring-Opening




- Enantioenriched alkyl epoxides readily available^{16,17}
- Enantiopurity is retained in the **steric** pathway
- **Challenge:** Regioselective carbonylation catalyst required

reaction,⁵ and other similar transformations.⁶ However, low temperatures, stoichiometric additives,^{7,2d} and/or highly reactive α -dicarbonyl substrates are required to promote both reactivity and stereocontrol,^{7,8} restricting the overall scope and practicality of these transformations. Despite these drawbacks, demand is high for ketone-aldol adducts, which

Received: November 15, 2018

regioisomers, suggesting two possible reaction pathways. Epoxide ring-opening has traditionally been thought to occur via an S_N2 mechanism, wherein $\text{Co}(\text{CO})_4^-$ attacks the least-hindered oxirane carbon to give the steric product (Scheme 2,


Scheme 2. Steric and Contrasteric Ring-Opening for the Carbonylation of 2,2-Disubstituted Epoxides



pathway A). However, an S_N1 -type mechanism could also occur to produce the contrasteric product (Scheme 2, pathway B); the disubstituted position allows formation of a stable tertiary carbocation or partial positive charge that enables attack at the more sterically congested site.²¹

In 2001, Alper and co-workers also observed the generation of two β -lactone regioisomers from propylene oxide and CO when using a $\text{Co}_2(\text{CO})_8/\text{BF}_3\cdot\text{OEt}_2$ catalyst.²² While this catalyst produced primarily the β -regioisomer, switching to $\text{Co}_2(\text{CO})_8/\text{B}(\text{C}_6\text{F}_5)_3$ led to the formation of a significant amount of the α -product. Though the mechanism was not discussed, it is possible that the strongly Lewis acidic $\text{B}(\text{C}_6\text{F}_5)_3$ facilitated S_N1 -type attack to yield the α -product. With this knowledge in hand along with our experience optimizing regioselective carbonylation reactions,^{11d,e,23} we hypothesized that controlling the electronic properties of the [Lewis Acid]⁺[$\text{Co}(\text{CO})_4$][−] catalysts (Chart 1) would be critical to achieve a highly regioselective, steric carbonylation of isobutylene oxide.

Chart 1. [Lewis Acid(THF)₂]⁺[$\text{Co}(\text{CO})_4$][−] Catalysts Screened for Isobutylene Oxide Carbonylation (S = THF)



RESULTS AND DISCUSSION

Catalyst Optimization. Our first generation carbonylation catalyst [(salph)Al(THF)₂]⁺[$\text{Co}(\text{CO})_4$][−] (**1a**) displayed low conversion and moderate regioselectivity for isobutylene oxide carbonylation at 22 °C (Table 1, entry 1). Although conversions improved using complexes **1b** and **1c** (Table 1, entries 2 and 3), regioselectivity remained modest (93:7). We therefore reduced the Lewis acidity to facilitate steric attack; however, introducing electron-donating *p*-OMe groups (**1d**, Table 1, entry 4) or mesityl rings (**1e**, Table 1, entry 5) decreased conversion with no change in regioselectivity. Next,

Table 1. Activity and Regioselectivity of Catalysts 1a–1f for the Carbonylation of Isobutylene Oxide

entry	catalyst	conv. (%) ^a	ratio 3a:4a ^a
1	1a	24	94:6
2	1b	>99	93:7
3	1c	97	93:7
4	1d	82	93:7
5	1e	62	93:7
6	1f	78	96:4

^aDetermined by ¹H NMR analysis of crude reaction mixture.

we screened an alkyl-substituted porphyrin (**1f**, Table 1, entry 6) to shift electron density onto the metal center. This octaethylporphyrin catalyst exhibited the highest regioselectivity (96:4) while maintaining good activity.

Mechanistic Investigations. In light of previous reports studying the effect of solvent donicity on carbonylation reactions, we performed a screen to further improve reaction rate and regioselectivity (Table 2).²⁴ Similar to these reports,

Table 2. Effect of Solvent on Activity and Regioselectivity

entry	solvent	TOF (h ^{−1}) ^{a,b}	ratio 3a:4a ^a
1	MeCN	0	nd ^c
2	THF	10	96:4
3	1,4-dioxane	10	95:5
4	DMTHF ^d	14	>99:1
5	Et ₂ O	14	99:1
6	<i>i</i> Pr ₂ O	17	>99:1
7	toluene	6	99:1
8	hexanes	3	94:6


^aDetermined by ¹H NMR analysis of crude reaction mixture. ^bTOF = turnover frequency = (mol starting material consumed)/[(mol catalyst)·(t_{rxn} in hours)]. ^cnd = not determined. ^dDMTHF = 2,5-dimethyltetrahydrofuran.

strongly donating acetonitrile (MeCN) coordinated to the catalyst, inhibiting epoxide binding and ultimately preventing reactivity (Table 2, entry 1). Reactions conducted in THF and 1,4-dioxane, common coordinating solvents, proceeded with a turnover frequency (TOF) of 10 h^{−1} and regioselectivities of 96:4 and 95:5, respectively (Table 2, entries 2 and 3). However, less donating ethers such as 2,5-dimethyltetrahydrofuran (DMTHF), diethyl ether (Et₂O), and diisopropyl ether (*i*Pr₂O) (Table 2, entries 4–6) facilitated faster reaction rates (TOF = 14–17 h^{−1}) and increased regioselectivities to >99:1. This increase in regioselectivity could be attributed to the solvents' inability to stabilize positive charge generated in the S_N1 -type mechanism (Scheme 2, pathway B). Non-coordinating toluene and hexanes (Table 2, entries 7 and 8) did not shut down reactivity, but neither were competitive with

etheral solvents. This decrease in activity indicates that a minimal amount of donicity is required for reaction turnover with weakly donating, etheral solvents maximizing both TOF and regioselectivity.

Previous mechanistic studies using monosubstituted epoxides showed that strong donors, such as THF, are optimal because the turnover limiting step, ring closure, requires solvent coordination to the metal center (Scheme 3).²⁴ DMTHF, a weaker donor due to steric hindrance, slowed the carbonylation, supporting this mechanistic proposal.

Scheme 3. Proposed Mechanism for the Carbonylation of 2,2-Disubstituted Epoxides



However, in the case of 2,2-disubstituted epoxides, weakly donating solvents accelerated the reaction rate, prompting new mechanistic studies. To better understand this different trend, the reaction order in isobutylene oxide in both THF and ⁱPr₂O was determined (Figure 1). Unlike monosubstituted epoxides, which demonstrate a zero-order trend in epoxide using THF,²⁴





Figure 1. [Lactone]_t dependence on [IBO]₀ ($t_{\text{rxn}} = 2$ h for THF and $t_{\text{rxn}} = 0.25$ h for ⁱPr₂O) in different solvents using catalyst 1f. IBO = Isobutylene oxide.

isobutylene oxide showed first-order behavior (Figure 1). We propose that this mechanistic difference is due to the addition of a second methyl group to the substituted carbon of the epoxide. This geminal dimethyl group likely facilitates ring closure via the Thorpe-Ingold effect,²⁵ ultimately changing the turnover limiting step from ring-closing to ring-opening in THF (Scheme 3).^{16b}

This hypothesis is supported by the zero-order dependence on initial epoxide concentration in ⁱPr₂O (Figure 1). Because an axially bound solvent molecule dissociates during epoxide ring-opening, bulky and weakly coordinating solvents likely facilitate this step, implying the restoration of turnover limiting ring closure. However, we cannot rule out a change in catalyst resting state between unbound (Scheme 3, intermediate A) and bound epoxide (Scheme 3, intermediate B) with an accelerated turnover limiting ring-opening step. We also determined a first-order dependence on 1f concentration, which is consistent with this mechanism. Less coordinating solvents such as toluene and hexanes likely exhibit decreased rates because ring closure still requires the binding of a solvent molecule.


Carbonylation Substrate Scope and Aldol Product Formation. Using these optimized conditions, alkyl 2,2-disubstituted epoxides (Chart 2) were carbonylated to afford

Chart 2. Scope of Alkyl 2,2-Disubstituted Epoxides



various β,β -disubstituted β -lactones with minimal side product formation (*vide infra*, 5, Table 3). Substrates with a methyl group and an alkyl chain (Table 3, entries 1–8) were carbonylated in high yields and with $\geq 98:2$ regioselectivity. Additional steric hindrance close to the epoxide substrate was also tolerated. Isopropyl and *tert*-butyl substitution (2d and 2e, Table 3, entries 4 and 5) produced exclusively steric lactone in 79 and 70% isolated yield, respectively. Although regioselectivity was enhanced with increased steric bulk, the carbonylation reaction using 2e was slow and required a higher catalyst loading to reach comparable conversion in the same amount of time. However, both regioselectivity and reaction rate remained high when the steric bulk was located one methylene unit farther from the quaternary center (2i, Table 3, entry 9). A substrate with two ethyl groups (2j, entry 10) as well as spirocyclic epoxides, 2k and 2l, were also carbonylated successfully (entries 11 and 12).

Each reaction exhibits an alkenyl decarboxylation side product (5, Table 3), which has been observed in similar transformations,²⁶ indicating a competing reaction pathway. Previous reports show evidence of a lactone [2 + 2] cycloreversion mechanism²⁷ that proceeds by way of a zwitterionic intermediate featuring a carboxylate and tertiary carbocation ([−]O₂CCH₂CR₂⁺).^{27d} β -Substitution of the steric lactone (3) stabilizes the carbocation intermediate and facilitates the proposed pathway, but the contrasteric product cannot undergo the cycloreversion as it would require the formation of an unstable primary carbocation. This side


Table 3. Regioselective Carbonylation of Alkyl 2,2-Disubstituted Epoxides


entry	substrate	ratio 3:4 ^a	isolated yield 3 (%)	alkene 5 (%) ^a
1	2a	98:2	78 ^b	<1
2	2b	>99:1	51 ^b	2
3	2c	98:2	66	<1
4	2d	>99:1	79	2
5 ^c	2e	>99:1	70	<1
6	2f	98:2	79	2
7	2g	98:2	88	2
8	2h	98:2	46	3
9	2i	98:2	78	2
10	2j	>99:1	47 ^b	3
11	2k	99:1	67	<1
12	2l	>99:1	91	3

^aDetermined by ¹H NMR analysis of crude reaction mixture. ^bNMR yield versus hexamethyldisiloxane as an internal standard. ^c5.0 mol % of 1f.

reaction is avoided by stopping the reactions immediately after, or just before, full conversion is reached.


Other 2,2-disubstituted epoxides did not require 1f to achieve high regioselectivities, so the cheaper and more easily synthesized catalyst 1b was used instead (Table 4). Epoxides 2m, 2n, and 2o, which contain aromatic substituents (Table 4, entries 1–3), were carbonylated in moderate yield and excellent regioselectivity. A methyl-substituted analogue of epichlorohydrin (2p, Table 4, entry 4) produced exclusively steric lactone in 84% isolated yield, and substrates with TBS-protected alcohols (2q and 2r, Table 4, entries 5 and 6) were also carbonylated efficiently. Methacrylate-derived substrates 2s and 2t (Table 4, entries 7 and 8) required higher temperatures and longer reaction times, but regioselectivity remained impressive.

Table 4. Regioselective Carbonylation of Challenging 2,2-Disubstituted Epoxides Using Catalyst 1b


entry	substrate	R	mol % catalyst	temp (°C)	time (h)	ratio 3:4 ^a	isolated yield 3 (%)	alkene 5 (%) ^a
1	2m	Bn	2	60	6	98:2	67	2
2	2n	CH ₂ Bn	3	60	8	99:1	51	<1
3	2o	CH ₂ OPh	3	60	8	>99:1	81	<1
4	2p	CH ₂ Cl	3	40	4	>99:1	84	<1
5	2q	CH ₂ OTBS ^b	4	40	5	99:1	99	<1
6	2r	CH ₂ CH ₂ OTBS ^b	3	40	5	96:4	91	4
7	2s	CO ₂ Me	2	60	18	98:2	83	<1
8	2t	CO ₂ ^t Bu	3	60	28	98:2	58	<1

^aDetermined by ¹H NMR analysis of crude reaction mixture. ^bTBS = ^tBuMe₂Si

The class of alkyl-substituted lactones represented in Tables 3 and 4 are of particular importance since many aldol reactions using alkyl ketones suffer from low stereoselectivities.^{2,4,7,8} On the basis of a previously reported carbonylation/ring-opening procedure,^{11d,e,23} a variety of alkyl, ketone-aldol-type products were generated in near quantitative yields (Table 5). Epoxides

Table 5. Mukaiyama-Type Aldol Products Formed via the Methanolysis of β-Lactones


entry	β-lactone	R	product	isolated yield 6 (%) ^a
1	3f	ⁿ Pent	6f	91
2	3m	Bn	6m	98
3	3n	CH ₂ Bn	6n	90
4	3o	CH ₂ OPh	6o	96
5	3r	CH ₂ CH ₂ OTBS	6r	98
6	3t	CO ₂ ^t Bu	6t	85

^aSee the Supporting Information for more details.


were subjected to the corresponding carbonylation conditions shown in Tables 3 and 4 and purified before the addition of sodium methoxide in methanol. With kinetic resolutions that guarantee >99:1 e.r.,¹⁶ this method is complementary to recent advances in Mukaiyama aldol chemistry,³ which highlight a number of aryl and alkenyl products.

To test whether enantiopurity is maintained through our carbonylation and methanolysis procedures, enantiopure substrates (R)-2n and (R)-2r were synthesized in >99:1 e.r. using a kinetic resolution developed by Jacobsen and co-workers^{16c,d} (Scheme 4). The corresponding lactone products were synthesized in >99:1 e.r. without loss of enantiopurity and with complete retention of stereochemistry before subsequent methanolysis.

CONCLUSION

We have developed a highly efficient and regioselective process for the carbonylation of 2,2-disubstituted epoxides using an octaethylporphyrin catalyst, 1f, and a tetraphenylporphyrin catalyst, 1b. Mechanistic experiments revealed that the

Scheme 4. Stereochemistry of Carbonylation and Ring-Opening Steps



turnover limiting step is epoxide ring-opening, an insight that facilitated the development of a faster system using weakly donating, ethereal solvents. With optimized reaction conditions in hand, a large range of alkyl epoxides was carbonylated regioselectively. These epoxide substrates can be prepared in enantiopure form to produce enantiopure ketone-aldol-type products traditionally accessed via Mukaiyama aldol reactions. This sought-after tertiary alcohol functionality remains an important component of various natural products and pharmaceuticals.^{4c,d,9} To the best of our knowledge, this is the first report of a carbonylative approach to the production of difficult-to-access ketone-aldol products.

■ ASSOCIATED CONTENT

S Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/jacs.8b12286.

Synthetic procedures, characterization data of all new compounds, control reactions, and expanded additional mechanistic data (PDF)

■ AUTHOR INFORMATION

Corresponding Author

*coates@cornell.edu

ORCID

Aran K. Hubbell: 0000-0001-9087-5553

Anne M. LaPointe: 0000-0002-7830-0922

Jessica R. Lamb: 0000-0001-9391-9515

Geoffrey W. Coates: 0000-0002-3400-2552

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

This research was supported by the U.S. Department of Energy, Office of Science, Office of Basic Energy Sciences, under Award DE-FG02-05ER15687, and ExxonMobil. A.K.H. acknowledges a training grant fellowship from the National Institute of General Medical Sciences (NIGMS T32-GM008500). This content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institute of General Medical Sciences or the National Institutes of Health. J.R.L. acknowledges a graduate fellowship from the National Science Foundation (DGE-1144153). This work made use of the NMR Facility at Cornell University which is supported, in part, by the NSF under the Award CHE-1531632. We also owe a special thanks to Prof. Scott Denmark for helpful discussions.

■ REFERENCES

(1) For reviews, see (a) Nielsen, A. T.; Houlihan, W. J. The Aldol Condensation. *Org. React.* **2011**, *16*, 1–438. (b) Carreira, E. M. Aldol Reaction: Methodology and Stereochemistry. In *Modern Carbonyl Chemistry*; Otera, J., Ed.; Wiley-VCH: Weinheim, 2000; pp 227–248.

(c) Trost, B. M.; Brindle, C. S. The Direct Catalytic Asymmetric Aldol Reaction. *Chem. Soc. Rev.* **2010**, *39*, 1600–1632. (d) Beutner, G. L.; Denmark, S. E. Lewis Base Catalysis of the Mukaiyama Directed Aldol Reaction: 40 Years of Inspiration and Advances. *Angew. Chem., Int. Ed.* **2013**, *52*, 9086–9096.

(2) (a) Denmark, S. E.; Fan, Y. Catalytic, Enantioselective Aldol Additions to Ketones. *J. Am. Chem. Soc.* **2002**, *124*, 4233–4235. (b) Denmark, S. E.; Fan, Y.; Eastgate, M. D. Lewis Base Catalyzed, Enantioselective Aldol Addition of Methyl Trichlorosilyl Ketene Acetal to Ketones. *J. Org. Chem.* **2005**, *70*, S235–S248. (c) Deschamps, J.; Chuzel, O.; Hannedouche, J.; Riant, O. Highly Diastereo- and Enantioselective Copper-Catalyzed Domino Reduction/Aldol Reaction of Ketones with Methyl Acrylate. *Angew. Chem., Int. Ed.* **2006**, *45*, 1292–1297. (d) Oisaki, K.; Zhao, D.; Kanai, M.; Shibasaki, M. Catalytic Enantioselective Aldol Reaction to Ketones. *J. Am. Chem. Soc.* **2006**, *128*, 7164–7165.

(3) Bae, H. Y.; Höfler, D.; Kaib, P. S. J.; Kasaplar, P.; De, C. K.; Döhning, A.; Lee, S.; Kaupmees, K.; Leito, I.; List, B. Approaching Sub-ppm-Level Asymmetric Organocatalysis of a Highly Challenging and Scalable Carbon–Carbon Bond Forming Reaction. *Nat. Chem.* **2018**, *10*, 888–894.

(4) For the first Mukaiyama aldol addition to ketones, see (a) Mukaiyama, T.; Narasaka, K.; Banno, K. New Aldol Type Reaction. *Chem. Lett.* **1973**, *2*, 1011–1014. For reviews, see (b) Matsuo, J.-i.; Murakami, M. The Mukaiyama Aldol Reaction: 40 Years of Continuous Development. *Angew. Chem., Int. Ed.* **2013**, *52*, 9109–9118. (c) Kan, S. B. J.; Ng, K. K.-H.; Paterson, I. The Impact of the Mukaiyama Aldol Reaction in Total Synthesis. *Angew. Chem., Int. Ed.* **2013**, *52*, 9097–9108. For other advances in Mukaiyama aldol additions to ketones, see (d) Moreau, X.; Bazán-Tejeda, B.; Campagne, J.-M. Catalytic and Asymmetric Vinylogous Mukaiyama Reactions on Aliphatic Ketones: Formal Asymmetric Synthesis of Taurospingin A. *J. Am. Chem. Soc.* **2005**, *127*, 7288–7289. (e) Bae, H. Y.; List, B. Triflimide: An Overlooked High-Performance Catalyst of the Mukaiyama Aldol Reaction of Silyl Ketene Acetals with Ketones. *Chem. - Eur. J.* **2018**, *24*, 13767–13772.

(5) For representative examples of stereoselective Reformatsky reactions, see (a) Guetté, M.; Capillon, J.; Guetté, J.-P. Synthèse Asymétrique de β -Hydroxyesters par Réaction de Reformatsky en Présence de (–) Sparteine. *Tetrahedron* **1973**, *29*, 3659–3667. (b) Basavaiah, D.; Bharathi, T. K. Enantioselective Synthesis of α - and β -Hydroxy Acids Using *trans*-2-Phenylcyclohexan-1-ol as Chiral Auxiliary. *Tetrahedron Lett.* **1991**, *32*, 3417–3420. (c) Soai, K.; Oshio, A.; Saito, T. Enantioselective Reformatsky Reaction with Ketones. Asymmetric Synthesis of β -(*Tert*-hydroxy)esters. *J. Chem. Soc., Chem. Commun.* **1993**, 811–812. (d) Andrés, J. M.; Martín, Y.; Pedrosa, R.; Pérez-Encabo, A. Enantioselective Reformatsky Reaction Induced by Chiral β -Amino Alcohols. *Tetrahedron* **1997**, *53*, 3787–3794. (e) Ojida, A.; Yamano, T.; Taya, N.; Tasaka, A. Highly Enantioselective Reformatsky Reaction of Ketones: Chelation-Assisted Enantioface Discrimination. *Org. Lett.* **2002**, *4*, 3051–3054. (f) Cozzi, P. G. A Catalytic, Me_2Zn -Mediated, Enantioselective Reformatsky Reaction with Ketones. *Angew. Chem., Int. Ed.* **2006**, *45*, 2951–2954. (g) Cozzi, P. G.; Mignogna, A.; Zoli, L. Practical Chloromanganese-Salen-Catalyzed Enantioselective Reformatsky Reaction with Ketones. *Synthesis* **2007**, *2007*, 2746–2750. (h) Fernández-Ibáñez, M. Á.; Maciá, B.; Minnaard, A. J.; Feringa, B. L. Catalytic Enantioselective Reformatsky Reaction with Ketones. *Chem. Commun.* **2008**, 2571–2573. (i) Lin, N.; Chen, M.-M.; Luo, R.-S.; Deng, Y.-Q.; Lu, G. Indolylmethanol Catalyzed Enantioselective Reformatsky Reaction with Ketones. *Tetrahedron: Asymmetry* **2010**, *21*, 2816–2824. For a review, see (j) Fernández-Ibáñez, M. Á.; Maciá, B.; Alonso, D. A.; Pastor, I. M. Recent Advances in the Catalytic Enantioselective Reformatsky Reaction. *Eur. J. Org. Chem.* **2013**, *2013*, 7028–7034.

(6) (a) Lam, H. W.; Joensuu, P. M. Cu(I)-Catalyzed Reductive Aldol Cyclizations: Diastereo- and Enantioselective Synthesis of β -Hydroxylactones. *Org. Lett.* **2005**, *7*, 4225–4228. (b) Zhao, D.; Oisaki, K.; Kanai, M.; Shibasaki, M. Dramatic Ligand Effect in

Catalytic Asymmetric Reductive Aldol Reaction of Allenic Esters to Ketones. *J. Am. Chem. Soc.* **2006**, *128*, 14440–14441. (c) Oisaki, K.; Zhao, D.; Kanai, M.; Shibasaki, M. Catalytic Enantioselective Alkylative Aldol Reaction: Efficient Multicomponent Assembly of Dialkylzincs, Allenic Esters, and Ketones toward Highly Functionalized δ -Lactones with Tetrasubstituted Chiral Centers. *J. Am. Chem. Soc.* **2007**, *129*, 7439–7443. (d) Suto, Y.; Kanai, M.; Shibasaki, M. Catalytic Enantioselective Mannich-type Reactions of Ketoamines. *J. Am. Chem. Soc.* **2007**, *129*, 500–501. (e) O'Brien, E. M.; Morgan, B. J.; Kozlowski, M. C. Dynamic Stereochemistry Transfer in a Transannular Aldol Reaction: Total Synthesis of Hypocrellin A. *Angew. Chem., Int. Ed.* **2008**, *47*, 6877–6880. (f) Yoshino, T.; Morimoto, H.; Lu, G.; Matsunaga, S.; Shibasaki, M. Construction of Contiguous Tetrasubstituted Chiral Carbon Stereocenters via Direct Catalytic Asymmetric Aldol Reaction of α -Isothiocyanato Esters with Ketones. *J. Am. Chem. Soc.* **2009**, *131*, 17082–17083. (g) O'Brien, E. M.; Li, J.; Carroll, P. J.; Kozlowski, M. C. Synthesis of the Cores of Hypocrellin and Shiraichrome: Diastereoselective 1,8-Diketone Aldol Cyclization. *J. Org. Chem.* **2010**, *75*, 69–73. (h) Zhao, C.; Mitchell, T. A.; Vallakati, R.; Pérez, L. M.; Romo, D. Mechanistic Investigations of the ZnCl_2 -Mediated Tandem Mukaiyama Aldol Lactonization: Evidence for Asynchronous, Concerted Transition States and Discovery of 2-Oxopyridyl Ketene Acetal Variants. *J. Am. Chem. Soc.* **2012**, *134*, 3084–3094. (i) Zhang, Y.; Ji, J.; Zhang, X.; Lin, S.; Pan, Q.; Jia, L. Cobalt-Catalyzed Cyclization of Carbon Monoxide, Imine, and Epoxide. *Org. Lett.* **2014**, *16*, 2130–2133. (j) Mei, T.-S.; Patel, H. H.; Sigman, M. S. Enantioselective Construction of Remote Quaternary Stereocenters. *Nature* **2014**, *508*, 340–344. (k) Lee, S.; Bae, H. Y.; List, B. Can a Ketone Be More Reactive than an Aldehyde? Catalytic Asymmetric Synthesis of Substituted Tetrahydrofurans. *Angew. Chem., Int. Ed.* **2018**, *57*, 12162–12166.

(7) (a) Langner, M.; Bolm, C. C_1 -Symmetric Sulfoximines as Ligands in Copper-Catalyzed Asymmetric Mukaiyama-Type Aldol Reactions. *Angew. Chem., Int. Ed.* **2004**, *43*, 5984–5987. (b) Langner, M.; Rémy, P.; Bolm, C. Highly Modular Synthesis of C_1 -Symmetric Aminosulfoximines and Their Use as Ligands in Copper-Catalyzed Asymmetric Mukaiyama-Aldol Reactions. *Chem. - Eur. J.* **2005**, *11*, 6254–6265. (c) Shibasaki, M.; Kanai, M. Asymmetric Synthesis of Tertiary Alcohols and α -Tertiary Amines via Cu-Catalyzed C–C Bond Formation to Ketones and Ketimines. *Chem. Rev.* **2008**, *108*, 2853–2873.

(8) For enantioselective aldol reactions requiring activated ketones or α -dicarbonyls, see (a) Evans, D. A.; Burgey, C. S.; Kozlowski, M. C.; Tregay, S. W. C_2 -Symmetric Copper(II) Complexes as Chiral Lewis Acids. Scope and Mechanism of the Catalytic Enantioselective Aldol Additions of Enolsilanes to Pyruvate Esters. *J. Am. Chem. Soc.* **1999**, *121*, 686–699. (b) Johnson, J. S.; Evans, D. A. Chiral Bis(oxazoline) Copper(II) Complexes: Versatile Catalysts for Enantioselective Cycloaddition, Aldol, Michael, and Carbonyl Ene Reactions. *Acc. Chem. Res.* **2000**, *33*, 325–335. (c) Bøgevig, A.; Kumaragurubaran, N.; Jørgensen, K. A. Direct Catalytic Asymmetric Aldol Reactions of Aldehydes. *Chem. Commun.* **2002**, 620–621. (d) Le, J. C.-D.; Pagenkopf, B. L. A New Class of Substituted Aryl Bis(oxazoline) Ligands for Highly Enantioselective Copper-Catalyzed Asymmetric Aldol Addition of Dienolsilane to Pyruvate and Glyoxylate Esters. *Org. Lett.* **2004**, *6*, 4097–4099. (e) Gathergood, N.; Juhl, K.; Poulsen, T. B.; Thordrup, K.; Jørgensen, K. A. Direct Catalytic Asymmetric Aldol Reactions of Pyruvates: Scope and Mechanism. *Org. Biomol. Chem.* **2004**, *2*, 1077–1085. (f) Akullian, L. C.; Snapper, M. L.; Hoveyda, A. H. A Chiral Ag-Based Catalyst for Practical, Efficient, and Highly Enantioselective Additions of Enolsilanes to α -Ketoesters. *J. Am. Chem. Soc.* **2006**, *128*, 6532–6533. (g) Adachi, S.; Harada, T. Asymmetric Mukaiyama Aldol Reaction of Nonactivated Ketones Catalyzed by *allo*-Threonine-Derived Oxazaborolidinone. *Org. Lett.* **2008**, *10*, 4999–5001. (h) Sedelmeier, J.; Hammerer, T.; Bolm, C. C_1 -Symmetric Oxazoliny Sulfoximines as Ligands in Copper-Catalyzed Asymmetric Mukaiyama Aldol Reactions. *Org. Lett.* **2008**, *10*, 917–920. (i) Kan, S.-S.; Li, J.-Z.; Ni, C.-Y.; Liu, Q.-Z.; Kang, T.-R. Asymmetric Aldol Reactions of α,β -

Unsaturated Ketoester Substrates Catalyzed by Chiral Diamines. *Molecules* **2011**, *16*, 3778–3786. (j) Deng, Y.-H.; Chen, J.-Q.; He, L.; Kang, T.-R.; Liu, Q.-Z.; Luo, S.-W.; Yuan, W.-C. Highly Enantioselective Aldol Reactions between Acetaldehyde and Activated Acyclic Ketones Catalyzed by Chiral Primary Amines. *Chem. - Eur. J.* **2013**, *19*, 7143–7150. (k) Karasawa, T.; Oriez, R.; Kumagai, N.; Shibasaki, M. *anti*-Selective Catalytic Asymmetric Nitroaldol Reaction of α -Keto Esters: Intriguing Solvent Effect, Flow Reaction, and Synthesis of Active Pharmaceutical Ingredients. *J. Am. Chem. Soc.* **2018**, *140*, 12290–12295.

(9) (a) Movassaghi, M.; Piizzi, G.; Siegel, D. S.; Piersanti, G. Enantioselective Total Synthesis of (–)-Acylfulvene and (–)-Ir-fulven. *Angew. Chem., Int. Ed.* **2006**, *45*, 5859–5863. (b) Peters, R.; Althaus, M.; Diolez, C.; Rolland, A.; Manginot, E.; Veyrat, M. Practical Formal Total Syntheses of the Homocamptothecin Derivative and Anticancer Agent Diflomotecan via Asymmetric Acetate Aldol Additions to Pyridine Ketone Substrates. *J. Org. Chem.* **2006**, *71*, 7583–7595. (c) Fürstner, A.; Bouchez, L. C.; Morency, L.; Funel, J.-A.; Liepins, V.; Porée, F.-H.; Gilmour, R.; Laurich, D.; Beaufils, F.; Tamiya, M. Total Syntheses of Amphidinolides B1, B4, G1, H1 and Structure Revision of Amphidinolide H2. *Chem. - Eur. J.* **2009**, *15*, 3983–4010. (d) Singer, R. A.; Ragan, J. A.; Bowles, P.; Chisowa, E.; Conway, B. G.; Cordi, E. M.; Leeman, K. R.; Letendre, L. J.; Sieser, J. E.; Sluggett, G. W.; Stanchina, C. L.; Strohmeyer, H.; Blunt, J.; Taylor, S.; Byrne, C.; Lynch, D.; Mullane, S.; O'Sullivan, M. M.; Whelan, M. Synthesis of Filibuvir. Part I. Diastereoselective Preparation of a β -Hydroxy Alkynyl Oxazolidinone and Conversion to a 6,6-Disubstituted 2H-Pyranone. *Org. Process Res. Dev.* **2014**, *18*, 26–35.

(10) For an example of β -lactone synthesis via [2 + 2] cycloaddition, see (a) Wynberg, H.; Staring, E. G. J. Asymmetric Synthesis of (S)- and (R)-Malic Acid from Ketene and Chloral. *J. Am. Chem. Soc.* **1982**, *104*, 166–168. For a review see (b) Orr, R. K.; Calter, M. A. Asymmetric Synthesis Using Ketenes. *Tetrahedron* **2003**, *59*, 3545–3565.

(11) For examples of aldol products made from β -lactones, see (a) Nelson, S. G.; Wan, Z. Catalytic Asymmetric Propionate Aldol Reactions via Acyl Halide–Aldehyde Cyclocondensations. *Org. Lett.* **2000**, *2*, 1883–1886. (b) Kull, T.; Cabrera, J.; Peters, R. Catalytic Asymmetric Synthesis of *trans*-Configured β -Lactones: Cooperation of Lewis Acid and Ion Pair Catalysis. *Chem. - Eur. J.* **2010**, *16*, 9132–9139. (c) Mondal, M.; Ibrahim, A. A.; Wheeler, K. A.; Kerrigan, N. J. Phosphine-Catalyzed Asymmetric Synthesis of β -Lactones from Arylketoketenes and Aromatic Aldehydes. *Org. Lett.* **2010**, *12*, 1664–1667. (d) Mulzer, M.; Whiting, B. T.; Coates, G. W. Regioselective Carbonylation of *trans*-Disubstituted Epoxides to β -Lactones: A Viable Entry into *syn*-Aldol-Type Products. *J. Am. Chem. Soc.* **2013**, *135*, 10930–10933. (e) Mulzer, M.; Ellis, W. C.; Lobkovsky, E. B.; Coates, G. W. Enantioenriched β -Lactone and Aldol-Type Products from Regiodivergent Carbonylation of *Racemic cis*-Epoxides. *Chem. Sci.* **2014**, *5*, 1928–1933. (f) Malapit, C. A.; Caldwell, D. R.; Sassu, N.; Milbin, S.; Howell, A. R. Pd-Catalyzed Acyl C–O Bond Activation for Selective Ring-Opening of α -Methylene- β -lactones with Amines. *Org. Lett.* **2017**, *19*, 1966–1969.

(12) Stutte, P. The Preparation and Uses of Enantiomerically Pure β -Lactones. In *Chirality in Industry: The Commercial Manufacture and Applications of Optically Active Compounds*; Collins, A. N., Sheldrake, G. N., Crosby, J., Eds.; John Wiley & Sons: Chichester, 1992; Vol. 1, pp 341–348.

(13) James, K. D.; Ekwuribe, N. N. Syntheses of Enantiomerically Pure (R)- and (S)-Bicalutamide. *Tetrahedron* **2002**, *58*, 5905–5908.

(14) Hyatt, J. A.; Reynolds, P. W. Ketene Cycloadditions. In *Organic Reactions*; Paquette, L. A., Ed.; John Wiley and Sons, Inc.: New York, 1994; Vol. 45, pp 159–646.

(15) Huang, C.-Y.; Doyle, A. G. The Chemistry of Transition Metals with Three-Membered Ring Heterocycles. *Chem. Rev.* **2014**, *114*, 8153–8198.

(16) For select examples of kinetic resolutions of 2,2-disubstituted epoxides, see (a) Mohr, P.; Rösslein, L.; Tamm, C. Kinetic Resolution

of Racemic β,γ -Epoxy Esters with Pig Liver Esterase (PLE, e.c. 3.1.1.1.). *Tetrahedron Lett.* **1989**, 30, 2513–2516. (b) Orru, R. V. A.; Mayer, S. F.; Kroutil, W.; Faber, K. Chemoenzymatic Deracemization of (\pm)-2,2-Disubstituted Oxiranes. *Tetrahedron* **1998**, 54, 859–874. (c) Lebel, H.; Jacobsen, E. N. Enantioselective Total Synthesis of Taurospongins. *J. Org. Chem.* **1998**, 63, 9624–9625. (d) Lebel, H.; Jacobsen, E. N. Chromium Catalyzed Kinetic Resolution of 2,2-Disubstituted Epoxides. *Tetrahedron Lett.* **1999**, 40, 7303–7306. (e) Ueberbacher, B. J.; Osprian, I.; Mayer, S. F.; Faber, K. A Chemoenzymatic, Enantioconvergent, Asymmetric Total Synthesis of (R)-Fridamycin. *Eur. J. Org. Chem.* **2005**, 2005, 1266–1270. (f) Elenkov, M. M.; Hoeffken, H. W.; Tang, L.; Hauer, B.; Janssen, D. B. Enzyme-Catalyzed Nucleophilic Ring Opening of Epoxides for the Preparation of Enantiopure Tertiary Alcohols. *Adv. Synth. Catal.* **2007**, 349, 2279–2285. (g) Elenkov, M. M.; Tang, L.; Meetsma, A.; Hauer, B.; Janssen, D. B. Formation of Enantiopure 5-Substituted Oxazolidinones through Enzyme-Catalyzed Kinetic Resolution of Epoxides. *Org. Lett.* **2008**, 10, 2417–2420. (h) Molinaro, C.; Guilbault, A.-A.; Kosjek, B. Resolution of 2,2-Disubstituted Epoxides via Biocatalytic Azidolysis. *Org. Lett.* **2010**, 12, 3772–3775.

(17) For select enantioselective syntheses of 2,2-disubstituted epoxides, see (a) Sone, T.; Yamaguchi, A.; Matsunaga, S.; Shibasaki, M. Catalytic Asymmetric Synthesis of 2,2-Disubstituted Terminal Epoxides via Dimethyloxosulfonium Methylide Addition to Ketones. *J. Am. Chem. Soc.* **2008**, 130, 10078–10079. (b) Sone, T.; Lu, G.; Matsunaga, S.; Shibasaki, M. Catalytic Asymmetric Synthesis of 2,2-Disubstituted Oxetanes from Ketones by Using a One-Pot Sequential Addition of Sulfur Ylide. *Angew. Chem., Int. Ed.* **2009**, 48, 1677–1680. (c) Sone, T.; Yamaguchi, A.; Matsunaga, S.; Shibasaki, M. Enantioselective Synthesis of 2,2-Disubstituted Terminal Epoxides via Catalytic Asymmetric Corey-Chaykovsky Epoxidation of Ketones. *Molecules* **2012**, 17, 1617–1634.

(18) Malapit, C. A.; Luvaga, I. K.; Caldwell, D. R.; Schipper, N. K.; Howell, A. R. Rh-Catalyzed Conjugate Addition of Aryl and Alkenyl Boronic Acids to α -Methylene- β -lactones: Stereoselective Synthesis of trans-3,4-Disubstituted β -Lactones. *Org. Lett.* **2017**, 19, 4460–4463.

(19) For the first carbonylation of isobutylene oxide, see (a) Drent, E.; Kragt, W. J. (Shell Internationale Research Maatschappij B, V., Neth.) Eur. Pat. Appl. EP 577206 A2, 1993. (b) *Chem. Abstr.* **1994**, 120, 191517c.

(20) For two early reports of isobutylene oxide carbonylation by our group, see (a) Getzler, Y. D. Y. L.; Mahadevan, V.; Lobkovsky, E. B.; Coates, G. W. Synthesis of β -Lactones: A Highly Active and Selective Catalyst for Epoxide Carbonylation. *J. Am. Chem. Soc.* **2002**, 124, 1174–1175. (b) Mahadevan, V.; Getzler, Y. D. Y. L.; Coates, G. W. [Lewis Acid]⁺[Co(CO)₄][−] Complexes: A Versatile Class of Catalysts for Carbonylative Ring Expansion of Epoxides and Aziridines. *Angew. Chem., Int. Ed.* **2002**, 41, 2781–2784.

(21) Church, T. L.; Getzler, Y. D. Y. L.; Byrne, C. M.; Coates, G. W. Carbonylation of Heterocycles by Homogeneous Catalysts. *Chem. Commun.* **2007**, 657–674.

(22) Lee, J. T.; Thomas, P. J.; Alper, H. Synthesis of β -Lactones by the Regioselective, Cobalt and Lewis Acid Catalyzed Carbonylation of Simple and Functionalized Epoxides. *J. Org. Chem.* **2001**, 66, 5424–5426.

(23) Mulzer, M.; Coates, G. W. Carbonylation of *cis*-Disubstituted Epoxides to *trans*- β -Lactones: Catalysts Displaying Steric and Contrasting Regioselectivity. *J. Org. Chem.* **2014**, 79, 11851–11862.

(24) (a) Church, T. L.; Getzler, Y. D. Y. L.; Coates, G. W. The Mechanism of Epoxide Carbonylation by [Lewis Acid]⁺[Co(CO)₄][−] Catalysts. *J. Am. Chem. Soc.* **2006**, 128, 10125–10133. (b) Rowley, J. M.; Lobkovsky, E. B.; Coates, G. W. Catalytic Double Carbonylation of Epoxides to Succinic Anhydrides: Catalyst Discovery, Reaction Scope, and Mechanism. *J. Am. Chem. Soc.* **2007**, 129, 4948–4960.

(25) (a) Beesley, R. M.; Ingold, C. K.; Thorpe, J. F. The Formation and Stability of *spiro*-Compounds. Part I. *spiro*-Compounds from Cyclohexane. *J. Chem. Soc., Trans.* **1915**, 107, 1080–1106. (b) Ingold, C. K. The Conditions Underlying the Formation of Unsaturated and of Cyclic Compounds from Halogenated Open-Chain Derivatives.

Part I. Products Derived from α -Halogenated Glutaric Acids. *J. Chem. Soc., Trans.* **1921**, 119, 305–329.

(26) (a) Chen, Q.; Mulzer, M.; Shi, P.; Beuning, P. J.; Coates, G. W.; O'Doherty, G. A. *De Novo* Asymmetric Synthesis of Fridamycin. *Org. Lett.* **2011**, 13, 6592–6595. (b) Chen, Q.; Zhong, Y.; O'Doherty, G. A. Convergent *De Novo* Synthesis of Vineomycinone B₂ Methyl Ester. *Chem. Commun.* **2013**, 49, 6806–6808.

(27) (a) Gresham, T. L.; Jansen, J. E.; Shaver, F. W.; Beears, W. L. β -Propiolactone. XIV. β -Isovalerolactone. *J. Am. Chem. Soc.* **1954**, 76, 486–488. (b) Liang, H. T.; Bartlett, P. D. Kinetics and Mechanism of the Reactions of β -Isovalerolactone in Water. *J. Am. Chem. Soc.* **1958**, 80, 3585–3590. (c) Noyce, D. S.; Banitt, E. H. The Stereochemistry of the Decarboxylation of β -Lactones to Form Olefins. *J. Org. Chem.* **1966**, 31, 4043–4047. (d) Mulzer, J.; Zippel, M.; Brüntrup, G. Thermal Decarboxylation of β -Lactones: Steric Hindrance of Mesomerism as Indication of a Zwitterionic Intermediate. *Angew. Chem., Int. Ed. Engl.* **1980**, 19, 465–466.