

Chromatography

Phases, Rf values, polarity and gas chromatography

Name _____ Class _____ Date _____

EQUATIONS

$R_f = \text{distance moved by spot} \div \text{distance moved by solvent front}$ (both from the baseline; no unit)

$\text{Area \%} = \text{peak area} \div \text{total area} \times 100 \%$ (equal detector response)

Mean, sample standard deviation ($n - 1$), $RSD = s \div \text{mean} \times 100 \%$

Show units on every quantity in every step.

A. Understanding

1. (a) What are the stationary phase and the mobile phase? (b) Explain why two substances in a mixture separate as the mobile phase moves.

B. Rf values

2. On a TLC plate the solvent front is 8.40 cm above the baseline. A mixture gives spots at 2.10 cm, 4.62 cm and 6.72 cm. Standards on the same plate: X, R_f 0.25; Y, R_f 0.55; Z, R_f 0.70. (a) Calculate the three R_f values. (b) Which standards are in the mixture? (c) What can you say about the third spot?

3. Explain each step: (a) the baseline is drawn in pencil (b) the solvent level is below the baseline (c) the container has a lid (d) the solvent front is marked as soon as the plate is removed.

C. Polarity

4. Hexane, propanone and ethanoic acid are run on a silica plate. (a) Predict the order of their R_f values, highest first, and explain. (b) How would a more polar solvent change the R_f values?

5. A sample gives a single spot on a TLC plate. Does this prove the sample is pure? Explain, and suggest how to check.

D. Gas chromatography

6. Standards give retention times: ethanol 2.4 min, propan-1-ol 4.1 min, butan-1-ol 6.8 min. A sample shows peaks at 2.4 min and 6.8 min. (a) What does it contain? (b) Suggest why ethanol comes out first.

7. Peak areas for a three-component mixture are 2310, 1540 and 3850. Assuming equal detector response, calculate the percentage of each component.

E. Replicates and quantities

8. An unknown dye is run five times (quintuplicate) beside standards. R_f values: 0.61, 0.63, 0.62, 0.60, 0.64. Standards: dye M, R_f 0.62; dye N, R_f 0.71. (a) Calculate the mean and the standard deviation. (b) Which dye is it? Justify using the standard deviation.

9. In a GC analysis, a 0.500 mg/mL standard gives a peak area of 2480. A sample is injected three times, giving areas of 1850, 1870 and 1860. (a) Find the mean area, the standard deviation and the RSD. (b) Assuming area is proportional to concentration, find the concentration in the sample. (c) What mass of the compound is in 20.0 mL of the sample?