

- [5] P. Lindner, *Int. J. Quantum Chem.* **4**, 217 (1970).
 [6] M. H. Alexander, *Phys. Rev.* **A1**, 1397 (1970).
 [7] R. E. Johnson, S. T. Epstein and W. J. Meath, *J. Chem. Phys.* **47**, 1271 (1967).
 [8] A. Dalgarno, *Advan. Phys.* **11**, 281 (1962).
 [9] W. Kolos, *Int. J. Quantum Chem.* **1**, 169 (1967).

Received September 27, 1971.

New Proof of the Minimum Principle for Excited States

RALPH H. YOUNG*

*Department of Chemistry, University of California,
Riverside, California 92502, USA*

Let H be a self-adjoint operator in Hilbert space. It is a well-known and frequently used fact that the n th root of a linear variational computation for H is an upper bound to the n th eigenvalue of H [1, 2, 3]. The invariably cited references [1] are incomplete in that they neglect questions of convergence for their sequence of successive approximations. Complete proofs have been based on maximum-of-minima arguments [2] or repeated application of the Rayleigh-Ritz principle [3]. Here is an alternative proof, which hopefully sheds additional light on the nature of this upper-bound principle.

Let f_1, f_2, \dots, f_n be n functions such that

$$(1) \quad \left. \begin{aligned} \langle f_i | f_j \rangle &= \delta_{ij} \\ \langle f_i | H | f_j \rangle &= \tilde{E}_i \delta_{ij} \end{aligned} \right\} \quad \text{for } i, j = 1, 2, \dots, n$$

(These may be the first n solutions to a linear variational problem.) We assume that H has a purely discrete spectrum from $-\infty$ past \tilde{E}_n (i.e. on $(-\infty, \tilde{E}_n + \epsilon]$ for some $\epsilon > 0$); let the ψ_k ($k = 1, 2, \dots$) be the orthonormal eigenfunctions of H corresponding to eigenvalues $E_k \leq \tilde{E}_n$. We wish to prove that there are at least n such ψ_k 's; or, equivalently, that the manifold \mathcal{M}_ψ spanned by these eigenfunctions has dimension $\geq n$.

The manifold spanned by the f_k 's, \mathcal{M}_f , resembles \mathcal{M}_ψ in that

$$(2) \quad \langle g | H | g \rangle \leq \tilde{E}_n \|g\|^2 \quad \text{if } g \in \mathcal{M}_f$$

It seems unreasonable that \mathcal{M}_ψ should have smaller dimension than \mathcal{M}_f . To confirm

* Present address: Eastman Kodak Research Laboratories, Rochester, New York 14650, U.S.A.

this suspicion, we show that the n functions Pf_k ,

$$(3) \quad Pf_k = \sum_{E_j \leq \tilde{E}_n} \langle \psi_j | f_k \rangle \psi_j$$

which clearly are in \mathcal{M}_ψ , are linearly independent. (P is the projector onto \mathcal{M}_ψ .) Were they not, there would be a choice of coefficients a_k , not all zero, such that

$$(4) \quad \sum_{k=1}^n a_k (Pf_k) = 0$$

$$(5) \quad \sum_{k=1}^n a_k f_k = (1 - P) \sum_{k=1}^n a_k f_k$$

in which case

$$(6) \quad \begin{aligned} \left\langle \sum_{k=1}^n a_k f_k \left| H \right| \sum_{k=1}^n a_k f_k \right\rangle &= \left\langle (1 - P) \sum_{k=1}^n a_k f_k \left| H \right| (1 - P) \sum_{k=1}^n a_k f_k \right\rangle \\ &> \tilde{E}_n \left\| (1 - P) \sum_{k=1}^n a_k f_k \right\|^2 \\ &= \tilde{E}_n \left\| \sum_{k=1}^n a_k f_k \right\|^2 \end{aligned}$$

Equation (6) contradicts (2), since $\sum_{k=1}^n a_k f_k \in \mathcal{M}_\psi$. Hence \mathcal{M}_ψ has dimension $\geq n$, H has at least n eigenvalues $\leq \tilde{E}_n$ (counting m -fold degenerate eigenvalues m times), and

$$(7) \quad E_n \leq \tilde{E}_n$$

Acknowledgement

The author wishes to thank Dr. Walter J. Deal and Dr. William H. Miller for helpful conversations on this subject, and the referee for an additional reference.

Bibliography

- [1] E. A. Hylleraas and B. Undheim, Z. Physik. **65**, 759 (1930); J. K. L. MacDonald, Phys. Rev. **43**, 830 (1933).
- [2] S. H. Gould, *Variational Methods for Eigenvalue Problems* (University of Toronto Press, Toronto, 1957); J. F. Perkins, J. Chem. Phys. **45**, 2156 (1966).
- [3] P.-O. Löwdin, Phys. Rev. **139**, A357 (1965).

Received March 11, 1971

Revised September 20, 1971