TUTORIAL 1

Complex Numbers and Induction

1.1 Course Logistics

NOTE

The current International Engineering syllabus and official announcements determine the actual class arrangements, submission deadlines, office hours, and assessment rules. In particular, WeBWorK is optional practice and contributes neither to the grade nor to a protective component in the international course.

1.1.1 Access to Tutorials

Tutorials may be held over Zoom, and recordings may be available. The way links and recordings are made available may differ between staff members; they will usually be found on Moodle. Technical problems may prevent a tutorial from being made available. The course team will try to avoid such problems, but students remain responsible for catching up on any material missed because of a technical failure. Follow the current course announcements for the arrangements that apply to your group.

1.1.2 Homework and Assessment

- For a homework sheet with two parts, the first contains two questions marked for submission, to be solved independently and submitted for grading. The second contains additional questions for independent practice, which are not submitted. Full solutions to some of these questions are provided so that you can assess your own work. Check the current assignment instructions for the questions you must actually submit.
- The homework is based on material from the lectures and/or tutorials.
- Submitting work gives you professional feedback. The additional questions help you consolidate the material and practise solving relevant problems. You are strongly encouraged to attempt as many questions as possible.
- One possible publication schedule is Tuesday morning, with submissions accepted until Saturday evening of the following week (a submission window labelled as 12 days). The current course announcements determine the deadlines for the international program.
- Where WeBWorK sheets are provided, they may be released approximately every week or at a longer interval and may be open for two weeks. International Engineering students may use available WeBWorK questions as optional practice. The material may be in Hebrew; WeBWorK scores do not contribute to the international course grade or to any protective component.

Created by Chen Gadi

 © 2026 Chen Gadi

- The final assessment rules are stated in the current syllabus. Homework instructions appear in the relevant course section. Direct questions about grades and examinations to the course lecturer.

1.1.3 Office Hours

- Office hours with the teaching assistant are intended for questions arising from the tutorial material and homework.
- The teaching assistant sets the time, location, and format of office hours. Office hours are not an additional tutorial and do not replace independent study.

Procedures may change; updates will be announced when necessary.

1.1.4 Study Recommendations

Mathematics is a language, and learning it requires practice. This course is an integral (and differential) part of the mathematical foundations that support your later academic studies. The following suggestions are intended to help you learn the material and manage the course.
Scope of this advice: The recommendations in this section do not make the author responsible for readers' study decisions, understanding, or application of the material. Each reader remains responsible for their own learning and use of the information.

- Attempt all questions marked for submission and all practice questions; they have been selected carefully.
- If you fall slightly behind, use the recordings and tutorials to catch up, and check that you can solve the homework questions on the same topic.
- If you fall substantially behind, prioritize the theoretical foundations and basic practice questions, and try to rejoin the class's pace as soon as possible.
- Consider making a ``three magic sheets'' summary: one sheet for definitions, one for propositions and theorems, and one for useful techniques.
- A competitive atmosphere does not help the group or the individual. Help one another. Group learning and support for classmates can strengthen both learning and resilience.

1.1.5 Clarifications, Personal Recommendations, and Expectations

- Send messages and questions through the website system at chengadi.science.

1. Logistics requests should include your full name, student ID, and course name. Requests lacking these details will not be processed. If you have received no response within three days, I may have missed your request; send a reminder through the system and include or quote the original request.
2. Questions about the material: in previous years and semesters, I received a very large number of questions (at least 20 requests per day). Questions may be sent through the website system, subject to the following guidelines:
   (a) First make an independent effort to understand the issue, using notes, books, the internet, and tools such as ChatGPT. Check mathematical claims independently.

Created by Chen Gadi

 © 2026 Chen Gadi

(b) Keep your message short and specific. Include the course name and a question that can be answered in one or two lines. Attach an image of the relevant material where possible, rather than referring only to a particular theorem.

(c) Do not send requests such as ``Is my method correct?'' or ``Check this part.'' I will not grade homework solutions through messages. You can discuss an approach during office hours or obtain feedback through the homework grader.

(d) You may ask for clarification of the wording of a homework question.

(e) I will try to answer within a few days. Messages that are too long, or that require more than a few minutes to read and answer, may not receive a response. Thank you for your understanding.

- I may upload videos of myself solving problems on YouTube. These are an additional resource for students, rather than part of my formal responsibilities; please do not send requests about their production or availability.
- During tutorials, I will not summarize all the theorems and conclusions covered in lectures. I recommend preparing your own summary. Tutorials are intended to practise the lecture material and work on somewhat more demanding questions that encourage thought and independent learning.
- I will also make sure to include basic exercises.
- Time will be set aside for questions during class. I will not repeatedly answer questions that have already been addressed.
- **The lectures determine the required material.**
  It is not possible to cover every topic in the tutorials or practise every type of question on each topic.
- I strongly recommend exploring the material through the resources available to you: books, online sources, earlier summaries (for example, materials from the students' association), office hours, and similar resources. **Each student is responsible for their own learning.**
- Tutorial pace and presentation: I will try to work at a reasonable pace, write clearly, and use headings. Please let me know if the presentation needs adjustment.
- WeBWorK is optional and recommended as additional practice with feedback. **For the International Engineering Program, WeBWorK does not count toward the final grade and does not provide protective credit.**
- I will try to provide examples, proofs, and counterexamples for each topic.
- I will conduct an anonymous mid-semester teaching survey, and I welcome feedback.
- I will take questions at the designated times rather than in the middle of solving an exercise.
- I will try to upload typed tutorial sheets before the tutorials, but they may contain errors or imprecise statements.
  The material presented in the lectures, both in writing and orally, remains authoritative.
- Come prepared: review the definitions and material covered in the lectures before each tutorial.

Created by Chen Gadi

 © 2026 Chen Gadi

MOTIVATION AND GUIDING IDEA

Cartesian form is well suited to addition and comparison of components; polar form is well suited to multiplication, powers, and roots. Before calculating, choose a representation that fits the requested operation. For induction, first choose the precise statement and the first value for which it is supposed to hold.

NOTATION AND CONVENTIONS

For $z = x+iy \in \mathbb{C}$, the symbols $\operatorname{Re} z = x$ and $\operatorname{Im} z = y$ denote real numbers; in particular, the imaginary part is y, not iy. Both z^* and $\overline{z}$ denote the conjugate $x - iy$. The modulus $|z|$ is a nonnegative real number. When $z \neq 0$, $\arg z$ refers to an argument modulo integer multiples of 2π, whereas $\operatorname{Arg} z \in [0, 2\pi)$ is the fixed choice in these notes. The notation $a \mid b$ means that $b = am$ for some $m \in \mathbb{Z}$. The expression $P(n) \Rightarrow P(n+1)$ is a logical implication; it does not by itself establish the base case.

DEFINITION 1.1

Existence, uniqueness, and conditions of use. Every $z \in \mathbb{C}$ has a unique representation $z = x + iy$ with $x, y \in \mathbb{R}$. Thus two complex numbers are equal if and only if their real parts are equal and their imaginary parts are equal. For $z \neq 0$, its inverse exists and is unique, since $zz^* = |z|^2 > 0$; check this condition before dividing by z. If $r > 0$ and $\theta \in \mathbb{R}$, de Moivre's formula gives

$$(re^{i\theta})^n = r^n e^{in\theta} \qquad (n \in \mathbb{Z}).$$

Negative powers require $r > 0$. For nonnegative integer powers, the formula also holds when $r = 0$, except for 0^0, which is not defined here. When finding nth roots of a nonzero number, for $n \geq 1$, list all n roots rather than selecting a single angle.

Created By Chen Gadi

How to use these notes

These notes are intended for students in all engineering disciplines. They combine definitions, guiding ideas, examples, and detailed solutions.

MOTIVATION AND GUIDING IDEA

Objectives. We will convert between Cartesian and polar forms, calculate powers and roots, and interpret operations in the complex plane. We will then formulate an induction proof, stating the domain on which its claim holds.

MOTIVATION AND GUIDING IDEA

A working habit. In every solution, state what is given, what must be established, which definition or result you use, and why every operation is permitted. Correct calculations and clear explanations complement one another.

 © 2026 Chen Gadi

1.2 Complex numbers: Cartesian form

DEFINITION 1.2

The imaginary unit i satisfies $i^2 = -1$.
A complex number has the form $z = x + iy$, where $x, y \in \mathbb{R}$.

1. If $\mathrm{Re}(z) = x = 0$, then z is purely imaginary.
2. If $\mathrm{Im}(z) = y = 0$, then z is real.

EXERCISE 1.1

Solve $z^2 - 4z + 13 = 0$ over $\mathbb{C}$. Factor the polynomial into linear factors over $\mathbb{C}$, and determine whether such a factorization exists over $\mathbb{R}$.

WORKED SOLUTION

The discriminant is $\Delta = (-4)^2 - 4 \cdot 1 \cdot 13 = 16 - 52 = -36$. Therefore

$$z_{1,2} = \frac{4 \pm \sqrt{-36}}{2} = \frac{4 \pm 6i}{2} = 2 \pm 3i.$$

Consequently,

$$z^2 - 4z + 13 = (z - (2 + 3i))(z - (2 - 3i)).$$

This is a factorization over $\mathbb{C}$. Over $\mathbb{R}$, the discriminant is negative, so there are no real roots and no factorization into two real linear factors.

DEFINITION 1.3

- **Complex conjugate:** for $z = x + iy$, write $z^* := x - iy$.
- **Modulus:** $|z| = \sqrt{x^2 + y^2} \in \mathbb{R}$.
- **Multiplicative inverse:** for $z \neq 0$, $z^{-1} = \dfrac{z^*}{|z|^2}$.

EXERCISE 1.2

Prove the following statements. **(a)** $\mathrm{Re}\, z = \dfrac{z + z^*}{2}$ and $\mathrm{Im}\, z = \dfrac{z - z^*}{2i}$. **(b)** $|z| \geq \mathrm{Im}\, z$ and $|z| \geq \mathrm{Re}\, z$ for every $z \in \mathbb{C}$. **(c)** (At home) $(z^*)^* = z$. **(d)** (At home) If z is real, then $z^* = z$.

WORKED SOLUTION

Write $z = x + iy$. Then

$$\frac{z + z^*}{2} = x = \mathrm{Re}\, z, \qquad \frac{z - z^*}{2i} = y = \mathrm{Im}\, z,$$

and

$$|z| = \sqrt{x^2 + y^2} \geq |x| \geq x = \mathrm{Re}\, z, \qquad |z| \geq |y| \geq y = \mathrm{Im}\, z.$$

Created by Chen Gadi

 © 2026 Chen Gadi

EXERCISE 1.2 CONTINUED

(c) $(z^*)^* = (x - iy)^* = x + iy = z$.
(d) If z is real, then $y = 0$, and hence $z^* = x - 0i = x = z$.

1.2.1 Useful properties

1. $|z| = |z^*|$.
2. $z \cdot z^* = |z|^2$ (multiplication by the conjugate gives a real number).
3. $|zw| = |z|\,|w|$.
4. $(zw)^* = z^* w^*$.

EXERCISE 1.3

For $z \neq 0$, prove that $(z^{-1})^* = (z^*)^{-1}$.

WORKED SOLUTION

Since $z \neq 0$, we also have $z^* \neq 0$ and $|z|^2 > 0$. The inverse formula gives

$$(z^{-1})^* = \left(\frac{z^*}{|z|^2}\right)^* = \frac{z}{|z|^2}, \qquad (z^*)^{-1} = \frac{(z^*)^*}{|z^*|^2} = \frac{z}{|z|^2}.$$

Both sides equal the same expression, so $(z^{-1})^* = (z^*)^{-1}$.

1.3 Polar form

1.3.1 Converting between Cartesian and polar forms

DEFINITION 1.4

For every $z = x + iy \neq 0$, there are $r = |z| > 0$ and an angle θ such that

$$z = r(\cos\theta + i\sin\theta) = re^{i\theta}, \qquad x = r\cos\theta, \quad y = r\sin\theta.$$

Conversely, $r = \sqrt{x^2 + y^2}$. If $x \neq 0$, then $\tan\theta = y/x$, but this ratio alone does not determine the quadrant. **Remarks:**

- The angle is determined up to adding $2\pi k$, where $k \in \mathbb{Z}$. For $z = 0$, we have $r = 0$, and no argument is defined.
- All angles in exponents $e^{i\theta}$ are measured in radians. Degrees are used only for numerical illustration.

EXERCISE 1.4

Convert $z = 3 + 3i$ and $w = -4 + 4i$ to polar form. Then write $a = 6e^{i\pi/3}$ in Cartesian form. For each polar representation, choose an angle in $[0, 2\pi)$.

Created by Chen Gadi

 © 2026 Chen Gadi

EXERCISE 1.4 CONTINUED

WORKED SOLUTION

For z, we have $|z| = \sqrt{3^2 + 3^2} = 3\sqrt{2}$. Its components are equal and positive, so the angle is $\pi/4$:

$$z = 3\sqrt{2}e^{i\pi/4}.$$

For w, we have $|w| = \sqrt{(-4)^2 + 4^2} = 4\sqrt{2}$. The point lies in the second quadrant, and the absolute values of its components have ratio 1. Therefore

$$w = 4\sqrt{2}e^{3i\pi/4}.$$

Finally, use Euler's identity and the trigonometric values at $\pi/3$:

$$a = 6\left(\cos\frac{\pi}{3} + i\sin\frac{\pi}{3}\right) = 6\left(\frac{1}{2} + i\frac{\sqrt{3}}{2}\right) = 3 + 3\sqrt{3}i.$$

The angles are in radians. Both given complex numbers have positive modulus, so the undefined argument at zero causes no difficulty.

1.3.2 Arithmetic operations




Figure 1.1. Cartesian and polar forms in the same plane: the modulus is the distance from the origin, and the argument is the direction angle.

DEFINITION 1.5

Multiplication: $(r_1e^{i\theta_1})(r_2e^{i\theta_2}) = r_1r_2\,e^{i(\theta_1+\theta_2)}$.

Division (when the denominator is nonzero): $\dfrac{r_1e^{i\theta_1}}{r_2e^{i\theta_2}} = \dfrac{r_1}{r_2}\,e^{i(\theta_1-\theta_2)}$.

Conjugation: $(re^{i\theta})^* = r\,e^{-i\theta}$.

Periodicity: $re^{i(\theta+2\pi k)} = re^{i\theta}$, $re^{i(\theta+\pi)} = -re^{i\theta}$.

Created by Chen Gadi

 © 2026 Chen Gadi

EXERCISE 1.5

Let $z = 1 + i$ and $w = -4 + 4i$. Calculate zw and w/z in two ways: using polar form and using Cartesian arithmetic. Check that the resulting moduli agree with the multiplication and division formulas.

WORKED SOLUTION

Write $z = \sqrt{2}e^{i\pi/4}$ and $w = 4\sqrt{2}e^{3i\pi/4}$. Then

$$zw = 8e^{i\pi} = -8, \qquad \frac{w}{z} = 4e^{i\pi/2} = 4i.$$

Cartesian arithmetic gives

$$(1+i)(-4+4i) = -4 + 4i - 4i + 4i^2 = -8,$$

$$\frac{-4+4i}{1+i} = \frac{(-4+4i)(1-i)}{(1+i)(1-i)} = \frac{-4+8i+4}{2} = 4i.$$

Checking the moduli, $|z||w| = \sqrt{2} \cdot 4\sqrt{2} = 8 = |-8|$, and $|w|/|z| = 4 = |4i|$. The two calculations agree.

1.3.3 Powers

EXERCISE 1.6

Calculate $(1+i)^{12}$ using polar form. Verify your answer using $(1+i)^4$.

WORKED SOLUTION

Since $1 + i = \sqrt{2}e^{i\pi/4}$,

$$(1+i)^{12} = (\sqrt{2})^{12} e^{12i\pi/4} = 2^6 e^{3i\pi} = -64.$$

For verification, $(1+i)^2 = 2i$, so $(1+i)^4 = (2i)^2 = -4$. Thus $(1+i)^{12} = ((1+i)^4)^3 = (-4)^3 = -64$.

1.3.4 Complex roots

DEFINITION 1.6

Let $0 \neq z \in \mathbb{C}$, and let $n \geq 1$ be an integer. The equation $w^n = z = re^{i\theta}$ has n distinct roots:

$$w_j = \sqrt[n]{r}\, e^{i\frac{\theta+2\pi j}{n}}, \quad j = 0, 1, \ldots, n-1.$$

Geometric meaning: the roots lie on the circle of radius $\sqrt[n]{r}$ at equal angular intervals of $2\pi/n$. For $n \geq 3$, they are the vertices of a regular polygon; for $n = 2$, they are two opposite points, and for $n = 1$, there is one point.

Created by Chen Gadi

 © 2026 Chen Gadi

EXERCISE 1.7

Find all solutions of $z^4 = -i$.

WORKED SOLUTION

Write $-i = e^{-i\pi/2}$. Therefore

$$z_j = e^{i\frac{-\pi/2+2\pi j}{4}} = e^{i\frac{\pi(4j-1)}{8}}, \quad j = 0, 1, 2, 3,$$

so the angles are $-\pi/8,\ 3\pi/8,\ 7\pi/8,\ 11\pi/8$, all at radius 1.

EXERCISE 1.8

Let $n \geq 2$. Find all nth roots of unity and prove that their sum is zero. What happens when $n = 1$?

WORKED SOLUTION

The roots are $w_j = e^{i\frac{2\pi j}{n}}$, $j = 0, \ldots, n-1$. Put $w = e^{i\frac{2\pi}{n}}$. Then $w_j = w^j$, and therefore

$$\sum_{j=0}^{n-1} w^j = \frac{w^n - 1}{w - 1} = 0,$$

because $w^n = 1$ and $w \neq 1$ when $n \geq 2$. For $n = 1$, the only root is 1, so the sum is 1, rather than zero.

EXERCISE 1.9

(At home) Consider $a_n x^n + a_{n-1}x^{n-1} + \cdots + a_1 x + a_0 = 0$, with real coefficients. Prove that if z is a solution, then z^* is also a solution.

WORKED SOLUTION

Use the identities $(z_1 + \cdots + z_m)^* = z_1^* + \cdots + z_m^*$ and $(z_1 \cdots z_m)^* = z_1^* \cdots z_m^*$, together with $a^* = a$ for real a. If $a_0 + a_1 z + \cdots + a_n z^n = 0$, conjugating gives $a_0 + a_1 z^* + \cdots + a_n (z^*)^n = 0$. Thus z^* is a solution.

Created by Chen Gadi

 © 2026 Chen Gadi

1.4 Further properties of complex numbers




Figure 1.2. The roots of unity are equally spaced around the unit circle; their sum is zero whenever their order is greater than one.

1.4.1 Multiplication by i

Multiplication by i rotates a complex number through $\frac{\pi}{2}$ counterclockwise.

EXERCISE 1.10

Prove that multiplication by i gives a rotation through $\frac{\pi}{2}$.

WORKED SOLUTION

Cartesian representation:
Let $z = a + bi$. Multiplication by i gives

$$z \cdot i = (a + bi)i = -b + ai.$$

The vector (a, b) becomes $(-b, a)$. This is exactly a counterclockwise rotation through $\pi/2$ about the origin: the new x coordinate is $-b$, and the new y coordinate is a.
Polar representation:
For $z \neq 0$, write $z = re^{i\theta}$, where $r = |z|$ and θ is an angle from the positive x axis to z. Then

$$z \cdot i = re^{i\theta} \cdot i = re^{i\theta} \cdot e^{i\frac{\pi}{2}} = re^{i(\theta+\frac{\pi}{2})}.$$

Multiplication by i adds $\frac{\pi}{2}$ to θ, so z rotates through $\frac{\pi}{2}$ about the origin, as required. The origin is fixed, since $0 \cdot i = 0$.

Created by Chen Gadi

 © 2026 Chen Gadi

1.4.2 Division in polar form

DEFINITION 1.7

If $z_1 = r_1 e^{i\theta_1}$ and $z_2 = r_2 e^{i\theta_2}$, with $r_2 \neq 0$, then

$$\frac{z_1}{z_2} = \frac{r_1}{r_2} e^{i(\theta_1 - \theta_2)}.$$

EXERCISE 1.11

Calculate $\frac{z_1}{z_2}$ for $z_1 = 4e^{i\frac{\pi}{3}}$ and $z_2 = 2e^{i\frac{\pi}{6}}$.

WORKED SOLUTION

We obtain

$$\frac{z_1}{z_2} = \frac{4}{2} e^{i\left(\frac{\pi}{3} - \frac{\pi}{6}\right)} = 2e^{i\frac{\pi}{6}}.$$

EXERCISE 1.12

Let $z_1, z_2 \neq 0$. Prove that an argument of z_1/z_2 is the difference of arguments of z_1 and z_2, up to an integer multiple of 2π.

WORKED SOLUTION

Write $z_1 = r_1 e^{i\theta_1}$ and $z_2 = r_2 e^{i\theta_2}$, where $r_1, r_2 > 0$. Division gives

$$\frac{z_1}{z_2} = \frac{r_1}{r_2} e^{i(\theta_1 - \theta_2)}.$$

Thus $\theta_1 - \theta_2$ is an argument of the quotient. If an argument in a prescribed interval is required, add a suitable integer multiple of 2π.

Created by Chen Gadi

1.4.3 Choosing an argument in a fixed interval

In these notes, $\mathrm{Arg}(z)$ denotes the unique argument in $[0, 2\pi)$, for $z \neq 0$.

RECALL AND TOOLS

The equalities $\mathrm{Arg}(z_1 \cdot z_2) = \mathrm{Arg}(z_1) + \mathrm{Arg}(z_2)$ and $\mathrm{Arg}\left(\frac{z_1}{z_2}\right) = \mathrm{Arg}(z_1) - \mathrm{Arg}(z_2)$ do not always hold, because the expression on the right may lie outside $[0, 2\pi)$.

EXERCISE 1.13

Calculate $\mathrm{Arg}(z_1 \cdot z_2)$ for $z_1 = e^{i\frac{3\pi}{4}}$ and $z_2 = e^{i\frac{5\pi}{4}}$.

WORKED SOLUTION

First calculate the product itself:

$$z_1 z_2 = e^{i3\pi/4} e^{i5\pi/4} = e^{i2\pi} = 1.$$

 © 2026 Chen Gadi

EXERCISE 1.13 CONTINUED

The angle 2π is an argument of 1, but it does not belong to $[0, 2\pi)$. Hence

$$\mathrm{Arg}(z_1 z_2) = \mathrm{Arg}(1) = 0, \qquad \mathrm{Arg}(z_1) + \mathrm{Arg}(z_2) = 2\pi.$$

These two values are unequal, but congruent modulo 2π.

1.4.4 Calculating arguments and identifying quadrants

DEFINITION 1.8

For a complex number $z = a + bi \neq 0$ with $a \neq 0$, every argument θ satisfies

$$\tan(\theta) = \frac{b}{a}.$$

To determine θ precisely, take the quadrant into account:

- If $a > 0, b > 0$, then $\theta = \arctan\left(\frac{b}{a}\right)$.
- If $a < 0, b > 0$, add π to obtain an angle in the second quadrant: $\theta = \pi + \arctan\left(\frac{b}{a}\right)$.
- If $a < 0, b < 0$, add π for the third quadrant: $\theta = \pi + \arctan\left(\frac{b}{a}\right)$.
- If $a > 0, b < 0$, add 2π to obtain a positive angle in the fourth quadrant: $\theta = 2\pi + \arctan\left(\frac{b}{a}\right)$.

If $b = 0, a > 0$, then $\mathrm{Arg}(z) = 0$; if $b = 0, a < 0$, then $\mathrm{Arg}(z) = \pi$. If $a = 0$, the angle depends only on b:

- If $b > 0$, then $\theta = \frac{\pi}{2}$.
- If $b < 0$, then $\theta = \frac{3\pi}{2}$.

EXERCISE 1.14

Find the angle θ for $z = -1 + \sqrt{3}i$.

WORKED SOLUTION

Here $a = -1, b = \sqrt{3}$. First calculate

$$\tan(\theta) = \frac{\sqrt{3}}{-1} = -\sqrt{3}.$$

Since $a < 0$ and $b > 0$, the number lies in the second quadrant. Therefore

$$\theta = \pi - \frac{\pi}{3} = \frac{2\pi}{3}.$$

Created by Chen Gadi

 © 2026 Chen Gadi

EXERCISE 1.15

Find the angle θ for $z = -5 - 5i$.

WORKED SOLUTION

Here $a = -5, b = -5$. First calculate

$$\tan(\theta) = \frac{b}{a} = \frac{-5}{-5} = 1.$$

Since $a < 0, b < 0$, the number lies in the third quadrant. Therefore

$$\theta = \pi + \arctan(1) = \pi + \frac{\pi}{4} = \frac{5\pi}{4}.$$

1.5 The induction principle

DEFINITION 1.9

The principle of mathematical (weak) induction:
Let n_0 be a nonnegative integer, and let $P(n)$ be a statement defined for every integer $n \geq n_0$. Suppose that:

1. **Base case:** $P(n_0)$ is true.
2. **Inductive step:** for every integer $n \geq n_0$, the assumption $P(n)$ implies $P(n+1)$.

Then $P(n)$ is true for every integer $n \geq n_0$.

Created by Chen Gadi

EXERCISE 1.16

Prove by induction that, for every $n \geq 1$,

$$1 + 2 + \cdots + n = \frac{n(n+1)}{2}.$$

WORKED SOLUTION

The statement: $P(n) : \sum_{i=1}^{n} i = \frac{n(n+1)}{2}$.
Base case: for $n = 1$, the left side is 1, and the right side is $\frac{1 \cdot 2}{2} = 1$.
Induction hypothesis: suppose $\sum_{i=1}^{k} i = \frac{k(k+1)}{2}$ for some $k \geq 1$.
Inductive step:

$$\sum_{i=1}^{k+1} i = \left(\sum_{i=1}^{k} i\right) + (k+1) = \frac{k(k+1)}{2} + (k+1) = \frac{(k+1)(k+2)}{2}.$$

This is the formula for $k+1$.
Conclusion: the statement holds for every $n \geq 1$.

 © 2026 Chen Gadi

EXERCISE 1.17

Prove that $7 \mid (8^n - 1)$ for every $n \geq 1$.

WORKED SOLUTION

The statement: $P(n):\ 7 \mid (8^n - 1)$.
Base case: for $n = 1$, $8^1 - 1 = 7$, and indeed $7 \mid 7$.
Induction hypothesis: suppose $7 \mid (8^k - 1)$ for some $k \geq 1$. Thus there is $m \in \mathbb{Z}$ such that $8^k - 1 = 7m$.
Inductive step:

$$8^{k+1} - 1 = 8 \cdot 8^k - 1 = 8(8^k - 1) + 7.$$

Using $8^k - 1 = 7m$, we get

$$8^{k+1} - 1 = 8 \cdot 7m + 7 = 7(8m + 1),$$

so $7 \mid (8^{k+1} - 1)$.
Conclusion: $7 \mid (8^n - 1)$ for every $n \geq 1$.
A shorter observation: $8 \equiv 1 \pmod 7 \Rightarrow 8^n \equiv 1 \pmod 7$.

EXERCISE 1.18

Prove that, for every $n \geq 4$,

$$2^n \geq n^2.$$

WORKED SOLUTION

We use induction on n.
Base case: for $n = 4$, $2^4 = 16 = 4^2$.
Induction hypothesis: suppose $2^k \geq k^2$ for some $k \geq 4$.
Inductive step:

$$2^{k+1} = 2 \cdot 2^k \ \geq\ 2k^2.$$

It suffices to show $2k^2 \geq (k+1)^2 \iff k^2 - 2k - 1 \geq 0$. For $k \geq 4$, $k^2 - 2k - 1 > 0$, so $2^{k+1} \geq (k+1)^2$.
By induction, $2^n \geq n^2$ for every $n \geq 4$.

1.6 Supplementary exercises: geometry and signal calibration

EXERCISE 1.19

Find and describe geometrically all $z \in \mathbb{C}$ satisfying $|z - 1| = |z + 1|$. Which of them also satisfy $|z| = 2$?

WORKED SOLUTION

Write $z = x + iy$. Both sides are nonnegative, so squaring preserves the equality:

$$|z - 1|^2 = (x - 1)^2 + y^2, \qquad |z + 1|^2 = (x + 1)^2 + y^2.$$

Created by Chen Gadi

 © 2026 Chen Gadi

EXERCISE 1.19 CONTINUED

Therefore

$$(x-1)^2+y^2=(x+1)^2+y^2 \iff x^2-2x+1=x^2+2x+1 \iff x=0.$$

This is the imaginary axis: the set of points equidistant from 1 and -1. The additional condition gives $y^2=4$, so the intersection consists of $z=2i$ and $z=-2i$.

EXERCISE 1.20

In a simple model of a signal with two components, a calibration system applies $T(z)=(1+i)z$. How does this change the modulus and angle? Calculate $T(1-i)$, and find an input whose output is $2i$.

WORKED SOLUTION

Since $1+i=\sqrt{2}e^{i\pi/4}$, every $z=re^{i\theta}\neq 0$ satisfies $T(z)=\sqrt{2}\,r\,e^{i(\theta+\pi/4)}$. Thus the modulus is multiplied by $\sqrt{2}$, and the direction rotates counterclockwise through $\pi/4$. We also have $T(0)=0$. Direct calculation gives $T(1-i)=(1+i)(1-i)=1-i^2=2$. To obtain $2i$, solve $(1+i)z=2i$:

$$z=\frac{2i}{1+i}=\frac{2i(1-i)}{(1+i)(1-i)}=\frac{2i+2}{2}=1+i.$$

Check: $(1+i)^2=1+2i+i^2=2i$.

Additional practice

EXERCISE 1.21 FOUNDATION

Let $z=-3+4i$. Calculate $|z|$, z^*, and z^{-1}. Solve $zw=5+10i$, and verify the solution by direct multiplication.

WORKED SOLUTION

We have $|z|=\sqrt{9+16}=5$, so $z^*=-3-4i$ and $z^{-1}=(-3-4i)/25$. Since $z\neq 0$, division by it is permitted and gives a unique solution:

$$w=\frac{5+10i}{-3+4i}=\frac{(5+10i)(-3-4i)}{25}=\frac{25-50i}{25}=1-2i.$$

For verification, $(-3+4i)(1-2i)=-3+6i+4i-8i^2=5+10i$. Uniqueness follows because multiplication by a nonzero number is invertible.

EXERCISE 1.22 CHALLENGE

Let $z\in\mathbb{C}\setminus\{-1\}$. Prove that $(z-1)/(z+1)$ is purely imaginary if and only if $|z|=1$. Then solve $(z-1)/(z+1)=2i$, and check that the solution satisfies the geometric condition.

Created by Chen Gadi

 © 2026 Chen Gadi

EXERCISE 1.22 CHALLENGE CONTINUED

WORKED SOLUTION

Multiply the numerator and denominator by the conjugate of the denominator. Since $z \neq -1$, the denominator $|z+1|^2$ is positive, and

$$\frac{z-1}{z+1} = \frac{(z-1)(z^*+1)}{|z+1|^2} = \frac{|z|^2 - 1 + z - z^*}{|z+1|^2}.$$

The expression $z - z^*$ is purely imaginary. Hence

$$\operatorname{Re}\left(\frac{z-1}{z+1}\right) = \frac{|z|^2 - 1}{|z+1|^2}.$$

The real part is zero if and only if $|z|^2 = 1$, equivalent to $|z| = 1$ because the modulus is nonnegative. This proves both directions and includes $z = 1$, whose quotient is 0. In the given equation, multiplying by $z + 1 \neq 0$ gives $z - 1 = 2iz + 2i$. Thus

$$(1-2i)z = 1+2i, \qquad z = \frac{1+2i}{1-2i} = \frac{(1+2i)^2}{5} = \frac{-3+4i}{5}.$$

This number differs from -1 and has modulus $\sqrt{9+16}/5 = 1$. Moreover,

$$\frac{z-1}{z+1} = \frac{-8+4i}{2+4i} = 2i.$$

It is therefore the unique solution and lies on the unit circle.

Created by Chen Gadi

 © 2026 Chen Gadi