

5B Prove Corollary 5.3 (to Pasch's theorem)

Theorem 5.2 (Pasch's theorem): Suppose $\triangle ABC$ is a triangle and ℓ is a line that does not contain any of the points A, B, C . If ℓ intersects one of the sides of $\triangle ABC$, then it also intersects another side.

Corollary 5.3: If $\triangle ABC$ is a triangle and ℓ is a line that does not contain any of the points A, B, C , then either ℓ intersects exactly two sides of $\triangle ABC$ or it intersects none of them.

ℓ is a line. It does not contain any of the points A, B, C , so it is not collinear to any of the segments $\overline{AB}, \overline{BC}, \overline{AC}$

Case: ℓ does not intersect any of the line segments.

Then ℓ does not intersect any of the line segments.

Case: ℓ intersects one of the line segments. Then by theorem 5.2, it intersects another side.

Then ℓ intersects exactly two sides of $\triangle ABC$

5C Suppose $\triangle ABC$ is a triangle and ℓ is a line (which might or might not contain one or more vertices). Is it possible for ℓ to intersect exactly one side of $\triangle ABC$? Exactly two? All three? In each case, either give an example or prove that it is impossible.

Case 1: Intersects exactly one side of $\triangle ABC$

If ℓ intersects at a vertex, for example A , then since $A \in \overline{AB}, \overline{AC}$, then it intersects more than one side.

Then assume that ℓ does not intersect at a vertex, and intersects at one side.

But by theorem 5.2, we know that if it intersects at one side, it must intersect another side.

Then it is not possible for ℓ to intersect the triangle on exactly one side.

Case 2: Intersects exactly two sides of $\triangle ABC$

Yes, it is possible: example picture provided (Picture is of triangle $\triangle ABC$, and line ℓ , where ℓ intersects through sides $\overline{AB}, \overline{AC}$, and touches none of the vertices.)

Case 3: Intersects exactly three sides of $\triangle ABC$

Yes, it is possible: example picture provided (Picture is of triangle $\triangle ABC$, and line ℓ , where ℓ is collinear to \overline{AB} , and touches both vertices A, B)

5D Prove Theorem 5.8 (The converse to the isosceles triangle theorem). [Hint: One way to proceed is to construct an indirect proof, like Euclid's proof of proposition 1.6. Another is to mimic Pappus's proof of the isosceles triangle theorem.]

Theorem 5.7: Isosceles Triangle theorem: If two sides of a triangle are congruent to each other, then the angles opposite those sides are congruent.

Theorem 5.8: If two angles of a triangle are congruent to each other, then the sides opposite those angles are congruent.

Let $\triangle ABC$ be a triangle with $\angle B \cong \angle C$

Then draw the mirrored image $\triangle A'B'C'$ under the correspondence $A \leftrightarrow B, B \leftrightarrow C$, and $C \leftrightarrow A$

The triangles $\triangle ABC, \triangle A'B'C'$ satisfy the hypothesis set by ASA congruence, since $\overline{BC} \cong \overline{B'C'}$

and $\angle B \equiv \angle B', \angle C \cong \angle C'$

Thus the corresponding sides $\overline{AB}, \overline{AC}$ are congruent

The figure shown is the same as Fig 5.7 from the book, mimicing Pappus's proof.

5E Prove Theorem 5.10 (The triangle copying theorem)

Suppose $\triangle ABC$ is a triangle and \overline{DE} is a segment congruent to \overline{AB} . On eah side of \overleftrightarrow{DE} there is a point F such that $\triangle DEF \cong \triangle ABC$

5F Prove Theorem 5.18 (The triangle inequality)

If A, B, C are noncollinear points, then $AC < AB + BC$

Construct a triangle with vertices A, B, C , call it $\triangle ABC$

For the line \overleftrightarrow{AB} , take the point D , such that $BD = BC$