

Óptica II

Chengyu Jin

5 de mayo de 2026

Índice general

1. Quantum Ideal Gas	2
1.1. Non-interacting Ising model	2
1.2. Ideal Quantum Gas	3

Capítulo 1

Quantum Ideal Gas

1.1. Non-interacting Ising model

$$H = -J \sum_i s_i \text{ where } s_i = \pm 1 \text{ (spin)}$$

We must to find:

$$Z$$

$$A$$

$$U = \langle H \rangle$$

$$\langle M \rangle = \langle \sum s_i \rangle / N$$

For interacting case

At T_c my particles are disordered. Below T_c , symmetry is broken (spontaneous magnetization).

$$Z = (z_i)^N$$

$$z_i = \sum_{s_i=\pm 1} e^{-\beta H_i} = e^{\beta J} + e^{-\beta J} = 2 \cosh(\beta J)$$

$$Z = (2 \cosh(\beta J))^N$$

$$A = -kT \ln Z = -NkT \ln(2 \cosh(\beta J))$$

$$U = \left(-\frac{\partial \ln Z}{\partial \beta} \right)_{N,V} = -\frac{\partial}{\partial \beta} (N \ln(2 \cosh(\beta J))) = -N \frac{2 \sinh(\beta J)}{2 \cosh(\beta J)} \cdot J = -NJ \tanh(\beta J)$$

In other way:

$$U = \langle H \rangle = -J \sum_i \langle s_i \rangle = -NJ \tanh(\beta J)$$
$$\langle s_i \rangle = 1 \cdot P(s_i = 1) - 1 \cdot P(s_i = -1) = \frac{2 \sinh(\beta J)}{2 \cosh(\beta J)} = \tanh(\beta J)$$

$$\text{where } P(s_i) = \frac{e^{-\beta J s_i}}{2 \cosh(\beta J)}$$
$$P(1) = \frac{e^{\beta J}}{2 \cosh(\beta J)}$$
$$P(-1) = \frac{e^{-\beta J}}{2 \cosh(\beta J)}$$

1.2. Ideal Quantum Gas

System made of N identical particles:

$$\Psi(r_1, r_2, \dots, r_N, t) \quad (1.1)$$

Its evolution is given by:

$$i\hbar \frac{\partial \Psi}{\partial t} = H\Psi \quad ; \quad H = \sum_{i=1}^N \frac{p_i^2}{2m} + V(r_1, r_2, \dots, r_N) \quad (1.2)$$

P_{ij} is the **permutation operator** which exchanges the labels of particles i and j :

$$P_{ij}\Psi(\dots, r_i, \dots, r_j, \dots, t) = \Psi(\dots, r_j, \dots, r_i, \dots, t) \quad (1.3)$$

$$[H, P_{ij}] = 0 \quad \forall i, j \quad (i \neq j)$$

If $H\phi_n = E_n\phi_n$:

$$\phi(r_1, \dots, r_N, t) = \sum a_n \phi_n(r_1, \dots, r_N) \quad ; \quad a_n = \langle \phi_n | \Psi \rangle \quad (1.4)$$

Now we apply P_{ij} to Ψ and then H :

$$\begin{aligned} HP_{ij}\Psi(\dots, r_i, \dots) &\rightarrow \\ H\Psi(\dots, r_j, \dots, r_i, \dots, t) &= \sum a_n E_n \phi_n(\dots, r_j, \dots, r_i, \dots, t) = \sum a_n E_n \underbrace{P_{ij}}_{\text{scalars}} \phi_n(\dots, r_i, \dots, r_j, \dots, t) \\ &= P_{ij} \sum a_n E_n \phi_n(\dots, r_i, \dots, r_j, \dots, t) = P_{ij} H\Psi \\ &\rightarrow \text{We have proved that } [H, P_{ij}] = 0 \quad \forall i, j \quad (i \neq j) \end{aligned}$$

Commuting observables have a complete set of common eigenfunctions:

$$P_{ij}H\phi = P_{ij}E\phi = E(P_{ij}\phi) \xrightarrow{[H, P_{ij}]=0} H(P_{ij}\phi) \quad (1.5)$$

This implies $P_{ij}\phi$ is also an eigenfunction of H with the eigenvalue E .

$$P_{ij}\phi = \Theta\phi$$

Value of Θ ?

$$\begin{aligned} P_{ij}\phi(\dots, r_i, \dots, r_j, \dots, t) &= \phi(\dots, r_j, \dots, r_i, \dots, t) \\ P_{ij}^2\phi(\dots, r_i, \dots, r_j, \dots, t) &= \phi(\dots, r_i, \dots, r_j, \dots, t) = I\phi \end{aligned}$$

Then $P_{ij}^2 = I$.

As $P_{ij}\phi = \Theta\phi$:

$$\begin{aligned} P_{ij}\phi(\dots, r_i, \dots, r_j, \dots, t) &= \Theta\phi(\dots, r_i, \dots, r_j, \dots, t) \\ P_{ij}^2\phi(\dots, r_i, \dots, r_j, \dots, t) &= \Theta^2\phi(\dots, r_i, \dots, r_j, \dots, t) \end{aligned}$$

As $P_{ij}^2 = I$, then $\Theta^2 = 1 \implies \Theta = \pm 1$.

- **Bosons:** $\Theta = +1 \rightarrow \text{Symmetric}$
- **Fermions:** $\Theta = -1 \rightarrow \text{Antisymmetric}$

What happens if two particles are in the same position ($r_i = r_j$) and I exchange them with P_{ij} ?

First case (Fermions): $P_{ij}\phi(\dots, r_i, \dots, r_i, \dots, t) = -\phi(\dots, r_i, \dots, r_i, \dots, t)$
 $\implies \phi(\dots, r_i, \dots, r_i, \dots, t) = 0 \rightarrow$ It is **forbidden** that two particles occupy the same position. These particles are known as **Fermions** (half-integer spin).

Second case (Bosons): $P_{ij}\phi(\dots, r_i, \dots, r_i, \dots, t) = \phi(\dots, r_i, \dots, r_i, \dots, t)$
 $\implies \phi(\dots, r_i, \dots, r_i, \dots, t) = \phi(\dots, r_i, \dots, r_i, \dots, t) \rightarrow$ It is **not forbidden** that two particles occupy the same position. These particles are known as **Bosons** (integer spin).

Examples:

Fermions	Bosons
Electron	Quark
Proton	Gluon
Neutron	α particle
$\vdots$	$\vdots$
ϕ is antisymmetric ($\Theta = -1$)	ϕ is symmetric ($\Theta = +1$)

$$\text{Ideal} \rightarrow H = \sum_i^N H_i$$

$$\begin{aligned} \sum_i n_i &= N \\ \sum_i n_i \epsilon_i &= E \end{aligned}$$

Fermions vs Bosons occupancy:

- **Fermions:** $n_j \in \{0, 1\}$ (Pauli exclusion principle).
- **Bosons:** $n_j \in \{0, 1, 2, \dots, N\}$ (No restriction).

$$Z(T, V) = \sum_{\{n_i\}} e^{-\beta \sum n_i \epsilon_i} = \sum_{\{n_i\}} e^{-\beta E(\{n_i\})} \delta_{N, \sum n_i} \quad (1.6)$$

→ This restriction ($\sum n_i = N$) does not allow us to perform the sum easily.

Boltzmann approximation → $z_N = \sum_{\{n_i\}} \frac{n_1! n_2! \dots n_i!}{N!} e^{-\beta \sum n_i \epsilon_i} \approx \frac{1}{N!} \sum_{\{n_i\}} e^{-\beta \sum n_i \epsilon_i}$

Ensembles Comparison:

- **Microcanonical:** (E, V, N) fixed. Isolated system.
- **Canonical:** $(\langle E \rangle, V, N)$ fixed. Thermal contact with reservoir at T .
- **Grand Canonical (Macrocanonical):** $(\langle E \rangle, V, \langle N \rangle)$ fixed. Thermal and particle contact (μ, T) .

Macrocanonical ensemble

$\Xi(\mu, T, V) = \sum_{N=0}^{\infty} e^{\beta \mu N} Z_N(T, V) \rightarrow \text{similar to } \int e^{tx} p(x) dx \rightarrow \text{moment generating function}$

$$\begin{aligned} \Xi(\mu, T, V) &= \sum_{N=0}^{\infty} e^{\beta \mu N} \sum_{\{n_i\}} e^{-\beta \sum n_i \epsilon_i} \delta_{N, \sum n_i} = \sum_{\{n_i\}} e^{\beta \mu \sum n_i} \sum_{N=0}^{\infty} e^{\beta \mu N} \delta_{N, \sum n_i} \\ &= \sum_{\{n_i\}} e^{-\beta \sum n_i \epsilon_i} e^{\beta \mu \sum n_i} = \sum_{\{n_i\}} e^{\beta \sum (\mu - \epsilon_i) n_i} = \sum_{\{n_i\}} \prod_i e^{\beta (\mu - \epsilon_i) n_i} \\ &= \prod_i \sum_{\{n_i\}} e^{\beta (\mu - \epsilon_i) n_i} \end{aligned}$$

Fermions

$$n_i = 0, 1 \quad \forall i$$

then

$$\Xi(\mu, T, V) = \prod_i (1 + e^{\beta (\mu - \epsilon_i)}) \quad (1.7)$$

where 1 is when $n_i = 0$ and the exponential is when $n_i = 1$.

Bosons

$$n_i = 0, 1, \dots, \infty \quad \forall i$$

then

$$\Xi(\mu, T, V) = \prod_i \sum_{n_i=0}^{\infty} (e^{\beta(\mu-\epsilon_i)})^{n_i} = \prod_i (1 - e^{\beta(\mu-\epsilon_i)})^{-1} \quad (1.8)$$

using $\sum x^n = \frac{1}{1-x}$ for $|x| < 1$.

I can write generally this:

$$\Xi(\mu, T, V) = \prod_i (1 - \Theta e^{\beta(\mu-\epsilon_i)})^{-1/\Theta} \quad (1.9)$$

where $\Theta = \begin{cases} +1 & \text{for bosons} \\ -1 & \text{for fermions} \end{cases}$

Canonical ensemble	Macrocanonical ensemble
$A = -kT \ln Z$	$J = -kT \ln \Xi$
$A = U - TS$	$J = U - TS - \mu N$

From the macrocanonical ensemble we are going to use:

- $J = -kT \ln \Xi$
- $PV = -J$
- $\langle N \rangle = - \left(\frac{\partial J}{\partial \mu} \right)_{T,V}$

From what we know:

$$J = -kT \ln \Xi = -kT \ln \left[\prod_i (1 - \Theta e^{\beta(\mu-\epsilon_i)})^{-1/\Theta} \right] = \frac{kT}{\Theta} \sum_i \ln(1 - \Theta e^{\beta(\mu-\epsilon_i)}) \quad (1.10)$$

For Bosons $\mu < 0$ For Fermions μ can be any value

$$\langle N \rangle = - \left(\frac{\partial J}{\partial \mu} \right)_{T,V} = - \frac{kT}{\Theta} \sum_i \frac{-\Theta \beta e^{\beta(\mu-\epsilon_i)}}{1 - \Theta e^{\beta(\mu-\epsilon_i)}} = \sum_i \frac{1}{e^{\beta(\epsilon_i-\mu)} - \Theta} \rightarrow \text{this remembers us the restriction} \quad (1.11)$$

$$\Rightarrow \langle N \rangle = \langle \sum n_i \rangle = \sum \langle n_i \rangle$$

So we can know the average of n_i for each energy level ϵ_i :

$$\langle n_i \rangle = \frac{1}{e^{\beta(\epsilon_i-\mu)} - \Theta} \quad (1.12)$$

Average occupation number of level i :

- **Fermi-Dirac Statistics (Fermions):** $\langle n_i \rangle = \frac{1}{e^{\beta(\epsilon_i - \mu)} + 1}$
- **Bose-Einstein Statistics (Bosons):** $\langle n_i \rangle = \frac{1}{e^{\beta(\epsilon_i - \mu)} - 1}$

$$J = \frac{kT}{\Theta} \sum_i \ln(1 - \Theta e^{\beta(\mu - \epsilon_i)}) \quad (1.13)$$

$$\lim_{\Theta \rightarrow 0} J = \lim_{\Theta \rightarrow 0} -kT \sum_i \ln(1 - \Theta e^{\beta(\mu - \epsilon_i)})^{1/\Theta} = -kT \sum_i e^{\beta(\mu - \epsilon_i)} \quad (1.14)$$

Using $e^x = \lim_{n \rightarrow \infty} (1 + \frac{x}{n})^n$

Remember that $z_{MB} = \frac{1}{N!} (z_{MB})^N$ so $\Xi(\mu, T, V) = \sum_{N=0}^{\infty} e^{\beta\mu N} z_{MB} = \sum_{N=0}^{\infty} \lambda^N \frac{z_1^N}{N!} = e^{\lambda z_1}$ where λ is the fugacity: $\lambda = e^{\beta\mu}$

we reached the same result as the boltzmann gas approximation

$$J = -kT \ln \Xi = -\lambda z_1 kT = -kT e^{\beta\mu} \sum_i e^{-\beta\epsilon_i} = -kT \sum_i e^{\beta(\mu - \epsilon_i)}$$

$$\text{Then } J(\mu, T, V) = \frac{kT}{\Theta} \sum_i \ln(1 - \Theta e^{\beta(\mu - \epsilon_i)}) \begin{cases} \Theta = 1 & \text{Bosons} \\ \Theta = -1 & \text{Fermions} \\ \Theta \rightarrow 0 & \text{MB} \end{cases} \rightarrow \text{this uni-}$$

fied the three statistics.

$$\langle n_i \rangle = \frac{1}{e^{\beta(\epsilon_i - \mu)} - \Theta}$$

Boltzmann approximation $\rightarrow$ We see that bosons and fermions converge to the same behavior. In this limit: $\langle n_i \rangle = e^{\beta(\mu - \epsilon_i)} \ll 1$. In 3D $\rightarrow \rho \Lambda^3 \ll 1$.

Let's demonstrate that the Boltzmann approximation (which means $\langle n_i \rangle = e^{\beta(\mu - \epsilon_i)} \ll 1$) corresponds to considering $\rho \Lambda^3 \ll 1$

For free particle in 3D: $z_1 = \frac{V}{\Lambda^3}$, then $Z = \frac{z_1^N}{N!}$ (where $L \gg \Lambda$)
 $\rightarrow$ as calculated before for the Boltzmann gas.

$$A(N, T, V) = -kT \ln Z = -kT \ln \left(\frac{V^N}{\Lambda^{3N} N!} \right)$$

$$\mu = \left(\frac{\partial A}{\partial N} \right)_{T,V} \rightarrow \text{The chemical potential is the cost of Free Energy of varying } N.$$

A system always wants to minimize the Helmholtz Free Energy: $A = U - TS$

- sometimes we just need to minimize $U \rightarrow$ Example: $T = 0$
- sometimes we just need to maximize $S \rightarrow$ Example: fixed U

What does μ mean to A ?

- When $\mu < 0$, this means that as we increase N , we get a lower A . In this situation, the system wants more N in order to decrease A .
- When $\mu > 0$, this means that as we decrease N , we get higher A . Then the system wants to reduce N to decrease A .

Boltzmann approximation: $e^{\beta(\mu-\epsilon)} \ll 1 \implies \beta(\mu-\epsilon) \ll 0 \implies \mu\beta - \epsilon\beta \ll 0 \xrightarrow{\epsilon \rightarrow 0} \ln(\rho\Lambda^3) < 0 \implies \rho\Lambda^3 \ll 1$ (If it's satisfied for $\epsilon = 0$, the most unfavorable case, it's satisfied for any ϵ).

Thus, the Boltzmann approximation is equivalent to considering $\rho \ll 1$ and/or $T \gg 1$.

Thermodynamic properties of Bosons and Fermions

$$J(\mu, T, V) = \frac{kT}{\Theta} \sum_i \ln(1 - \Theta e^{\beta(\mu - \epsilon_i)})$$

We are going to consider **free particles in a 3D box**. What is the spectrum in this case?

$$\begin{aligned} H\psi_{n_x, n_y, n_z} &= \epsilon_{n_x, n_y, n_z} \psi_{n_x, n_y, n_z} \\ H &= \sum_{i=1}^N -\frac{\hbar^2}{2m} \nabla_i^2 \\ \epsilon_{n_x, n_y, n_z} &= \epsilon_{n_x} + \epsilon_{n_y} + \epsilon_{n_z} \\ \epsilon_{n_x} &= \frac{\hbar^2 \pi^2}{2m L^2} n_x^2 \quad \text{where } n = x, y, z \\ \psi_{n_x, n_y, n_z} &= \left(\frac{2}{L}\right)^{3/2} \sin\left(\frac{n_x \pi x}{L}\right) \sin\left(\frac{n_y \pi y}{L}\right) \sin\left(\frac{n_z \pi z}{L}\right) \end{aligned}$$

As $L \rightarrow \infty$, we are going to perform the sum over n_x, n_y, n_z instead of ϵ :
 $J(\mu, T, V) = \frac{kT}{\Theta} \sum_{n_x=1}^{\infty} \sum_{n_y=1}^{\infty} \sum_{n_z=1}^{\infty} \ln(1 - \Theta e^{\beta(\mu - \epsilon_{n_x, n_y, n_z})})$

$$\xrightarrow{\text{change of variable}} p_{n_x} = \frac{\hbar \pi n_x}{L} \rightarrow \epsilon_{n_x, n_y, n_z} = \frac{p_{n_x}^2 + p_{n_y}^2 + p_{n_z}^2}{2m}$$

$$J(\mu, T, V) = \frac{kT}{\theta} \sum_{p_x=\frac{\hbar}{2L}} \sum_{p_y=\frac{\hbar}{2L}} \sum_{p_z=\frac{\hbar}{2L}} \ln(1 - \theta e^{\beta(\mu - \frac{(p_x^2 + p_y^2 + p_z^2)}{2m})})$$

when $L \gg 1$

$$\sum_{p_x} \sum_{p_y} \sum_{p_z} \rightarrow \frac{gL^3}{h^3} \int_0^{\infty} dp_x \int_0^{\infty} dp_y \int_0^{\infty} dp_z f(p) \rightarrow \frac{gL^3}{2^3 h^3} \int_{-\infty}^{\infty} dp_x \int_{-\infty}^{\infty} dp_y \int_{-\infty}^{\infty} dp_z f(p)$$

where $f(p) = f(-p)$ is an even function.

Using $dp_x dp_y dp_z = 4\pi p^2 dp$:

$$\rightarrow \frac{4\pi V}{h^3} \int_0^\infty p^2 f(p) dp = \int_0^\infty D(p) f(p) dp \quad \text{where} \quad \boxed{D(p) = \frac{V}{h^3} 4\pi p^2} \text{ density of states}$$

We are going to see this many times. $D(p)dp \rightarrow$ number of states in $[p, p + dp]$.

Since we have a free particle: $\varepsilon = \frac{p^2}{2m}$, I can express $D(p)$ as $D(\varepsilon)$. $D(p)dp = D(\varepsilon)d\varepsilon$ must be the same, the number of states couldn't change when I change the variable.

$$\begin{aligned} D(\varepsilon)d\varepsilon &= D(p)dp \rightarrow D(\varepsilon)\frac{d\varepsilon}{dp} = D(p) \rightarrow D(\varepsilon)\frac{p}{m} = \frac{V}{h^3} 4\pi 2m\varepsilon \rightarrow D(\varepsilon)\frac{\sqrt{2m\varepsilon}}{m} = \frac{V}{h^3} 4\pi 2m\varepsilon \\ &\rightarrow D(\varepsilon) = \frac{V}{h^3} \frac{m}{\sqrt{2m\varepsilon}} 4\pi 2m\varepsilon = \frac{V}{h^3} 4\pi m \sqrt{2m\varepsilon} = \frac{V}{h^3} 4\pi m^{3/2} \sqrt{2\varepsilon} \end{aligned}$$

$$\hookrightarrow D(\varepsilon) = D(p)\frac{dp}{d\varepsilon} = \dots \text{ same result.}$$

$$J(\mu, T, V) = \frac{kT}{\theta} \frac{V}{h^3} 4\pi \int_0^\infty p^2 \ln(1 - \theta e^{\beta(\mu - p^2/2m)}) dp = \frac{kT}{\theta} \int_0^\infty D(p) \ln(1 - \theta e^{\beta(\mu - p^2/2m)}) dp = \frac{kT}{\theta} \int_0^\infty D(\varepsilon) \ln(1 - \theta e^{\beta(\mu - \varepsilon)}) d\varepsilon$$

As factor of energy $\rightarrow$ we changed $D(p)dp = D(\varepsilon)d\varepsilon$ so we have not quantum constant.

$$\frac{kT}{\theta} \int_0^\infty D(\varepsilon) \ln(1 - \theta e^{\beta(\mu - \varepsilon)}) d\varepsilon = \frac{kT}{\theta} \frac{V}{h^3} 4\pi m^{3/2} \sqrt{2} \int_0^\infty \sqrt{\varepsilon} \ln(1 - \theta e^{\beta(\mu - \varepsilon)}) d\varepsilon \rightarrow \text{in-} \\ \text{tegration by parts}$$

$$\begin{aligned} u &= \ln(1 - \theta e^{\beta(\mu - \varepsilon)}) \quad du = \frac{\theta e^{\beta(\mu - \varepsilon)}}{1 - \theta e^{\beta(\mu - \varepsilon)}} \beta \quad v = \frac{2}{3} \varepsilon^{3/2} \\ &= \left(\ln(1 - \theta e^{\beta(\mu - \varepsilon)}) \frac{2}{3} \varepsilon^{3/2} \right) \Big|_0^\infty - \int_0^\infty \frac{2}{3} \varepsilon^{3/2} \frac{\theta e^{\beta(\mu - \varepsilon)}}{1 - \theta e^{\beta(\mu - \varepsilon)}} \beta d\varepsilon \left[\frac{kT}{\theta} \frac{V}{h^3} 4\pi m^{3/2} \sqrt{2} \right] \\ &= -\frac{2}{3} \frac{V}{h^3} 4\pi m^{3/2} \sqrt{2} \int_0^\infty \varepsilon^{3/2} \frac{\beta \theta e^{\beta(\mu - \varepsilon)}}{1 - \theta e^{\beta(\mu - \varepsilon)}} d\varepsilon = d\varepsilon \int_0^\infty \varepsilon^{3/2} \frac{1}{e^{\beta(\varepsilon - \mu)} - \theta} d\varepsilon \end{aligned}$$

$<$ this is the average occupation number $\langle n \rangle$

$$J = -PV \rightarrow P(T, \mu) = \frac{2}{3} \frac{V}{\Lambda^3} \frac{1}{\Gamma(3/2)} \int_0^\infty \varepsilon^{3/2} \frac{1}{e^{\beta(\varepsilon - \mu)} - \theta} d\varepsilon$$

This integral is very difficult, the way going to study it is specific cases.

$$\langle N \rangle = - \left(\frac{\partial J}{\partial \mu} \right)_{T, V} = V \left(\frac{\partial P}{\partial \mu} \right)_T \approx \frac{V}{\Lambda^3} (kT)^{3/2} \int_0^\infty E^{1/2} \langle n(E) \rangle_\theta d\varepsilon$$

$$\langle N \rangle = \sum_i \langle n(\varepsilon_i) \rangle_\theta \rightarrow \text{in the continuum limit} \rightarrow \text{Derive this I got the same as} \\ = \frac{V}{h^3} \frac{8\pi}{\sqrt{2}} m^{3/2} \int_0^\infty \varepsilon^{1/2} \langle n(\varepsilon) \rangle_\theta d\varepsilon$$

$$\rho(T, \mu) = \frac{\langle N \rangle}{V} = \frac{8\pi m^{3/2}}{h^3 \sqrt{2}} \int_0^\infty \varepsilon^{1/2} \langle n(\varepsilon) \rangle_\theta d\varepsilon$$

$$\langle E \rangle = \sum_i \varepsilon_i \langle n(\varepsilon_i) \rangle_\theta$$

$$\langle E \rangle = \int_\varepsilon D(\varepsilon) \varepsilon \langle n(\varepsilon) \rangle_\theta d\varepsilon \rightarrow \text{in the continuum limit} = \frac{V 8\pi m^{3/2}}{h^3 \sqrt{2}} \int_0^\infty \varepsilon^{3/2} \langle n(\varepsilon) \rangle_\theta d\varepsilon = -\frac{3}{2} J = \frac{3}{2} PV \rightarrow \text{It not depends if the particles are bosons or fermions.}$$

$$\left. \begin{aligned} PV &= \frac{2}{3} \langle E \rangle \\ \langle E \rangle &= \frac{3}{2} PV \end{aligned} \right\} \text{this is equal for bosons and fermions.}$$

$$\text{For Boltzmann gas: } \left. \begin{aligned} PV &= NkT \\ u = \langle E \rangle &= \frac{3}{2} NkT \end{aligned} \right\} PV = \frac{2}{3} u \rightarrow \text{It is also satisfied for Boltzmann gas.}$$

Weakly degenerate limit

$$\text{Classical limit: } \langle n(\epsilon) \rangle_\theta \ll 1 \rightarrow \lambda \text{ (fugacity)} \ll 1 \quad \lambda = e^{\beta\mu}$$

$$\text{Boltzmann approximation is OK as } \left(\frac{\partial P}{\partial \lambda} \right)_T = \text{ctan}(\rho(\epsilon)) \rightarrow \lambda \rho \Lambda^3 \text{ is the same value obtained in Boltzmann approximation.}$$

$$\text{In this case, } \frac{\lambda}{\Lambda^3} \text{ connects to the ideal gas law since I recover that } P = \rho kT \text{ as } PV = NkT \quad D(\epsilon) \text{ is } g \frac{V}{h^3} (1 - \theta e^{-\beta(\epsilon-\mu)})^{-1}$$

$$\text{For fermions we can check that when } \theta = -1 \rightarrow \text{higher } P$$

$$P(T, \mu) = \frac{g}{\theta} \frac{kT}{\Lambda^3} \frac{V}{h^3} \int_0^\infty \epsilon^{1/2} \langle n(\epsilon) \rangle_\theta d\epsilon \quad P(3/2) = \frac{3}{4} \sqrt{\pi}$$

$$\int_0^\infty \frac{\epsilon^{3/2}}{e^{\beta(\epsilon-\mu)} - \theta} d\epsilon = \int_0^\infty \frac{\epsilon^{3/2}}{e^{\beta\epsilon} \lambda^{-1} - \theta} d\epsilon \rightarrow \text{since } \theta \lambda e^{-\beta\epsilon} < 1 \rightarrow \int_0^\infty \epsilon^{3/2} \lambda e^{-\beta\epsilon} \sum_{n=0}^\infty (\theta \lambda e^{-\beta\epsilon})^n = \int_0^\infty \epsilon^{3/2} \lambda e^{-\beta\epsilon} (1 + \theta \lambda e^{-\beta\epsilon} + \theta^2 \lambda^2 e^{-2\beta\epsilon} + \dots) = \lambda \Gamma(5/2) \beta^{-5/2} + \lambda^2 \theta \Gamma(5/2) 2^{-5/2} \beta^{-5/2} + \lambda^3 \theta^2 \Gamma(5/2) 3^{-5/2} \beta^{-5/2} + \dots = \lambda \frac{3}{4} \sqrt{\pi} \beta^{-5/2} (1 + \theta \lambda 2^{-5/2} + \theta^2 \lambda^2 3^{-5/2} + \dots)$$

$$\text{Using this: } \int_0^\infty dx x^n e^{-ax} = \frac{\Gamma(n+1)}{a^{n+1}}$$

$$\text{Then } P(T, \mu) = \frac{g}{\Lambda^3} \frac{kT}{\theta} \frac{1}{\theta} (\lambda \frac{3}{4} \sqrt{\pi} \beta^{5/2} \beta^{-5/2} (1 + \theta \lambda 2^{-5/2} + \theta^2 \lambda^2 3^{-5/2} + \dots)) = \frac{gkT}{\Lambda^3} (\lambda + \theta \lambda^2 2^{-5/2} + \theta^2 \lambda^3 3^{-5/2} + \dots) \quad \text{Our goal is to express this } P = P(T, \rho). \text{ Check this in line.}$$

$$\rho(T, \mu) = \frac{\langle N \rangle}{V} = \frac{1}{V} \left(\frac{\partial J}{\partial \mu} \right)_{T, V} = \left(\frac{\partial P}{\partial \mu} \right)_T \Rightarrow \rho(T, \lambda) = \left(\frac{\partial P}{\partial \lambda} \right)_T \left(\frac{\partial \lambda}{\partial \mu} \right)_T = \lambda \left(\frac{\partial P}{\partial \lambda} \right)_T = \frac{g}{\Lambda^3} (1 + \theta \lambda 2^{-3/2} + \theta^2 \lambda^2 3^{-3/2} + \dots)$$

$$\text{Next challenge: } \lambda(\rho, T) \rightarrow \lambda \propto \rho \rightarrow \text{then substitute in } P \text{ to obtain } P = P(\rho, T) \text{ From } \rho(T, \mu) = \frac{g}{\Lambda^3} (\lambda + \theta \lambda^2 2^{-3/2} + \theta^2 \lambda^3 3^{-3/2} + \dots) \Rightarrow \lambda(\rho, T) = \Lambda^3 \rho - \theta 2^{-3/2} (\Lambda^3 \rho)^2 - \left(\frac{1}{3^{3/2}} - \frac{1}{2^{3/2}} \right) (\Lambda^3 \rho)^3 + \dots$$

$$\text{Lagrange theorem} \rightarrow (\text{Inverse } y = \rho \text{ and } x = \lambda) \text{ We found } y = ax + bx^2 + cx^3 + \dots \text{ (eq. 1) We want } x = sy + ty^2 + uy^3 + \dots \text{ Step 1: Substitute (we want until second order in } y \text{)}$$

order, y^2 so) $x(y)$ into the rhs (right hand side) of eq. (1) $y = a(sy+ty^2+\dots)+b(sy+ty^2)^2+c(sy+ty^2)^3+\dots$. Let's compute this up to the second order $O(y^2)$ First term: $a(sy+ty^2) = asy+aty^2$ Second term: $b(sy)^2 = bs^2y^2 \rightarrow y = asy+aty^2+bs^2y^2+O(y^3)$ (eq. 2)

Step 2: Equal term by term in eq. (2) $y = asy + (aty + bs^2)y^2 + O(y^3)$ $y = asy \Rightarrow s = 1/a$ $aty^2 + bs^2y^2 = 0 \rightarrow at + bs^2 = 0 \rightarrow at + \frac{b}{a^2} = 0 \rightarrow t = -b/a^3$

$$\text{So } x(y) = \frac{y}{a} - \frac{b}{a^3}y^2$$

$\Rightarrow P(T, \lambda(T, \rho)) = kT\rho(1 - \theta(2^{-5/2} - 2^{-3/2})\Lambda^3\rho + O((\Lambda^3\rho)^2)) \uparrow$ virial expansion in terms of ρ With $y = \rho$, $x = \lambda$, $a = \Lambda^{-3}$, $b = \theta 2^{-3/2} \hookrightarrow \lambda(\rho, T) = \Lambda^3\rho - \Lambda^3\theta 2^{-3/2}\rho^2 + \dots \approx \Lambda^3\rho(1 - \theta 2^{-3/2}\Lambda^3\rho + \dots)$ $\frac{\Lambda^3\rho}{g} < 1$ Boltzmann approximation. Is weakly degenerate limit.