

# Solving Linear Equations

R4 Cheng

May 10, 2024

## Matrix Operations

$$\begin{bmatrix} 1 & 2 & -4 \\ -2 & 3 & 1 \\ 4 & 1 & 2 \end{bmatrix} \begin{bmatrix} x_1 & x_2 & x_3 \\ y_1 & y_2 & y_3 \\ z_1 & z_2 & z_3 \end{bmatrix} = \begin{bmatrix} \underline{c_1} & \underline{c_2} & \underline{c_3} \end{bmatrix}$$

$$\underline{c_2} = x_2 \begin{bmatrix} 1 \\ -2 \\ 4 \end{bmatrix} + y_2 \begin{bmatrix} 2 \\ 3 \\ 1 \end{bmatrix} + z_2 \begin{bmatrix} -4 \\ 1 \\ 2 \end{bmatrix}$$

$$\begin{bmatrix} a & b & c \end{bmatrix} \begin{bmatrix} 1 & 2 & -4 \\ -2 & 3 & 1 \\ 4 & 1 & 2 \end{bmatrix} = a \begin{bmatrix} 1 & 2 & -4 \end{bmatrix} + b \begin{bmatrix} -2 & 3 & 1 \end{bmatrix} + c \begin{bmatrix} 4 & 1 & 2 \end{bmatrix}$$

## Properties of Matrices

$A(BC) = (AB)C$  (Associative law holds)

$AB \neq BA$  (Commutative law does not hold)

$C(A + B) = CA + CB$  or  $(A + B)C = AC + BC$  (Distributive laws hold)

**Remark.** *We can change Gaussian Elimination to Matrix multiplication*

## Identity Matrix

The identity matrix is a square matrix with ones on the main diagonal and zeros elsewhere. It is denoted by  $I$  or  $I_n$  for an  $n \times n$  matrix.  $AI = IA = A$ , for any  $n \times n$  matrix  $A$ .

$$I = \begin{bmatrix} 1 & 0 & \cdots & 0 \\ 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 1 \end{bmatrix}$$

## Inverse Matrix

The inverse of a square matrix  $A$ , denoted as  $A^{-1}$ , is a matrix such that  $AA^{-1} = A^{-1}A = I$ , where  $I$  is the identity matrix. The inverse matrix can be found using the formula:

$$A^{-1} = \frac{1}{\det(A)} \cdot \text{adj}(A)$$

where  $\det(A)$  is the determinant of matrix  $A$  and  $\text{adj}(A)$  is the adjugate of matrix  $A$ .

For example, to find the inverse of a  $2 \times 2$  matrix:

$$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

the inverse matrix is given by:

$$A^{-1} = \frac{1}{ad - bc} \cdot \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$$

To find the inverse of a  $3 \times 3$  matrix, you can use the formula:

$$A^{-1} = \frac{1}{\det(A)} \cdot \text{adj}(A)$$

where  $\text{adj}(A)$  is the adjugate of matrix  $A$ . The adjugate of a  $3 \times 3$  matrix is given by:

$$\text{adj}(A) = \begin{bmatrix} A_{11} & A_{21} & A_{31} \\ A_{12} & A_{22} & A_{32} \\ A_{13} & A_{23} & A_{33} \end{bmatrix}$$

where  $A_{ij}$  is the cofactor of element  $a_{ij}$  in matrix  $A$ .

Note that not all matrices have an inverse. A matrix is invertible if and only if its determinant is non-zero.

### Attributes

- It is unique.
- The inverse of  $A^{-1}$  is  $A$  itself.

---

**Claim.** Suppose  $A$  is invertible. Then its inverse is unique.

**Proof.** Suppose  $B$  and  $C$  are both inverses of  $A$ . Then  $B = BI = B(AC) = (BA)C = IC = C$ .

**Remark.** *left inverse = right inverse = inverse*

**Claim.** The inverse of  $A^{-1}$  is  $A$  itself.

**Proof.**  $AA^{-1} = I$  and  $A^{-1}A = I$ .

**Claim.** If  $A$  is invertible, then the one and only solution to  $A\underline{x} = \underline{b}$  is  $\underline{x} = A^{-1}\underline{b}$ .

**Proof.**  $A\underline{x} = \underline{b} \Rightarrow A^{-1}A\underline{x} = A^{-1}\underline{b} \Rightarrow \underline{x} = A^{-1}\underline{b}$ .

**Claim.** Suppose there is a nonzero solution  $\underline{x}$  to  $A\underline{x} = \underline{0}$  (homogeneous equation). Then  $A$  is not invertible.

**Proof.** If  $A$  is invertible, then  $A^{-1}$  exists. Then  $A^{-1}A\underline{x} = A^{-1}\underline{0} \Rightarrow \underline{x} = \underline{0}$ .

**Claim.** A diagonal matrix has an inverse provided no diagonal entries are zero.

**Proof.**

If

$$A = \begin{bmatrix} d_1 & 0 & \cdots & 0 \\ 0 & d_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & d_n \end{bmatrix}$$

then

$$A^{-1} = \begin{bmatrix} \frac{1}{d_1} & 0 & \cdots & 0 \\ 0 & \frac{1}{d_2} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \frac{1}{d_n} \end{bmatrix}$$

**Claim.** If  $A$  and  $B$  are invertible, then  $AB$  is invertible and  $(AB)^{-1} = B^{-1}A^{-1}$ .

**Proof.**

$$(B^{-1}A^{-1})(AB) = B^{-1}(A^{-1}A)B = B^{-1}IB = I$$

$$(AB)(B^{-1}A^{-1}) = A(BB^{-1})A^{-1} = AIA^{-1} = I$$

**Remark.**  $(ABC)^{-1} = C^{-1}B^{-1}A^{-1}$

## Gauss-Jordan Elimination

Given  $A$ , we want to find its inverse  $A^{-1}$ .  $AA^{-1} = I$

$$A \begin{bmatrix} \underline{col}_1 & \underline{col}_2 & \underline{col}_3 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} \underline{e}_1 & \underline{e}_2 & \underline{e}_3 \end{bmatrix}$$

$$A = \begin{bmatrix} 2 & -1 & 0 \\ -1 & 2 & -1 \\ 0 & -1 & 2 \end{bmatrix}$$

$$\begin{bmatrix} \boxed{2} & -1 & 0 & \vdots & 1 & 0 & 0 \\ -1 & 2 & -1 & \vdots & 0 & 1 & 0 \\ 0 & -1 & 2 & \vdots & 0 & 0 & 1 \end{bmatrix}$$

$\Rightarrow$

$$\begin{bmatrix} 2 & -1 & 0 & \vdots & 1 & 0 & 0 \\ 0 & \boxed{\frac{3}{2}} & -1 & \vdots & \frac{1}{2} & 1 & 0 \\ 0 & -1 & 2 & \vdots & 0 & 0 & 1 \end{bmatrix}$$

$\Rightarrow$

$$\begin{bmatrix} 2 & -1 & 0 & \vdots & 1 & 0 & 0 \\ 0 & \frac{3}{2} & -1 & \vdots & \frac{1}{2} & 1 & 0 \\ 0 & 0 & \boxed{\frac{4}{3}} & \vdots & \frac{1}{3} & \frac{2}{3} & 1 \end{bmatrix}$$

Until here is Gauss

$\Rightarrow$

$$\begin{bmatrix} 2 & -1 & 0 & \vdots & 1 & 0 & 0 \\ 0 & \frac{3}{2} & 0 & \vdots & \frac{3}{4} & \frac{3}{2} & \frac{3}{4} \\ 0 & 0 & \boxed{\frac{4}{3}} & \vdots & \frac{1}{3} & \frac{2}{3} & 1 \end{bmatrix}$$

$\Rightarrow$

$$\begin{bmatrix} 2 & 0 & 0 & \vdots & \frac{3}{2} & 1 & \frac{1}{2} \\ 0 & \boxed{\frac{3}{2}} & 0 & \vdots & \frac{3}{4} & \frac{3}{2} & \frac{3}{4} \\ 0 & 0 & \boxed{\frac{4}{3}} & \vdots & \frac{1}{3} & \frac{2}{3} & 1 \end{bmatrix}$$

$\Rightarrow$

$$\begin{bmatrix} 1 & 0 & 0 & \vdots & \frac{3}{4} & \frac{1}{2} & \frac{1}{4} \\ 0 & 1 & 0 & \vdots & \frac{1}{2} & 1 & \frac{1}{2} \\ 0 & 0 & 1 & \vdots & \frac{1}{4} & \frac{1}{2} & \frac{3}{4} \end{bmatrix}$$

Until here is Jordan

$$col_1 = \begin{bmatrix} \frac{3}{4} \\ \frac{1}{2} \\ \frac{1}{4} \end{bmatrix}, col_2 = \begin{bmatrix} \frac{1}{2} \\ 1 \\ \frac{1}{2} \end{bmatrix}, col_3 = \begin{bmatrix} \frac{1}{4} \\ \frac{1}{2} \\ \frac{3}{4} \end{bmatrix}, A^{-1} = \begin{bmatrix} \frac{3}{4} & \frac{1}{2} & \frac{1}{4} \\ \frac{1}{2} & 1 & \frac{1}{2} \\ \frac{1}{4} & \frac{1}{2} & \frac{3}{4} \end{bmatrix}$$

**Claim.** A matrix is invertible if and only if (iff.) it is nonsingular.

## Elimination = Factorization: $A = LU$

TODO

**Claim.** If  $A = L_1 D_1 U_1$  and  $A = L_2 D_2 U_2$ , where the  $L$ 's are lower-triangular with unit diagonal, the  $U$ 's are upper-triangular with unit diagonal, and the  $D$ 's are diagonal matrices with no zeros on the diagonal, then  $L_1 = L_2$ ,  $D_1 = D_2$ , and  $U_1 = U_2$ .