

## Calculus Stewart Ch11 Sec 2

50. A sequence of terms is defined by

$$a_1 = 1, a_n = (5 - n)a_{n-1}$$

Calculate  $\sum_{n=1}^{\infty} a_n$ .

Solution:

$$a_1 = 1, a_2 = 3, a_3 = 2 \cdot 3 = 6, a_4 = 1 \cdot 6 = 6, a_5 = 0 \cdot 6 = 0$$

$$\Rightarrow a_k = 0 \text{ for } k \geq 5$$

$$\sum_{n=1}^{\infty} a_n = 1 + 3 + 6 + 6 = 16$$

73. Find value of  $c$  if

$$\sum_{n=2}^{\infty} (1+c)^{-n} = 2$$

Solution:

$$\sum_{n=2}^{\infty} (1+c)^{-n} = \frac{\left(\frac{1}{1+c}\right)^2}{1 - \frac{1}{1+c}} = 2$$

$$\Rightarrow 2c^2 + 2c - 1 = 0 \Rightarrow c = \frac{1 - \sqrt{3}}{2} < 1$$

83. If  $\sum a_n$  is convergent and  $\sum b_n$  is divergent. Show that the series  $\sum(a_n + b_n)$  is divergent.

Proof:

Suppose  $\sum(a_n + b_n)$  is convergent. Then

$$\sum(a_n + b_n) - \sum a_n \text{ is convergent}$$

$$= \sum(a_n + b_n) - a_n = \sum b_n \text{ is convergent (* contradicts)}$$

$$\therefore \sum(a_n + b_n) \text{ is divergent}$$

## Calculus Stewart Ch11 Sec 3

42.  $\sum_{n=1}^{\infty} \frac{(\ln n)^2}{n^2}$

$$\lim_{n \rightarrow \infty} \frac{\ln n}{\sqrt[4]{n}} = \lim_{n \rightarrow \infty} \frac{\frac{1}{n}}{\frac{1}{4\sqrt[4]{n^3}}} = \lim_{n \rightarrow \infty} \frac{4}{\sqrt[4]{n}} = 0$$

$\therefore \ln n < \sqrt[4]{n}$  for large  $n$

$$\frac{(\ln n)^2}{n^2} < \frac{(\sqrt[4]{n})^2}{n^2} = \frac{\sqrt{n}}{n^2} = \frac{1}{n^{\frac{3}{2}}}$$

$$\Rightarrow \sum \frac{(\ln n)^2}{n^2} < \sum \frac{1}{n^{\frac{3}{2}}} \text{ converges}$$

45. For all positive values of  $b$  for which the series  $\sum_n b^{\ln n}$  converges.

Solution:

$$b^{\ln n} = n^{\ln b}$$

$$\sum_n b^{\ln n} \text{ converges iff. } -\ln b > 1 \Leftrightarrow 0 < b < \frac{1}{e}$$

#### Calculus Stewart Ch11 Sec 4

44. Show that if  $a_n > 0$  and  $\sum a_n$  is convergent, then  $\sum \ln(1 + a_n)$  is convergent.

Hint:

$$x > \ln(1 + x)$$

45. If  $\sum a_n$  is a convergent series with positive terms, is  $\sum \sin(a_n)$  also converges?

Solution:

$$\because \sum a_n \text{ converges} \therefore \lim_{n \rightarrow \infty} a_n = 0$$

$$\text{By Comparison Test, } \lim_{n \rightarrow \infty} \frac{|\sin(a_n)|}{|a_n|} = 1 > 0$$

$$\therefore \sum \sin(a_n) \text{ converges.}$$

#### Calculus Stewart Ch11 Sec 5

31. The terms of a series are defined recursively by the equations

$$a_1 = 2, a_{n+1} = \frac{5n+1}{4n+3} a_n$$

Determine whether  $\sum a_n$  converges or diverges.

Solution:

Since  $a_n$  is positive definite, by ratio test,

$$\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \frac{5}{4} > 1$$

$\therefore \sum a_n$  diverges.

# Calculus Stewart Ch11 Sec 7

23.  $\sum_{n=1}^{\infty} \tan\left(\frac{1}{n}\right)$

Solution:

By Comparison test,  $\lim_{n \rightarrow \infty} \frac{\tan\left(\frac{1}{n}\right)}{\frac{1}{n}} = \lim_{n \rightarrow \infty} \frac{\sin\left(\frac{1}{n}\right)}{\frac{1}{n}} \frac{1}{\cos\left(\frac{1}{n}\right)} = 1 > 0$ .

$\therefore \sum_{n=1}^{\infty} \tan\left(\frac{1}{n}\right)$  diverges

27.  $\sum_{k=1}^{\infty} \frac{k \ln k}{(k+1)^3}$

Solution:

$$\lim_{k \rightarrow \infty} \frac{\ln k}{\sqrt{k}} = 0$$

$\therefore \ln k < \sqrt{k}$  for large  $k$

$$\therefore \frac{k \ln k}{(k+1)^3} < \frac{k\sqrt{k}}{(k+1)^3} < \frac{k\sqrt{k}}{k^3} = \frac{1}{k^{\frac{3}{2}}}$$

$\therefore \sum_{k=1}^{\infty} \frac{k \ln k}{(k+1)^3}$  converges

36.  $\sum_{k=1}^{\infty} \frac{1}{(\ln k)^{\ln k}}$

Solution:

$\ln k > e^2$  for large  $k$

$$\frac{1}{(\ln k)^{\ln k}} < \frac{1}{(e^2)^{\ln k}} = \frac{1}{k^2}$$

$\therefore \sum_{k=1}^{\infty} \frac{1}{(\ln k)^{\ln k}}$  converges