
Homework 1
Due date: 2018.10.15

Problem 1. The critical density of the Universe is

$$\rho_c = \frac{3H_0^2}{8\pi G_N},$$

where H_0 is the present day Hubble expansion rate, given as $H_0 = 100 h \text{ km s}^{-1} \text{ Mpc}^{-1}$, where h is a dimensionless number. In the following questions, use $G_N = 6.674 \times 10^{-11} \text{ m}^3 \text{ kg}^{-1} \text{ s}^{-2}$, $1 \text{ Mpc} = 3.086 \times 10^{22} \text{ m}$, the Solar mass $M_\odot = 1.988 \times 10^{30} \text{ kg}$, and take $h = 0.7$.

Please round your results to two significant figures for 1), 2), 3) and 5), and don't worry about uncertainties.

- 1) What is the value of $1/H_0$ in Gyr? (1 Gyr = 10^9 year) [1 point]
- 2) What is the value of ρ_c in g/cm^3 ? [1 point]
- 3) What is the value of ρ_c in M_\odot/Mpc^3 ? [1 point]
- 4) In SI units, the Planck density (i.e., mass per unit volume), ρ_{Pl} , can be constructed as $\hbar^{n_1} c^{n_2} G_N^{n_3}$. Find n_1, n_2 and n_3 . [1 point]
- 5) In $\hbar = c = 1$ units, what are the values of ρ_{Pl} and ρ_c in GeV^4 ? [1 point] (Please note that we have given in class the relations between m & s & kg in SI units and the power of MeV or GeV in $\hbar = c = 1$ units. However, at that time we only kept one or two significant figures. Therefore, you might want to re-calculate them keeping more significant figures in order to achieve the correct two significant figure results of ρ_{Pl} and ρ_c .)

Problem 2. Given two four-vectors, $a^\mu = (2, 0, 1, 8)$ and $b^\mu = (0, 9, 2, 7)$, find: a_μ , b_μ , $\vec{a} \cdot \vec{a}$, $\vec{a} \cdot \vec{b}$, a^2 , $a \cdot b$, $(a + b)^2$. [2 points]

Note: please use the convention $g_{00} = -g_{11} = -g_{22} = -g_{33} = 1$ & $g_{\mu\nu} = 0$ (when $\mu \neq \nu$) for the calculations. A three-vector, e.g., \vec{a} , is the one appearing in $a^\mu = (a^0, \vec{a})$. A four-vector dot product, e.g., $a \cdot b$, is defined as $a \cdot b \equiv a^\mu b_\mu$. In particular, $a^2 \equiv a \cdot a$.

Problem 3. In a two-body scattering event, $A + B \rightarrow C + D$, it is convenient to introduce the *Mandelstam variables* $s \equiv (p_A + p_B)^2$, $t \equiv (p_A - p_C)^2$ and $u \equiv (p_A - p_D)^2$ (note that the square of a four-momentum should be always understood as, for example, $a^2 \equiv a \cdot a \equiv a^\mu a_\mu$). **The virtue of the Mandelstam variables is that they are Lorentz invariants, with the same value in any inertial system.**

- 1) Show that $s + t + u = m_A^2 + m_B^2 + m_C^2 + m_D^2$. [1 point]
- 2) Find the energy of A in the center-of-momentum (CM) frame, in terms of s, m_A and m_B . [2 points]
- 3) Find the energy of A in B's rest frame, in terms of s, m_A and m_B . [2 points]

Problem 4. In the perfect vacuum, an electron and a positron can annihilate into two or more photons ($e^+ + e^- \rightarrow n \gamma$, with $n \geq 2$). From the view of conservation of energy and momentum, please explain why the single photon creation process cannot happen. [1 point]