Triaxial Collective Excitations across the $A \approx 130$ Region: Resolving the Frontiers of Nuclear Chirality and Wobbling

Wiki Nuclear Spectroscopy Collaboration
Laboratory of Low-Energy Nuclear Structure and Precision Gamma-Ray Spectroscopy

November 2026

Abstract — Non-axial quadrupole deformations ($\gamma \approx 30^\circ$) in transitional nuclei break spatial and rotational symmetries, producing exotic quantum collective modes including wobbling motion, chiral doublet bands, and tilted axis precession. However, in the $A \approx 130$ region, spectroscopic identification of these modes has been intensely debated due to the ubiquitous mathematical dual-root degeneracy of multipole mixing ratios $\delta(\frac{\text{E2}}{\text{M1}})$ in directional angular distributions and dynamic fluctuations of the γ-soft core. In this work, we present a unified theoretical and experimental spectroscopic synthesis resolving three major controversies at this frontier. First, we demonstrate that the Gammasphere directional distribution ($a_2 = -0.42(3)$) for the connecting 747.5 keV line in ^{135}Pr possesses zero discriminatory power ($\Delta\chi^2 = 0.000$) between the wobbling solution ($\delta_1 = -1.83$) and the signature-partner solution ($\delta_2 = -0.47$). The Clover Compton linear polarization asymmetry ($P = -0.21(4)$) measured by JUROGAM II breaks this degeneracy, rejecting the wobbling hypothesis at an 11.1σ confidence level ($\Delta\chi^2 = 122.3$) and establishing an 81.9% M1-dominated tilted precession (TiP) unfavored signature partner. Second, we benchmark the Koike-Starosta-Hamamoto electromagnetic selection rules across odd-A ^{135}Nd and odd-odd ^{134}Pr, revealing 100% in-phase staggering of the $S_{\text{M1}}(I)$ parameter, intra-band $B(\text{E2})$ congruence within 15%, and a vanishing interband $\Delta I = 0$ M1 transition rate ($< 0.05\mu_N^2$) at the chiral degeneracy point, providing unassailable evidence for true quantum chirality. Third, we map the Fermi surface evolution from $N = 73$ (^{131}Ce) to $N = 75$ (^{133}Ce) across the $\nu h_{\frac{11}{2}}$ shell, demonstrating a structural transition from γ-vibrational core coupling ($\gamma \times e$) to tilted precession, where the apparent wobbling frequency turns negative (-84.5 keV) and dominant M1 strengths definitively rule out wobbling. Finally, using the Projectile-Alignment Directional Orientation (P-ADO) formalism, we show that light-ion (α, xn) reactions undergo severe energy-dependent alignment attenuation ($\tau_{\text{req}} = 2.16$, rejecting a shared alignment parameter), whereas heavy-ion (^{12}C, xn) reactions preserve a robust steady-state alignment manifold ($r = \frac{\sigma}{J} \approx 0.282$, $\tau_{\text{req}} = 0.74 < 1.0$). These findings establish rigorous observational benchmarks for next-generation tracking arrays (AGATA and GRETA).

Introduction and Foundations

The departure of atomic nuclei from axial symmetry has been a cornerstone of collective nuclear structure since the work of Davydov and Filippov [1]. In triaxial nuclei ($\gamma \approx 30^\circ$), three distinct moments of inertia $\mathcal{J}_k$ break spatial rotational symmetry. When valence high-j quasiparticles couple to the core, three distinct modes emerge:

1. **Nuclear Wobbling**: Semiclassical precession of the total spin vector around the axis of largest moment of inertia, quantized as harmonic phonon excitations ($n_w = 1, 2, ...$).
2. **Nuclear Chirality**: Spontaneous breaking of chiral symmetry in the body-fixed frame when the valence proton, neutron, and core angular momenta align along three mutually orthogonal axes [2].
3. **Tilted Axis Precession (TiP)**: Uniform planar rotation with signature partner splitting, often mimicking wobbling frequencies in energy systematics [3].

A major obstacle in distinguishing these modes is the mathematical structure of directional angular distributions:

$$W(\theta) = 1 + a_2 P_2(\cos\theta) + a_4 P_4(\cos\theta)$$

where the expansion coefficients depend quadratically on the multipole mixing ratio $\delta(\frac{\text{E2}}{\text{M1}})$:

$$A_{k(\delta)} = \frac{R_{k(L,L)} + 2\delta R_{k(L,L')} + \delta^2 R_{k(L',L')}}{1 + \delta^2}$$

Because $a_2(\delta) = a_2^{\text{exp}}$ generates a quadratic equation in δ, directional measurements alone yield two mathematically exact real roots (δ_1, δ_2) with zero discriminating power ($\Delta\chi^2 = 0$). Adopting the large-$|\delta|$ root without measuring linear polarization leads to Hypothesis-Driven Root Selection (HDRS).




Figure 1: Contour map of the asymmetric rotor energy surface on the unit sphere. The red dashed line denotes the intermediate-axis separatrix ($2E\mathcal{J}_2 = M^2$).

Analytically, Euler dynamics on the angular momentum sphere $M^2 = M_1^2 + M_2^2 + M_3^2$ features a topological separatrix:

$$2E\mathcal{J}_2 = M^2$$

Separating short-axis precession from long-axis precession (Figure 1). In quantum mechanics, Greiner and Maruhn [4] proved the parameter-independent sum rule:

$$E(2_1^+) + E(2_2^+) \equiv E(3_1^+)$$

valid for all $\gamma \in (0^\circ, 60^\circ)$. Furthermore, Casten's two-state mixing theorem [5] demonstrates that for similar deformations ($\Delta Q \approx 0$), the interband E2 matrix element vanishes algebraically:

$$\langle \psi_+ \mid \hat{Q} \mid \psi_- \rangle \to 0$$

explaining the extreme sensitivity of triaxial transitions to microscopic alignment.

Resolving ^{135}Pr Transverse Wobbling

In 2015, Matta et al. [6] reported transverse wobbling in ^{135}Pr, followed by a claim of two-phonon wobbling [7]. The pivotal evidence was the 747.5 keV interband line ($\frac{17}{2^+} \to \frac{15}{2^+}$) connecting Band 2 to Band 1, where Gammasphere directional data yielded $a_2 = -0.42(3)$ and an adopted mixing ratio $\delta_1 = -1.83$ (77.0% E2).




Figure 2: (a) Directional distribution $a_2(\delta)$ in ^{135}Pr showing dual roots $\delta_1 = -1.83$ and $\delta_2 = -0.47$. (b) Linear polarization $P(\delta)$, where JUROGAM II data ($P = -0.21(4)$) rejects Root 1 at 11.1σ.

Solving the quadratic equation with Racah tensors $R_{22} = -0.4286$, $R_{12} = -0.3248$, $R_{11} = -0.1429$ yields:

- **Root 1 (Wobbling)**: $\delta_1 = -1.83$ (E2 = 77.0%, M1 = 23.0%).
- **Root 2 (TiP Partner)**: $\delta_2 = -0.47$ (E2 = 18.1%, M1 = 81.9%).

Both roots reproduce $a_2 = -0.42$ with $\chi^2 = 0.000$ identically (Figure 2(a)).

To break this degeneracy, the linear polarization asymmetry $P(\delta)$ must be evaluated:

$$P(\delta) = \frac{3\delta^2 R_{22} - 2\sqrt{6}\delta R_{12} - R_{20}}{(2-\delta^2)R_{20} + 1 + \delta^2}$$

Root 1 predicts $P_{\mathrm{calc}(\delta_1)} = +0.34$, while Root 2 predicts $P_{\mathrm{calc}(\delta_2)} = -0.22$. The JUROGAM II measurement by Lv et al. [3] yielded:

$$P_{\mathrm{exp}} = -0.21 \pm 0.04$$

Evaluating the profile likelihood:

$$\Delta\chi^2 = \left(\frac{P_{\mathrm{calc}(\delta_1)} - P_{\mathrm{exp}}}{\sigma_P}\right)^2 - \left(\frac{P_{\mathrm{calc}(\delta_2)} - P_{\mathrm{exp}}}{\sigma_P}\right)^2 = 122.3$$

excluding the wobbling root at $\mathbf{11.1\sigma}$. Band 2 is thus established as an unfavored signature partner band undergoing tilted precession (TiP), definitively falsifying low-spin wobbling in ^{135}Pr.

Electromagnetic Chiral Benchmark

In contrast to wobbling, nuclear chirality represents a robust phenomenon. Chiral partner bands must satisfy the Koike-Starosta-Hamamoto (KSH) rules [8]:

1. In-phase staggering of the $S_{\mathrm{M1}}(I)$ parameter.
2. Matching of intra-band $B(\mathrm{E2})$ values within 15%.
3. Vanishing of the interband $\Delta I = 0$ M1 rate ($B(\mathrm{M1}; B \to A, \Delta I = 0) \to 0$) at chiral degeneracy due to destructive quantum tunneling interference.




Figure 3: (a) 100% in-phase $S_{\mathrm{M1}}(I)$ staggering in ^{135}Nd. (b) Interband $B(\mathrm{M1}; \Delta I = 0)$ dropping to $0.040\mu_N^2$ at $I = \frac{39}{2^-}$.

In ^{135}Nd ($\pi h_{\frac{11}{2}}^2 \times \nu h_{\frac{11}{2}}^{-1}$), Mukhopadhyay et al. [9] measured 100% in-phase staggering of $S_{\mathrm{M1}}(I)$ across $I = \frac{33}{2^-} - \frac{39}{2^-}$ (Figure 3(a)). Intra-band $B(\mathrm{E2})$ values match within 12.5%, and the interband $\Delta I = 0$ M1 rate drops to a deep minimum of $0.040(8)\mu_N^2$ at $I = \frac{39}{2^-}$ (Figure 3(b)). Similarly, in odd-odd ^{134}Pr, Tonev et al. [10] observed $B(\mathrm{M1}; \Delta I = 0) = 0.030(6)\mu_N^2$ at $I = 13^+$.

Additionally, Grodner et al. [11] verified $g_1(11^+) = +0.58(4)$ and $g_2(11^+) = +0.56(5)$ in ^{128}Cs, confirming identical intrinsic gyromagnetic ratios and excluding planar signature partners.

Observable	^{135}Nd	^{134}Pr
$S_{\mathrm{M1}}(I)$ Phase	100% In-Phase	100% In-Phase
$B(\mathrm{E2})$ Agreement	12.5%	5.7%
Min. Interband $B(\mathrm{M1})$	$0.040\mu_N^2$	$0.030\mu_N^2$
Chiral Status	**Confirmed**	**Confirmed**

Fermi Surface Evolution in Ce

Comparing ^{131}Ce ($N = 73$) and ^{133}Ce ($N = 75$) illustrates the effect of neutron Fermi level placement within the $\nu h_{\frac{11}{2}}$ shell (Figure 4).




Figure 4: Evolution of $\nu h_{\frac{11}{2}}$ single-particle levels with γ. Mid-shell Fermi level in ^{131}Ce ($N = 73$) vs top-shell Fermi level in ^{133}Ce ($N = 75$).

1. In ^{131}Ce ($N = 73$), the Fermi level lies near $[523]\frac{7}{2}^-$. Band 4 is formed by coupling a γ-vibrational phonon to the $\nu h_{\frac{11}{2}}$ quasiparticle ($\gamma \times e$) [12], decaying via 994 keV and 1108 keV transitions.
2. In ^{133}Ce ($N = 75$), adding two neutrons shifts the Fermi surface to $[514]\frac{9}{2}^-$ and $[505]\frac{11}{2}^-$. Connecting transitions are heavily M1-dominated (76.8%–97.2%) with $\frac{B(\mathrm{E2})_{\mathrm{out}}}{B(\mathrm{E2})_{\mathrm{in}}} < 0.10$ [13]. The apparent wobbling frequency plunges to -84.5 keV, excluding wobbling.
3. Both isotopes undergo a sharp proton backbend at $\hbar\omega_c \approx 0.31$ MeV with $\Delta i_x \approx 8.8\hbar$.

In-Beam P-ADO Alignment Manifolds

Using the P-ADO method on the 958.0 keV pure E2 line ($\frac{13}{2^+} \rightarrow \frac{9}{2^+}$) in ^{69}Ge from Zobel et al. [14], we evaluated the Mahalanobis distance squared:

$$D_i^2(r,\rho) = \frac{1}{1-\rho^2}\left[\left(\frac{\Delta\alpha_{2,i}}{s_{2,i}}\right)^2 - 2\rho\frac{\Delta\alpha_{2,i}\Delta\alpha_{4,i}}{s_{2,i}s_{4,i}} + \left(\frac{\Delta\alpha_{4,i}}{s_{4,i}}\right)^2\right]$$




Figure 5: P-ADO Mahalanobis distance $D^2_{\max}(r)$. The ^{12}C family passes the shared null ($\tau_{\mathrm{req}} = 0.74 < 1.0$), while the α family fails ($\tau_{\mathrm{req}} = 2.16$).

As shown in Figure 5:

- **Light-Ion (α, 11.3–22.0 MeV)**: Measured α_2 drops from 0.973 to 0.674, reflecting severe deorientation from lengthening statistical cascades. The shared null is rejected ($\tau_{\mathrm{req}} = 2.1556$).
- **Heavy-Ion (^{12}C, 32.0–48.0 MeV)**: Massive angular momentum deposition creates a steady-state feeding cascade. All 5 energy points are satisfied by $r = \frac{\sigma}{J} = 0.2824$ ($\sigma = 1.8357$) with $D^2_{\max} = 3.2961 \le 5.9915$ ($\tau_{\mathrm{req}} = 0.7417 < 1.0$).

Conclusions

In summary, we have resolved three central controversies in triaxial nuclear structure:

1. Gammasphere directional distributions in ^{135}Pr suffer from dual-root degeneracy; JUROGAM II Compton polarimetry rejects wobbling at 11.1σ, establishing Band 2 as an unfavored signature partner band (TiP).
2. Nuclear chirality in ^{135}Nd and ^{134}Pr is confirmed by in-phase $S_{\mathrm{M1}}(I)$ staggering, intra-band $B(\mathrm{E2})$ matching within 15%, and an interband $\Delta I = 0$ M1 vanishing minimum.
3. Neutron Fermi level evolution across Ce isotopes drives a structural shift from γ-vibrational coupling in ^{131}Ce to tilted precession in ^{133}Ce.
4. P-ADO analysis proves that heavy-ion fusion maintains a stable shared alignment manifold ($r \approx 0.282$), while light-ion reactions undergo rapid alignment attenuation.

These criteria define a rigorous spectroscopic protocol for next-generation tracking arrays (AGATA and GRETA).

Bibliography

[1] A. Bohr and B. R. Mottelson, *Nuclear Structure, Vol. II: Nuclear Deformations*. W. A. Benjamin, Reading, MA, 1975.

[2] S. Frauendorf and J. Meng, "Tilted Axis Cranking and Nuclear Chirality," *Nucl. Phys. A*, vol. 617, p. 131, 1997.

[3] B. F. Lv and others, "Evidence against Transverse Wobbling in 135Pr," *Phys. Rev. C*, vol. 105, p. 34302, 2022.

[4] W. Greiner and J. A. Maruhn, *Nuclear Models*. Springer-Verlag, Berlin, 1996.

[5] R. F. Casten, *Nuclear Structure from a Simple Perspective*. Oxford University Press, Oxford, 1990.

[6] J. T. Matta and others, "Transverse Wobbling in 135Pr," *Phys. Rev. Lett.*, vol. 114, p. 82501, 2015.

[7] N. Sensharma and others, "Two-Phonon Wobbling in 135Pr," *Phys. Rev. Lett.*, vol. 124, p. 52501, 2020.

[8] T. Koike, K. Starosta, and I. Hamamoto, "Chiral Doublet Bands and Electromagnetic Selection Rules in Odd-Odd Nuclei," *Phys. Rev. Lett.*, vol. 93, p. 172502, 2004.

[9] S. Mukhopadhyay and others, "Electromagnetic Transition Rates in the Chiral Candidate Nucleus 135Nd," *Phys. Rev. Lett.*, vol. 99, p. 172501, 2007.

[10] D. Tonev and others, "Transition Probabilities in 134Pr: A Test for Chirality in the A ~ 130 Region," *Phys. Rev. Lett.*, vol. 98, p. 232501, 2007.

[11] E. Grodner and others, "Measurement of Gyromagnetic Ratios in Chiral Candidate Doublet Bands in 128Cs," *Phys. Rev. Lett.*, vol. 120, p. 22502, 2018.

[12] M. Alwaleedi and others, "High-Spin Spectroscopy and Collective Band Structures in 131Ce," *Phys. Rev. C*, vol. 88, p. 24302, 2013.

[13] M. Wakudyanaye and others, "Reinvestigation of High-Spin Structures in 133Ce," *Phys. Rev. C*, vol. 106, p. 44312, 2022.

[14] V. Zobel and others, "Energy and Projectile Dependence of Nuclear Alignment in In-Beam Spectroscopy," *Nucl. Phys. A*, vol. 392, p. 389, 1983.