A coverage-calibrated audit of alignment transfer in oriented-state gamma-ray angular distributions: the public ^{67}Ga benchmark

Author information to be completed[1*]

[1*]Affiliation and collaboration information to be completed.

Corresponding author(s). E-mail(s): corresponding.author@to-be-completed;

Abstract

Projectile-alignment directional orientation (P-ADO) analyses can transfer an alignment proxy between reaction conditions when extracting multipole mixing ratios. We study that transfer with the public ^{67}**Ga 958.0**-keV $\mathbf{13/2^+ \to 9/2^+}$ pure-$\boldsymbol{E2}$ benchmark of Zobel et al. The ten published $\boldsymbol{(\alpha_2, \alpha_4)}$ pairs are analyzed under an explicitly independent-row Gaussian table contract with fixed quoted marginal errors and a scanned common row correlation $\boldsymbol{\rho}$. At $\boldsymbol{\rho = 0}$, the shared proxy gives $\boldsymbol{r = 0.2801}$; projectile-family and energy-trend alternatives do not improve the descriptive information criteria, while a separate global convex-mixture control is weakly identified and is not interpreted as a feeding fraction. The observed profile-preserving maximum-row discrepancy is **22.18**. Repeating the same profile operation in 1000 pseudo-experiments per true-$\boldsymbol{\rho}$ value gives empirical 95% quantiles of **8.47**–**8.79** for true $\boldsymbol{\rho = -0.6, 0, 0.6}$. The result is a conditional table-level calibration warning, rather than an experimental p-value or a general invalidation of P-ADO.

Keywords: nuclear alignment, P-ADO, angular distributions, covariance, coverage, feeding

1 Introduction

Angular distributions from oriented nuclear states are used to determine transition multipolarities and mixing ratios in in-beam gamma-ray spectroscopy. The measured coefficients are not determined by the transition operator alone: they also depend on the population of magnetic substates, the feeding history, detector geometry, and the treatment of the fitted angular-distribution covariance [1, 2]. This dependence becomes consequential when a mixing-ratio branch is later used as a collective-mode discriminator in the A$\approx$ 130 region, where wobbling, tilted precession, signature-partner bands, and gamma-soft particle-core coupling can produce overlapping energy fingerprints but different electromagnetic transition patterns [3–6].

The P-ADO approach is useful because a pure-$E2$ reference transition can provide an alignment anchor for a neighbouring transition. The corresponding shortcut is to treat the inferred $r = \sigma/J$ as transferable between reaction conditions. Earlier work already established the qualitative limits of that shortcut: alignment coefficients depend on the feeding history and a Gaussian population is a conditional one-parameter approximation [7, 8]; Zobel et al. subsequently showed that attenuation changes with projectile and beam energy and discussed the role of direct and cascade feeding [9]. The contribution sought here is narrower and quantitative: we provide a reproducible audit

workflow that asks when a shared alignment proxy is statistically compatible with published coefficient pairs under an explicit covariance contract.

We revisit the ten-row ^{67}Ga table as a controlled, reproducible calibration case study. The purpose is not to reanalyse the original event data, which are unavailable in the source package, and not to claim that the P-ADO method is invalid. The A$\approx$ 130 connection is therefore a downstream analysis gate: before a P-ADO branch is used to support a collective-mode assignment, its alignment transfer and statistical coverage should satisfy the tests defined here. We quantify three boundaries that are often hidden in a compact mixing-ratio analysis: the distinction between a shared and a condition-specific alignment response, the effect of an unknown row-internal α_2–α_4 covariance, and the coverage of a maximum-row discrepancy statistic.

2 Benchmark and evidence boundary

The benchmark is the 958.0-keV $13/2^+ \to 9/2^+$ pure-$E2$ transition in ^{67}Ga. The ten rows contain five α-induced conditions from 11.3 to 22 MeV and five ^{12}C-induced conditions from 32 to 48 MeV. The reactions are $^{64}\mathrm{Zn}(\alpha, p\gamma)^{67}\mathrm{Ga}$ and $^{57}\mathrm{Fe}(^{12}\mathrm{C}, pn\gamma)^{67}\mathrm{Ga}$, respectively [9]. The values used here are the published fit coefficients and quoted statistical uncertainties transcribed from Table 1; they are not raw event counts.

The earlier Zobel study contains related ^{67}Ga values and is retained as a dependent source lineage rather than merged as an independent experiment [7]. We use the 1983 Table 1 as the canonical fixture because its reaction labels and locator are explicit; resolving whether the two publications share acquisition or fitting inputs is outside the present table-level analysis.

The source does not supply the row-internal covariance between α_2 and α_4, nor does it supply the complete response matrices or event-level feeding information. Its discussion of the α_2–α_4 plane mentions error ellipses but does not print their numerical parameters (printed p.393/PDF p.5). We therefore treat the correlation ρ as a sensitivity parameter rather than as a measured quantity. The declared contract is block-diagonal, $\Sigma_{\mathrm{table}} = \mathrm{blockdiag}(\Sigma_1, \ldots, \Sigma_{10})$: rows are treated as independent, the quoted $s_{2,i}$ and $s_{4,i}$ are treated as fixed marginal standard errors, and one common ρ is scanned. These are analysis assumptions, not source measurements.

3 Models and test statistics

For a state of spin J, a Gaussian magnetic-substate population is written

$$P(m \mid r) \propto \exp\left[-\frac{m^2}{2(Jr)^2}\right], \qquad r = \frac{\sigma}{J}. \tag{1}$$

The predicted vector $\boldsymbol{\mu}(r) = (\alpha_2(r), \alpha_4(r))$ is obtained by averaging the normalized Yamazaki tensor polynomials over $P(m \mid r)$. For each table row i, the covariance contract is

$$\Sigma_i(\rho) = \begin{pmatrix} s_{2,i}^2 & \rho s_{2,i} s_{4,i} \\ \rho s_{2,i} s_{4,i} & s_{4,i}^2 \end{pmatrix}, \tag{2}$$

where $s_{2,i}$ and $s_{4,i}$ are the quoted marginal statistical errors. The finite-J response map is evaluated at $J = 13/2$, with the Gaussian width proxy restricted to $0.02 \le r \le 1$. This map is a model calculation under the Yamazaki convention; it is not a direct inversion of an independently measured alignment tensor. An independent attenuation-table replay is retained in the machine-readable response-map validation output; it checks the finite-J Gaussian convention but does not validate the Zobel detector response, event counts, covariance or feeding history.

We compare three response descriptions sharing the common submodel:

$$H_{\mathrm{shared}} : r_i = r_0, \tag{3}$$

$$H_{\mathrm{family}} : r_i = r_\alpha \text{ or } r_{^{12}\mathrm{C}}, \tag{4}$$

$$H_{\mathrm{energy}} : r_i = r_0 + r_1(E_i - \bar{E})/10. \tag{5}$$

The global convex-mixture control is analyzed separately as a sensitivity model; it uses one common mixture prediction for all rows and has no condition-specific feeding weights. The Gaussian profile objective is

$$-2\log L = \sum_i \Big[(\mathbf{y}_i - \boldsymbol{\mu}_i)^{\mathsf{T}} \Sigma_i^{-1} (\mathbf{y}_i - \boldsymbol{\mu}_i) + \log\det\Sigma_i + 2\log(2\pi)\Big]. \tag{6}$$

For future event-level work, the count likelihood should be written in a Poisson form rather than treating a background-subtracted spectrum as a set of independent Gaussian observations; asymptotic likelihood-ratio thresholds also require coverage checks when nuisance parameters are weakly identified. As a complementary discrepancy diagnostic, we calculate

$$D^2_{\max} = \max_i (\mathbf{y}_i - \boldsymbol{\mu}_i)^\mathsf{T} \Sigma_i^{-1} (\mathbf{y}_i - \boldsymbol{\mu}_i) \qquad (7)$$

and report $\tau_{\max} = \sqrt{D^2_{\max}/q^{(2)}_{0.95}}$, with $q^{(2)}_{0.95} = 5.9915$. This threshold is deliberately reported as a declared two-dimensional reference contract, not as a calibrated ten-row family-wise p-value. The profiled discrepancy used for the covariance study is a separate statistic,

$$T_{\text{prof}}(\mathbf{y}) = \min_{\rho \in \mathcal{R}} \max_i d_i^2(\widehat{r}_\rho, \rho)\,, \qquad (8)$$

where $\widehat{r}_\rho$ minimizes the shared-model Gaussian objective at fixed ρ. It is not a profile-likelihood-ratio statistic; the same r fit, ρ scan and row maximum are repeated in each pseudo-experiment.

4 Results

4.1 Shared and extended alignment profiles

At $\rho = 0$, the shared model gives $r_0 = 0.2801$. The family model gives $r_\alpha = 0.2887$ and $r_{^{12}\mathrm{C}} = 0.2758$, while the energy model gives $r_0 = 0.2785$ and $r_1 = 0.0053$ per 10 MeV. The energy slope is descriptive only: the projectile families occupy disjoint energy ranges in this table. A global convex-mixture control lowers the descriptive objective, but its component parameters are weakly identified and are not interpreted as direct- or cascade-feeding fractions. The extra response parameters do not improve AIC or BIC under the fixed $\rho = 0$ contract. Thus the table does not identify a unique condition-specific or feeding-specific replacement model.

The shared fit gives $\tau_{\max} = 2.28$ at $\rho = 0$. The statistic used for the covariance sensitivity is instead the minimum over the declared ρ grid of the fitted maximum-row discrepancy; the corresponding observed $D^2_{\max}$ minimum is reported below and is calibrated with the same profiling operation in a separate toy study.

4.2 Covariance sensitivity

Figure 1 shows the $\tau_{\max}$ profile over the declared sensitivity range $-0.8 \le \rho \le 0.8$. The shared model remains above unity throughout this finite grid. The energy model has the smallest maximum discrepancy over much of the interval, whereas the family model does not dominate the profile. This is not evidence for a unique energy law; it shows that the table-level data produce a large discrepancy for a single response under the declared contract, while the alternative model form remains unresolved.

4.3 Coverage of the maximum-row threshold

We generated 1000 pseudo-experiments from the shared model with $r = 0.28$, $\rho = 0$, and the row-specific quoted marginal errors. In each pseudo-experiment, r was refitted and the maximum row-wise D^2 was calculated. The single-row two-dimensional threshold $q^{(2)}_{0.95} = 5.9915$ was satisfied in 65.4% of pseudo-experiments (Monte Carlo standard error 0.015). The empirical 95th percentile of the maximum statistic was 10.01, rather than 5.99.

This result does not by itself calibrate the profiled statistic used for the observed table. It changes the interpretation of a nominal threshold: applying a single-row 95% contour to the maximum over ten rows is not a 95% family-wise test. A publication-level rejection or acceptance statement requires a predeclared family-wise statistic, an event-level likelihood, or a coverage-calibrated test.

4.4 Profile-preserving calibration

To calibrate the statistic used for the observed covariance sensitivity, we repeated the full ρ-profile and maximum-row operation in each pseudo-experiment. With 1000 pseudo-experiments for each true $\rho \in \{-0.6, 0, 0.6\}$, the empirical 95th quantiles of the profiled statistic were 8.79, 8.56, and 8.47, respectively. Bootstrap Monte Carlo 95% intervals for these quantiles were recorded in the machine-readable output. The observed profiled value is 22.18 at $\rho = 0.64$.

Table 1 Table-level fits at $\rho = 0$. $D^2_{\max}$ is the largest row-wise discrepancy. BIC uses the ten independent row vectors as its sample units.

Model	Fitted response	AIC	BIC	$D^2_{\max}$
Shared	$r_0 = 0.2801$	32.04	32.34	31.26
Family	$r_\alpha = 0.2887;\ r_{12C} = 0.2758$	32.63	33.23	36.24
Energy	$r_0 = 0.2785;\ r_1 = 0.0053$	32.63	33.24	26.96
Mixture	$r_d = 0.1952;\ r_c = 0.3659;\ w = 0.5304$	33.56	34.47	30.71




Fig. 1 (a) Published α_2 values and quoted marginal errors for the two projectile families, with the three response models and the separate mixture control. (b) Maximum-row discrepancy scale $\tau_{\max}$ as a function of the assumed common row correlation ρ. (c) Quantiles of the maximum-row D^2 statistic in shared-null pseudo-experiments.

This comparison is still conditional on the independent-row Gaussian contract and the finite profile grid; it is not an experimental p-value or a supremum calibration over all covariance structures.

As a descriptive influence control, the same profiled statistic was recomputed after removing each row in turn. The leave-one-row-out values range from 15.81 to 24.44, so the observed discrepancy is not numerically produced by a single row in this fixture. Because the number of rows changes, these values are not assigned a separate calibrated null distribution.

4.5 Coverage sensitivity to the true alignment and covariance

We extended the fixed-ρ sensitivity study over five true values of r from 0.10 to 0.60 and seven true row correlations from -0.8 to 0.8, with 1000 pseudo-experiments per cell. When the fitted covariance matched the generating covariance, the empirical coverage of the single-row threshold was approximately 0.62–0.68. When the data were generated with a nonzero correlation but the fit forced $\rho = 0$, the coverage fell as low as 0.47 at the edge of the scanned negative-correlation range and was also below 0.53 in several other cells. The nominal threshold is therefore not stable under either the ten-row maximum operation or modest covariance misspecification.

For comparison, the smallest observed shared-model $D^2_{\max}$ across the profile grid $-0.8 \le \rho \le 0.8$ with spacing $\Delta\rho = 0.02$ is 22.18, at $\rho = 0.64$. The fixed-ρ toy 95th percentiles range from about 9.60 to 10.59 across the expanded sensitivity grid. This is a finite-grid sensitivity result; the profile-preserving calibration above is the relevant calibration for the observed profiled statistic.

5 Implications for A≈ 130 collective-mode spectroscopy

The benchmark has a direct methodological consequence for A≈ 130 studies. ADO, DCO, polarization, and mixing-ratio analyses often combine




Fig. 2 Coverage of the single-row two-dimensional threshold for the maximum over ten rows in a 5×7 fixed-ρ toy grid. The left panel uses the generating correlation in the fit; the right panel forces $\rho = 0$ while varying the generating correlation. Values are empirical fractions from 1000 toys per cell.

information from different reaction conditions or inherit an alignment width from a reference transition. Zobel's experiment provides the physical reason to question that transfer: direct feeding and cascade feeding can change with projectile and beam energy [9]. The present calculation adds a quantitative layer by showing that the shared-response assumption can fail at the level of a multi-row observable vector, while a naive single-row threshold can have poor family-wise coverage.

The result does not select wobbling, tilted precession, chirality, or signature-partner bands in any particular A≈ 130 nucleus. It supports a pre-analysis diagnostic: a mixing-ratio branch used in collective-mode interpretation should retain condition-specific alignment nuisance, the α_2–α_4 covariance, and a response or feeding control. Independent polarization or conversion-electron information remains necessary when the angular-distribution map has two real mixing-ratio roots [1, 3, 5].

6 Limitations and next tests

The missing row covariance is the dominant limitation. The table contains fitted coefficients, not raw spectra, so the present analysis cannot recover the covariance from first principles. The Gaussian population is also only one response model; a direct-plus-cascade mixture could explain part of the row-to-row variation, but the global convex mixture used here has no condition-specific feeding weights and does not validate a feeding mechanism. The energy-dependent and family models are descriptive alternatives, and the energy trend is confounded with projectile family in this fixture.

The next decisive test is a blind, event-level analysis with at least one pure-$E2$ calibration line and several mixed transitions from the same acquisition. It should use the raw angular counts, on/off or sideband background terms, detector response, and a holdout transition. Pseudo-experiment coverage must be checked before a likelihood-ratio threshold is interpreted as a mode-selection significance. The present grid is a calibration sensitivity study, not a substitute for that event-level test.

7 Conclusions

The public ^{67}Ga P-ADO table produces a large maximum-row discrepancy for a single shared alignment response under the stated table-level contract. The data do not identify whether a family-specific or energy-dependent response is the correct replacement, and the nominal two-dimensional single-row threshold is not calibrated for the maximum over ten rows. The transferable result is therefore a response and coverage requirement for mixing-ratio analyses that are later used to identify A≈ 130 collective modes.

Data availability

The normalized ten-row benchmark table and all derived analysis outputs are included in this local manuscript package. A public release should deposit the normalized CSV, scripts, JSON outputs and a version identifier in a repository with a persistent archive record. The source paper and its exact locator are recorded in the local evidence record ‘knowledge/sources/zobel-1983-energy-projectile-alignment.md‘; the publisher PDF and raw event counts are not redistributed here. The source does not provide row-internal or cross-row covariance, and no unavailable raw data are claimed to have been reconstructed.

Code availability

The profile, fixed-ρ coverage, profile-preserving calibration and coverage-grid scripts are included under ‘analysis/‘ in this manuscript package. They use Python, NumPy and SciPy; the fixed seeds, covariance contract and BIC row-unit convention are recorded in the JSON results and README. The response-map validation output records the independent finite-J table replay. The outputs are table-level analyses and are not an event-level detector-response reanalysis. Generative AI assistance was used locally for literature triage, code review, manuscript consistency checks and language editing; the accountable authors must verify all claims, locators, declarations, permissions and the final submission version. Author names, affiliations, corresponding-author details, funding, conflicts and contribution roles remain to be completed by the author group.

References

[1] Hamilton, W.D.: The Electromagnetic Interaction in Nuclear Spectroscopy. North-Holland, Amsterdam (1975)

[2] Yamazaki, T.: Tables of coefficients for angular distributions of gamma rays from aligned nuclei. Nuclear Data, Section A **3**, 1–23 (1967) https://doi.org/10.1016/S0550-306X(67)80002-8

[3] Matta, J.T., *et al.*: Transverse wobbling in ^{135}Pr. Physical Review Letters **114**, 082501 (2015) https://doi.org/10.1103/PhysRevLett.114.082501

[4] Lawrie, E.A., Shirinda, O., Petrache, C.M.: Tilted precession and wobbling in triaxial nuclei. Physical Review C **101**, 034306 (2020) https://doi.org/10.1103/PhysRevC.101.034306

[5] Lv, B.-F., *et al.*: Evidence against the wobbling nature of low-spin bands in ^{135}Pr. Physics Letters B **824**, 136840 (2022) https://doi.org/10.1016/j.physletb.2021.136840

[6] Nomura, K., Petrache, C.M.: Questioning the wobbling interpretation of low-spin bands in γ-soft nuclei within the interacting boson-fermion model. Physical Review C **105**, 024320 (2022) https://doi.org/10.1103/PhysRevC.105.024320

[7] Zobel, V., Cleemann, L., Eberth, J., Hellmeister, H.P., Neumann, W., Wiehl, N.: The shape of magnetic substate distributions in fusion-evaporation reactions. Nuclear Instruments and Methods **171**, 223–231 (1980) https://doi.org/10.1016/0029-554X(80)90494-2

[8] Ionescu, V.A., Kern, J., Nordmann, C., Olbrich, S., Reichart, W.: Improved analysis of gamma-ray angular distributions in (particle, xn) experiments. Nuclear Instruments and Methods in Physics Research **190**(1), 19–27 (1981) https://doi.org/10.1016/0029-554X(81)90200-7

[9] Zobel, V., Cleemann, L., Eberth, J., Heck, T., Huesing, K., Neumann, W., Nolte, M.: Energy and projectile dependence of nuclear alignment in fusion-evaporation reactions. Nuclear Instruments and Methods in Physics Research **207**, 389–394 (1983) https://doi.org/10.1016/0167-5087(83)90649-X