Same-parent consistency checks for the 365-keV transition in ^{129}Ba

Author information to be completed[1*]

[1*]Affiliation and collaboration information to be completed.

Corresponding author(s). E-mail(s): corresponding.author@to-be-completed;

Abstract

The mixing ratio of the 365-keV transition in **^{129}Ba** has been reconsidered in a proposed wobbling interpretation. We examine what its same-parent 461.6-keV companion can add to that inference. Specializing standard angular-distribution theory gives $\boldsymbol{A_{4,t} = -(8/3) f_{E2} s_4 A_{4,r}}$, where $\boldsymbol{f_{E2} = \delta^2/(1+\delta^2)}$ and $\boldsymbol{s_4}$ is a finite-response ratio. This relation constrains a product; a calibrated response is required to infer the magnitude of $\boldsymbol{\delta}$. The evaluated central coefficients require a negative product, while a rank-2 check also has a central mismatch for equal responses. These observations motivate a fit in coefficient space rather than a division by a weak reference coefficient. Under a declared diagonal Gaussian covariance, the profiled distances for the small- and large-magnitude domains differ by only 0.0136. Their ordering changes within a specified covariance sensitivity family. A provisional error-scale comparison using newly obtained primary-source OCR also removes a previously apparent two-margin interval mismatch. The analysis identifies an unresolved companion-line consistency question, without assigning a new mixing ratio or a collective mode.

Keywords: gamma-ray angular distribution, multipole mixing ratio, nuclear alignment, ^{129}Ba

1 Physical question and existing method

The electromagnetic character of interband transitions is important when testing wobbling interpretations in the $A \approx 130$ region. A quasiparticle–triaxial-rotor calculation relates spin-dependent energies to electromagnetic matrix elements, but an energy trend alone does not fix the transition operator or the experimental mixing ratio [1]. In ^{129}Ba, Chakraborty et al. reexamined legacy angular-distribution coefficients and reported a minimum near $\delta = -2$, with an approximately 80(4)% radiative $E2$ fraction for the 365-keV line. They proposed a wobbling interpretation and requested further measurements [2].

A useful reference already appears in the level scheme: the same $13/2^-$ parent decays through the 364.7-keV mixed $M1/E2$ line and a 461.6-keV line assigned $E2$ [3]. Can this companion independently constrain the branch choice? Using a pure transition from the same state to constrain alignment is an established technique, explicitly discussed by Diamond et al. [4]. The statistical-tensor and radiation-coefficient formalism is also standard [5, 6]. We make no claim to originate either principle. The present study applies them to a specific legacy-data consistency problem and examines the dependence on response and uncertainty assumptions.

A single rank-2 angular coefficient can admit two mixing-ratio magnitudes. A reliable rank-4 coefficient can distinguish them when alignment and response are known. With unknown alignment, even the two coefficients need not identify a unique magnitude. A same-parent reference is informative only if it samples the same selected parent ensemble and its multipolarity and response are established. These conditions are the subject of this study, rather than consequences of a level-scheme label.

2 Source identity and numerical inputs

We retain the evaluated ENSDF input in Table 1. The selected reaction dataset belongs to the 2014 $A = 129$ evaluation [3]; it attributes the paired angular coefficients to Gizon et al. [7]. The two records share the 643.6-keV parent. The rounded 365-keV label is not used to match unrelated transitions.

The publisher's full-text API supplies OCR covering Gizon et al., pp.464–476. It describes angular-distribution measurements at seven angles and gives Table 1 intensities at 55° to the beam [7]. Its discussion of the 365.0- and 461.8-keV branches supports the same-parent identity, with the energies subsequently revised in the evaluation. The accompanying PDF access provided only the first page. The Table 1 scan has therefore not been visually checked, and the damaged OCR cannot certify missing signs or every table column.

The OCR indicates error scales of 0.05 for the target A_2 and 0.04/0.06 for the reference A_2/A_4, differing from the evaluated values. We preserve this as a version question and use those scales only in a labeled sensitivity scenario. We have not established whether the difference reflects later revision, transcription or OCR. The 1978 polarization article remains unavailable in full in the present archive. No raw angular yields or covariance have been recovered, and no combination of the two source versions is treated as a new measurement.

3 Same-parent angular constraints

For an axially aligned ensemble and a response that permits scalar angular-rank folding, write

$$A_{k,x} = \rho_{k,x} Q_{k,x} F_{k,x}(\delta_x), \qquad x = t, r. \tag{1}$$

Here ρ_k describes nuclear alignment, Q_k the finite response, and F_k the radiation coefficient in a specified phase convention. Shared selected-parent populations imply $\rho_{k,t} = \rho_{k,r}$. Different gates or time selections may violate this equality even when the state identity agrees.

For an $M1/E2$ target, the rank-4 $M1/M1$ and interference terms vanish by angular-momentum coupling. A pure-$E2$ reference consequently gives

$$\begin{aligned} A_{4,t} &= -\frac{8}{3} f_{E2} s_4 A_{4,r}, \\ f_{E2} &= \frac{\delta^2}{1+\delta^2}, \qquad s_4 = \frac{Q_{4,t}}{Q_{4,r}}. \end{aligned} \tag{2}$$

The signed radiation factors for this spin sequence are

$$F_{4,t} = \frac{32\sqrt{2431}}{3003}, \qquad F_{4,r} = -\frac{4\sqrt{2431}}{1001}. \tag{3}$$

Their ratio is $-8/3$. An exact factorial calculation and an independent standard Wigner-symbol calculation agree. Direct photon-amplitude projections for all 14 parent magnetic substates also verify the cancellation for common diagonal populations; no Gaussian alignment assumption is needed.

Equation (2) identifies only $f_{E2}s_4$ when s_4 is unknown. With a known nonzero s_4, the allowed f_{E2} determines $|\delta|$, but $f_{E2}(\delta) = f_{E2}(-\delta)$ leaves the sign undetermined. If the reference coefficient vanishes, ratio inversion is undefined. A known negative s_4 is mathematically usable; nonnegative detector efficiency does not by itself guarantee a positive rank-4 attenuation factor.

For different selected-parent populations a separate factor $g_4 = \rho_{4,t}/\rho_{4,r}$ remains in Eq. (2). It is not an instrumental Q_4. Reference multipole contamination changes the radiation coefficient, while a general angular response can mix ranks and require a fuller treatment. These effects cannot all be absorbed into a negative s_4.

Table 1 Canonical evaluated input. Energies are in keV; quoted margins are preserved. The bottom row gives an explicitly provisional sensitivity scenario, with the evaluated means and signs retained. It is not a replacement measurement.

Line	$J_i \to J_f$	E_γ	A_2	A_4
Mixed target	$13/2^- \to 11/2^-$	364.7	$-0.70(3)$	$+0.16(5)$
$E2$ reference candidate	$13/2^- \to 9/2^-$	461.6	$+0.22(2)$	$+0.08(5)$
Provisional OCR error scales, target/reference			0.05/0.04	0.05/0.06

The rank-2 companion relation supplies a second check on the same observations. In the declared phase convention,

$$\frac{A_{2,t}}{A_{2,r}} = s_2 q(\delta), \qquad q(\delta) = \frac{-7 + 14\sqrt{11}\delta - \delta^2}{10(1+\delta^2)}. \tag{4}$$

Its two-dimensional quadratic-form eigenvalues give the exact range

$$\frac{-2-\sqrt{137}}{5} \le q(\delta) \le \frac{-2+\sqrt{137}}{5}. \tag{5}$$

The central ratio $-0.70/0.22 = -3.1818$ is below the lower limit -2.74094 for $s_2 = 1$. Allowing a positive s_2 requires at least 1.16085 at these central values. This amplitude check needs a response magnitude, whereas the rank-4 opposite-sign condition needs only $s_4 > 0$. Neither is a statistical rejection of noisy data.

4 Profiling the original coefficients

The reference A_4 is only 1.6 times its evaluated margin. Rather than approximate its ratio as Gaussian, we work with $y = (A_{2,t}, A_{4,t}, A_{2,r}, A_{4,r})^T$. For fixed δ, s_2, s_4, let $c_2 = s_2 q(\delta)$, $c_4 = -\frac{8}{3} s_4 f_{E2}$ and

$$B_\delta = \begin{pmatrix} c_2 & 0 \\ 0 & c_4 \\ 1 & 0 \\ 0 & 1 \end{pmatrix}. \tag{6}$$

The unknown true reference coefficients form $b = (b_2, b_4)^T$. For a fixed positive-definite Gaussian covariance V, the profiled residual distance is

$$D(\delta) = \min_b (y - B_\delta b)^T V^{-1} (y - B_\delta b) \tag{7}$$

$$= (L_\delta y)^T (L_\delta V L_\delta^T)^{-1} (L_\delta y), \tag{8}$$

$$L_\delta = \begin{pmatrix} 1 & 0 & -c_2 & 0 \\ 0 & 1 & 0 & -c_4 \end{pmatrix}. \tag{9}$$

This is Gaussian nuisance profiling with fixed V, not marginalization; no determinant of the projected covariance is added. Allowing any real b relaxes physical population-positivity constraints, so a fit does not certify a realizable alignment distribution.

We use $s_2 = s_4 = 1$ as a diagnostic scenario. With the four quoted margins collected in S, the covariance sensitivity family is

$$V = S\,[R(\rho_c) \otimes R(\rho_w)]\,S, \qquad R(\rho) = \begin{pmatrix} 1 & \rho \\ \rho & 1 \end{pmatrix}. \tag{10}$$

Here ρ_w couples angular ranks within each line and ρ_c couples lines. We scan each over $-0.8, -0.4, 0, 0.4, 0.8$; these are chosen correlations, not measured values or an exhaustive covariance model. Minimization uses $\theta = \arctan\delta$, with the pure-$E2$ endpoints included. Small- and large-magnitude domains share the $|\delta| = 1$ boundary in their closures. Distances and contrasts receive no p-value, confidence-interval or branch-probability interpretation.

5 Results and sensitivity

The rank-4 central coefficients give $f_{E2} s_4 = -0.75$. For $s_4 > 0$, their same signs cannot be exact means under the shared-parent and pure-reference assumptions. Their one-margin boxes are also both positive. However, the illustrative two-margin comparison at $s_4 = 1$ is fragile: the evaluated reference box maps to an upper target value 0.0533, just below the target lower value 0.06. Raising s_4 to 1.125 permits contact. Using the provisional OCR reference margin 0.06 gives an upper image of 0.1067 and hence overlap. These are literal marginal boxes, not joint confidence regions.

Table 2 Conditional diagonal-covariance results with $s_2 = s_4 = 1$. The minimizers describe the diagnostic; they are not revised experimental mixing ratios.

Quantity	Evaluated	OCR scales
δ at minimum	-1.0270	-1.2412
$D_{\min}$	12.8797	8.2357
$D_{\text{small}} - D_{\text{large}}$	0.0136	0.4685

The joint coefficient calculation in Table 2 does not provide a stable large-versus-small distinction. For evaluated margins and $\rho_c = 0$, varying ρ_w from -0.8 to $+0.8$ changes $D_{\text{small}} - D_{\text{large}}$ from $+3.530$ to -1.931. The ordering thus reverses inside this specified family. The provisional OCR-scale calculation changes the distances again (Fig. 1). None of these assumed covariances has been established for the source experiment. The calculation demonstrates dependence on missing inputs, rather than measuring the correct covariance or reproducing the 2024 fit.

Numerical checks compare Eq. (8) with a direct whitened least-squares fit, agreeing within 1.2×10^{-12}. Doubling the angular search grid from 1025 to 2049 points before refinement changes the diagonal minima by less than 9×10^{-15}. An isotropic zero-reference control has no δ sensitivity, as required. These checks establish implementation consistency, not experimental coverage. Coverage would require repeated sampling under a justified observation model.

6 Implications

The same-parent line is useful because it adds a falsifiable consistency requirement to a mixing-ratio analysis. It has not yet supplied a new experimental root. Resolving the ^{129}Ba case requires a visual check of the primary coefficient table and its later revisions, the response ratio, reference purity, and the joint fit or calibration covariance. The primary source's discussion of single-spectrum angular measurements narrows the experimental question; speculative gate differences cannot substitute for evidence about that acquisition.

The 2024 paper reports a joint angular-coefficient minimum. Our additional companion-reference assumptions are not a reconstruction of its fitting procedure, and the present results do not establish why its branch differs from historical values. Electromagnetic fractions also do not fix absolute collectivity: lifetime and branching information would still be required for an absolute $B(E2)$ test. The analysis therefore leaves wobbling, signature-partner and tilted-precession interpretations to comparisons with complete electromagnetic evidence.

7 Conclusions

Specializing established angular-distribution theory gives exact same-parent rank-4 and rank-2 consistency conditions for the ^{129}Ba 365/462-keV pair. Their evaluated central values raise a reference-transfer question, but weak reference precision, unverified response ratios and a source-version discrepancy prevent a unique inference. Coefficient-space profiling makes that limitation quantitative: the small- and large-magnitude domains are nearly coincident under diagonal covariance, and their ordering changes in a declared correlation scan. The useful outcome is a specific companion-line and covariance validation target, rather than a new collective-mode assignment. The originality and empirical sufficiency of this application remain to be established before journal submission.

Data and code availability

The draft includes the normalized evaluated coefficients, a separate input-contract record, exact angular-momentum calculations, coefficient-space profiling and figure-generation code. The publisher OCR and first-page PDFs are locally archived for source checking and are excluded from redistribution. Permanent public repository and license information remain to be completed.

Declarations

Author identities, contributions, affiliations, funding and competing-interest statements remain to be completed. AI-assisted tools were used to develop this draft, code and consistency checks; final scientific verification and responsibility remain with the eventual authors. This document is a research draft and has not passed final claim-specific review.




Fig. 1 Declared two-rank coefficient-space diagnostics. (a) Profile distance above each diagonal-covariance minimum; the dashed line is $|\delta| = 1$. (b) Small-minus-large domain distance along the chosen $\rho_c = 0$ slice. The two curves retain the same evaluated means and signs but use different error scales. No line represents a statistical rejection threshold.

References

[1] Frauendorf, S., Dönau, F.: Transverse wobbling: A collective mode in odd-A triaxial nuclei. Physical Review C **89**, 014322 (2014) https://doi.org/10.1103/PhysRevC.89.014322

[2] Chakraborty, S., Bhattacharyya, S., Mukherjee, G., Majumder, C.: Exploring the possibility of wobbling motion in ^{129}Ba. Physical Review C **110**, 024324 (2024) https://doi.org/10.1103/PhysRevC.110.024324

[3] Timar, J., Elekes, Z., Singh, B.: Nuclear data sheets for A=129. Nuclear Data Sheets **121**, 143–394 (2014) https://doi.org/10.1016/j.nds.2014.09.002

[4] Diamond, R.M., Matthias, E., Newton, J.O., Stephens, F.S.: Nuclear alignment in heavy-ion reactions. Physical Review Letters **16**, 1205–1207 (1966) https://doi.org/10.1103/PhysRevLett.16.1205

[5] Yamazaki, T.: Tables of coefficients for angular distributions of gamma rays from aligned nuclei. Nuclear Data, Section A **3**, 1–23 (1967) https://doi.org/10.1016/S0550-306X(67)80002-8

[6] Der Mateosian, E., Sunyar, A.W.: Tables of attenuation coefficients for angular distribution of gamma rays from partially aligned nuclei. Atomic Data and Nuclear Data Tables **13**, 391–406 (1974) https://doi.org/10.1016/0092-640X(74)90007-2

[7] Gizon, J., Gizon, A., Meyer-ter-Vehn, J.: Triaxial shape and onset of high-spin bands in ^{129}Ba. Nuclear Physics A **277**(3), 464–476 (1977) https://doi.org/10.1016/0375-9474(77)90714-X