

# The Gauss-Bonnet Theorem

Chen Lin

University of Chicago

October 7, 2021

# Statement of the theorem


## Theorem (Gauss-Bonnet Theorem)

If  $R$  is a region on a surface  $S$ , then

# Statement of the theorem

## Theorem (Gauss-Bonnet Theorem)

If  $R$  is a region on a surface  $S$ , then




# Statement of the theorem

## Theorem (Gauss-Bonnet Theorem)

If  $R$  is a region on a surface  $S$ , then

$$\int_{\partial R} k_g ds + \iint_R K dA + \sum_{i=1}^p \theta_i = 2\pi\chi(R).$$




# Statement of the theorem

## Theorem (Gauss-Bonnet Theorem)

If  $R$  is a region on a surface  $S$ , then

$$\int_{\partial R} k_g ds + \iint_R K dA + \sum_{i=1}^p \theta_i = 2\pi\chi(R).$$

- $k_g$  is geodesic curvature on  $\partial R$




# Statement of the theorem

## Theorem (Gauss-Bonnet Theorem)

If  $R$  is a region on a surface  $S$ , then

$$\int_{\partial R} k_g ds + \iint_R K dA + \sum_{i=1}^p \theta_i = 2\pi\chi(R).$$




- $k_g$  is geodesic curvature on  $\partial R$
- $K$  is Gaussian curvature of  $S$

# Statement of the theorem

## Theorem (Gauss-Bonnet Theorem)

If  $R$  is a region on a surface  $S$ , then

$$\int_{\partial R} k_g ds + \iint_R K dA + \sum_{i=1}^p \theta_i = 2\pi\chi(R).$$




- $k_g$  is geodesic curvature on  $\partial R$
- $K$  is Gaussian curvature of  $S$
- $\theta_1, \dots, \theta_p$  are the exterior angles of  $R$

# Statement of the theorem

## Theorem (Gauss-Bonnet Theorem)

If  $R$  is a region on a surface  $S$ , then

$$\int_{\partial R} k_g ds + \iint_R K dA + \sum_{i=1}^p \theta_i = 2\pi\chi(R).$$



- $k_g$  is geodesic curvature on  $\partial R$
- $K$  is Gaussian curvature of  $S$
- $\theta_1, \dots, \theta_p$  are the exterior angles of  $R$
- $\chi(R)$  is the Euler characteristic of  $R$

Why is it awesome?




# Statement of the theorem

## Theorem (Gauss-Bonnet Theorem)

If  $R$  is a region on a surface  $S$ , then

$$\int_{\partial R} k_g ds + \iint_R K dA + \sum_{i=1}^p \theta_i = 2\pi\chi(R).$$



- $k_g$  is geodesic curvature on  $\partial R$
- $K$  is Gaussian curvature of  $S$
- $\theta_1, \dots, \theta_p$  are the exterior angles of  $R$
- $\chi(R)$  is the Euler characteristic of  $R$

Why is it awesome?

- $k_g$ ,  $K$ , and  $\theta_i$  are geometric, but  $\chi(R)$  is topological!

# Outline

# Outline

## ① The geometry

# Outline

- ① The geometry
- ② The topology

# Outline

- ① The geometry
- ② The topology
- ③ Sketch of proof

# Outline

- ① The geometry
- ② The topology
- ③ Sketch of proof
- ④ Some corollaries

# What is a surface?

# What is a surface?



$S \subset \mathbb{R}^3$  is a **surface** if it is 'smooth' and looks flat locally.



# What is a surface?

$S \subset \mathbb{R}^3$  is a **surface** if it is 'smooth' and looks flat locally.



- Tangent planes exist at each point



# What is a surface?

$S \subset \mathbb{R}^3$  is a **surface** if it is 'smooth' and looks flat locally.

- Tangent planes exist at each point
- Identify the tangent plane with a unit normal vector




# What is curvature?

# What is curvature?

**From calculus:** for a path  $\alpha : I \rightarrow \mathbb{R}^3$  with  $\|\alpha'\| \equiv 1$ , curvature is  $\|\alpha''\|$ .


# What is curvature?

**From calculus:** for a path  $\alpha : I \rightarrow \mathbb{R}^3$  with  $\|\alpha'\| \equiv 1$ , curvature is  $\|\alpha''\|$ .



# What is curvature?


**From calculus:** for a path  $\alpha : I \rightarrow \mathbb{R}^3$  with  $\|\alpha'\| \equiv 1$ , curvature is  $\|\alpha''\|$ .



What does curvature mean for a surface?

# What is curvature?

**From calculus:** for a path  $\alpha : I \rightarrow \mathbb{R}^3$  with  $\|\alpha'\| \equiv 1$ , curvature is  $\|\alpha''\|$ .



What does curvature mean for a surface?


- We'll start with paths on surfaces

# What is curvature? (continued)



## What is curvature? (continued)


Suppose  $\alpha : I \rightarrow S$  is a path and  $\|\alpha'\| \equiv 1$ . We can study:



## What is curvature? (continued)

Suppose  $\alpha : I \rightarrow S$  is a path and  $\|\alpha'\| \equiv 1$ . We can study:


- 1 how 'straight'  $\alpha$  is on  $S$



## What is curvature? (continued)

Suppose  $\alpha : I \rightarrow S$  is a path and  $\|\alpha'\| \equiv 1$ . We can study:


- 1 how 'straight'  $\alpha$  is on  $S$  (geodesic curvature  $k_g$ )



## What is curvature? (continued)

Suppose  $\alpha : I \rightarrow S$  is a path and  $\|\alpha'\| \equiv 1$ . We can study:


- 1 how 'straight'  $\alpha$  is on  $S$  (geodesic curvature  $k_g$ )
- 2 the concavity of  $S$  along  $\alpha$



## What is curvature? (continued)

Suppose  $\alpha : I \rightarrow S$  is a path and  $\|\alpha'\| \equiv 1$ . We can study:


- ① how 'straight'  $\alpha$  is on  $S$  (geodesic curvature  $k_g$ )
- ② the concavity of  $S$  along  $\alpha$  (normal curvature  $k_n$ )



# What is curvature? (continued)

Suppose  $\alpha : I \rightarrow S$  is a path and  $\|\alpha'\| \equiv 1$ . We can study:

- ① how 'straight'  $\alpha$  is on  $S$  (geodesic curvature  $k_g$ )
- ② the concavity of  $S$  along  $\alpha$  (normal curvature  $k_n$ )




$k_g$ : component of  $\alpha''$  on tangent plane

## What is curvature? (continued)

Suppose  $\alpha : I \rightarrow S$  is a path and  $\|\alpha'\| \equiv 1$ . We can study:

- ① how 'straight'  $\alpha$  is on  $S$  (geodesic curvature  $k_g$ )
- ② the concavity of  $S$  along  $\alpha$  (normal curvature  $k_n$ )




$k_g$ : component of  $\alpha''$  on tangent plane

$k_n$ : component of  $\alpha''$  along  $N$

# What is curvature? (continued)

Suppose  $\alpha : I \rightarrow S$  is a path and  $\|\alpha'\| \equiv 1$ . We can study:

- 1 how 'straight'  $\alpha$  is on  $S$  (geodesic curvature  $k_g$ )
- 2 the concavity of  $S$  along  $\alpha$  (normal curvature  $k_n$ )



$k_g$ : component of  $\alpha''$  on tangent plane

$k_n$ : component of  $\alpha''$  along  $N$



# What is Gaussian curvature?

# What is Gaussian curvature?

- The **(Gaussian) curvature** of  $S$  at the point is

$$K = (\max k_n) \cdot (\min k_n).$$

# What is Gaussian curvature?

- The **(Gaussian) curvature** of  $S$  at the point is

$$K = (\max k_n) \cdot (\min k_n).$$


- Might be helpful to look at the sign of  $K$ :

# What is Gaussian curvature?


- The **(Gaussian) curvature** of  $S$  at the point is

$$K = (\max k_n) \cdot (\min k_n).$$


- Might be helpful to look at the sign of  $K$ :



$K < 0$



$K > 0$




$K = 0$

# What are exterior angles?



# What are exterior angles?

Think of a region  $R$  as a 'nice' subset of  $S$



# What are exterior angles?

Think of a region  $R$  as a 'nice' subset of  $S$



Denote exterior angles with  $\theta_1, \dots, \theta_p$  and interior as  $\varphi_i := \pi - \theta_i$ .

# What is the Euler characteristic?




# What is the Euler characteristic?

We can always “triangulate”  $R$ .



# What is the Euler characteristic?

We can always “triangulate”  $R$ .





# What is the Euler characteristic?

We can always “triangulate”  $R$ .



# What is the Euler characteristic?



We can always “triangulate”  $R$ .



For any triangulation of  $R$ , we compute

# What is the Euler characteristic?

We can always “triangulate”  $R$ .





For any triangulation of  $R$ , we compute

$$\underbrace{\chi(R) = V - E + F}_{\text{Euler characteristic of } R}$$

# What is the Euler characteristic?

We can always “triangulate”  $R$ .





For any triangulation of  $R$ , we compute

$$\underbrace{\chi(R) = V - E + F}_{\text{Euler characteristic of } R}$$

# What is the Euler characteristic?

We can always “triangulate”  $R$ .





For any triangulation of  $R$ , we compute

$$\underbrace{\chi(R) = V - E + F}_{\text{Euler characteristic of } R}$$

- $V$ : number of vertices

# What is the Euler characteristic?

We can always “triangulate”  $R$ .



For any triangulation of  $R$ , we compute



$$\underbrace{\chi(R) = V - E + F}_{\text{Euler characteristic of } R}$$

- $V$ : number of vertices
- $E$ : number of edges



# What is the Euler characteristic?

We can always “triangulate”  $R$ .





For any triangulation of  $R$ , we compute

$$\underbrace{\chi(R) = V - E + F}_{\text{Euler characteristic of } R}$$

- $V$ : number of vertices
- $E$ : number of edges
- $F$ : number of faces

# What is the Euler characteristic?

We can always “triangulate”  $R$ .



For any triangulation of  $R$ , we compute

$$\underbrace{\chi(R) = V - E + F}_{\text{Euler characteristic of } R}$$

- $V$ : number of vertices
- $E$ : number of edges
- $F$ : number of faces

$\chi(R)$  does not depend on triangulation!

# Proving the Gauss-Bonnet Theorem

## Theorem (Gauss-Bonnet Theorem)

If  $R$  is a region on a surface  $S$ , then

$$\int_{\partial R} k_g \, ds + \iint_R K \, dA + \sum_{i=1}^p \theta_i = 2\pi\chi(R).$$

# Proving the Gauss-Bonnet Theorem

## Theorem (Gauss-Bonnet Theorem)

If  $R$  is a region on a surface  $S$ , then

$$\int_{\partial R} k_g \, ds + \iint_R K \, dA + \sum_{i=1}^p \theta_i = 2\pi\chi(R).$$

① Show the fact for triangles:

$$\int_{\partial R} k_g \, ds + \iint_R K \, dA + \sum_{i=1}^3 \theta_i = 2\pi.$$

# Proving the Gauss-Bonnet Theorem

## Theorem (Gauss-Bonnet Theorem)

If  $R$  is a region on a surface  $S$ , then

$$\int_{\partial R} k_g ds + \iint_R K dA + \sum_{i=1}^p \theta_i = 2\pi\chi(R).$$

① Show the fact for triangles:

$$\int_{\partial R} k_g ds + \iint_R K dA + \sum_{i=1}^3 \theta_i = 2\pi.$$

Uses Stokes' Theorem:  $\oint_{\partial R} \mathbf{F} \cdot d\mathbf{r} = \iint_R \text{curl} \mathbf{F} \cdot \mathbf{N} dA$

## Theorem (Gauss-Bonnet Theorem)

If  $R$  is a region on a surface  $S$ , then

$$\int_{\partial R} k_g \, ds + \iint_R K \, dA + \sum_{i=1}^p \theta_i = 2\pi\chi(R).$$

## Theorem (Gauss-Bonnet Theorem)

If  $R$  is a region on a surface  $S$ , then

$$\int_{\partial R} k_g ds + \iint_R K dA + \sum_{i=1}^p \theta_i = 2\pi\chi(R).$$

② Triangulate the surface, apply

$$\int_{\partial R} k_g ds + \iint_R K dA + \sum_{i=1}^3 \theta_i = 2\pi.$$

for each triangle.

## Theorem (Gauss-Bonnet Theorem)

If  $R$  is a region on a surface  $S$ , then

$$\int_{\partial R} k_g ds + \iint_R K dA + \sum_{i=1}^p \theta_i = 2\pi\chi(R).$$

- ② Triangulate the surface, apply

$$\int_{\partial R} k_g ds + \iint_R K dA + \sum_{i=1}^3 \theta_i = 2\pi.$$

for each triangle.

- ③ Sum everything up and  $\chi(R)$  will pop out.



# Why care?

# Why care?

The Gauss-Bonnet Theorem unites geometry and topology.

# Why care?

The Gauss-Bonnet Theorem unites geometry and topology.

- If  $S$  is a closed surface, then

# Why care?

The Gauss-Bonnet Theorem unites geometry and topology.

- If  $S$  is a closed surface, then

$$\iint_S K \, dA = 2\pi\chi(S),$$

# Why care?

The Gauss-Bonnet Theorem unites geometry and topology.

- If  $S$  is a closed surface, then

$$\begin{aligned}\iint_S K \, dA &= 2\pi\chi(S), \\ &= 2(2 - 2g).\end{aligned}$$

- $g$ : how many holes  $S$  has


# Why care?

The Gauss-Bonnet Theorem unites geometry and topology.


- If  $S$  is a closed surface, then

$$\begin{aligned}\iint_S K \, dA &= 2\pi\chi(S), \\ &= 2(2 - 2g).\end{aligned}$$

- $g$ : how many holes  $S$  has



$$g = 0$$



$$g = 1$$




$$g = 2$$

# Why care? (continued)

## Why care? (continued)


- Take some sphere.





## Why care? (continued)


- Take some sphere.



- Draw a triangle whose sides are 'straight-lines'.

## Why care? (continued)

- Take some sphere.




- Draw a triangle whose sides are 'straight-lines'.
- The Gauss-Bonnet Theorem will tell us

$$\varphi_1 + \varphi_2 + \varphi_3 > \pi.$$

## Why care? (continued)

- Take some sphere.



- Draw a triangle whose sides are 'straight-lines'.
- The Gauss-Bonnet Theorem will tell us

$$\varphi_1 + \varphi_2 + \varphi_3 > \pi.$$

- I think that's pretty epic.

**THANK YOU!**