

# Chapter 4

## *Divide and Conquer*

The slides for this course are based on the course textbook: Cormen, Leiserson, Rivest, and Stein, Introduction to Algorithms, 3rd edition, The MIT Press, McGraw-Hill, 2010.

Many of the slides were provided by the publisher for use with the textbook. They are copyrighted, 2010.

These slides are for classroom use only, and may be used only by students in this specific course this semester. They are NOT a substitute for reading the textbook!

# Chapter 4 Topics

- Maximum subarray
- The substitution method
- The recursion-tree method
- The master method

# Designing Algorithms

- There are a number of design paradigms for algorithms that have proven useful for many types of problems
- Insertion sort – incremental approach
- Other examples of design approaches
  - divide and conquer
  - greedy algorithms
  - dynamic programming

# Divide and Conquer

- A good divide and conquer algorithm generally implies an easy recursive version of the algorithm
- Three steps
  - Divide the problem into a number of subproblems
  - Conquer the subproblems by solving them recursively. When the subproblem size is small enough, just solve the subproblem.
  - Combine - the solutions of subproblems to form the solution of the original problem

# Merge Sort

- Divide
  - divide an  $n$ -element sequence into two  $n/2$  element sequences
- Conquer
  - if the resulting list is of length 1 it is sorted
  - else call the merge sort recursively
- Combine
  - merge the two sorted sequences

MERGE-SORT ( $A, p, r$ )

1     **if**  $p < r$


2             **then**  $q \leftarrow \lfloor (p+r)/2 \rfloor$

3             *MERGE-SORT*( $A, p, q$ )

4             *MERGE-SORT*( $A, q+1, r$ )

5             *MERGE*( $A, p, q, r$ )

To sort  $A[1..n]$ , invoke MERGE-SORT with  
*MERGE-SORT*( $A, 1, \text{length}(A)$ )



# Recurrences

Definition –

a recurrence is an equation or inequality that describes a function in terms of its value on smaller inputs



# Recurrence for Divide and Conquer Algorithms

$$T(n) = \begin{cases} \Theta(1) & \text{Base case} \\ aT(n/b) + D(n) + C(n) & \end{cases}$$

*Conquer cost*      *Divide cost*      *Combine cost*

The diagram illustrates the recurrence relation for Divide and Conquer algorithms. The recurrence is given as  $T(n) = \begin{cases} \Theta(1) & \text{Base case} \\ aT(n/b) + D(n) + C(n) & \end{cases}$ . Annotations with red arrows point to the components of the recurrence: 'Base case' points to  $\Theta(1)$ , 'Conquer cost' points to  $aT(n/b)$ , 'Divide cost' points to  $D(n)$ , and 'Combine cost' points to  $C(n)$ .

# Analysis of Merge-Sort

Here is what we got as the running time:

$$T(n) = \begin{cases} \Theta(1) & \text{if } n = 1 \\ 2T(n/2) + \Theta(1) + \Theta(n) & \text{if } n > 1 \end{cases}$$

We can ignore the  $\Theta(1)$  factor, as it is irrelevant compared to  $\Theta(n)$ , and we can rewrite this recurrence as:

$$T(n) = \begin{cases} \Theta(1) & \text{if } n = 1 \\ 2T(n/2) + \Theta(n) & \text{if } n > 1 \end{cases}$$

# Why Recurrences?

- The complexity of many interesting algorithms is easily expressed as a recurrence – especially divide and conquer algorithms
- The complexity of recursive algorithms is readily expressed as a recurrence.

# Why solve recurrences?

- To make it easier to compare the complexity of two algorithms
- To make it easier to compare the complexity of the algorithm to standard reference functions.

# Example Recurrences for Algorithms

- Insertion sort
- Linear search of a list

# Recurrences for Algorithms, continued

- Binary search

# “Casual” About Some Details

- Boundary conditions
  - These are usually constant for small  $n$
- Floors and ceilings
  - Usually makes no difference in solution
  - Usually assume  $n$  is an “appropriate” integer (i.e., a power of 2) and assume that the function behaves the same way if floors and ceilings were taken into consideration

# Merge Sort Assumptions

- The actual recurrence describing the worst-case running time for merge sort is:


$$T(n) = \begin{cases} \Theta(1) & \text{for } n \leq 1 \\ T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + \Theta(n) & \text{otherwise} \end{cases}$$

- But we typically assume that  $n = 2^k$  where  $k$  is an integer and use the simpler recurrence.



# Maximum-subarray Problem


- Stock investment: Buy one unit of stock only one time and then sell it at a later date
- Goal: to maximize the profit



Day	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
Price	100	113	110	85	105	102	86	63	81	101	94	106	101	79	94	90	97
Change		13	-3	-25	20	-3	-16	-23	18	20	-7	12	-5	-22	15	-4	7

# One Potential Solution

- Find the highest price and search left to find the lowest prior price
- Find the lowest price and search right to find the highest later price
- Take the pair with the greater difference
- Do not work! See counterexample below.




Day	0	1	2	3	4
Price	10	11	7	10	6
Change		1	-4	3	-4

# Maximum Subarray



- Consider the daily change in price
- Maximum subarray problem: find the non-empty, contiguous subarray of A whose values have the largest sum.

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
A	13	-3	-25	20	-3	-16	-23	18	20	-7	12	-5	-22	15	-4	7

  
maximum subarray

# Divide and Conquer

- Suppose we want to find a maximum subarray of  $A[\text{low}..\text{high}]$
- Divide and conquer will find the midpoint, say  $\text{mid}$ , of the subarray, and consider the subarrays  $A[\text{low}..\text{mid}]$  and  $A[\text{mid}+1..\text{high}]$
- Any contiguous subarray  $A[i..j]$  must lie in one area out of three possibilities



# Find Max Crossing Subarray

- First, it is easy to find a maximum subarray crossing the midpoint
- We just need to find maximum subarrays of the form  $A[i..mid]$  and  $A[mid+1..j]$  and combine them

FIND-MAX-CROSSING-SUBARRAY( $A, low, mid, high$ )

// Find a maximum subarray of the form  $A[i \dots mid]$ .

$left\text{-}sum = -\infty$

$sum = 0$

**for**  $i = mid$  **downto**  $low$

$sum = sum + A[i]$

**if**  $sum > left\text{-}sum$

$left\text{-}sum = sum$

$max\text{-}left = i$

// Find a maximum subarray of the form  $A[mid + 1 \dots j]$ .

$right\text{-}sum = -\infty$

$sum = 0$

**for**  $j = mid + 1$  **to**  $high$

$sum = sum + A[j]$

**if**  $sum > right\text{-}sum$

$right\text{-}sum = sum$

$max\text{-}right = j$

// Return the indices and the sum of the two subarrays.

**return** ( $max\text{-}left, max\text{-}right, left\text{-}sum + right\text{-}sum$ )

# Find Maximum subarray

- We can then write a divide and conquer algorithm to solve the maximum subarray problem.
- Divide into three cases, and choose the best solution
  - Left subarray
  - Crossing subarray
  - Right subarray

FIND-MAXIMUM-SUBARRAY( $A, low, high$ )

**if**  $high == low$

**return** ( $low, high, A[low]$ )                      // base case: only one element

**else**  $mid = \lfloor (low + high) / 2 \rfloor$

    ( $left-low, left-high, left-sum$ ) =

        FIND-MAXIMUM-SUBARRAY( $A, low, mid$ )

    ( $right-low, right-high, right-sum$ ) =

        FIND-MAXIMUM-SUBARRAY( $A, mid + 1, high$ )

    ( $cross-low, cross-high, cross-sum$ ) =

        FIND-MAX-CROSSING-SUBARRAY( $A, low, mid, high$ )

**if**  $left-sum \geq right-sum$  and  $left-sum \geq cross-sum$

**return** ( $left-low, left-high, left-sum$ )

**elseif**  $right-sum \geq left-sum$  and  $right-sum \geq cross-sum$

**return** ( $right-low, right-high, right-sum$ )

**else return** ( $cross-low, cross-high, cross-sum$ )



# Analyzing the algorithm

- So the total running time is?

$$T(n) = 2 \cdot T\left(\frac{n}{2}\right) + \underbrace{D(n)} + \underbrace{Q(n)}$$

$$T(n) = 2T\left(\frac{n}{2}\right) + O(1) + \underline{n}$$

# Methods for Solving Recurrences

- Constructive induction
- Iterative substitution
  - Recurrence trees
- Master Theorem

# Constructive Induction

- Use mathematical induction to derive an answer
- Steps
  1. Guess the form of the solution
  2. Use mathematical induction to find constants or show that they can be found and to prove that the answer is correct

# Constructive induction

- Goal
  - Derive a function of  $n$  (or other variables used to express the size of the problem) that is not a recurrence so we can establish an upper and/or lower bound on the recurrence
  - We may get an exact solution or we may just get upper or lower bounds on the solution

# Constructive Induction

- Suppose  $T$  includes a parameter  $n$  and  $n$  is a natural number (positive integer)
- Instead of proving directly that  $T$  holds for all values of  $n$ , prove

- $T$  holds for a base case  $b$  (often  $n = 1$ )
- For every  $n > b$ , if  $T$  holds for  $n-1$ , then  $T$  holds for  $n$ .

» Assume  $T$  holds for  $n-1$

» Prove that  $T$  holds for  $n$  follows from this assumption

$$T(n) ? T(n-1)$$

$$T(n) ? T(n/2), \text{ if } T \text{ holds for } n/2, \text{ then } T \text{ holds for } n$$

# Example 1

- Given

$$T(n) = \begin{cases} 1 & \text{for } n \leq 1 \\ T(n-1) + n & \text{otherwise} \end{cases}$$

- Prove  $T(n) \in O(n^2)$

- Note that this is the recurrence for insertion sort and we have already shown that this is  $O(n^2)$  using other methods

$$T(n) = \sum_{i=1}^n i = \frac{n(n+1)}{2} \in O(n^2)$$

# Proof of Example 1

$$T(n) = an^2 + bn + c, \therefore T(n) = \frac{1}{2}n^2 + \frac{1}{2}n \quad \text{①}$$

base:

$$T(1) = a \cdot 1^2 + b \cdot 1 + c = 1 \Rightarrow a + b + c = 1$$

induction's step: <sup>suppose</sup>  $T(n-1) = a(n-1)^2 + b(n-1) + c$

(We need to show:

$$T(n) = T(n-1) + n$$

$$\Rightarrow an^2 + bn + c = a(n-1)^2 + b(n-1) + c + n$$

$$\Rightarrow an^2 + bn + c = an^2 - 2an + a + bn - b + c + n$$
$$= an^2 + (b+1-2a)n + a-b+c$$

$\Rightarrow c=0$

$$\text{② } \cancel{a} = \cancel{a} + 1 - 2a \Rightarrow a = \frac{1}{2} \quad \text{③ } \cancel{c} = a - b + \cancel{c} \Rightarrow b = a = \frac{1}{2}$$

## Example 2 – Establishing an Upper Bound

Recurrence :  $T(n) = 4T(n/2) + n$

Guess :  $T(n) \in O(n^3)$

$$\underline{T(n) \leq C \cdot n^3}$$

*loose upper bound*  
 ~~$T(n) = an^3 + bn^2 + cn + d$~~

base :  $T(2) = 4T(1) + 2 \leq C \cdot 2^3 \Rightarrow C \geq \frac{6}{8}$

inductive step : Assume  $T(n/2) \leq C \cdot (\frac{n}{2})^3$

We need to prove  $T(n) \leq C \cdot n^3$

$$T(n) = 4 \cdot T(n/2) + n \leq 4 \cdot C \cdot (\frac{n}{2})^3 + n$$

$$\frac{C}{2} n^3 + n$$

$$\text{So } \frac{C}{2} n^3 + n \leq C n^3 \Rightarrow \frac{C}{2} n^3 + n = \frac{C}{2} n^3 + n$$

$$C \geq \sqrt[3]{2/n} \geq 2$$



## Ex. 3 – Fallacious Argument

Recurrence:  $T(n) = 4T(n/2) + n$

Guess:  $T(n) \in O(n^2)$

$$\underline{T(n)} \leq Cn^2$$

base: skipped

Inductive step: Assume  $T(\frac{n}{2}) \leq C(\frac{n}{2})^2$

We need to prove  $\underline{T(n) \leq Cn^2}$

$$T(n) = 4T(n/2) + n \leq 4 \cdot C(\frac{n}{2})^2 + n$$

$$= Cn^2 + n \quad \underline{\underline{\text{failed!}}}$$

## Example 3 – Try again

Assume  
 $T(n) \leq$  "desired term" – "positive term"

$$\underline{T(n) \leq C_1 n^2 - C_2 n} \Rightarrow \underline{T(n) \in O(n^2)}$$

base case:  $n=2$   
 $\underline{T(n) = C_1 \cdot 2^2 - C_2 \cdot 2 \geq 1}$

inductive step: Assumption:  $\underline{T(\frac{n}{2}) \leq C_1 (\frac{n}{2})^2 - C_2 \frac{n}{2}}$   
We need to prove:  $\underline{T(n) \leq C_1 n^2 - C_2 n}$  ( $\forall C_2 \geq 1$ )  
$$\begin{aligned} T(n) &= 4T(\frac{n}{2}) + n \leq 4 \cdot C_1 (\frac{n}{2})^2 - 4C_2 \frac{n}{2} + n \quad T(n) \leq C_1 n^2 - C_2 n \\ &= C_1 n^2 - 2C_2 n + n = C_1 n^2 - C_2 n - (C_2 - 1)n \end{aligned}$$

# Boundary Conditions

- Boundary conditions are not usually important because we don't need an actual  $c$  value (if polynomially bounded)
- But sometimes it makes a big difference
  - Exponential solutions
  - Suppose we are searching for a solution to:  
$$T(n) = T(n/2)^2$$
  - and we find the partial solution:  
$$T(n) = c^n$$

# Boundary Conditions, cont.

If the boundary condition is

$$T(n) = 2$$

this implies that  $T(n) \in \Theta(2^n)$ .

But if the boundary condition is

$$T(n) = 3$$

this implies that  $T(n) \in \Theta(3^n)$ ,

and  $\Theta(3^n) \neq \Theta(2^n)$ .

The results are even more dramatic if  $T(1) = 1$

$$T(1) = 1 \Rightarrow T(n) = \Theta(1^n) = \Theta(1)$$

# Boundary Conditions

The solutions to the recurrences below have very different upper bounds:

$$T(n) = \begin{cases} 1 & \text{for } n = 1 \\ T(n/2)^2 & \text{otherwise} \end{cases}$$
$$T(n) = \begin{cases} 2 & \text{for } n = 1 \\ T(n/2)^2 & \text{otherwise} \end{cases}$$
$$T(n) = \begin{cases} 3 & \text{for } n = 1 \\ T(n/2)^2 & \text{otherwise} \end{cases}$$

# Iterating the Recurrence

- Called *iterative substitution*
- Sometimes referred to as *plug and chug*.
- In iterative substitution we substitute the original form of the recurrence everywhere  $T$  occurs on the right side of the recurrence equation.
- Repeat as needed until a pattern appears.
- We can use this method to get an estimate that we can use for the substitution method.

# Iterating the Recurrence

Look at the recurrence relation:

$$T(n) = \begin{cases} 0 & \text{if } n = 0 \\ T(n - 1) + n & \text{if } n > 0 \end{cases}$$

Substituting  $n - 1$  for  $n$  in the relation above we get:

$$T(n - 1) = T(n - 2) + (n - 1)$$

Substitute for  $n - 1$  in the original relation:

$$T(n) = (T(n - 2) + (n - 1)) + n$$

We know that

$$T(n - 2) = T(n - 3) + (n - 2)$$

So substitute this for  $T(n - 2)$  above:

$$T(n) = (T(n - 3) + (n - 2)) + (n - 1) + n$$

# Iterating the Recurrence

We see the following pattern:

$$T(n) = T(n - 1) + n$$

$$T(n) = (T(n - 2) + (n - 1)) + n$$

$$T(n) = (T(n - 3) + (n - 2)) + (n - 1) + n$$

...

$$T(n) = T(n - (n - 2)) + 2 + 3 + \dots + (n - 2) + (n - 1) + n$$

$$T(n) = T(n - (n - 1)) + 2 + 3 + \dots + (n - 2) + (n - 1) + n$$

$$T(n) = T(n - (n - 0)) + 2 + 3 + \dots + (n - 2) + (n - 1) + n$$

We can rewrite  $(n - (n - 0))$  as  $(n - n)$  or as  $(0)$ , thus:

$$T(n) = T(0) + 1 + 2 + 3 + \dots + (n - 2) + (n - 1) + n$$

But we know that  $T(0) = 0$  is the base case, so:

$$T(n) = 0 + 1 + 2 + 3 + \dots + (n - 2) + (n - 1) + n$$

$$\frac{n(n+1)}{2}$$



# Iterating the Recurrence

The summation of

$$T(n) = 0 + 1 + 2 + 3 + \dots + (n - 2) + (n - 1) + n$$

is

$$T(n) = (n(n + 1) / 2) = \frac{1}{2} n^2 + \frac{1}{2} n$$

which we recognize as  $O(n^2)$ .

# Iterating the Recurrence

$$T\left(\frac{n}{2}\right) = 2T\left(\frac{n}{2^2}\right) + c\frac{n}{2}$$

Let's look at the recurrence equation for Merge Sort again:

$$T(n) = \begin{cases} c & \text{if } n = 1 \\ 2T(n/2) + cn & \text{if } n > 1 \end{cases}$$

$$T(n) = 2T\left(\frac{n}{2}\right) + cn$$

$$= 2\left(2T\left(\frac{n}{2^2}\right) + c\frac{n}{2}\right) + cn$$

$$= 2^2 T\left(\frac{n}{2^2}\right) + cn + cn$$

$$= 2^2 \left(2T\left(\frac{n}{2^3}\right) + c\frac{n}{2^2}\right) + cn + cn$$

$$= 2^3 T\left(\frac{n}{2^3}\right) + cn + cn + cn$$

$$? = 2^{\log_2 n} T\left(\frac{n}{2^{\log_2 n}}\right) + \underbrace{cn + \dots + cn}_{\log_2 n} \approx \underline{\underline{(\log_2 n) \cdot n + cn}}$$

$$\begin{aligned} n &\rightarrow n=2^k \\ \frac{n}{2^k} &= 1 \\ k &= \log_2 n \end{aligned}$$

$$O(n \log n)$$

$$= n(1 + 2 + 2^2 + \dots + 2^{\log_2 n}) + 4n^2$$

## Example 4

$$T(n) = n + 4T(n/2) \quad T(n/2) = \frac{n}{2} + 4T(n/4)$$

$$= n + 4\left(\frac{n}{2} + 4T\left(\frac{n}{4}\right)\right) \quad T\left(\frac{n}{4}\right) = \frac{n}{4} + 4T\left(\frac{n}{8}\right)$$

$$= n + 4 \cdot \frac{n}{2} + 4^2 T\left(\frac{n}{4}\right)$$

$$= n + 4 \cdot \frac{n}{2} + 4^2 \left(\frac{n}{4} + 4T\left(\frac{n}{8}\right)\right)$$

$$= n + 4 \cdot \frac{n}{2} + 4^2 \frac{n}{4} + 4^3 T\left(\frac{n}{8}\right)$$

$$= n + 4 \frac{n}{2} + 4^2 \frac{n}{4} + 4^3 \frac{n}{8} + \dots$$

$$= \underbrace{n + 2n + 2^2 n + 2^3 n + \dots}_{\log_2 n} + \underbrace{n \cdot \frac{4^{\log_2 n}}{2^{\log_2 n}}}_{n \cdot 1} + \dots + \underbrace{4^{\log_2 n}}_{n^2} + \dots + \underbrace{I(1)}_{1}$$

$$4^{\log_2 n} = 2^{2 \log_2 n} = 2^{\log_2 n^2} = n^2$$

$$1 = \frac{n}{2^k}$$

$$k = \log_2 n$$

# Example 5


Recurrence:  $T(n) = 4T(n/3) + n$

Guess:  $T(n) \in O(n^2)$


# Recurrence Trees

- Allow you to visualize the process of iterating the recurrence
- Allows you make a good guess for the substitution method
- Or to organize the bookkeeping for iterating the recurrence
- Example


$$T(n) = T(n/4) + T(n/2) + n^2$$



Since the values decrease geometrically, the total is at most a constant factor more than the largest term and hence the solution is  $\Theta(n^2)$




**Figure 4.1** The construction of a recursion tree for the recurrence  $T(n) = 3T(n/4) + cn^2$ . Part (a) shows  $T(n)$ , which is progressively expanded in (b)–(d) to form the recursion tree. The fully expanded tree in part (d) has height  $\log_4 n$  (it has  $\log_4 n + 1$  levels).




**Figure 4.2** A recursion tree for the recurrence  $T(n) = T(n/3) + T(2n/3) + cn$ .





**Figure 4.3** The recursion tree generated by  $T(n) = aT(n/b) + f(n)$ . The tree is a complete  $a$ -ary tree with  $n^{\log_b a}$  leaves and height  $\log_b n$ . The cost of each level is shown at the right, and their sum is given in equation (4.6).



**Figure 4.4** The recursion tree generated by  $T(n) = aT(\lceil n/b \rceil) + f(n)$ . The recursive argument  $n_j$  is given by equation (4.12).

# The master method

Provides a cookbook method for solving recurrences of the form

$$T(n) = aT(n/b) + f(n)$$

where  $a \geq 1$  and  $b > 1$  and  $f(n)$  is an asymptotically positive function.

# Divide and Conquer Algorithms

- The form of the master theorem is very convenient because divide and conquer algorithms have recurrences of the form

$$T(n) = aT(n/b) + D(n) + C(n)$$

where

$a$  is the number of subproblem s at each step

$n/b$  is the size of each subproblem

$D(n)$  is the cost of dividing into subproblem s

$C(n)$  is the cost of combining the solutions to  
subproblem s

# Form of the Master Theorem

- Combines  $D(n)$  and  $C(n)$  into  $f(n)$
- For example, in Merge-Sort

$$T(n) = \begin{cases} \Theta(1) & \text{if } n = 1 \\ 2T(n/2) + \Theta(n) & \text{if } n > 1 \end{cases}$$

$$a = 2, b = 2$$

$$f(n) = \Theta(n)$$

- We will ignore floors and ceilings. The proof of the Master Theorem includes a proof that this is ok.

# Form of the Master Theorem

- Combines  $D(n)$  and  $C(n)$  into  $f(n)$
- For example, in Merge-Sort

$$T(n) = \begin{cases} \Theta(1) & \text{for } n = 1 \\ 2T(n/2) + \Theta(n) & \text{for } n > 1 \end{cases}$$

$$a = 2, b = 2$$

$$f(n) = \Theta(n)$$

We will ignore floors and ceilings. The proof of the Master Theorem includes a proof that this is ok.

# Form of the Master Theorem

- The Master Method is used for recurrence equations of the form:

$$T(n) = \begin{cases} c & \text{for } n < d \\ aT(n/b) + f(n) & \text{for } n \geq 1 \end{cases}$$

# Master theorem

Let  $a \geq 1$  and  $b > 1$  be constants, let  $f(n)$  be a function, and let  $T(n)$  be defined on the non-negative integers by the recurrence

$$T(n) = aT(n/b) + f(n)$$

where we interpret  $n/b$  to mean either the floor or ceiling of  $n/b$ . Then  $T(n)$  can be bounded asymptotically as follows:



# Master theorem

Case 1 : if  $f(n) = O\left(n^{\log_b a - \varepsilon}\right)$  for some constant  $\varepsilon > 0$ , then

$$T(n) = \Theta\left(n^{\log_b a}\right)$$

Case 2: if  $f(n) = \Theta\left(n^{\log_b a}\right)$ , then

$$T(n) = \Theta\left(n^{\log_b a} \lg n\right)$$

Case 3 : if  $f(n) = \Omega\left(n^{\log_b a + \varepsilon}\right)$  for some constant  $\varepsilon > 0$ , and

if  $af(n/b) \leq cf(n)$  for some constant  $c < 1$  and all sufficiently large  $n$ , then

$$T(n) = \Theta\left(f(n)\right)$$

## 3 cases

1. If there is a small constant  $\varepsilon > 0$ , such that

$$\underline{f(n) = O(n^{\log_b a - \varepsilon})} \Rightarrow f(n) = o(n^{\log_b a})$$

then  $T(n)$  is

$$\Theta(n^{\log_b a})$$

Here  $f(n)$  is polynomially smaller than the  
special function  $n^{\log_b a}$

## 3 cases

2. If

$$f(n) = \Theta\left(n^{\log_b a}\right)$$

then  $T(n)$  is

$$\Theta\left(n^{\log_b a} \lg n\right)$$

Here  $f(n)$  is asymptotically equal to the special  
function  $n^{\log_b a}$

3 cases  $f(n) \sim n^{\log_b a}$

3. If there are small constants  $\varepsilon > 0$  and  $c < 1$ ,  
such that  $af(n/b) \leq cf(n)$

$$f(n) = \Omega\left(n^{\log_b a + \varepsilon}\right) \Leftrightarrow f(n) \not\sim n^{\log_b a}$$

for all sufficiently large  $n$ , then  $T(n)$  is

$$\Theta(f(n))$$

Here  $f(n)$  is polynomially larger than the special  
function  $n^{\log_b a}$

# What does the master theorem say?

Compare two functions:

$$f(n) \quad \text{and} \quad n^{\log_b a}$$

When  $f(n)$  grows asymptotically slower (Case 1)

$$T(n) = \Theta(n^{\log_b a})$$

When the growth rates are the same (Case 2)

$$T(n) = \Theta(f(n) \lg n) = \Theta(n^{\log_b a} \lg n)$$

When  $f(n)$  grows asymptotically faster (Case 3)

$$T(n) = \Theta(f(n))$$

# Using the Master Method

Using the master method, solve the recurrence

$$T(n) = 4T(n/2) + n$$

$$a = 4$$

$$b = 2$$

$$f(n) = n$$

$$f(n) = n < n^{\log_2 4} = n^2$$

$$T(n) \in \Theta(n^2)$$

# Using the Master Method

$$T(n) = 64T(n/4) + n$$

$a=64$   
 $b=4$   
 $f(n)=n$

$n^{\log_b a} = n^3$

$T(n) \in \Theta(n^3)$

# Using the Master Method

Using the master method, solve the recurrence

$$T(n) = T(2n/3) + 1$$

$$a = 1$$

$$b = 3/2$$

$$n^{\log_b a} = n^{\log_{3/2} 1} = n^0 = 1$$

$$\underline{f(n) = 1}$$

$$T(n) = \Theta(\log n)$$



# Using the Master Method

$$T(n) = T(3n / 4) + 1$$

# Using the Master Method

Using the master method, solve the recurrence

$$T(n) = T(n/3) + n$$

$$a = 1$$

$$b = 3$$

$$\underline{\underline{f(n) = n}}$$

$$\log_b a = 1$$

$$T(n) \in \underline{\underline{\Theta(n)}}$$

# Using the Master Method

②  $T(n) = 4T(n/3) + n \lg n$

①  $T(n) = 3T(n/4) + n \lg n$

$a = 3$

$b = 4$



$n^{\log_b a} = n^{\log_4 3}$

$n^{\log_3 4}$

$f(n) = n \lg n$

$\frac{n \cdot n^{(\log_3 4 - 1)}}{c}$

$\lg n ? n^c T(n) = n \lg n$

$T(n) = n^{\log_3 4}$

# Conclusion

- We talked about:
  - ✓ The substitution method (2 types)
  - ✓ The recursion-tree method
  - ✓ The master method
- Be able to solve recurrences using all three of these methods.

# The Master Theorem

Let  $a \geq 1$  and  $b > 1$  be constants, let  $f(n)$  be a function, and let  $T(n)$  be defined on the nonnegative integers by the recurrence  $T(n) = aT(n/b) + f(n)$

where  $n/b$  can be either  $\lfloor n/b \rfloor$  or  $\lceil n/b \rceil$

Then  $T(n)$  can be bounded asymptotically as follows:

1. If  $f(n) = O(n^{\log_b a - \varepsilon})$  for some constant  $\varepsilon > 0$ ,  
then  $T(n) = \Theta(n^{\log_b a})$
2. If  $f(n) = \Theta(n^{\log_b a})$ , then  $T(n) = \Theta(n^{\log_b a} \lg n)$
3. If  $f(n) = \Omega(n^{\log_b a - \varepsilon})$  for some constant  $\varepsilon > 0$ , and  
if  $af(n/b) \leq cf(n)$  for some constant  $c < 1$  and all  
sufficiently large  $n$ , then  $T(n) = \Theta(f(n))$