

Topics in Analysis and Linear Algebra

Le Chen

le.chen@emory.edu

Emory University

Atlanta GA

Last updated on

July 27, 2021

Summer Bootcamp for
Emory Biostatistics and Bioinformatics
PhD Program

July 22 - 28, 2021

Chapter 3. Real Number System and Calculus

This part is mostly based on Chapter 2 of

*J. McDonald and N. Weiss, **A course in real analysis***, Academic Press, 2005.

Chapter 3. Real Number System and Calculus

§ 3.1 Real number system

§ 3.2 Sequences of real numbers

§ 3.3 Open and closed sets

§ 3.4 Real-valued functions

§ 3.5 Liminf and limsup of sets

§ 3.6 Some techniques in calculus

Chapter 3. Real Number System and Calculus

§ 3.1 Real number system

§ 3.2 Sequences of real numbers

§ 3.3 Open and closed sets

§ 3.4 Real-valued functions

§ 3.5 Liminf and limsup of sets

§ 3.6 Some techniques in calculus

What is a real number?



1

¹Image from Wikipedia.



2

²Image from

<https://geeksoutofthebox.com/2019/03/15/simons-real-numbers-diagram/>

Real number system can be formulated in three groups of axioms

(F) Field Axioms

(O) Order Axioms

(C) Completeness Axioms

Real number system can be formulated in three groups of axioms

(F) Field Axioms

(O) Order Axioms

(C) Completeness Axioms

Real number system can be formulated in three groups of axioms

(F) Field Axioms

(O) Order Axioms

(C) Completeness Axioms

Field Axioms

Let $x, y, z \in \mathbb{R}$. Then we have that

(F1) $x + y = y + x$ and $xy = yx$. (Commutative)

(F2) $(x + y) + z = x + (y + z)$ and $(xy)z = x(yz)$. (Associative)

(F3) $x(y + z) = xy + xz$. (Distributive)

(F4) There exist $0, 1 \in \mathbb{R}$ with $0 \neq 1$ such that for all $x \in \mathbb{R}$

$x + 0 = x$ and $x \cdot 1 = x$. (Identities)

(F5) For each $x \in \mathbb{R}$, there exists a $-x \in \mathbb{R}$ such that $x + (-x) = 0$ and, if $x \neq 0$, there exists an $x^{-1} \in \mathbb{R}$ such that $xx^{-1} = 1$. (Inverses)

Field Axioms

Let $x, y, z \in \mathbb{R}$. Then we have that

(F1) $x + y = y + x$ and $xy = yx$. (Commutative)

(F2) $(x + y) + z = x + (y + z)$ and $(xy)z = x(yz)$. (Associative)

(F3) $x(y + z) = xy + xz$. (Distributive)

(F4) There exist $0, 1 \in \mathbb{R}$ with $0 \neq 1$ such that for all $x \in \mathbb{R}$

$x + 0 = x$ and $x \cdot 1 = x$. (Identities)

(F5) For each $x \in \mathbb{R}$, there exists a $-x \in \mathbb{R}$ such that $x + (-x) = 0$ and, if $x \neq 0$, there exists an $x^{-1} \in \mathbb{R}$ such that $xx^{-1} = 1$. (Inverses)

Field Axioms

Let $x, y, z \in \mathbb{R}$. Then we have that

$$(F1) \quad x + y = y + x \text{ and } xy = yx. \quad (\text{Commutative})$$

$$(F2) \quad (x + y) + z = x + (y + z) \text{ and } (xy)z = x(yz). \quad (\text{Associative})$$

$$(F3) \quad x(y + z) = xy + xz. \quad (\text{Distributive})$$

$$(F4) \quad \text{There exist } 0, 1 \in \mathbb{R} \text{ with } 0 \neq 1 \text{ such that for all } x \in \mathbb{R}$$

$$x + 0 = x \quad \text{and} \quad x \cdot 1 = x. \quad (\text{Identities})$$

$$(F5) \quad \text{For each } x \in \mathbb{R}, \text{ there exists a } -x \in \mathbb{R} \text{ such that } x + (-x) = 0 \text{ and, if } x \neq 0, \text{ there exists an } x^{-1} \in \mathbb{R} \text{ such that } xx^{-1} = 1. \quad (\text{Inverses})$$

Field Axioms

Let $x, y, z \in \mathbb{R}$. Then we have that

$$(F1) \quad x + y = y + x \text{ and } xy = yx. \quad (\text{Commutative})$$

$$(F2) \quad (x + y) + z = x + (y + z) \text{ and } (xy)z = x(yz). \quad (\text{Associative})$$

$$(F3) \quad x(y + z) = xy + xz. \quad (\text{Distributive})$$

$$(F4) \quad \text{There exist } 0, 1 \in \mathbb{R} \text{ with } 0 \neq 1 \text{ such that for all } x \in \mathbb{R}$$

$$x + 0 = x \quad \text{and} \quad x \cdot 1 = x. \quad (\text{Identities})$$

$$(F5) \quad \text{For each } x \in \mathbb{R}, \text{ there exists a } -x \in \mathbb{R} \text{ such that } x + (-x) = 0 \text{ and, if } x \neq 0, \text{ there exists an } x^{-1} \in \mathbb{R} \text{ such that } xx^{-1} = 1. \quad (\text{Inverses})$$

Field Axioms

Let $x, y, z \in \mathbb{R}$. Then we have that

$$(F1) \quad x + y = y + x \text{ and } xy = yx. \quad (\text{Commutative})$$

$$(F2) \quad (x + y) + z = x + (y + z) \text{ and } (xy)z = x(yz). \quad (\text{Associative})$$

$$(F3) \quad x(y + z) = xy + xz. \quad (\text{Distributive})$$

$$(F4) \quad \text{There exist } 0, 1 \in \mathbb{R} \text{ with } 0 \neq 1 \text{ such that for all } x \in \mathbb{R}$$

$$x + 0 = x \quad \text{and} \quad x \cdot 1 = x. \quad (\text{Identities})$$

$$(F5) \quad \text{For each } x \in \mathbb{R}, \text{ there exists a } -x \in \mathbb{R} \text{ such that } x + (-x) = 0 \text{ and, if } x \neq 0, \text{ there exists an } x^{-1} \in \mathbb{R} \text{ such that } xx^{-1} = 1. \quad (\text{Inverses})$$

Order Axioms

Let $x, y, z \in \mathbb{R}$. Then we have that

(O1) $x < y$ and $y < z$ implies that $x < z$. (Transitive)

(O2) $x < y$ implies that $x + z < y + z$.

(O3) $x < y$ and $z > 0$ implies that $xz < yz$.

(O4) Exactly one of $x = y$, $x < y$, and $x > y$ holds. (Trichotomous)

Order Axioms

Let $x, y, z \in \mathbb{R}$. Then we have that

(O1) $x < y$ and $y < z$ implies that $x < z$. (Transitive)

(O2) $x < y$ implies that $x + z < y + z$.

(O3) $x < y$ and $z > 0$ implies that $xz < yz$.

(O4) Exactly one of $x = y$, $x < y$, and $x > y$ holds. (Trichotomous)

Order Axioms

Let $x, y, z \in \mathbb{R}$. Then we have that

(O1) $x < y$ and $y < z$ implies that $x < z$. (Transitive)

(O2) $x < y$ implies that $x + z < y + z$.

(O3) $x < y$ and $z > 0$ implies that $xz < yz$.

(O4) Exactly one of $x = y$, $x < y$, and $x > y$ holds. (Trichotomous)

Order Axioms


Let $x, y, z \in \mathbb{R}$. Then we have that

- (O1) $x < y$ and $y < z$ implies that $x < z$. (Transitive)
- (O2) $x < y$ implies that $x + z < y + z$.
- (O3) $x < y$ and $z > 0$ implies that $xz < yz$.
- (O4) Exactly one of $x = y$, $x < y$, and $x > y$ holds. (Trichotomous)

Completeness Axiom

Axiom A nonempty subset of real numbers that is **bounded above** has a **least upper bound**, which is denoted as


$$\sup A, \quad \sup_{x \in A} x, \quad \text{or} \quad \sup\{x : x \in A\}.$$



Completeness Axiom

Axiom A nonempty subset of real numbers that is **bounded above** has a **least upper bound**, which is denoted as


$$\sup A, \quad \sup_{x \in A} x, \quad \text{or} \quad \sup\{x : x \in A\}.$$



Completeness Axiom

Axiom A nonempty subset of real numbers that is **bounded above** has a **least upper bound**, which is denoted as


$$\sup A, \quad \sup_{x \in A} x, \quad \text{or} \quad \sup\{x : x \in A\}.$$



Completeness Axiom

Axiom A nonempty subset of real numbers that is **bounded above** has a **least upper bound**, which is denoted as


$$\sup A, \quad \sup_{x \in A} x, \quad \text{or} \quad \sup\{x : x \in A\}.$$



Completeness Axiom

Axiom A nonempty subset of real numbers that is **bounded above** has a **least upper bound**, which is denoted as


$$\sup A, \quad \sup_{x \in A} x, \quad \text{or} \quad \sup\{x : x \in A\}.$$



Completeness Axiom

Axiom A nonempty subset of real numbers that is **bounded above** has a **least upper bound**, which is denoted as


$$\sup A, \quad \sup_{x \in A} x, \quad \text{or} \quad \sup\{x : x \in A\}.$$



Completeness Axiom

Axiom A nonempty subset of real numbers that is **bounded above** has a **least upper bound**, which is denoted as


$$\sup A, \quad \sup_{x \in A} x, \quad \text{or} \quad \sup\{x : x \in A\}.$$



Completeness Axiom

Axiom A nonempty subset of real numbers that is **bounded above** has a **least upper bound**, which is denoted as


$$\sup A, \quad \sup_{x \in A} x, \quad \text{or} \quad \sup\{x : x \in A\}.$$



Completeness Axiom

Axiom A nonempty subset of real numbers that is **bounded above** has a **least upper bound**, which is denoted as


$$\sup A, \quad \sup_{x \in A} x, \quad \text{or} \quad \sup\{x : x \in A\}.$$



Completeness Axiom

Axiom A nonempty subset of real numbers that is **bounded above** has a **least upper bound**, which is denoted as


$$\sup A, \quad \sup_{x \in A} x, \quad \text{or} \quad \sup\{x : x \in A\}.$$



Completeness Axiom

Axiom A nonempty subset of real numbers that is **bounded above** has a **least upper bound**, which is denoted as


$$\sup A, \quad \sup_{x \in A} x, \quad \text{or} \quad \sup\{x : x \in A\}.$$



Completeness Axiom

Axiom A nonempty subset of real numbers that is **bounded above** has a **least upper bound**, which is denoted as


$$\sup A, \quad \sup_{x \in A} x, \quad \text{or} \quad \sup\{x : x \in A\}.$$



Completeness Axiom

Axiom A nonempty subset of real numbers that is **bounded above** has a **least upper bound**, which is denoted as


$$\sup A, \quad \sup_{x \in A} x, \quad \text{or} \quad \sup\{x : x \in A\}.$$



Completeness Axiom

Axiom A nonempty subset of real numbers that is **bounded above** has a **least upper bound**, which is denoted as


$$\sup A, \quad \sup_{x \in A} x, \quad \text{or} \quad \sup\{x : x \in A\}.$$



Completeness Axiom

Axiom A nonempty subset of real numbers that is **bounded above** has a **least upper bound**, which is denoted as


$$\sup A, \quad \sup_{x \in A} x, \quad \text{or} \quad \sup\{x : x \in A\}.$$



Completeness Axiom

Axiom A nonempty subset of real numbers that is **bounded above** has a **least upper bound**, which is denoted as

$$\sup A, \quad \sup_{x \in A} x, \quad \text{or} \quad \sup\{x : x \in A\}.$$



Let $\{x_n\}_{n \in \mathbb{N}}$ be a sequence and $a \in \mathbb{R}$. Then

$$\sup_n x_n \leq a \iff \forall n, x_n \leq a$$


$$\sup_n x_n < a \iff \forall n, x_n < a$$

$$a < \sup_n x_n \iff \exists n, x_n > a$$

$$a \leq \sup_n x_n \iff \exists n, x_n \geq a$$


Corr. A nonempty subset of real numbers that is **bounded below** has a **greatest lower bound**, which is denoted as

$$\inf A, \quad \inf_{x \in A} x, \quad \text{or} \quad \inf\{x : x \in A\}.$$




Corr. A nonempty subset of real numbers that is **bounded below** has a **greatest lower bound**, which is denoted as

$$\inf A, \quad \inf_{x \in A} x, \quad \text{or} \quad \inf\{x : x \in A\}.$$




Corr. A nonempty subset of real numbers that is **bounded below** has a **greatest lower bound**, which is denoted as

$$\inf A, \quad \inf_{x \in A} x, \quad \text{or} \quad \inf\{x : x \in A\}.$$




Corr. A nonempty subset of real numbers that is **bounded below** has a **greatest lower bound**, which is denoted as

$$\inf A, \quad \inf_{x \in A} x, \quad \text{or} \quad \inf\{x : x \in A\}.$$




Corr. A nonempty subset of real numbers that is **bounded below** has a **greatest lower bound**, which is denoted as

$$\inf A, \quad \inf_{x \in A} x, \quad \text{or} \quad \inf\{x : x \in A\}.$$




Corr. A nonempty subset of real numbers that is **bounded below** has a **greatest lower bound**, which is denoted as

$$\inf A, \quad \inf_{x \in A} x, \quad \text{or} \quad \inf\{x : x \in A\}.$$




Corr. A nonempty subset of real numbers that is **bounded below** has a **greatest lower bound**, which is denoted as

$$\inf A, \quad \inf_{x \in A} x, \quad \text{or} \quad \inf\{x : x \in A\}.$$




Corr. A nonempty subset of real numbers that is **bounded below** has a **greatest lower bound**, which is denoted as

$$\inf A, \quad \inf_{x \in A} x, \quad \text{or} \quad \inf\{x : x \in A\}.$$




Corr. A nonempty subset of real numbers that is **bounded below** has a **greatest lower bound**, which is denoted as

$$\inf A, \quad \inf_{x \in A} x, \quad \text{or} \quad \inf\{x : x \in A\}.$$




Corr. A nonempty subset of real numbers that is **bounded below** has a **greatest lower bound**, which is denoted as

$$\inf A, \quad \inf_{x \in A} x, \quad \text{or} \quad \inf\{x : x \in A\}.$$




Corr. A nonempty subset of real numbers that is **bounded below** has a **greatest lower bound**, which is denoted as

$$\inf A, \quad \inf_{x \in A} x, \quad \text{or} \quad \inf\{x : x \in A\}.$$




Corr. A nonempty subset of real numbers that is **bounded below** has a **greatest lower bound**, which is denoted as

$$\inf A, \quad \inf_{x \in A} x, \quad \text{or} \quad \inf\{x : x \in A\}.$$




Corr. A nonempty subset of real numbers that is **bounded below** has a **greatest lower bound**, which is denoted as

$$\inf A, \quad \inf_{x \in A} x, \quad \text{or} \quad \inf\{x : x \in A\}.$$




Corr. A nonempty subset of real numbers that is **bounded below** has a **greatest lower bound**, which is denoted as

$$\inf A, \quad \inf_{x \in A} x, \quad \text{or} \quad \inf\{x : x \in A\}.$$



Corr. A nonempty subset of real numbers that is **bounded below** has a **greatest lower bound**, which is denoted as

$$\inf A, \quad \inf_{x \in A} x, \quad \text{or} \quad \inf\{x : x \in A\}.$$



Let $\{x_n\}_{n \in \mathbb{N}}$ be a sequence and $a \in \mathbb{R}$. Then

$$a \leq \inf_n x_n \iff \forall n, x_n \geq a$$

$$a < \inf_n x_n \iff \forall n, x_n > a$$

$$\inf_n x_n < a \iff \exists n, x_n < a$$

$$\inf_n x_n \leq a \iff \exists n, x_n \leq a$$

E.g. $\sup[0, 1) = 1$ and $\inf[0, 1) = 0$.

\mathbb{N} has no least upper bound, but $\inf \mathbb{N} = 1$.

Let $A = \{x : x^2 < 3\}$. Then

$$\sup_{x \in A} x = \sqrt{3} \quad \text{and} \quad \inf_{x \in A} x = -\sqrt{3}.$$

E.g. $\sup[0, 1) = 1$ and $\inf[0, 1) = 0$.

\mathbb{N} has no least upper bound, but $\inf \mathbb{N} = 1$.

Let $A = \{x : x^2 < 3\}$. Then

$$\sup_{x \in A} x = \sqrt{3} \quad \text{and} \quad \inf_{x \in A} x = -\sqrt{3}.$$

E.g. $\sup[0, 1) = 1$ and $\inf[0, 1) = 0$.

\mathbb{N} has no least upper bound, but $\inf \mathbb{N} = 1$.

Let $A = \{x : x^2 < 3\}$. Then

$$\sup_{x \in A} x = \sqrt{3} \quad \text{and} \quad \inf_{x \in A} x = -\sqrt{3}.$$

Properties

1. Archimedean principle

For each $x \in \mathbb{R}$, there is an $n \in \mathbb{N}$ such that $n > x$.

2. Density of the irrational numbers

Between any two real numbers there is an irrational number.

3. Density of the rational numbers

Between any two real numbers there is a rational number.

Proof As exercises.



Properties

1. Archimedean principle

For each $x \in \mathbb{R}$, there is an $n \in \mathbb{N}$ such that $n > x$.

2. Density of the irrational numbers

Between any two real numbers there is an irrational number.

3. Density of the rational numbers

Between any two real numbers there is a rational number.

Proof As exercises.



Properties

1. Archimedean principle

For each $x \in \mathbb{R}$, there is an $n \in \mathbb{N}$ such that $n > x$.

2. Density of the irrational numbers

Between any two real numbers there is an irrational number.

3. Density of the rational numbers

Between any two real numbers there is a rational number.

Proof As exercises.



Properties

1. Archimedean principle

For each $x \in \mathbb{R}$, there is an $n \in \mathbb{N}$ such that $n > x$.

2. Density of the irrational numbers

Between any two real numbers there is an **irrational** number.

3. Density of the rational numbers

Between any two real numbers there is an rational number.

Proof As exercises.



Properties

1. Archimedean principle

For each $x \in \mathbb{R}$, there is an $n \in \mathbb{N}$ such that $n > x$.

2. Density of the irrational numbers

Between any two real numbers there is an **irrational** number.

3. Density of the rational numbers

Between any two real numbers there is an rational number.

Proof As exercises.



Properties

1. Archimedean principle

For each $x \in \mathbb{R}$, there is an $n \in \mathbb{N}$ such that $n > x$.

2. Density of the irrational numbers

Between any two real numbers there is an **irrational** number.

3. Density of the rational numbers

Between any two real numbers there is an **rational** number.

Proof As exercises.



Properties

1. Archimedean principle

For each $x \in \mathbb{R}$, there is an $n \in \mathbb{N}$ such that $n > x$.

2. Density of the irrational numbers

Between any two real numbers there is an **irrational** number.

3. Density of the rational numbers

Between any two real numbers there is an **rational** number.

Proof As exercises.



Extended Real Number System

Def. The *extended real numbers* $\mathbb{R}^* := \mathbb{R} \cup \{-\infty, \infty\}$.

$x \in \mathbb{R}$	$x + \infty = \infty + x = \infty$	$x - \infty = -\infty + x = -\infty$
$x > 0$	$x \cdot \infty = \infty \cdot x = \infty$	$x \cdot (-\infty) = (-\infty) \cdot x = -\infty$
$x = 0$	$0 \cdot \infty = \infty \cdot 0 = 0$	$0 \cdot (-\infty) = (-\infty) \cdot 0 = 0$
$x < 0$	$x \cdot \infty = \infty \cdot x = -\infty$	$x \cdot (-\infty) = (-\infty) \cdot x = \infty$
$x = \infty$	$\infty + \infty = \infty$ $\infty \cdot \infty = \infty$	$(-\infty) + (-\infty) = -\infty$ $(-\infty) \cdot (-\infty) = \infty$
	$\infty \cdot (-\infty) = (-\infty) \cdot \infty = -\infty$	

$\infty - \infty$ cannot be defined (HW).

Extended Real Number System

Def. The *extended real numbers* $\mathbb{R}^* := \mathbb{R} \cup \{-\infty, \infty\}$.

$x \in \mathbb{R}$	$x + \infty = \infty + x = \infty$	$x - \infty = -\infty + x = -\infty$
$x > 0$	$x \cdot \infty = \infty \cdot x = \infty$	$x \cdot (-\infty) = (-\infty) \cdot x = -\infty$
$x = 0$	$0 \cdot \infty = \infty \cdot 0 = 0$	$0 \cdot (-\infty) = (-\infty) \cdot 0 = 0$
$x < 0$	$x \cdot \infty = \infty \cdot x = -\infty$	$x \cdot (-\infty) = (-\infty) \cdot x = \infty$
$x = \infty$	$\infty + \infty = \infty$ $\infty \cdot \infty = \infty$	$(-\infty) + (-\infty) = -\infty$ $(-\infty) \cdot (-\infty) = \infty$
	$\infty \cdot (-\infty) = (-\infty) \cdot \infty = -\infty$	

$\infty - \infty$ cannot be defined (HW).

Extended Real Number System

Def. The *extended real numbers* $\mathbb{R}^* := \mathbb{R} \cup \{-\infty, \infty\}$.

$x \in \mathbb{R}$	$x + \infty = \infty + x = \infty$	$x - \infty = -\infty + x = -\infty$
$x > 0$	$x \cdot \infty = \infty \cdot x = \infty$	$x \cdot (-\infty) = (-\infty) \cdot x = -\infty$
$x = 0$	$0 \cdot \infty = \infty \cdot 0 = 0$	$0 \cdot (-\infty) = (-\infty) \cdot 0 = 0$
$x < 0$	$x \cdot \infty = \infty \cdot x = -\infty$	$x \cdot (-\infty) = (-\infty) \cdot x = \infty$
$x = \infty$	$\infty + \infty = \infty$ $\infty \cdot \infty = \infty$	$(-\infty) + (-\infty) = -\infty$ $(-\infty) \cdot (-\infty) = \infty$
	$\infty \cdot (-\infty) = (-\infty) \cdot \infty = -\infty$	

$\infty - \infty$ cannot be defined (HW).

Def. Let a and b be extended real numbers such that $a < b$. Then the *intervals on \mathbb{R}^** with *endpoints* a and b are as follows:

$$(a, b) = \{x \in \mathbb{R}^* : a < x < b\}$$

$$[a, b) = \{x \in \mathbb{R}^* : a \leq x < b\}$$

$$(a, b] = \{x \in \mathbb{R}^* : a < x \leq b\}$$

$$[a, b] = \{x \in \mathbb{R}^* : a \leq x \leq b\}$$

If both a and b are in \mathbb{R} , these intervals are the *bounded intervals* in \mathbb{R} .

Otherwise, if either $a = -\infty$ or $b = \infty$, then these intervals are *unbounded intervals*.

Def. Let a and b be extended real numbers such that $a < b$. Then the *intervals on \mathbb{R}^** with *endpoints* a and b are as follows:

$$(a, b) = \{x \in \mathbb{R}^* : a < x < b\}$$

$$[a, b) = \{x \in \mathbb{R}^* : a \leq x < b\}$$

$$(a, b] = \{x \in \mathbb{R}^* : a < x \leq b\}$$

$$[a, b] = \{x \in \mathbb{R}^* : a \leq x \leq b\}$$

If both a and b are in \mathbb{R} , these intervals are the *bounded intervals* in \mathbb{R} .

Otherwise, if either $a = -\infty$ or $b = \infty$, then these intervals are *unbounded intervals*.

Def. Let a and b be extended real numbers such that $a < b$. Then the *intervals on \mathbb{R}^** with *endpoints* a and b are as follows:

$$(a, b) = \{x \in \mathbb{R}^* : a < x < b\}$$

$$[a, b) = \{x \in \mathbb{R}^* : a \leq x < b\}$$

$$(a, b] = \{x \in \mathbb{R}^* : a < x \leq b\}$$

$$[a, b] = \{x \in \mathbb{R}^* : a \leq x \leq b\}$$

If both a and b are in \mathbb{R} , these intervals are the *bounded intervals* in \mathbb{R} .

Otherwise, if either $a = -\infty$ or $b = \infty$, then these intervals are *unbounded intervals*.

Thm Every subset A of \mathbb{R}^* has both a least upper bound and greatest lower bound. Moreover,

- a) If $A = \emptyset$, then $\sup A = -\infty$ and $\inf A = \infty$.
- b) If A is bounded above in \mathbb{R} , then $\sup A \in \mathbb{R}$; otherwise, $\sup A = \infty$.
- c) If A is bounded below in \mathbb{R} , then $\inf A \in \mathbb{R}$; otherwise, $\inf A = -\infty$.

Thm Every subset A of \mathbb{R}^* has both a least upper bound and greatest lower bound. Moreover,

- a) If $A = \emptyset$, then $\sup A = -\infty$ and $\inf A = \infty$.
- b) If A is bounded above in \mathbb{R} , then $\sup A \in \mathbb{R}$; otherwise, $\sup A = \infty$.
- c) If A is bounded below in \mathbb{R} , then $\inf A \in \mathbb{R}$; otherwise, $\inf A = -\infty$.

Thm Every subset A of \mathbb{R}^* has both a least upper bound and greatest lower bound. Moreover,

- a) If $A = \emptyset$, then $\sup A = -\infty$ and $\inf A = \infty$.
- b) If A is bounded above in \mathbb{R} , then $\sup A \in \mathbb{R}$; otherwise, $\sup A = \infty$.
- c) If A is bounded below in \mathbb{R} , then $\inf A \in \mathbb{R}$; otherwise, $\inf A = -\infty$.

Thm Every subset A of \mathbb{R}^* has both a least upper bound and greatest lower bound. Moreover,

- a) If $A = \emptyset$, then $\sup A = -\infty$ and $\inf A = \infty$.
- b) If A is bounded above in \mathbb{R} , then $\sup A \in \mathbb{R}$; otherwise, $\sup A = \infty$.
- c) If A is bounded below in \mathbb{R} , then $\inf A \in \mathbb{R}$; otherwise, $\inf A = -\infty$.

E.g.

a) $\inf \mathbb{N} = 1$ and $\sup \mathbb{N} = \infty$.

b) $\inf \mathbb{Z} = -\infty$ and $\sup \mathbb{Z} = \infty$.

c) If I is an interval in \mathbb{R}^* with endpoints a and b , $a \leq b$. Then $\inf I = a$ and $\sup I = b$.

E.g.

a) $\inf \mathbb{N} = 1$ and $\sup \mathbb{N} = \infty$.

b) $\inf \mathbb{Z} = -\infty$ and $\sup \mathbb{Z} = \infty$.

c) If I is an interval in \mathbb{R}^* with endpoints a and b , $a \leq b$. Then $\inf I = a$ and $\sup I = b$.

E.g.

a) $\inf \mathbb{N} = 1$ and $\sup \mathbb{N} = \infty$.

b) $\inf \mathbb{Z} = -\infty$ and $\sup \mathbb{Z} = \infty$.

c) If I is an interval in \mathbb{R}^* with endpoints a and b , $a \leq b$. Then $\inf I = a$ and $\sup I = b$.

E.g.

- a) $\inf \mathbb{N} = 1$ and $\sup \mathbb{N} = \infty$.
- b) $\inf \mathbb{Z} = -\infty$ and $\sup \mathbb{Z} = \infty$.
- c) If I is an interval in \mathbb{R}^* with endpoints a and b , $a \leq b$. Then $\inf I = a$ and $\sup I = b$.

HW Ex. 2.10 and 2.11 on p. 43.

Chapter 3. Real Number System and Calculus

§ 3.1 Real number system

§ 3.2 Sequences of real numbers

§ 3.3 Open and closed sets

§ 3.4 Real-valued functions

§ 3.5 Liminf and limsup of sets

§ 3.6 Some techniques in calculus

Def. Let $\{x_n\}_{n=1}^{\infty}$ be a sequence of real numbers. We say that the real number $L \in \mathbb{R}$ is the *limit* of this sequence³, namely,

$$\lim_{n \rightarrow \infty} x_n = L$$

if and only if for every real number $\epsilon > 0$, there exists a natural number N such that for all $n > N$, we have $|x_n - L| < \epsilon$.

Def'.

$$\lim_{n \rightarrow \infty} x_n = L \iff \forall \epsilon > 0 \exists N \in \mathbb{N} \forall n \in \mathbb{N} \text{ s.t. } (n \geq N \rightarrow |x_n - L| < \epsilon)$$

³In this case, we say that $\{x_n\}_{n=1}^{\infty}$ is *convergent*. Otherwise, we say that $\{x_n\}_{n=1}^{\infty}$ is *divergent*.

Def. Let $\{x_n\}_{n=1}^{\infty}$ be a sequence of real numbers. We say that the real number $L \in \mathbb{R}$ is the *limit* of this sequence³, namely,

$$\lim_{n \rightarrow \infty} x_n = L$$

if and only if for every real number $\epsilon > 0$, there exists a natural number N such that for all $n > N$, we have $|x_n - L| < \epsilon$.

Def'.

$$\lim_{n \rightarrow \infty} x_n = L \iff \forall \epsilon > 0 \exists N \in \mathbb{N} \forall n \in \mathbb{N} \text{ s.t. } (n \geq N \rightarrow |x_n - L| < \epsilon)$$

³In this case, we say that $\{x_n\}_{n=1}^{\infty}$ is *convergent*. Otherwise, we say that $\{x_n\}_{n=1}^{\infty}$ is *divergent*.

E.g. $\{(n-1)/n\}_{n=1}^{\infty}$ is convergent and $\lim_{n \rightarrow \infty} (n-1)/n = 1$.

$\{(-1)^n\}_{n=1}^{\infty}$ is divergent.

$\{n^2\}_{n=1}^{\infty}$ is divergent.

E.g. $\{(n-1)/n\}_{n=1}^{\infty}$ is convergent and $\lim_{n \rightarrow \infty} (n-1)/n = 1$.

$\{(-1)^n\}_{n=1}^{\infty}$ is divergent.

$\{n^2\}_{n=1}^{\infty}$ is divergent.

E.g. $\{(n-1)/n\}_{n=1}^{\infty}$ is convergent and $\lim_{n \rightarrow \infty} (n-1)/n = 1$.

$\{(-1)^n\}_{n=1}^{\infty}$ is divergent.

$\{n^2\}_{n=1}^{\infty}$ is divergent.

Recall that $\mathbb{R}^* := \mathbb{R} \cup \{-\infty, +\infty\}$ is the *extended real line*.

Def. A sequence of real numbers $\{a_1, a_2, \dots\}$ is said to *converge in* \mathbb{R}^* if one of the following three conditions hold:

- (i) The sequence converges to a finite real number as in the previous definition.
- (ii) For each $M \in \mathbb{R}$, there exists an $N \in \mathbb{N}$ such that for all $n \geq N$, $x_n > M$.
- (iii) For each $M \in \mathbb{R}$, there exists an $N \in \mathbb{N}$ such that for all $n \geq N$, $x_n < M$.

- (i) We say that the sequence converges in \mathbb{R} or the limit exists and is finite.
- (ii) We say the sequence converges to ∞ and write $\lim_{n \rightarrow \infty} x_n = \infty$.
- (iii) We say the sequence converges to $-\infty$ and write $\lim_{n \rightarrow \infty} x_n = -\infty$.

Recall that $\mathbb{R}^* := \mathbb{R} \cup \{-\infty, +\infty\}$ is the *extended real line*.

Def. A sequence of real numbers $\{a_1, a_2, \dots\}$ is said to *converge in \mathbb{R}^** if one of the following three conditions hold:

- (i) The sequence converges to a finite real number as in the previous definition.
- (ii) For each $M \in \mathbb{R}$, there exists an $N \in \mathbb{N}$ such that for all $n \geq N$, $x_n > M$.
- (iii) For each $M \in \mathbb{R}$, there exists an $N \in \mathbb{N}$ such that for all $n \geq N$, $x_n < M$.

- (i) We say that the sequence converges in \mathbb{R} or the limit exists and is finite.
- (ii) We say the sequence converges to ∞ and write $\lim_{n \rightarrow \infty} x_n = \infty$.
- (iii) We say the sequence converges to $-\infty$ and write $\lim_{n \rightarrow \infty} x_n = -\infty$.

Recall that $\mathbb{R}^* := \mathbb{R} \cup \{-\infty, +\infty\}$ is the *extended real line*.

Def. A sequence of real numbers $\{a_1, a_2, \dots\}$ is said to *converge in \mathbb{R}^** if one of the following three conditions hold:

- (i) The sequence converges to a finite real number as in the previous definition.
- (ii) For each $M \in \mathbb{R}$, there exists an $N \in \mathbb{N}$ such that for all $n \geq N$, $x_n > M$.
- (iii) For each $M \in \mathbb{R}$, there exists an $N \in \mathbb{N}$ such that for all $n \geq N$, $x_n < M$.

- (i) We say that the sequence converges in \mathbb{R} or the limit exists and is finite.
- (ii) We say the sequence converges to ∞ and write $\lim_{n \rightarrow \infty} x_n = \infty$.
- (iii) We say the sequence converges to $-\infty$ and write $\lim_{n \rightarrow \infty} x_n = -\infty$.

Recall that $\mathbb{R}^* := \mathbb{R} \cup \{-\infty, +\infty\}$ is the *extended real line*.

Def. A sequence of real numbers $\{a_1, a_2, \dots\}$ is said to *converge in \mathbb{R}^** if one of the following three conditions hold:

- (i) The sequence converges to a finite real number as in the previous definition.
 - (ii) For each $M \in \mathbb{R}$, there exists an $N \in \mathbb{N}$ such that for all $n \geq N$, $x_n > M$.
 - (iii) For each $M \in \mathbb{R}$, there exists an $N \in \mathbb{N}$ such that for all $n \geq N$, $x_n < M$.
-

- (i) We say that the sequence converges in \mathbb{R} or the limit exists and is finite.
- (ii) We say the sequence converges to ∞ and write $\lim_{n \rightarrow \infty} x_n = \infty$.
- (iii) We say the sequence converges to $-\infty$ and write $\lim_{n \rightarrow \infty} x_n = -\infty$.

Recall that $\mathbb{R}^* := \mathbb{R} \cup \{-\infty, +\infty\}$ is the *extended real line*.

Def. A sequence of real numbers $\{a_1, a_2, \dots\}$ is said to *converge in \mathbb{R}^** if one of the following three conditions hold:

- (i) The sequence converges to a finite real number as in the previous definition.
 - (ii) For each $M \in \mathbb{R}$, there exists an $N \in \mathbb{N}$ such that for all $n \geq N$, $x_n > M$.
 - (iii) For each $M \in \mathbb{R}$, there exists an $N \in \mathbb{N}$ such that for all $n \geq N$, $x_n < M$.
-

- (i) We say that the sequence converges in \mathbb{R} or the limit exists and is finite.
- (ii) We say the sequence converges to ∞ and write $\lim_{n \rightarrow \infty} x_n = \infty$.
- (iii) We say the sequence converges to $-\infty$ and write $\lim_{n \rightarrow \infty} x_n = -\infty$.

Recall that $\mathbb{R}^* := \mathbb{R} \cup \{-\infty, +\infty\}$ is the *extended real line*.

Def. A sequence of real numbers $\{a_1, a_2, \dots\}$ is said to *converge in \mathbb{R}^** if one of the following three conditions hold:

- (i) The sequence converges to a finite real number as in the previous definition.
 - (ii) For each $M \in \mathbb{R}$, there exists an $N \in \mathbb{N}$ such that for all $n \geq N$, $x_n > M$.
 - (iii) For each $M \in \mathbb{R}$, there exists an $N \in \mathbb{N}$ such that for all $n \geq N$, $x_n < M$.
-

- (i) We say that the sequence converges in \mathbb{R} or the limit exists and is finite.
- (ii) We say the sequence converges to ∞ and write $\lim_{n \rightarrow \infty} x_n = \infty$.
- (iii) We say the sequence converges to $-\infty$ and write $\lim_{n \rightarrow \infty} x_n = -\infty$.

Recall that $\mathbb{R}^* := \mathbb{R} \cup \{-\infty, +\infty\}$ is the *extended real line*.

Def. A sequence of real numbers $\{a_1, a_2, \dots\}$ is said to *converge in \mathbb{R}^** if one of the following three conditions hold:

- (i) The sequence converges to a finite real number as in the previous definition.
 - (ii) For each $M \in \mathbb{R}$, there exists an $N \in \mathbb{N}$ such that for all $n \geq N$, $x_n > M$.
 - (iii) For each $M \in \mathbb{R}$, there exists an $N \in \mathbb{N}$ such that for all $n \geq N$, $x_n < M$.
-

- (i) We say that the sequence converges in \mathbb{R} or the limit exists and is finite.
- (ii) We say the sequence converges to ∞ and write $\lim_{n \rightarrow \infty} x_n = \infty$.
- (iii) We say the sequence converges to $-\infty$ and write $\lim_{n \rightarrow \infty} x_n = -\infty$.

Recall that $\mathbb{R}^* := \mathbb{R} \cup \{-\infty, +\infty\}$ is the *extended real line*.

Def. A sequence of real numbers $\{a_1, a_2, \dots\}$ is said to *converge in \mathbb{R}^** if one of the following three conditions hold:

- (i) The sequence converges to a finite real number as in the previous definition.
 - (ii) For each $M \in \mathbb{R}$, there exists an $N \in \mathbb{N}$ such that for all $n \geq N$, $x_n > M$.
 - (iii) For each $M \in \mathbb{R}$, there exists an $N \in \mathbb{N}$ such that for all $n \geq N$, $x_n < M$.
-

- (i) We say that the sequence converges in \mathbb{R} or the limit exists and is finite.
- (ii) We say the sequence converges to ∞ and write $\lim_{n \rightarrow \infty} x_n = \infty$.
- (iii) We say the sequence converges to $-\infty$ and write $\lim_{n \rightarrow \infty} x_n = -\infty$.

E.g. $\{(n-1)/n\}_{n=1}^{\infty}$ converges in \mathbb{R} .

$\{(-1)^n\}_{n=1}^{\infty}$ does not converge in \mathbb{R}^* .

$\{n^2\}_{n=1}^{\infty}$ converges in \mathbb{R}^* and $\lim_{n \rightarrow \infty} n^2 = \infty$.

E.g. $\{(n-1)/n\}_{n=1}^{\infty}$ converges in \mathbb{R} .

$\{(-1)^n\}_{n=1}^{\infty}$ does not converge in \mathbb{R}^* .

$\{n^2\}_{n=1}^{\infty}$ converges in \mathbb{R}^* and $\lim_{n \rightarrow \infty} n^2 = \infty$.

E.g. $\{(n-1)/n\}_{n=1}^{\infty}$ converges in \mathbb{R} .

$\{(-1)^n\}_{n=1}^{\infty}$ does not converge in \mathbb{R}^* .

$\{n^2\}_{n=1}^{\infty}$ converges in \mathbb{R}^* and $\lim_{n \rightarrow \infty} n^2 = \infty$.

Monotone sequence

Def. If $x_1 \leq x_2 \leq \dots$, then $\{x_n\}_{n=1}^{\infty}$ is said to be *nondecreasing*.

If $x_1 \geq x_2 \geq \dots$, then $\{x_n\}_{n=1}^{\infty}$ is said to be *nonincreasing*.

$\{x_n\}_{n=1}^{\infty}$ is said to be *monotone* if it is either nondecreasing or nonincreasing.

Monotone sequence

Def. If $x_1 \leq x_2 \leq \dots$, then $\{x_n\}_{n=1}^{\infty}$ is said to be *nondecreasing*.

If $x_1 \geq x_2 \geq \dots$, then $\{x_n\}_{n=1}^{\infty}$ is said to be *nonincreasing*.

$\{x_n\}_{n=1}^{\infty}$ is said to be *monotone* if it is either nondecreasing or nonincreasing.

Monotone sequence

Def. If $x_1 \leq x_2 \leq \dots$, then $\{x_n\}_{n=1}^{\infty}$ is said to be *nondecreasing*.

If $x_1 \geq x_2 \geq \dots$, then $\{x_n\}_{n=1}^{\infty}$ is said to be *nonincreasing*.

$\{x_n\}_{n=1}^{\infty}$ is said to be *monotone* if it is either nondecreasing or nonincreasing.

E.g. $\{(n-1)/n\}_{n=1}^{\infty}$ is monotone and it is nondecreasing.

$\{(-1)^n\}_{n=1}^{\infty}$ is not monotone.

$\{n^2\}_{n=1}^{\infty}$ is monotone and it is nondecreasing.

E.g. $\{(n-1)/n\}_{n=1}^{\infty}$ is monotone and it is nondecreasing.

$\{(-1)^n\}_{n=1}^{\infty}$ is not monotone.

$\{n^2\}_{n=1}^{\infty}$ is monotone and it is nondecreasing.

E.g. $\{(n-1)/n\}_{n=1}^{\infty}$ is monotone and it is nondecreasing.

$\{(-1)^n\}_{n=1}^{\infty}$ is not monotone.

$\{n^2\}_{n=1}^{\infty}$ is monotone and it is nondecreasing.

Thm Any monotone sequence $\{x_n\}_{n=1}^{\infty}$ of real numbers converges in \mathbb{R}^* .

Moreover, we have the following:

a) If $\{x_n\}_{n=1}^{\infty}$ is nondecreasing, then

$$\lim_{n \rightarrow \infty} x_n = \sup\{x_n : n \in \mathbb{N}\}.$$

In particular, $\{x_n\}_{n=1}^{\infty}$ converges in \mathbb{R} if it is bounded above and is ∞ otherwise.

b) If $\{x_n\}_{n=1}^{\infty}$ is nonincreasing, then

$$\lim_{n \rightarrow \infty} x_n = \inf\{x_n : n \in \mathbb{N}\}.$$

In particular, $\{x_n\}_{n=1}^{\infty}$ converges in \mathbb{R} if it is bounded below and is $-\infty$ otherwise.

Thm Any monotone sequence $\{x_n\}_{n=1}^{\infty}$ of real numbers converges in \mathbb{R}^* .

Moreover, we have the following:

a) If $\{x_n\}_{n=1}^{\infty}$ is nondecreasing, then

$$\lim_{n \rightarrow \infty} x_n = \sup\{x_n : n \in \mathbb{N}\}.$$

In particular, $\{x_n\}_{n=1}^{\infty}$ converges in \mathbb{R} if it is bounded above and is ∞ otherwise.

b) If $\{x_n\}_{n=1}^{\infty}$ is nonincreasing, then

$$\lim_{n \rightarrow \infty} x_n = \inf\{x_n : n \in \mathbb{N}\}.$$

In particular, $\{x_n\}_{n=1}^{\infty}$ converges in \mathbb{R} if it is bounded below and is $-\infty$ otherwise.

Thm Any monotone sequence $\{x_n\}_{n=1}^{\infty}$ of real numbers converges in \mathbb{R}^* .

Moreover, we have the following:

a) If $\{x_n\}_{n=1}^{\infty}$ is nondecreasing, then

$$\lim_{n \rightarrow \infty} x_n = \sup\{x_n : n \in \mathbb{N}\}.$$

In particular, $\{x_n\}_{n=1}^{\infty}$ converges in \mathbb{R} if it is bounded above and is ∞ otherwise.

b) If $\{x_n\}_{n=1}^{\infty}$ is nonincreasing, then

$$\lim_{n \rightarrow \infty} x_n = \inf\{x_n : n \in \mathbb{N}\}.$$

In particular, $\{x_n\}_{n=1}^{\infty}$ converges in \mathbb{R} if it is bounded below and is $-\infty$ otherwise.

Thm Any monotone sequence $\{x_n\}_{n=1}^{\infty}$ of real numbers converges in \mathbb{R}^* .

Moreover, we have the following:

a) If $\{x_n\}_{n=1}^{\infty}$ is nondecreasing, then

$$\lim_{n \rightarrow \infty} x_n = \sup\{x_n : n \in \mathbb{N}\}.$$

In particular, $\{x_n\}_{n=1}^{\infty}$ converges in \mathbb{R} if it is **bounded above** and is ∞ otherwise.

b) If $\{x_n\}_{n=1}^{\infty}$ is nonincreasing, then

$$\lim_{n \rightarrow \infty} x_n = \inf\{x_n : n \in \mathbb{N}\}.$$

In particular, $\{x_n\}_{n=1}^{\infty}$ converges in \mathbb{R} if it is bounded below and is $-\infty$ otherwise.

Thm Any monotone sequence $\{x_n\}_{n=1}^{\infty}$ of real numbers converges in \mathbb{R}^* .

Moreover, we have the following:

a) If $\{x_n\}_{n=1}^{\infty}$ is nondecreasing, then

$$\lim_{n \rightarrow \infty} x_n = \sup\{x_n : n \in \mathbb{N}\}.$$

In particular, $\{x_n\}_{n=1}^{\infty}$ converges in \mathbb{R} if it is **bounded above** and is ∞ otherwise.

b) If $\{x_n\}_{n=1}^{\infty}$ is nonincreasing, then

$$\lim_{n \rightarrow \infty} x_n = \inf\{x_n : n \in \mathbb{N}\}.$$

In particular, $\{x_n\}_{n=1}^{\infty}$ converges in \mathbb{R} if it is bounded below and is $-\infty$ otherwise.

Thm Any monotone sequence $\{x_n\}_{n=1}^{\infty}$ of real numbers converges in \mathbb{R}^* .

Moreover, we have the following:

a) If $\{x_n\}_{n=1}^{\infty}$ is nondecreasing, then

$$\lim_{n \rightarrow \infty} x_n = \sup\{x_n : n \in \mathbb{N}\}.$$

In particular, $\{x_n\}_{n=1}^{\infty}$ converges in \mathbb{R} if it is **bounded above** and is ∞ otherwise.

b) If $\{x_n\}_{n=1}^{\infty}$ is nonincreasing, then

$$\lim_{n \rightarrow \infty} x_n = \inf\{x_n : n \in \mathbb{N}\}.$$

In particular, $\{x_n\}_{n=1}^{\infty}$ converges in \mathbb{R} if it is **bounded below** and is $-\infty$ otherwise.

Proof. We will prove the case when $\{x_n\}_{n=1}^{\infty}$ is nondecreasing. The nonincreasing case can be proved in a similar way.

As $\sup_n x_n$ always exists in \mathbb{R}^* , we need to consider two cases:

Case I: $\sup_n x_n \in \mathbb{R}$

Case II: $\sup_n x_n = \infty$

Let's prove Case I here. Let $x = \sup_n x_n$. In order to show that $\lim_n x_n = x$, by monotonicity, we need to show that

$$\forall \epsilon > 0 \exists N \forall n \text{ s.t. } (n \geq N) \rightarrow (x - a_n \leq \epsilon).$$

Proof. We will prove the case when $\{x_n\}_{n=1}^\infty$ is nondecreasing. The nonincreasing case can be proved in a similar way.

As $\sup_n x_n$ always exists in \mathbb{R}^* , we need to consider two cases:

Case I: $\sup_n x_n \in \mathbb{R}$

Case II: $\sup_n x_n = \infty$

Let's prove Case I here. Let $x = \sup_n x_n$. In order to show that $\lim_n x_n = x$, by monotonicity, we need to show that

$$\forall \epsilon > 0 \exists N \forall n \text{ s.t. } (n \geq N) \rightarrow (x - a_n \leq \epsilon).$$

Proof. We will prove the case when $\{x_n\}_{n=1}^{\infty}$ is nondecreasing. The nonincreasing case can be proved in a similar way.

As $\sup_n x_n$ always exists in \mathbb{R}^* , we need to consider two cases:

Case I: $\sup_n x_n \in \mathbb{R}$

Case II: $\sup_n x_n = \infty$

Let's prove Case I here. Let $x = \sup_n x_n$. In order to show that $\lim_n x_n = x$, by monotonicity, we need to show that

$$\forall \epsilon > 0 \exists N \forall n \text{ s.t. } (n \geq N) \rightarrow (x - a_n \leq \epsilon).$$

Proof. We will prove the case when $\{x_n\}_{n=1}^{\infty}$ is nondecreasing. The nonincreasing case can be proved in a similar way.

As $\sup_n x_n$ always exists in \mathbb{R}^* , we need to consider two cases:

Case I: $\sup_n x_n \in \mathbb{R}$

Case II: $\sup_n x_n = \infty$

Let's prove Case I here. Let $x = \sup_n x_n$. In order to show that $\lim_n x_n = x$, by monotonicity, we need to show that

$$\forall \epsilon > 0 \exists N \forall n \text{ s.t. } (n \geq N) \rightarrow (x - a_n \leq \epsilon).$$

Proof. We will prove the case when $\{x_n\}_{n=1}^{\infty}$ is nondecreasing. The nonincreasing case can be proved in a similar way.

As $\sup_n x_n$ always exists in \mathbb{R}^* , we need to consider two cases:

Case I: $\sup_n x_n \in \mathbb{R}$

Case II: $\sup_n x_n = \infty$

Let's prove Case I here. Let $x = \sup_n x_n$. In order to show that $\lim_n x_n = x$, by monotonicity, we need to show that

$$\forall \epsilon > 0 \exists N \forall n \text{ s.t. } (n \geq N) \rightarrow (x - a_n \leq \epsilon).$$

Proof. We will prove the case when $\{x_n\}_{n=1}^{\infty}$ is nondecreasing. The nonincreasing case can be proved in a similar way.

As $\sup_n x_n$ always exists in \mathbb{R}^* , we need to consider two cases:

Case I: $\sup_n x_n \in \mathbb{R}$

Case II: $\sup_n x_n = \infty$

Let's prove Case I here. Let $x = \sup_n x_n$. In order to show that $\lim_n x_n = x$, by monotonicity, we need to show that

$$\forall \epsilon > 0 \exists N \forall n \text{ s.t. } (n \geq N) \rightarrow (x - a_n \leq \epsilon).$$

Proof. (Continued) Otherwise,

$$\exists \epsilon > 0 \forall N \exists n \text{ s.t. } (n \geq N) \wedge (x - a_n > \epsilon).$$

Hence, there are infinitely many terms falling below $x - \epsilon$.

Since $\{x_n\}$ is nondecreasing, this implies all a_n fall below $x - \epsilon$, i.e.,

$$a_n < x - \epsilon, \quad \text{for all } n \geq 1.$$

which is equivalent to $\sup_n x_n < x - \epsilon$. This contradicts with the fact that $\sup_n x_n = x$.

Therefore,

$$\lim_n x_n = x = \sup_n x_n.$$

□

Proof. (Continued) Otherwise,

$$\exists \epsilon > 0 \forall N \exists n \text{ s.t. } (n \geq N) \wedge (x - a_n > \epsilon).$$

Hence, there are infinitely many terms falling below $x - \epsilon$.

Since $\{x_n\}$ is nondecreasing, this implies all a_n fall below $x - \epsilon$, i.e.,

$$a_n < x - \epsilon, \quad \text{for all } n \geq 1.$$

which is equivalent to $\sup_n x_n < x - \epsilon$. This contradicts with the fact that $\sup_n x_n = x$.

Therefore,

$$\lim_n x_n = x = \sup_n x_n.$$

□

Proof. (Continued) Otherwise,

$$\exists \epsilon > 0 \forall N \exists n \text{ s.t. } (n \geq N) \wedge (x - a_n > \epsilon).$$

Hence, there are infinitely many terms falling below $x - \epsilon$.

Since $\{x_n\}$ is nondecreasing, this implies all a_n fall below $x - \epsilon$, i.e.,

$$a_n < x - \epsilon, \quad \text{for all } n \geq 1.$$

which is equivalent to $\sup_n x_n < x - \epsilon$. This contradicts with the fact that $\sup_n x_n = x$.

Therefore,

$$\lim_n x_n = x = \sup_n x_n.$$

□

Proof. (Continued) Otherwise,

$$\exists \epsilon > 0 \forall N \exists n \text{ s.t. } (n \geq N) \wedge (x - a_n > \epsilon).$$

Hence, there are infinitely many terms falling below $x - \epsilon$.

Since $\{x_n\}$ is nondecreasing, this implies all a_n fall below $x - \epsilon$, i.e.,

$$a_n < x - \epsilon, \quad \text{for all } n \geq 1.$$

which is equivalent to $\sup_n x_n < x - \epsilon$. This contradicts with the fact that $\sup_n x_n = x$.

Therefore,

$$\lim_n x_n = x = \sup_n x_n.$$

□

E.g. $\{(n-1)/n\}_{n=1}^{\infty}$ is nondecreasing and converges in \mathbb{R} . It is bounded above.

$\{(-1)^n\}_{n=1}^{\infty}$ is not monotone.

$\{n^2\}_{n=1}^{\infty}$ is nondecreasing, does not converge in \mathbb{R} , converges in \mathbb{R}^* .

E.g. $\{(n-1)/n\}_{n=1}^{\infty}$ is nondecreasing and converges in \mathbb{R} . It is bounded above.

$\{(-1)^n\}_{n=1}^{\infty}$ is not monotone.

$\{n^2\}_{n=1}^{\infty}$ is nondecreasing, does not converge in \mathbb{R} , converges in \mathbb{R}^* .

E.g. $\{(n-1)/n\}_{n=1}^{\infty}$ is nondecreasing and converges in \mathbb{R} . It is bounded above.

$\{(-1)^n\}_{n=1}^{\infty}$ is not monotone.

$\{n^2\}_{n=1}^{\infty}$ is nondecreasing, does not converge in \mathbb{R} , converges in \mathbb{R}^* .

Cluster points

Def. Let $\{x_n\}_{n=1}^{\infty}$ be a sequence of real numbers.

- a) A real number x is said to be a *cluster point* of $\{x_n\}_{n=1}^{\infty}$ if for each $\epsilon > 0$ and $N \in \mathbb{N}$, there exists an $n \geq N$ such that $|x - x_n| < \epsilon$.
- b) ∞ is a *cluster point* of $\{x_n\}_{n=1}^{\infty}$ if for each $M \in \mathbb{R}$ and $N \in \mathbb{N}$, there exists an $n \geq N$ such that $x_n > M$.
- c) $-\infty$ is a *cluster point* of $\{x_n\}_{n=1}^{\infty}$ if for each $M \in \mathbb{R}$ and $N \in \mathbb{N}$, there exists an $n \geq N$ such that $x_n < M$.

- a) $x \in \mathbb{R}$ is a cluster point of $\{x_n\}$ if

$$\forall \epsilon \forall N \exists n \quad (n \geq N) \rightarrow (|x - x_n| < \epsilon). \quad (1)$$

Cluster points

Def. Let $\{x_n\}_{n=1}^{\infty}$ be a sequence of real numbers.

- a) A real number x is said to be a **cluster point** of $\{x_n\}_{n=1}^{\infty}$ if for each $\epsilon > 0$ and $N \in \mathbb{N}$, there exists an $n \geq N$ such that $|x - x_n| < \epsilon$.
- b) ∞ is a *cluster point* of $\{x_n\}_{n=1}^{\infty}$ if for each $M \in \mathbb{R}$ and $N \in \mathbb{N}$, there exists an $n \geq N$ such that $x_n > M$.
- c) $-\infty$ is a *cluster point* of $\{x_n\}_{n=1}^{\infty}$ if for each $M \in \mathbb{R}$ and $N \in \mathbb{N}$, there exists an $n \geq N$ such that $x_n < M$.

a) $x \in \mathbb{R}$ is a cluster point of $\{x_n\}$ if

$$\forall \epsilon \forall N \exists n \quad (n \geq N) \rightarrow (|x - x_n| < \epsilon). \quad (1)$$

Cluster points

Def. Let $\{x_n\}_{n=1}^{\infty}$ be a sequence of real numbers.

- a) A real number x is said to be a **cluster point** of $\{x_n\}_{n=1}^{\infty}$ if for each $\epsilon > 0$ and $N \in \mathbb{N}$, there exists an $n \geq N$ such that $|x - x_n| < \epsilon$.
- b) ∞ **is a cluster point** of $\{x_n\}_{n=1}^{\infty}$ if for each $M \in \mathbb{R}$ and $N \in \mathbb{N}$, there exists an $n \geq N$ such that $x_n > M$.
- c) $-\infty$ **is a cluster point** of $\{x_n\}_{n=1}^{\infty}$ if for each $M \in \mathbb{R}$ and $N \in \mathbb{N}$, there exists an $n \geq N$ such that $x_n < M$.

- a) $x \in \mathbb{R}$ is a cluster point of $\{x_n\}$ if

$$\forall \epsilon \forall N \exists n \quad (n \geq N) \rightarrow (|x - x_n| < \epsilon). \quad (1)$$

Cluster points

Def. Let $\{x_n\}_{n=1}^{\infty}$ be a sequence of real numbers.

- a) A real number x is said to be a **cluster point** of $\{x_n\}_{n=1}^{\infty}$ if for each $\epsilon > 0$ and $N \in \mathbb{N}$, there exists an $n \geq N$ such that $|x - x_n| < \epsilon$.
 - b) ∞ **is a cluster point** of $\{x_n\}_{n=1}^{\infty}$ if for each $M \in \mathbb{R}$ and $N \in \mathbb{N}$, there exists an $n \geq N$ such that $x_n > M$.
 - c) $-\infty$ **is a cluster point** of $\{x_n\}_{n=1}^{\infty}$ if for each $M \in \mathbb{R}$ and $N \in \mathbb{N}$, there exists an $n \geq N$ such that $x_n < M$.
-

a) $x \in \mathbb{R}$ is a cluster point of $\{x_n\}$ if

$$\forall \epsilon \forall N \exists n \quad (n \geq N) \rightarrow (|x - x_n| < \epsilon). \quad (1)$$

Cluster points

Def. Let $\{x_n\}_{n=1}^{\infty}$ be a sequence of real numbers.

- a) A real number x is said to be a **cluster point** of $\{x_n\}_{n=1}^{\infty}$ if for each $\epsilon > 0$ and $N \in \mathbb{N}$, there exists an $n \geq N$ such that $|x - x_n| < \epsilon$.
 - b) ∞ **is a cluster point** of $\{x_n\}_{n=1}^{\infty}$ if for each $M \in \mathbb{R}$ and $N \in \mathbb{N}$, there exists an $n \geq N$ such that $x_n > M$.
 - c) $-\infty$ **is a cluster point** of $\{x_n\}_{n=1}^{\infty}$ if for each $M \in \mathbb{R}$ and $N \in \mathbb{N}$, there exists an $n \geq N$ such that $x_n < M$.
-

a) $x \in \mathbb{R}$ is a cluster point of $\{x_n\}$ if

$$\forall \epsilon \forall N \exists n \quad (n \geq N) \rightarrow (|x - x_n| < \epsilon). \quad (1)$$

Cluster points

Def. Let $\{x_n\}_{n=1}^{\infty}$ be a sequence of real numbers.

- a) A real number x is said to be a **cluster point** of $\{x_n\}_{n=1}^{\infty}$ if for each $\epsilon > 0$ and $N \in \mathbb{N}$, there exists an $n \geq N$ such that $|x - x_n| < \epsilon$.
 - b) ∞ **is a cluster point** of $\{x_n\}_{n=1}^{\infty}$ if for each $M \in \mathbb{R}$ and $N \in \mathbb{N}$, there exists an $n \geq N$ such that $x_n > M$.
 - c) $-\infty$ **is a cluster point** of $\{x_n\}_{n=1}^{\infty}$ if for each $M \in \mathbb{R}$ and $N \in \mathbb{N}$, there exists an $n \geq N$ such that $x_n < M$.
-

a) $x \in \mathbb{R}$ is a cluster point of $\{x_n\}$ if

$$\forall \epsilon \forall N \exists n \quad (n \geq N) \rightarrow (|x - x_n| < \epsilon). \quad (1)$$

E.g.1 $\{(n-1)/n\}_{n=1}^{\infty}$ has one cluster point: 1.

$\{(-1)^n\}_{n=1}^{\infty}$ has two cluster points: -1 and $+1$.

$\{n^2\}_{n=1}^{\infty}$ has one cluster point: $+\infty$.

E.g.1 $\{(n-1)/n\}_{n=1}^{\infty}$ has one cluster point: 1.

$\{(-1)^n\}_{n=1}^{\infty}$ has two cluster points: -1 and $+1$.

$\{n^2\}_{n=1}^{\infty}$ has one cluster point: $+\infty$.

E.g.1 $\{(n-1)/n\}_{n=1}^{\infty}$ has one cluster point: 1.

$\{(-1)^n\}_{n=1}^{\infty}$ has two cluster points: -1 and $+1$.

$\{n^2\}_{n=1}^{\infty}$ has one cluster point: $+\infty$.

E.g.2 Consider the sequence $\{2, 1, 0, 2, 2, \frac{1}{2}, 2, 3, \frac{2}{3}, 2, 4, \frac{3}{4}, \dots\}$, that is,

$$x_n = \begin{cases} (n-3)/n & \text{if } n \equiv 0 \pmod{3} \\ 2 & \text{if } n \equiv 1 \pmod{3} \\ (n+1)/3 & \text{if } n \equiv 2 \pmod{3} \end{cases}$$

It has three cluster points: 1, 2, ∞ .

E.g.3 Let $\{r_n\}_{n=1}^{\infty}$ be an enumeration of the rational numbers. By the density of the rational numbers, every extended real number is a cluster point of the sequence $\{r_n\}_{n=1}^{\infty}$.

Thm A convergent sequence has exactly one cluster point, namely, its limit.
Thus, a sequence having more than one cluster point cannot converge.

Proof Suppose that $\{x_n\}$ is a convergent sequence and let x be its limit, namely, $\lim_{n \rightarrow \infty} x_n = x$. We need to prove:

- (1) x is a cluster point.
- (2) x is the only cluster point of $\{x_n\}$.

We also need to consider two cases:

Case I: $x \in \mathbb{R}$.

Case II: $x = \infty$ or $-\infty$.

We will focus on Case I only.

Now we first prove (1).

Thm A convergent sequence has exactly one cluster point, namely, its limit.
Thus, a sequence having more than one cluster point cannot converge.

Proof Suppose that $\{x_n\}$ is a convergent sequence and let x be its limit, namely, $\lim_{n \rightarrow \infty} x_n = x$. We need to prove:

- (1) x is a cluster point.
- (2) x is the only cluster point of $\{x_n\}$.

We also need to consider two cases:

Case I: $x \in \mathbb{R}$.

Case II: $x = \infty$ or $-\infty$.

We will focus on Case I only.

Now we first prove (1).

Thm A convergent sequence has exactly one cluster point, namely, its limit.
Thus, a sequence having more than one cluster point cannot converge.

Proof Suppose that $\{x_n\}$ is a convergent sequence and let x be its limit, namely, $\lim_{n \rightarrow \infty} x_n = x$. We need to prove:

- (1) x is a cluster point.
- (2) x is the only cluster point of $\{x_n\}$.

We also need to consider two cases:

Case I: $x \in \mathbb{R}$.

Case II: $x = \infty$ or $-\infty$.

We will focus on Case I only.

Now we first prove (1).

Thm A convergent sequence has exactly one cluster point, namely, its limit.
Thus, a sequence having more than one cluster point cannot converge.

Proof Suppose that $\{x_n\}$ is a convergent sequence and let x be its limit, namely, $\lim_{n \rightarrow \infty} x_n = x$. We need to prove:

- (1) x is a cluster point.
- (2) x is the only cluster point of $\{x_n\}$.

We also need to consider two cases:

Case I: $x \in \mathbb{R}$.

Case II: $x = \infty$ or $-\infty$.

We will focus on Case I only.

Now we first prove (1).

Thm A convergent sequence has exactly one cluster point, namely, its limit.
Thus, a sequence having more than one cluster point cannot converge.

Proof Suppose that $\{x_n\}$ is a convergent sequence and let x be its limit, namely, $\lim_{n \rightarrow \infty} x_n = x$. We need to prove:

- (1) x is a cluster point.
- (2) x is the only cluster point of $\{x_n\}$.

We also need to consider two cases:

Case I: $x \in \mathbb{R}$.

Case II: $x = \infty$ or $-\infty$.

We will focus on Case I only.

Now we first prove (1).

Thm A convergent sequence has exactly one cluster point, namely, its limit.
Thus, a sequence having more than one cluster point cannot converge.

Proof Suppose that $\{x_n\}$ is a convergent sequence and let x be its limit, namely, $\lim_{n \rightarrow \infty} x_n = x$. We need to prove:

- (1) x is a cluster point.
- (2) x is the only cluster point of $\{x_n\}$.

We also need to consider two cases:

Case I: $x \in \mathbb{R}$.

Case II: $x = \infty$ or $-\infty$.

We will focus on Case I only.

Now we first prove (1).

Thm A convergent sequence has exactly one cluster point, namely, its limit.
Thus, a sequence having more than one cluster point cannot converge.

Proof Suppose that $\{x_n\}$ is a convergent sequence and let x be its limit, namely, $\lim_{n \rightarrow \infty} x_n = x$. We need to prove:

- (1) x is a cluster point.
- (2) x is the only cluster point of $\{x_n\}$.

We also need to consider two cases:

Case I: $x \in \mathbb{R}$.

Case II: $x = \infty$ or $-\infty$.

We will focus on Case I only.

Now we first prove (1).

Thm A convergent sequence has exactly one cluster point, namely, its limit.
Thus, a sequence having more than one cluster point cannot converge.

Proof Suppose that $\{x_n\}$ is a convergent sequence and let x be its limit, namely, $\lim_{n \rightarrow \infty} x_n = x$. We need to prove:

- (1) x is a cluster point.
- (2) x is the only cluster point of $\{x_n\}$.

We also need to consider two cases:

Case I: $x \in \mathbb{R}$.

Case II: $x = \infty$ or $-\infty$.

We will focus on Case I only.

Now we first prove (1).

Thm A convergent sequence has exactly one cluster point, namely, its limit.
Thus, a sequence having more than one cluster point cannot converge.

Proof Suppose that $\{x_n\}$ is a convergent sequence and let x be its limit, namely, $\lim_{n \rightarrow \infty} x_n = x$. We need to prove:

- (1) x is a cluster point.
- (2) x is the only cluster point of $\{x_n\}$.

We also need to consider two cases:

Case I: $x \in \mathbb{R}$.

Case II: $x = \infty$ or $-\infty$.

We will focus on Case I only.

Now we first prove (1).

	Once for all rest
	$\underbrace{\hspace{1cm}}$
$\lim_n x_n = x$	$\forall \epsilon \exists N \forall n \quad (n \geq N) \rightarrow (x_n - x < \epsilon)$
$\Downarrow ??$	
x is a cluster point of $\{x_n\}$	$\forall \epsilon \forall \tilde{N} \exists \tilde{n} \quad (\tilde{n} \geq \tilde{N}) \wedge (x_{\tilde{n}} - x < \epsilon)$
	$\underbrace{\hspace{1cm}}$
	Infinite games

(1) is proved by choosing any $\tilde{n} \geq \max \{ \tilde{N}, N \}$

	Once for all rest
	$\underbrace{\hspace{1cm}}$
$\lim_n x_n = x$	$\forall \epsilon \exists N \forall n \quad (n \geq N) \rightarrow (x_n - x < \epsilon)$
$\Downarrow ??$	
x is a cluster point of $\{x_n\}$	$\forall \epsilon \forall \tilde{N} \exists \tilde{n} \quad (\tilde{n} \geq \tilde{N}) \wedge (x_{\tilde{n}} - x < \epsilon)$
	$\underbrace{\hspace{1cm}}$
	Infinite games

(1) is proved by choosing any $\tilde{n} \geq \max \{ \tilde{N}, N \}$

	Once for all rest
	$\underbrace{\hspace{1.5cm}}$
$\lim_n x_n = x$	$\forall \epsilon \exists N \forall n \quad (n \geq N) \rightarrow (x_n - x < \epsilon)$
$\Downarrow ??$	
x is a cluster point of $\{x_n\}$	$\forall \epsilon \forall \tilde{N} \exists \tilde{n} \quad (\tilde{n} \geq \tilde{N}) \wedge (x_{\tilde{n}} - x < \epsilon)$
	$\underbrace{\hspace{1.5cm}}$
	Infinite games

(1) is proved by choosing any $\tilde{n} \geq \max \{ \tilde{N}, N \}$

	Once for all rest
	\frown
$\lim_n x_n = x$	$\forall \epsilon \exists N \forall n \quad (n \geq N) \rightarrow (x_n - x < \epsilon)$
$\Downarrow ??$	
x is a cluster point of $\{x_n\}$	$\forall \epsilon \forall \tilde{N} \exists \tilde{n} \quad (\tilde{n} \geq \tilde{N}) \wedge (x_{\tilde{n}} - x < \epsilon)$
	\smile
	Infinite games

(1) is proved by choosing any $\tilde{n} \geq \max\{\tilde{N}, N\}$


	Once for all rest
	$\underbrace{\hspace{1cm}}$
$\lim_n x_n = x$	$\forall \epsilon \exists N \forall n \quad (n \geq N) \rightarrow (x_n - x < \epsilon)$
$\Downarrow ??$	
x is a cluster point of $\{x_n\}$	$\forall \epsilon \forall \tilde{N} \exists \tilde{n} \quad (\tilde{n} \geq \tilde{N}) \wedge (x_{\tilde{n}} - x < \epsilon)$
	$\underbrace{\hspace{1cm}}$
	Infinite games

(1) is proved by choosing any $\tilde{n} \geq \max\{\tilde{N}, N\}$

	Once for all rest
	$\underbrace{\hspace{1cm}}$
$\lim_n x_n = x$	$\forall \epsilon \exists N \forall n \ (n \geq N) \rightarrow (x_n - x < \epsilon)$
$\Downarrow ??$	
x is a cluster point of $\{x_n\}$	$\forall \epsilon \forall \tilde{N} \exists \tilde{n} \ (\tilde{n} \geq \tilde{N}) \wedge (x_{\tilde{n}} - x < \epsilon)$
	$\underbrace{\hspace{1cm}}$
	Infinite games

(1) is proved by choosing any $\tilde{n} \geq \max(\tilde{N}, N)$.

As for (2), suppose y is another cluster point.



By choosing any $\epsilon < |x - y|/2$, we see that


1. In the ϵ -neighborhood of y , there are infinitely many terms.
2. In the ϵ -neighborhood of x , all but finite many terms are here.

Contradiction!

Therefore, there exists only one cluster point.

□

As for (2), suppose y is another cluster point.



By choosing any $\epsilon < |x - y|/2$, we see that


1. In the ϵ -neighborhood of y , there are infinitely many terms.
2. In the ϵ -neighborhood of x , all but finite many terms are here.

Contradiction!

Therefore, there exists only one cluster point.

□

As for (2), suppose y is another cluster point.



By choosing any $\epsilon < |x - y|/2$, we see that


1. In the ϵ -neighborhood of y , there are infinitely many terms.
2. In the ϵ -neighborhood of x , all but finite many terms are here.

Contradiction!

Therefore, there exists only one cluster point.

□

As for (2), suppose y is another cluster point.



By choosing any $\epsilon < |x - y|/2$, we see that


1. In the ϵ -neighborhood of y , there are infinitely many terms.
2. In the ϵ -neighborhood of x , all but finite many terms are here.

Contradiction!

Therefore, there exists only one cluster point.

□

As for (2), suppose y is another cluster point.



By choosing any $\epsilon < |x - y|/2$, we see that


1. In the ϵ -neighborhood of y , there are infinitely many terms.
2. In the ϵ -neighborhood of x , all but finite many terms are here.

Contradiction!

Therefore, there exists only one cluster point.

□

As for (2), suppose y is another cluster point.



By choosing any $\epsilon < |x - y|/2$, we see that

1. In the ϵ -neighborhood of y , there are infinitely many terms.
2. In the ϵ -neighborhood of x , all but finite many terms are here.

Contradiction!

Therefore, there exists only one cluster point.

□

A few more properties

1. A sequence is convergent iff each subsequence is convergent.
2. Sandwich theorem: If $x_n \leq c_n \leq b_n$ for all $n > N$ and $x_n \rightarrow L$ and $b_n \rightarrow L$, then $c_n \rightarrow L$.

A few more properties

1. A sequence is convergent iff each subsequence is convergent.
2. **Sandwich theorem:** If $x_n \leq c_n \leq b_n$ for all $n > N$ and $x_n \rightarrow L$ and $b_n \rightarrow L$, then $c_n \rightarrow L$.

Limit superior and limit inferior

Def. The *limit inferior* of a sequence $\{x_n\}_{n=1}^{\infty}$ is defined as

$$\liminf_{n \rightarrow \infty} x_n := \sup_n \left(\inf_{m \geq n} x_m \right) \in \mathbb{R}^*.$$

The *limit superior* of $\{x_n\}_{n=1}^{\infty}$ is defined as

$$\limsup_{n \rightarrow \infty} x_n := \inf_n \left(\sup_{m \geq n} x_m \right) \in \mathbb{R}^*.$$

Limit superior and limit inferior

Def. The *limit inferior* of a sequence $\{x_n\}_{n=1}^{\infty}$ is defined as

$$\liminf_{n \rightarrow \infty} x_n := \sup_n \left(\inf_{m \geq n} x_m \right) \in \mathbb{R}^*.$$

The *limit superior* of $\{x_n\}_{n=1}^{\infty}$ is defined as

$$\limsup_{n \rightarrow \infty} x_n := \inf_n \left(\sup_{m \geq n} a_m \right) \in \mathbb{R}^*.$$

Remark Since the sequences $\{y_n\}_{n=1}^{\infty}$ and $\{z_n\}_{n=1}^{\infty}$ defined as

$$y_n := \sup_{m \geq n} x_m \quad \text{and} \quad z_n := \inf_{m \geq n} x_m,$$

are, respectively, nonincreasing and nondecreasing, we see that

$$\begin{aligned} \inf_n y_n &= \inf_n \sup_{m \geq n} x_m = \limsup_{x \rightarrow \infty} x_n \\ \sup_n z_n &= \sup_n \inf_{m \geq n} x_m = \liminf_{x \rightarrow \infty} x_n. \end{aligned}$$

Hence,

$$\limsup_{n \rightarrow \infty} x_n = \lim_{n \rightarrow \infty} \sup_{m \geq n} x_m \quad \text{and} \quad \liminf_{n \rightarrow \infty} x_n = \lim_{n \rightarrow \infty} \inf_{m \geq n} x_m.$$

Remark Since the sequences $\{y_n\}_{n=1}^{\infty}$ and $\{z_n\}_{n=1}^{\infty}$ defined as

$$y_n := \sup_{m \geq n} x_m \quad \text{and} \quad z_n := \inf_{m \geq n} x_m,$$

are, respectively, nonincreasing and nondecreasing, we see that

$$\begin{aligned} \inf_n y_n &= \inf_n \sup_{m \geq n} x_m = \limsup_{x \rightarrow \infty} x_n \\ \sup_n z_n &= \sup_n \inf_{m \geq n} x_m = \liminf_{x \rightarrow \infty} x_n. \end{aligned}$$

Hence,

$$\limsup_{n \rightarrow \infty} x_n = \lim_{n \rightarrow \infty} \sup_{m \geq n} x_m \quad \text{and} \quad \liminf_{n \rightarrow \infty} x_n = \lim_{n \rightarrow \infty} \inf_{m \geq n} x_m.$$

Remark Since the sequences $\{y_n\}_{n=1}^{\infty}$ and $\{z_n\}_{n=1}^{\infty}$ defined as


$$y_n := \sup_{m \geq n} x_m \quad \text{and} \quad z_n := \inf_{m \geq n} x_m,$$

are, respectively, nonincreasing and nondecreasing, we see that

$$\begin{aligned} \inf_n y_n &= \inf_n \sup_{m \geq n} x_m = \limsup_{x \rightarrow \infty} x_n \\ \sup_n z_n &= \sup_n \inf_{m \geq n} x_m = \liminf_{x \rightarrow \infty} x_n. \end{aligned}$$

Hence,

$$\limsup_{n \rightarrow \infty} x_n = \lim_{n \rightarrow \infty} \sup_{m \geq n} x_m \quad \text{and} \quad \liminf_{n \rightarrow \infty} x_n = \lim_{n \rightarrow \infty} \inf_{m \geq n} x_m.$$



4

⁴Image from Wikipedia.

Characterization of the limsup and liminf.

Thm Let $\{x_n\}_{n=1}^{\infty}$ be a sequence of real numbers. Then

- a) $\limsup x_n = x \in \mathbb{R}$ iff for each $\epsilon > 0$,
 - i) there is an $N \in \mathbb{N}$ such that $x_n \leq x + \epsilon$ for all $n \geq N$, and
 - ii) for each $n \in \mathbb{N}$, there is an $m \geq n$ such that $x_m > x - \epsilon$.
 - b) $\limsup x_n = \infty$ iff for each $M \in \mathbb{R}$ and $N \in \mathbb{N}$, there is an $n \geq N$ such that $x_n > M$; in other words, iff the sequence is unbounded from above.
 - c) Similarly, $\limsup x_n = -\infty$ if and only if $\lim x_n = -\infty$.
-

- a) $\limsup x_n = x \in \mathbb{R}$ iff for each $\epsilon > 0$,
 - i) the ϵ -neighborhood of x has been visited infinitely many times; and
 - ii) only finite many terms are greater than $x + \epsilon$.

Characterization of the limsup and liminf.

Thm Let $\{x_n\}_{n=1}^{\infty}$ be a sequence of real numbers. Then

- a) $\limsup x_n = x \in \mathbb{R}$ iff for each $\epsilon > 0$,
 - i) there is an $N \in \mathbb{N}$ such that $x_n \leq x + \epsilon$ for all $n \geq N$, and
 - ii) for each $n \in \mathbb{N}$, there is an $m \geq n$ such that $x_m > x - \epsilon$.
 - b) $\limsup x_n = \infty$ iff for each $M \in \mathbb{R}$ and $N \in \mathbb{N}$, there is an $n \geq N$ such that $x_n > M$; in other words, iff the sequence is unbounded from above.
 - c) Similarly, $\limsup x_n = -\infty$ if and only if $\lim x_n = -\infty$.
-

- a) $\limsup x_n = x \in \mathbb{R}$ iff for each $\epsilon > 0$,
 - i) the ϵ -neighborhood of x has been visited infinitely many times; and
 - ii) only finite many terms are greater than $x + \epsilon$.

Characterization of the limsup and liminf.

Thm Let $\{x_n\}_{n=1}^{\infty}$ be a sequence of real numbers. Then

- a) $\limsup x_n = x \in \mathbb{R}$ iff for each $\epsilon > 0$,
 - i) there is an $N \in \mathbb{N}$ such that $x_n \leq x + \epsilon$ for all $n \geq N$, and
 - ii) for each $n \in \mathbb{N}$, there is an $m \geq n$ such that $x_m > x - \epsilon$.
 - b) $\limsup x_n = \infty$ iff for each $M \in \mathbb{R}$ and $N \in \mathbb{N}$, there is an $n \geq N$ such that $x_n > M$; in other words, iff the sequence is unbounded from above.
 - c) Similarly, $\limsup x_n = -\infty$ if and only if $\lim x_n = -\infty$.
-

- a) $\limsup x_n = x \in \mathbb{R}$ iff for each $\epsilon > 0$,
 - i) the ϵ -neighborhood of x has been visited infinitely many times; and
 - ii) only finite many terms are greater than $x + \epsilon$.

Characterization of the limsup and liminf.

Thm Let $\{x_n\}_{n=1}^{\infty}$ be a sequence of real numbers. Then

- a) $\limsup x_n = x \in \mathbb{R}$ iff for each $\epsilon > 0$,
 - i) there is an $N \in \mathbb{N}$ such that $x_n \leq x + \epsilon$ for all $n \geq N$, and
 - ii) for each $n \in \mathbb{N}$, there is an $m \geq n$ such that $x_m > x - \epsilon$.
 - b) $\limsup x_n = \infty$ iff for each $M \in \mathbb{R}$ and $N \in \mathbb{N}$, there is an $n \geq N$ such that $x_n > M$; in other words, iff the sequence is unbounded from above.
 - c) Similarly, $\limsup x_n = -\infty$ if and only if $\lim x_n = -\infty$.
-

- a) $\limsup x_n = x \in \mathbb{R}$ iff for each $\epsilon > 0$,
 - i) the ϵ -neighborhood of x has been visited infinitely many times; and
 - ii) only finite many terms are greater than $x + \epsilon$.

Characterization of the limsup and liminf.

Thm Let $\{x_n\}_{n=1}^{\infty}$ be a sequence of real numbers. Then

- a) $\limsup x_n = x \in \mathbb{R}$ iff for each $\epsilon > 0$,
 - i) there is an $N \in \mathbb{N}$ such that $x_n \leq x + \epsilon$ for all $n \geq N$, and
 - ii) for each $n \in \mathbb{N}$, there is an $m \geq n$ such that $x_m > x - \epsilon$.
 - b) $\limsup x_n = \infty$ iff for each $M \in \mathbb{R}$ and $N \in \mathbb{N}$, there is an $n \geq N$ such that $x_n > M$; in other words, iff the sequence is unbounded from above.
 - c) Similarly, $\limsup x_n = -\infty$ if and only if $\lim x_n = -\infty$.
-

- a) $\limsup x_n = x \in \mathbb{R}$ iff for each $\epsilon > 0$,
 - i) the ϵ -neighborhood of x has been visited infinitely many times; and
 - ii) only finite many terms are greater than $x + \epsilon$.

Characterization of the limsup and liminf.

Thm Let $\{x_n\}_{n=1}^{\infty}$ be a sequence of real numbers. Then

- a) $\limsup x_n = x \in \mathbb{R}$ iff for each $\epsilon > 0$,
 - i) there is an $N \in \mathbb{N}$ such that $x_n \leq x + \epsilon$ for all $n \geq N$, and
 - ii) for each $n \in \mathbb{N}$, there is an $m \geq n$ such that $x_m > x - \epsilon$.
 - b) $\limsup x_n = \infty$ iff for each $M \in \mathbb{R}$ and $N \in \mathbb{N}$, there is an $n \geq N$ such that $x_n > M$; in other words, iff the sequence is unbounded from above.
 - c) Similarly, $\limsup x_n = -\infty$ if and only if $\lim x_n = -\infty$.
-

- a) $\limsup x_n = x \in \mathbb{R}$ iff for each $\epsilon > 0$,
 - i) the ϵ -neighborhood of x has been visited infinitely many times; and
 - ii) only finite many terms are greater than $x + \epsilon$.

Characterization of the limsup and liminf.

Thm Let $\{x_n\}_{n=1}^{\infty}$ be a sequence of real numbers. Then

- a) $\limsup x_n = x \in \mathbb{R}$ iff for each $\epsilon > 0$,
 - i) there is an $N \in \mathbb{N}$ such that $x_n \leq x + \epsilon$ for all $n \geq N$, and
 - ii) for each $n \in \mathbb{N}$, there is an $m \geq n$ such that $x_m > x - \epsilon$.
 - b) $\limsup x_n = \infty$ iff for each $M \in \mathbb{R}$ and $N \in \mathbb{N}$, there is an $n \geq N$ such that $x_n > M$; in other words, iff the sequence is unbounded from above.
 - c) Similarly, $\limsup x_n = -\infty$ if and only if $\lim x_n = -\infty$.
-

- a) $\limsup x_n = x \in \mathbb{R}$ iff for each $\epsilon > 0$,
 - i) the ϵ -neighborhood of x has been visited infinitely many times; and
 - ii) only finite many terms are greater than $x + \epsilon$.

Characterization of the limsup and liminf.

Thm Let $\{x_n\}_{n=1}^{\infty}$ be a sequence of real numbers. Then

- a) $\limsup x_n = x \in \mathbb{R}$ iff for each $\epsilon > 0$,
 - i) there is an $N \in \mathbb{N}$ such that $x_n \leq x + \epsilon$ for all $n \geq N$, and
 - ii) for each $n \in \mathbb{N}$, there is an $m \geq n$ such that $x_m > x - \epsilon$.
 - b) $\limsup x_n = \infty$ iff for each $M \in \mathbb{R}$ and $N \in \mathbb{N}$, there is an $n \geq N$ such that $x_n > M$; in other words, iff the sequence is unbounded from above.
 - c) Similarly, $\limsup x_n = -\infty$ if and only if $\lim x_n = -\infty$.
-

- a) $\limsup x_n = x \in \mathbb{R}$ iff for each $\epsilon > 0$,
 - i) the ϵ -neighborhood of x has been visited infinitely many times; and
 - ii) only finite many terms are greater than $x + \epsilon$.

Characterization of the limsup and liminf.

Thm Let $\{x_n\}_{n=1}^{\infty}$ be a sequence of real numbers. Then

- a) $\limsup x_n = x \in \mathbb{R}$ iff for each $\epsilon > 0$,
 - i) there is an $N \in \mathbb{N}$ such that $x_n \leq x + \epsilon$ for all $n \geq N$, and
 - ii) for each $n \in \mathbb{N}$, there is an $m \geq n$ such that $x_m > x - \epsilon$.
 - b) $\limsup x_n = \infty$ iff for each $M \in \mathbb{R}$ and $N \in \mathbb{N}$, there is an $n \geq N$ such that $x_n > M$; in other words, iff the sequence is unbounded from above.
 - c) Similarly, $\limsup x_n = -\infty$ if and only if $\lim x_n = -\infty$.
-

- a) $\limsup x_n = x \in \mathbb{R}$ iff for each $\epsilon > 0$,
 - i) the ϵ -neighborhood of x has been visited infinitely many times; and
 - ii) only finite many terms are greater than $x + \epsilon$.

Characterization of the limsup and liminf.

Thm Let $\{x_n\}_{n=1}^{\infty}$ be a sequence of real numbers. Then

- a) $\limsup x_n = x \in \mathbb{R}$ iff for each $\epsilon > 0$,
 - i) there is an $N \in \mathbb{N}$ such that $x_n \leq x + \epsilon$ for all $n \geq N$, and
 - ii) for each $n \in \mathbb{N}$, there is an $m \geq n$ such that $x_m > x - \epsilon$.
 - b) $\limsup x_n = \infty$ iff for each $M \in \mathbb{R}$ and $N \in \mathbb{N}$, there is an $n \geq N$ such that $x_n > M$; in other words, iff the sequence is unbounded from above.
 - c) Similarly, $\limsup x_n = -\infty$ if and only if $\lim x_n = -\infty$.
-

- a) $\limsup x_n = x \in \mathbb{R}$ iff for each $\epsilon > 0$,
 - i) the ϵ -neighborhood of x has been visited infinitely many times; and
 - ii) only finite many terms are greater than $x + \epsilon$.

Characterization of the limsup and liminf.

Thm Let $\{x_n\}_{n=1}^{\infty}$ be a sequence of real numbers. Then

- a) $\limsup x_n = x \in \mathbb{R}$ iff for each $\epsilon > 0$,
 - i) there is an $N \in \mathbb{N}$ such that $x_n \leq x + \epsilon$ for all $n \geq N$, and
 - ii) for each $n \in \mathbb{N}$, there is an $m \geq n$ such that $x_m > x - \epsilon$.
 - b) $\limsup x_n = \infty$ iff for each $M \in \mathbb{R}$ and $N \in \mathbb{N}$, there is an $n \geq N$ such that $x_n > M$; in other words, iff the sequence is unbounded from above.
 - c) Similarly, $\limsup x_n = -\infty$ if and only if $\lim x_n = -\infty$.
-

- a) $\limsup x_n = x \in \mathbb{R}$ iff for each $\epsilon > 0$,
 - i) the ϵ -neighborhood of x has been visited infinitely many times; and
 - ii) only finite many terms are greater than $x + \epsilon$.

Proof. (Sketch) We only prove part (a). Let $y_n = \sup_{k \geq n} x_k$.

$$\begin{aligned} \limsup x_n = x &\iff \lim_{n \rightarrow \infty} \sup_{k \geq n} x_k = x \\ &\iff \forall \epsilon \exists N \forall n \ (n \geq N) \rightarrow \left(\sup_{k \geq n} x_k \in (x - \epsilon, x + \epsilon) \right) \end{aligned}$$

$$\sup_{k \geq n} x_k < x + \epsilon \iff \text{all terms starting from } n \text{ fall below } x + \epsilon$$

$$\sup_{k \geq n} x_k > x - \epsilon \iff \exists k \geq n \text{ s.t. } x_k > x - \epsilon$$

□

Proof. (Sketch) We only prove part (a). Let $y_n = \sup_{k \geq n} x_k$.

$$\begin{aligned} \limsup x_n = x &\iff \lim_{n \rightarrow \infty} \sup_{k \geq n} x_k = x \\ &\iff \forall \epsilon \exists N \forall n \ (n \geq N) \rightarrow \left(\sup_{k \geq n} x_k \in (x - \epsilon, x + \epsilon) \right) \end{aligned}$$

$$\sup_{k \geq n} x_k < x + \epsilon \iff \text{all terms starting from } n \text{ fall below } x + \epsilon$$

$$\sup_{k \geq n} x_k > x - \epsilon \iff \exists k \geq n \text{ s.t. } x_k > x - \epsilon$$

□

Thm Let $\{x_n\}_{n=1}^{\infty}$ be a sequence of real numbers. Then

- a) $\liminf x_n = x \in \mathbb{R}$ iff for each $\epsilon > 0$,
 - i) there is an $N \in \mathbb{N}$ such that $x_n \geq x - \epsilon$ for all $n \geq N$, and
 - ii) for each $n \in \mathbb{N}$, there is an $m \geq n$ such that $x_m < x + \epsilon$.
- b) $\liminf x_n = -\infty$ iff for each $M \in \mathbb{R}$ and $N \in \mathbb{N}$, there is an $n \geq N$ such that $x_n < M$; in other words, iff the sequence is unbounded from below.
- c) Similarly, $\liminf x_n = \infty$ if and only $\lim x_n = \infty$.

Thm Let $\{x_n\}_{n=1}^{\infty}$ be a sequence of real numbers. Then

- a) $\liminf x_n = x \in \mathbb{R}$ iff for each $\epsilon > 0$,
 - i) there is an $N \in \mathbb{N}$ such that $x_n \geq x - \epsilon$ for all $n \geq N$, and
 - ii) for each $n \in \mathbb{N}$, there is an $m \geq n$ such that $x_m < x + \epsilon$.
- b) $\liminf x_n = -\infty$ iff for each $M \in \mathbb{R}$ and $N \in \mathbb{N}$, there is an $n \geq N$ such that $x_n < M$; in other words, iff the sequence is unbounded from below.
- c) Similarly, $\liminf x_n = \infty$ if and only $\lim x_n = \infty$.

Thm Let $\{x_n\}_{n=1}^{\infty}$ be a sequence of real numbers. Then

- a) $\liminf x_n = x \in \mathbb{R}$ iff for each $\epsilon > 0$,
 - i) there is an $N \in \mathbb{N}$ such that $x_n \geq x - \epsilon$ for all $n \geq N$, and
 - ii) for each $n \in \mathbb{N}$, there is an $m \geq n$ such that $x_m < x + \epsilon$.
- b) $\liminf x_n = -\infty$ iff for each $M \in \mathbb{R}$ and $N \in \mathbb{N}$, there is an $n \geq N$ such that $x_n < M$; in other words, iff the sequence is unbounded from below.
- c) Similarly, $\liminf x_n = \infty$ if and only $\lim x_n = \infty$.

Thm Let $\{x_n\}_{n=1}^{\infty}$ be a sequence of real numbers. Then

- a) $\liminf x_n = x \in \mathbb{R}$ iff for each $\epsilon > 0$,
 - i) there is an $N \in \mathbb{N}$ such that $x_n \geq x - \epsilon$ for all $n \geq N$, and
 - ii) for each $n \in \mathbb{N}$, there is an $m \geq n$ such that $x_m < x + \epsilon$.
- b) $\liminf x_n = -\infty$ iff for each $M \in \mathbb{R}$ and $N \in \mathbb{N}$, there is an $n \geq N$ such that $x_n < M$; in other words, iff the sequence is unbounded from below.
- c) Similarly, $\liminf x_n = \infty$ if and only $\lim x_n = \infty$.

Thm Let $\{x_n\}_{n=1}^{\infty}$ be a sequence of real numbers. Then

- a) $\liminf x_n = x \in \mathbb{R}$ iff for each $\epsilon > 0$,
 - i) there is an $N \in \mathbb{N}$ such that $x_n \geq x - \epsilon$ for all $n \geq N$, and
 - ii) for each $n \in \mathbb{N}$, there is an $m \geq n$ such that $x_m < x + \epsilon$.
- b) $\liminf x_n = -\infty$ iff for each $M \in \mathbb{R}$ and $N \in \mathbb{N}$, there is an $n \geq N$ such that $x_n < M$; in other words, iff the sequence is unbounded from below.
- c) Similarly, $\liminf x_n = \infty$ if and only $\lim x_n = \infty$.

Thm Let $\{x_n\}_{n=1}^{\infty}$ be a sequence of real numbers. Then

- a) $\liminf x_n = x \in \mathbb{R}$ iff for each $\epsilon > 0$,
 - i) there is an $N \in \mathbb{N}$ such that $x_n \geq x - \epsilon$ for all $n \geq N$, and
 - ii) for each $n \in \mathbb{N}$, there is an $m \geq n$ such that $x_m < x + \epsilon$.
- b) $\liminf x_n = -\infty$ iff for each $M \in \mathbb{R}$ and $N \in \mathbb{N}$, there is an $n \geq N$ such that $x_n < M$; in other words, iff the sequence is unbounded from below.
- c) Similarly, $\liminf x_n = \infty$ if and only if $\lim x_n = \infty$.

Still other characterization (as an exercise):

Thm $\limsup_{n \rightarrow \infty} x_n = x \in \mathbb{R}$ if and only if x is the smallest real number such that for any positive real number $\epsilon > 0$, there exists a natural number N such that $x_n < x + \epsilon$ for all $n > N$.

$\liminf_{n \rightarrow \infty} x_n = x \in \mathbb{R}$ if and only if x is the largest real number such that for any positive real number $\epsilon > 0$, there exists a natural number N such that $x_n > x - \epsilon$ for all $n > N$.

Proof. HW for motivated students.

□

Still other characterization (as an exercise):

Thm $\limsup_{n \rightarrow \infty} x_n = x \in \mathbb{R}$ if and only if x is the **smallest** real number such that for any positive real number $\epsilon > 0$, there exists a natural number N such that $x_n < x + \epsilon$ for all $n > N$.

$\liminf_{n \rightarrow \infty} x_n = x \in \mathbb{R}$ if and only if x is the largest real number such that for any positive real number $\epsilon > 0$, there exists a natural number N such that $x_n > x - \epsilon$ for all $n > N$.

Proof. HW for motivated students.



Still other characterization (as an exercise):

Thm $\limsup_{n \rightarrow \infty} x_n = x \in \mathbb{R}$ if and only if x is the **smallest** real number such that for any positive real number $\epsilon > 0$, there exists a natural number N such that $x_n < x + \epsilon$ for all $n > N$.

$\liminf_{n \rightarrow \infty} x_n = x \in \mathbb{R}$ if and only if x is the **largest** real number such that for any positive real number $\epsilon > 0$, there exists a natural number N such that $x_n > x - \epsilon$ for all $n > N$.

Proof. HW for motivated students.



Still other characterization (as an exercise):

Thm $\limsup_{n \rightarrow \infty} x_n = x \in \mathbb{R}$ if and only if x is the **smallest** real number such that for any positive real number $\epsilon > 0$, there exists a natural number N such that $x_n < x + \epsilon$ for all $n > N$.

$\liminf_{n \rightarrow \infty} x_n = x \in \mathbb{R}$ if and only if x is the **largest** real number such that for any positive real number $\epsilon > 0$, there exists a natural number N such that $x_n > x - \epsilon$ for all $n > N$.

Proof. HW for motivated students.



E.g.1 Let $x_n = (-1)^n$. Then $\{x_n\}_{n=1}^{\infty}$ has two cluster points: ± 1 , amount which

$$\liminf_{n \rightarrow \infty} x_n = -1 \quad \text{and} \quad \limsup_{n \rightarrow \infty} x_n = 1.$$

Variations

$$x_n = (-1)^n \frac{n+5}{n}$$

$$x_n = 3 + \sin(n\pi) \frac{n^2}{n^2 + 8}$$

Similar examples

$$x_n = \sin\left(\frac{n\pi}{3}\right)$$

E.g.1 Let $x_n = (-1)^n$. Then $\{x_n\}_{n=1}^{\infty}$ has two cluster points: ± 1 , amount which

$$\liminf_{n \rightarrow \infty} x_n = -1 \quad \text{and} \quad \limsup_{n \rightarrow \infty} x_n = 1.$$

Variations

$$x_n = (-1)^n \frac{n+5}{n}$$

$$x_n = 3 + \sin(n\pi) \frac{n^2}{n^2 + 8}$$

Similar examples

$$x_n = \sin\left(\frac{n\pi}{3}\right)$$

E.g.1 Let $x_n = (-1)^n$. Then $\{x_n\}_{n=1}^{\infty}$ has two cluster points: ± 1 , amount which

$$\liminf_{n \rightarrow \infty} x_n = -1 \quad \text{and} \quad \limsup_{n \rightarrow \infty} x_n = 1.$$

Variations

$$x_n = (-1)^n \frac{n+5}{n}$$

$$x_n = 3 + \sin(n\pi) \frac{n^2}{n^2 + 8}$$

Similar examples

$$x_n = \sin\left(\frac{n\pi}{3}\right)$$

E.g.1 Let $x_n = (-1)^n$. Then $\{x_n\}_{n=1}^{\infty}$ has two cluster points: ± 1 , amount which

$$\liminf_{n \rightarrow \infty} x_n = -1 \quad \text{and} \quad \limsup_{n \rightarrow \infty} x_n = 1.$$

Variations

$$x_n = (-1)^n \frac{n+5}{n}$$

$$x_n = 3 + \sin(n\pi) \frac{n^2}{n^2 + 8}$$

Similar examples

$$x_n = \sin\left(\frac{n\pi}{3}\right)$$

E.g.1 Let $x_n = (-1)^n$. Then $\{x_n\}_{n=1}^{\infty}$ has two cluster points: ± 1 , amount which

$$\liminf_{n \rightarrow \infty} x_n = -1 \quad \text{and} \quad \limsup_{n \rightarrow \infty} x_n = 1.$$

Variations

$$x_n = (-1)^n \frac{n+5}{n}$$

$$x_n = 3 + \sin(n\pi) \frac{n^2}{n^2 + 8}$$

Similar examples

$$x_n = \sin\left(\frac{n\pi}{3}\right)$$

E.g.1 Let $x_n = (-1)^n$. Then $\{x_n\}_{n=1}^{\infty}$ has two cluster points: ± 1 , amount which

$$\liminf_{n \rightarrow \infty} x_n = -1 \quad \text{and} \quad \limsup_{n \rightarrow \infty} x_n = 1.$$

Variations

$$x_n = (-1)^n \frac{n+5}{n}$$

$$x_n = 3 + \sin(n\pi) \frac{n^2}{n^2 + 8}$$

Similar examples

$$x_n = \sin\left(\frac{n\pi}{3}\right)$$

E.g.2 Consider the sequence $\{2, 1, 0, 2, 2, \frac{1}{2}, 2, 3, \frac{2}{3}, 2, 4, \frac{3}{4}, \dots\}$, that is,

$$x_n = \begin{cases} (n-3)/n & \text{if } n \equiv 0 \pmod{3} \\ 2 & \text{if } n \equiv 1 \pmod{3} \\ (n+1)/3 & \text{if } n \equiv 2 \pmod{3} \end{cases}$$

It has three cluster points: 1, 2, ∞ , among which

$$\liminf_{n \rightarrow \infty} x_n = 1 \quad \text{and} \quad \limsup_{n \rightarrow \infty} x_n = \infty.$$

E.g.3 Let $\{r_n\}_{n=1}^{\infty}$ be an enumeration of the rational numbers. By the density of the rational numbers, every extended real number is a cluster point of the sequence $\{r_n\}_{n=1}^{\infty}$, amount which

$$\liminf_{n \rightarrow \infty} r_n = -\infty \quad \text{and} \quad \limsup_{n \rightarrow \infty} r_n = +\infty.$$

E.g.2 Consider the sequence $\{2, 1, 0, 2, 2, \frac{1}{2}, 2, 3, \frac{2}{3}, 2, 4, \frac{3}{4}, \dots\}$, that is,

$$x_n = \begin{cases} (n-3)/n & \text{if } n \equiv 0 \pmod{3} \\ 2 & \text{if } n \equiv 1 \pmod{3} \\ (n+1)/3 & \text{if } n \equiv 2 \pmod{3} \end{cases}$$

It has three cluster points: 1, 2, ∞ , among which

$$\liminf_{n \rightarrow \infty} x_n = 1 \quad \text{and} \quad \limsup_{n \rightarrow \infty} x_n = \infty.$$

E.g.3 Let $\{r_n\}_{n=1}^{\infty}$ be an enumeration of the rational numbers. By the density of the rational numbers, every extended real number is a cluster point of the sequence $\{r_n\}_{n=1}^{\infty}$, amount which

$$\liminf_{n \rightarrow \infty} r_n = -\infty \quad \text{and} \quad \limsup_{n \rightarrow \infty} r_n = +\infty.$$

The above examples suggest that

Prop. Let $\{x_n\}_{n=1}^{\infty}$ be a sequence of real numbers. Then

- a) $\limsup x_n$ is the largest cluster point of $\{x_n\}_{n=1}^{\infty}$.
- b) $\liminf x_n$ is the smallest cluster point of $\{x_n\}_{n=1}^{\infty}$.

The above examples suggest that

Prop. Let $\{x_n\}_{n=1}^{\infty}$ be a sequence of real numbers. Then

- a) $\limsup x_n$ is the largest cluster point of $\{x_n\}_{n=1}^{\infty}$.
- b) $\liminf x_n$ is the smallest cluster point of $\{x_n\}_{n=1}^{\infty}$.

The above examples suggest that

Prop. Let $\{x_n\}_{n=1}^{\infty}$ be a sequence of real numbers. Then

- a) $\limsup x_n$ is the largest cluster point of $\{x_n\}_{n=1}^{\infty}$.
- b) $\liminf x_n$ is the smallest cluster point of $\{x_n\}_{n=1}^{\infty}$.

The above examples suggest that

Prop. Let $\{x_n\}_{n=1}^{\infty}$ be a sequence of real numbers. Then

- a) $\limsup x_n$ is the largest cluster point of $\{x_n\}_{n=1}^{\infty}$.
- b) $\liminf x_n$ is the smallest cluster point of $\{x_n\}_{n=1}^{\infty}$.

Proof. We only prove (a). (b) can be proved in a similar way.

Let $x = \limsup x_n$. We have seen that x is a cluster point.

It remains to prove that x is the largest cluster point. The case when $x = \pm\infty$ is left for the motivated students.

Now assume that $x \in \mathbb{R}$.

Only finite many terms exceed $x + 1$, hence, ∞ is not a cluster point.

Let $y \in \mathbb{R}$ s.t. $x < y$. Set $\epsilon = (y - x)/2$.

Only finite many terms exceed $x + \epsilon$.

Since $y - \epsilon = x + \epsilon$, the ϵ -neighborhood of y has been visited only finitely many times.

Hence, y cannot be a cluster point. □

Proof. We only prove (a). (b) can be proved in a similar way.

Let $x = \limsup x_n$. We have seen that x is a cluster point.

It remains to prove that x is the largest cluster point. The case when $x = \pm\infty$ is left for the motivated students.

Now assume that $x \in \mathbb{R}$.

Only finite many terms exceed $x + 1$, hence, ∞ is not a cluster point.

Let $y \in \mathbb{R}$ s.t. $x < y$. Set $\epsilon = (y - x)/2$.

Only finite many terms exceed $x + \epsilon$.

Since $y - \epsilon = x + \epsilon$, the ϵ -neighborhood of y has been visited only finitely many times.

Hence, y cannot be a cluster point. □

Proof. We only prove (a). (b) can be proved in a similar way.

Let $x = \limsup x_n$. We have seen that x is a cluster point.

It remains to prove that x is the largest cluster point. The case when $x = \pm\infty$ is left for the motivated students.

Now assume that $x \in \mathbb{R}$.

Only finite many terms exceed $x + 1$, hence, ∞ is not a cluster point.

Let $y \in \mathbb{R}$ s.t. $x < y$. Set $\epsilon = (y - x)/2$.

Only finite many terms exceed $x + \epsilon$.

Since $y - \epsilon = x + \epsilon$, the ϵ -neighborhood of y has been visited only finitely many times.

Hence, y cannot be a cluster point. □

Proof. We only prove (a). (b) can be proved in a similar way.

Let $x = \limsup x_n$. We have seen that x is a cluster point.

It remains to prove that x is the largest cluster point. The case when $x = \pm\infty$ is left for the motivated students.

Now assume that $x \in \mathbb{R}$.

Only finite many terms exceed $x + 1$, hence, ∞ is not a cluster point.

Let $y \in \mathbb{R}$ s.t. $x < y$. Set $\epsilon = (y - x)/2$.

Only finite many terms exceed $x + \epsilon$.

Since $y - \epsilon = x + \epsilon$, the ϵ -neighborhood of y has been visited only finitely many times.

Hence, y cannot be a cluster point. □

Proof. We only prove (a). (b) can be proved in a similar way.

Let $x = \limsup x_n$. We have seen that x is a cluster point.

It remains to prove that x is the largest cluster point. The case when $x = \pm\infty$ is left for the motivated students.

Now assume that $x \in \mathbb{R}$.

Only finite many terms exceed $x + 1$, hence, ∞ is not a cluster point.

Let $y \in \mathbb{R}$ s.t. $x < y$. Set $\epsilon = (y - x)/2$.

Only finite many terms exceed $x + \epsilon$.

Since $y - \epsilon = x + \epsilon$, the ϵ -neighborhood of y has been visited only finitely many times.

Hence, y cannot be a cluster point. □

Proof. We only prove (a). (b) can be proved in a similar way.

Let $x = \limsup x_n$. We have seen that x is a cluster point.

It remains to prove that x is the largest cluster point. The case when $x = \pm\infty$ is left for the motivated students.

Now assume that $x \in \mathbb{R}$.

Only finite many terms exceed $x + 1$, hence, ∞ is not a cluster point.

Let $y \in \mathbb{R}$ s.t. $x < y$. Set $\epsilon = (y - x)/2$.

Only finite many terms exceed $x + \epsilon$.

Since $y - \epsilon = x + \epsilon$, the ϵ -neighborhood of y has been visited only finitely many times.

Hence, y cannot be a cluster point. □

Proof. We only prove (a). (b) can be proved in a similar way.

Let $x = \limsup x_n$. We have seen that x is a cluster point.

It remains to prove that x is the largest cluster point. The case when $x = \pm\infty$ is left for the motivated students.

Now assume that $x \in \mathbb{R}$.

Only finite many terms exceed $x + 1$, hence, ∞ is not a cluster point.

Let $y \in \mathbb{R}$ s.t. $x < y$. Set $\epsilon = (y - x)/2$.

Only finite many terms exceed $x + \epsilon$.

Since $y - \epsilon = x + \epsilon$, the ϵ -neighborhood of y has been visited only finitely many times.

Hence, y cannot be a cluster point. □

Proof. We only prove (a). (b) can be proved in a similar way.

Let $x = \limsup x_n$. We have seen that x is a cluster point.

It remains to prove that x is the largest cluster point. The case when $x = \pm\infty$ is left for the motivated students.

Now assume that $x \in \mathbb{R}$.

Only finite many terms exceed $x + 1$, hence, ∞ is not a cluster point.

Let $y \in \mathbb{R}$ s.t. $x < y$. Set $\epsilon = (y - x)/2$.

Only finite many terms exceed $x + \epsilon$.

Since $y - \epsilon = x + \epsilon$, the ϵ -neighborhood of y has been visited only finitely many times.

Hence, y cannot be a cluster point. □

Proof. We only prove (a). (b) can be proved in a similar way.

Let $x = \limsup x_n$. We have seen that x is a cluster point.

It remains to prove that x is the largest cluster point. The case when $x = \pm\infty$ is left for the motivated students.

Now assume that $x \in \mathbb{R}$.

Only finite many terms exceed $x + 1$, hence, ∞ is not a cluster point.

Let $y \in \mathbb{R}$ s.t. $x < y$. Set $\epsilon = (y - x)/2$.

Only finite many terms exceed $x + \epsilon$.

Since $y - \epsilon = x + \epsilon$, the ϵ -neighborhood of y has been visited only finitely many times.

Hence, y cannot be a cluster point. □

Properties

1.

$$\inf_n x_n \leq \liminf_{n \rightarrow \infty} x_n \leq \limsup_{n \rightarrow \infty} x_n \leq \sup_n x_n$$

2. A sequence $\{x_n\}_{n=1}^{\infty}$ of real numbers converges in \mathbb{R}^* if and only if it has exactly one cluster point. In such cases, the limit of the sequence is the unique cluster point. In other words,

$$\liminf_{n \rightarrow \infty} x_n = \limsup_{n \rightarrow \infty} x_n = C \iff \lim_{n \rightarrow \infty} x_n = C.$$


Properties

1.

$$\inf_n x_n \leq \liminf_{n \rightarrow \infty} x_n \leq \limsup_{n \rightarrow \infty} x_n \leq \sup_n x_n$$

2. A sequence $\{x_n\}_{n=1}^{\infty}$ of real numbers **converges in \mathbb{R}^*** if and only if it has exactly one cluster point. In such cases, the limit of the sequence is the unique cluster point. In other words,

$$\liminf_{n \rightarrow \infty} x_n = \limsup_{n \rightarrow \infty} x_n = c \iff \lim_{n \rightarrow \infty} x_n = c.$$



5

E.g. For all $\epsilon > 0$, the interval

$$\left(\liminf_{n \rightarrow \infty} x_n - \epsilon, \limsup_{n \rightarrow \infty} x_n + \epsilon \right)$$

contains *all but finitely many* numbers in $\{x_n\}$.

⁵Image is from Wikipedia.

E.g. (Continued) Similarly, if

$$\liminf_{n \rightarrow \infty} x_n < \limsup_{n \rightarrow \infty} x_n,$$

then for all ϵ with

$$0 < \epsilon < \frac{1}{2} \left(\limsup_{n \rightarrow \infty} x_n - \liminf_{n \rightarrow \infty} x_n \right),$$

infinitely many numbers in $\{x_n\}$ fall outside of the interval

$$\left(\liminf_{n \rightarrow \infty} x_n + \epsilon, \limsup_{n \rightarrow \infty} x_n - \epsilon \right).$$

Cauchy criterion

As we have seen that

A sequence of real numbers converges in \mathbb{R}^* if and only if it has exactly one cluster point.

There is another famous criterion for a sequence to converge in \mathbb{R} :

Cauchy Criterion

Cauchy criterion

As we have seen that

A sequence of real numbers **converges in \mathbb{R}^*** if and only if it has exactly one cluster point.

There is another famous criterion for a sequence to converge in \mathbb{R} :

Cauchy Criterion

Cauchy criterion


As we have seen that

A sequence of real numbers **converges in \mathbb{R}^*** if and only if it has exactly one cluster point.


There is another famous criterion for a sequence to **converge in \mathbb{R}** :

Cauchy Criterion

Cauchy sequence



Non-Cauchy sequence



⁶Images from Wikipedia.

Def. A sequence $\{x_n\}_{n=1}^{\infty}$ of real numbers is called a *Cauchy sequence* if for each $\epsilon > 0$, there exists an $N \in \mathbb{N}$ such that for all $m, n \geq N$, we have $|x_n - x_m| < \epsilon$.

Def.' A sequence $\{x_n\}_{n=1}^{\infty}$ is *Cauchy* iff

$$\forall \epsilon > 0 \exists N \in \mathbb{N} \forall m, n \geq N \{ |x_n - x_m| < \epsilon \}.$$

Thm (Cauchy Criterion)

A sequence of real numbers converges in \mathbb{R} iff it is Cauchy.

Def. A sequence $\{x_n\}_{n=1}^{\infty}$ of real numbers is called a *Cauchy sequence* if for each $\epsilon > 0$, there exists an $N \in \mathbb{N}$ such that for all $m, n \geq N$, we have $|x_n - x_m| < \epsilon$.

Def.' A sequence $\{x_n\}_{n=1}^{\infty}$ is *Cauchy* iff

$$\forall \epsilon > 0 \exists N \in \mathbb{N} \forall m, n \geq N \{|x_n - x_m| < \epsilon\}.$$

Thm (Cauchy Criterion)

A sequence of real numbers converges in \mathbb{R} iff it is Cauchy.

Def. A sequence $\{x_n\}_{n=1}^{\infty}$ of real numbers is called a *Cauchy sequence* if for each $\epsilon > 0$, there exists an $N \in \mathbb{N}$ such that for all $m, n \geq N$, we have $|x_n - x_m| < \epsilon$.

Def.' A sequence $\{x_n\}_{n=1}^{\infty}$ is *Cauchy* iff

$$\forall \epsilon > 0 \exists N \in \mathbb{N} \forall m, n \geq N \{|x_n - x_m| < \epsilon\}.$$

Thm (Cauchy Criterion)

A sequence of real numbers converges in \mathbb{R} iff it is Cauchy.

Def. A sequence $\{x_n\}_{n=1}^{\infty}$ of real numbers is called a *Cauchy sequence* if for each $\epsilon > 0$, there exists an $N \in \mathbb{N}$ such that for all $m, n \geq N$, we have $|x_n - x_m| < \epsilon$.

Def.' A sequence $\{x_n\}_{n=1}^{\infty}$ is *Cauchy* iff

$$\forall \epsilon > 0 \exists N \in \mathbb{N} \forall m, n \geq N \{|x_n - x_m| < \epsilon\}.$$

Thm (Cauchy Criterion)

A sequence of real numbers *converges in \mathbb{R}* iff it is Cauchy.

Proof. “ \Rightarrow ” Easy!

“ \Leftarrow ”: ...



Proof. " \Rightarrow " Easy!

" \Leftarrow ": ...



E.g.1 Let $a_n = \sqrt{n}$. Show that

- (i) The consecutive terms become arbitrarily close to each other as $n \rightarrow \infty$.
- (ii) The sequence is not Cauchy.

Sol. (i) This is because

$$a_{n+1} - a_n = \sqrt{n+1} - \sqrt{n} = \frac{1}{\sqrt{n+1} + \sqrt{n}} < \frac{1}{2\sqrt{n}} \rightarrow 0, \quad \text{as } n \rightarrow \infty.$$

(ii) To show $\{a_n\}_{n=1}^{\infty}$ is not Cauchy, let's negate the statement as follows:

$$\begin{aligned} & \neg (\forall \epsilon > 0 \exists N \in \mathbb{N} \forall m, n \geq N \{|x_n - x_m| < \epsilon\}) \\ \iff & \exists \epsilon > 0 \forall N \in \mathbb{N} \exists m, n \geq N \{|x_n - x_m| > \epsilon\} \end{aligned}$$

E.g.1 Let $a_n = \sqrt{n}$. Show that

- (i) The consecutive terms become arbitrarily close to each other as $n \rightarrow \infty$.
- (ii) The sequence is not Cauchy.

Sol. (i) This is because

$$a_{n+1} - a_n = \sqrt{n+1} - \sqrt{n} = \frac{1}{\sqrt{n+1} + \sqrt{n}} < \frac{1}{2\sqrt{n}} \rightarrow 0, \quad \text{as } n \rightarrow \infty.$$

(ii) To show $\{a_n\}_{n=1}^{\infty}$ is not Cauchy, let's negate the statement as follows:

$$\begin{aligned} & \neg (\forall \epsilon > 0 \exists N \in \mathbb{N} \forall m, n \geq N \{|x_n - x_m| < \epsilon\}) \\ \iff & \exists \epsilon > 0 \forall N \in \mathbb{N} \exists m, n \geq N \{|x_n - x_m| > \epsilon\} \end{aligned}$$

E.g.1 Let $a_n = \sqrt{n}$. Show that

- (i) The consecutive terms become arbitrarily close to each other as $n \rightarrow \infty$.
- (ii) The sequence is not Cauchy.

Sol. (i) This is because

$$a_{n+1} - a_n = \sqrt{n+1} - \sqrt{n} = \frac{1}{\sqrt{n+1} + \sqrt{n}} < \frac{1}{2\sqrt{n}} \rightarrow 0, \quad \text{as } n \rightarrow \infty.$$

(ii) To show $\{a_n\}_{n=1}^{\infty}$ is not Cauchy, let's negate the statement as follows:

$$\begin{aligned} & \neg (\forall \epsilon > 0 \exists N \in \mathbb{N} \forall m, n \geq N \{|x_n - x_m| < \epsilon\}) \\ \iff & \exists \epsilon > 0 \forall N \in \mathbb{N} \exists m, n \geq N \{|x_n - x_m| > \epsilon\} \end{aligned}$$

E.g.1 Let $a_n = \sqrt{n}$. Show that

- (i) The consecutive terms become arbitrarily close to each other as $n \rightarrow \infty$.
- (ii) The sequence is not Cauchy.

Sol. (i) This is because

$$a_{n+1} - a_n = \sqrt{n+1} - \sqrt{n} = \frac{1}{\sqrt{n+1} + \sqrt{n}} < \frac{1}{2\sqrt{n}} \rightarrow 0, \quad \text{as } n \rightarrow \infty.$$

(ii) To show $\{a_n\}_{n=1}^{\infty}$ is not Cauchy, let's negate the statement as follows:

$$\begin{aligned} & \neg (\forall \epsilon > 0 \exists N \in \mathbb{N} \forall m, n \geq N \{|x_n - x_m| < \epsilon\}) \\ \iff & \exists \epsilon > 0 \forall N \in \mathbb{N} \exists m, n \geq N \{|x_n - x_m| > \epsilon\} \end{aligned}$$

E.g.1 Let $a_n = \sqrt{n}$. Show that

- (i) The consecutive terms become arbitrarily close to each other as $n \rightarrow \infty$.
- (ii) The sequence is not Cauchy.

Sol. (i) This is because

$$a_{n+1} - a_n = \sqrt{n+1} - \sqrt{n} = \frac{1}{\sqrt{n+1} + \sqrt{n}} < \frac{1}{2\sqrt{n}} \rightarrow 0, \quad \text{as } n \rightarrow \infty.$$

(ii) To show $\{a_n\}_{n=1}^{\infty}$ is not Cauchy, let's negate the statement as follows:

$$\begin{aligned} & \neg (\forall \epsilon > 0 \exists N \in \mathbb{N} \forall m, n \geq N \{|x_n - x_m| < \epsilon\}) \\ \iff & \exists \epsilon > 0 \forall N \in \mathbb{N} \exists m, n \geq N \{|x_n - x_m| > \epsilon\} \end{aligned}$$

Sol. (Continued) Let's choose $\epsilon = 1$. For any $N \in \mathbb{N}$, we need to find $m, n \geq N$ such that

$$|a_n - a_m| \geq 1.$$

Indeed, let's choose $m = N$ and $n = 4N$

$$|a_n - a_m| = \sqrt{4N} - \sqrt{N} = \sqrt{N}(\sqrt{4} - 1) = \sqrt{N} \geq 1 = \epsilon.$$

□

Sol. (Continued) Let's choose $\epsilon = 1$. For any $N \in \mathbb{N}$, we need to find $m, n \geq N$ such that

$$|a_n - a_m| \geq 1.$$

Indeed, let's choose $m = N$ and $n = 4N$

$$|a_n - a_m| = \sqrt{4N} - \sqrt{N} = \sqrt{N}(\sqrt{4} - 1) = \sqrt{N} \geq 1 = \epsilon.$$

□

HW Ex. 2.15, 2.17 (a), 2.18, 2.19, 2.22, 2.29.

Chapter 3. Real Number System and Calculus

§ 3.1 Real number system

§ 3.2 Sequences of real numbers

§ 3.3 Open and closed sets

§ 3.4 Real-valued functions

§ 3.5 Liminf and limsup of sets

§ 3.6 Some techniques in calculus

Open sets

Def. A subset $O \subset \mathbb{R}$ is said to be an *open set* if for each $x \in O$, there exists an $r > 0$ such that $(x - r, x + r) \subset O$.

E.g. (a, b) with $-\infty \leq a < b \leq \infty$ is an open set, which are called *open interval intervals*.

$(0, 1)$ is not an open set.

Let K be a nonempty countable subset of \mathbb{R} . Then K cannot be an open set. For example, \mathbb{N} , \mathbb{Q} , \mathbb{Z} are not open sets.

\mathbb{Q}^c – the set of irrational numbers – is not an open set.

E.g. (a, b) with $-\infty \leq a < b \leq \infty$ is an open set, which are called *open interval intervals*.

$(0, 1)$ is not an open set.

Let K be a nonempty countable subset of \mathbb{R} . Then K cannot be an open set. For example, \mathbb{N} , \mathbb{Q} , \mathbb{Z} are not open sets.

\mathbb{Q}^c = the set of irrational numbers = is not an open set.

E.g. (a, b) with $-\infty \leq a < b \leq \infty$ is an open set, which are called *open interval intervals*.

$(0, 1)$ is not an open set.

Let K be a nonempty countable subset of \mathbb{R} . Then K cannot be an open set. For example, \mathbb{N} , \mathbb{Q} , \mathbb{Z} are not open sets.

\mathbb{Q}^c – the set of irrational numbers – is not an open set.

E.g. (a, b) with $-\infty \leq a < b \leq \infty$ is an open set, which are called *open interval intervals*.

$(0, 1]$ is not an open set.

Let K be a nonempty countable subset of \mathbb{R} . Then K cannot be an open set. For example, \mathbb{N} , \mathbb{Q} , \mathbb{Z} are not open sets.

\mathbb{Q}^c = the set of irrational numbers = is not an open set.

E.g. (a, b) with $-\infty \leq a < b \leq \infty$ is an open set, which are called *open interval intervals*.

$(0, 1]$ is not an open set.

Let K be a nonempty countable subset of \mathbb{R} . Then K cannot be an open set. For example, \mathbb{N} , \mathbb{Q} , \mathbb{Z} are not open sets.

\mathbb{Q}^c = the set of irrational numbers = is not an open set.

E.g. (a, b) with $-\infty \leq a < b \leq \infty$ is an open set, which are called *open interval intervals*.

$(0, 1]$ is not an open set.

Let K be a nonempty countable subset of \mathbb{R} . Then K cannot be an open set. For example, \mathbb{N} , \mathbb{Q} , \mathbb{Z} are not open sets.

\mathbb{Q}^c – the set of irrational numbers – is not an open set.

E.g. (a, b) with $-\infty \leq a < b \leq \infty$ is an open set, which are called *open interval intervals*.

$(0, 1]$ is not an open set.

Let K be a nonempty countable subset of \mathbb{R} . Then K cannot be an open set. For example, \mathbb{N} , \mathbb{Q} , \mathbb{Z} are not open sets.

\mathbb{Q}^c = the set of irrational numbers = is not an open set.

E.g. (a, b) with $-\infty \leq a < b \leq \infty$ is an open set, which are called *open interval intervals*.

$(0, 1]$ is not an open set.

Let K be a nonempty countable subset of \mathbb{R} . Then K cannot be an open set. For example, \mathbb{N} , \mathbb{Q} , \mathbb{Z} are not open sets.

\mathbb{Q}^c = the set of irrational numbers = is not an open set.

E.g. (a, b) with $-\infty \leq a < b \leq \infty$ is an open set, which are called *open interval intervals*.

$(0, 1]$ is not an open set.

Let K be a nonempty countable subset of \mathbb{R} . Then K cannot be an open set. For example, \mathbb{N} , \mathbb{Q} , \mathbb{Z} are not open sets.

Q5: The set of irrational numbers is not an open set.

E.g. (a, b) with $-\infty \leq a < b \leq \infty$ is an open set, which are called *open interval intervals*.

$(0, 1]$ is not an open set.

Let K be a nonempty countable subset of \mathbb{R} . Then K cannot be an open set. For example, \mathbb{N} , \mathbb{Q} , \mathbb{Z} are not open sets.

\mathbb{Q}^c – the set of irrational numbers – is not an open set.

E.g. (a, b) with $-\infty \leq a < b \leq \infty$ is an open set, which are called *open interval intervals*.

$(0, 1]$ is not an open set.

Let K be a nonempty countable subset of \mathbb{R} . Then K cannot be an open set. For example, \mathbb{N} , \mathbb{Q} , \mathbb{Z} are not open sets.

\mathbb{Q}^c – the set of irrational numbers – is not an open set.

E.g. (a, b) with $-\infty \leq a < b \leq \infty$ is an open set, which are called *open interval intervals*.

$(0, 1]$ is not an open set.

Let K be a nonempty countable subset of \mathbb{R} . Then K cannot be an open set. For example, \mathbb{N} , \mathbb{Q} , \mathbb{Z} are not open sets.

\mathbb{Q}^c – the set of irrational numbers – is not an open set.

Properties of open sets

1. \mathbb{R} and \emptyset are open sets.
2. If A and B are open sets, so is $A \cap B$. (finite intersection)
3. If $\{O_i\}_{i \in I}$ is a collection of open sets, then $\bigcup_{i \in I} O_i$ is open. (arbitrary union)

Proof. Exercise.



Let $Q_n = (-1/n, 1/n)$. Then $\bigcap_{n \in \mathbb{N}} Q_n = \{0\}$ is not an open set.

Properties of open sets

1. \mathbb{R} and \emptyset are open sets.
2. If A and B are open sets, so is $A \cap B$. (finite intersection)
3. If $\{O_i\}_{i \in I}$ is a collection of open sets, then $\bigcup_{i \in I} O_i$ is open. (arbitrary union)

Proof. Exercise.



Let $Q_n = (-1/n, 1/n)$. Then $\bigcap_{n \in \mathbb{N}} Q_n = \{0\}$ is not an open set.

Properties of open sets

1. \mathbb{R} and \emptyset are open sets.
2. If A and B are open sets, so is $A \cap B$. (finite intersection)
3. If $\{O_i\}_{i \in I}$ is a collection of open sets, then $\bigcup_{i \in I} O_i$ is open. (arbitrary union)

Proof. Exercise.



Let $Q_n = (-1/n, 1/n)$. Then $\bigcap_{n \in \mathbb{N}} Q_n = \{0\}$ is not an open set.

Properties of open sets

1. \mathbb{R} and \emptyset are open sets.
2. If A and B are open sets, so is $A \cap B$. (finite intersection)
3. If $\{O_i\}_{i \in I}$ is a collection of open sets, then $\bigcup_{i \in I} O_i$ is open. (arbitrary union)

Proof. Exercise.



Let $Q_n = (-1/n, 1/n)$. Then $\bigcap_{n \in \mathbb{N}} Q_n = \{0\}$ is not an open set.

Properties of open sets

1. \mathbb{R} and \emptyset are open sets.
2. If A and B are open sets, so is $A \cap B$. (finite intersection)
3. If $\{O_i\}_{i \in I}$ is a collection of open sets, then $\bigcup_{i \in I} O_i$ is open. (arbitrary union)

Proof. Exercise.



Let $Q_n = (-1/n, 1/n)$. Then $\bigcap_{n \in \mathbb{N}} Q_n = \{0\}$ is not an open set.

Properties of open sets

1. \mathbb{R} and \emptyset are open sets.
2. If A and B are open sets, so is $A \cap B$. (finite intersection)
3. If $\{O_i\}_{i \in I}$ is a collection of open sets, then $\bigcup_{i \in I} O_i$ is open. (arbitrary union)

Proof. Exercise.



Let $Q_n = (-1/n, 1/n)$. Then $\bigcap_{n \in \mathbb{N}} Q_n = \{0\}$ is not an open set.

Properties of open sets

1. \mathbb{R} and \emptyset are open sets.
2. If A and B are open sets, so is $A \cap B$. (finite intersection)
3. If $\{O_i\}_{i \in I}$ is a collection of open sets, then $\bigcup_{i \in I} O_i$ is open. (arbitrary union)

Proof. Exercise.



Let $Q_n = (-1/n, 1/n)$. Then $\bigcap_{n \in \mathbb{N}} Q_n = \{0\}$ is not an open set.

Def. For $a, b \in \mathbb{R}^*$ with $a < b$, (a, b) is an open set, which is called an *open interval*.

Thm. Each open set O is a countable union of disjoint open intervals.

Moreover, the representation is unique in the sense that if \mathcal{C} and \mathcal{D} are two pairwise disjoint collections of open intervals whose union is O , then $\mathcal{C} = \mathcal{D}$.

Def. For $a, b \in \mathbb{R}^*$ with $a < b$, (a, b) is an open set, which is called an *open interval*.

Thm. Each open set O is a countable union of disjoint open intervals.

Moreover, the representation is unique in the sense that if \mathcal{C} and \mathcal{D} are two pairwise disjoint collections of open intervals whose union is O , then $\mathcal{C} = \mathcal{D}$.

Def. For $a, b \in \mathbb{R}^*$ with $a < b$, (a, b) is an open set, which is called an *open interval*.

Thm. Each open set O is a countable union of disjoint open intervals.

Moreover, the representation is unique in the sense that if \mathcal{C} and \mathcal{D} are two pairwise disjoint collections of open intervals whose union is O , then $\mathcal{C} = \mathcal{D}$.

Closed sets

Def. Let $E \subset \mathbb{R}$. A real number x is called a *limit point* of E if for each $\epsilon > 0$, there is a $y \in E$ such that $|y - x| < \epsilon$.

The set of all limit point of E , denoted \bar{E} , is called the *closure* of E .

Closed sets

Def. Let $E \subset \mathbb{R}$. A real number x is called a *limit point* of E if for each $\epsilon > 0$, there is a $y \in E$ such that $|y - x| < \epsilon$.

The set of all limit point of E , denoted \bar{E} , is called the *closure* of E .

E.g. $\overline{\mathbb{R}} = \mathbb{R}$ and $\overline{\emptyset} = \emptyset$.

Let $a, b \in \mathbb{R}$ such that $a < b$. Then

$$\overline{(a, b)} = \overline{(a, b]} = \overline{[a, b)} = \overline{[a, b]} = [a, b].$$

$$\overline{\mathbb{N}} = \mathbb{N} \text{ and } \overline{\mathbb{Z}} = \mathbb{Z}.$$

$$\overline{\mathbb{Q}} = \mathbb{R} \text{ and } \overline{\mathbb{Q}^c} = \mathbb{R}.$$

If A is a finite subset of \mathbb{R} , then $\overline{A} = A$.

E.g. $\overline{\mathbb{R}} = \mathbb{R}$ and $\overline{\emptyset} = \emptyset$.

Let $a, b \in \mathbb{R}$ such that $a < b$. Then

$$\overline{(a, b)} = \overline{(a, b]} = \overline{[a, b)} = \overline{[a, b]} = [a, b].$$

$$\overline{\mathbb{N}} = \mathbb{N} \text{ and } \overline{\mathbb{Z}} = \mathbb{Z}.$$

$$\overline{\mathbb{Q}} = \mathbb{R} \text{ and } \overline{\mathbb{Q}^c} = \mathbb{R}.$$

If A is a finite subset of \mathbb{R} , then $\overline{A} = A$.

E.g. $\overline{\mathbb{R}} = \mathbb{R}$ and $\overline{\emptyset} = \emptyset$.

Let $a, b \in \mathbb{R}$ such that $a < b$. Then

$$\overline{(a, b)} = \overline{(a, b]} = \overline{[a, b)} = \overline{[a, b]} = [a, b].$$

$$\overline{\mathbb{N}} = \mathbb{N} \text{ and } \overline{\mathbb{Z}} = \mathbb{Z}.$$

$$\overline{\mathbb{Q}} = \mathbb{R} \text{ and } \overline{\mathbb{Q}^c} = \mathbb{R}.$$

If A is a finite subset of \mathbb{R} , then $\overline{A} = A$.

E.g. $\overline{\mathbb{R}} = \mathbb{R}$ and $\overline{\emptyset} = \emptyset$.

Let $a, b \in \mathbb{R}$ such that $a < b$. Then

$$\overline{(a, b)} = \overline{(a, b]} = \overline{[a, b)} = \overline{[a, b]} = [a, b].$$

$$\overline{\mathbb{N}} = \mathbb{N} \text{ and } \overline{\mathbb{Z}} = \mathbb{Z}.$$

$$\overline{\mathbb{Q}} = \mathbb{R} \text{ and } \overline{\mathbb{Q}^c} = \mathbb{R}.$$

If A is a finite subset of \mathbb{R} , then $\overline{A} = A$.

E.g. $\overline{\mathbb{R}} = \mathbb{R}$ and $\overline{\emptyset} = \emptyset$.

Let $a, b \in \mathbb{R}$ such that $a < b$. Then

$$\overline{(a, b)} = \overline{(a, b]} = \overline{[a, b)} = \overline{[a, b]} = [a, b].$$

$$\overline{\mathbb{N}} = \mathbb{N} \text{ and } \overline{\mathbb{Z}} = \mathbb{Z}.$$

$$\overline{\mathbb{Q}} = \mathbb{R} \text{ and } \overline{\mathbb{Q}^c} = \mathbb{R}.$$

If A is a finite subset of \mathbb{R} , then $\overline{A} = A$.

Def. A subset $F \subset \mathbb{R}$ is said to be a *closed set* if $\overline{F} = F$, i.e, F contains all its limit points.

E.g. \mathbb{R} and \emptyset are both open and closed.

Intervals such as $[a, b]$, $[a, \infty)$, $(-\infty, b]$ with $a, b \in \mathbb{R}$ are closed sets. They are called *closed intervals*.

\mathbb{N} and \mathbb{Z} are closed sets.

The sets of rationals \mathbb{Q} and irrationals \mathbb{Q}^c are neither open nor close.

If A is a finite subset of \mathbb{R} , then A is a close set.

E.g. \mathbb{R} and \emptyset are both open and closed.

Intervals such as $[a, b]$, $[a, \infty)$, $(-\infty, b]$ with $a, b \in \mathbb{R}$ are closed sets. They are called *closed intervals*.

\mathbb{N} and \mathbb{Z} are closed sets.

The sets of rationals \mathbb{Q} and irrationals \mathbb{Q}^c are neither open nor close.

If A is a finite subset of \mathbb{R} , then A is a close set.

E.g. \mathbb{R} and \emptyset are both open and closed.

Intervals such as $[a, b]$, $[a, \infty)$, $(-\infty, b]$ with $a, b \in \mathbb{R}$ are closed sets. They are called *closed intervals*.

\mathbb{N} and \mathbb{Z} are closed sets.

The sets of rationals \mathbb{Q} and irrationals \mathbb{Q}^c are neither open nor close.

If A is a finite subset of \mathbb{R} , then A is a close set.

E.g. \mathbb{R} and \emptyset are both open and closed.

Intervals such as $[a, b]$, $[a, \infty)$, $(-\infty, b]$ with $a, b \in \mathbb{R}$ are closed sets. They are called *closed intervals*.

\mathbb{N} and \mathbb{Z} are closed sets.

The sets of rationals \mathbb{Q} and irrationals \mathbb{Q}^c are neither open nor close.

If A is a finite subset of \mathbb{R} , then A is a close set.

E.g. \mathbb{R} and \emptyset are both open and closed.

Intervals such as $[a, b]$, $[a, \infty)$, $(-\infty, b]$ with $a, b \in \mathbb{R}$ are closed sets. They are called *closed intervals*.

\mathbb{N} and \mathbb{Z} are closed sets.

The sets of rationals \mathbb{Q} and irrationals \mathbb{Q}^c are neither open nor close.

If A is a finite subset of \mathbb{R} , then A is a close set.

Thm. A set is open if and only if its complement is closed.

Or equivalently, a set is closed if and only if its complement is open.

Thm. A set is open if and only if its complement is closed.

Or equivalently, a set is closed if and only if its complement is open.

Properties of closed sets

1. \mathbb{R} and \emptyset are **closed** sets.
2. If A and B are closed sets, so is $A \cup B$. (finite union)
3. If $\{F_i\}_{i \in I}$ is a collection of closed sets, then $\bigcap_{i \in I} F_i$ is closed. (arbitrary intersection)

Let $Q_n = [-1 + \frac{1}{n}, 1 - \frac{1}{n}]$. Then $\bigcup_{i \in \mathbb{N}} Q_n = (0, 1)$ is an open set.

Properties of closed sets

1. \mathbb{R} and \emptyset are **closed** sets.
2. If A and B are **closed** sets, so is $A \cup B$. (finite **union**)
3. If $\{F_i\}_{i \in I}$ is a collection of closed sets, then $\bigcap_{i \in I} F_i$ is closed.
(arbitrary intersection)

Let $Q_n = [-1 + \frac{1}{n}, 1 - \frac{1}{n}]$. Then $\bigcup_{i \in \mathbb{N}} Q_n = (0, 1)$ is an open set.

Properties of closed sets

1. \mathbb{R} and \emptyset are **closed** sets.
2. If A and B are **closed** sets, so is $A \cup B$. (finite **union**)
3. If $\{F_i\}_{i \in I}$ is a collection of **closed** sets, then $\bigcap_{i \in I} F_i$ is **closed**.
(arbitrary **intersection**)

Let $Q_n = [-1 + \frac{1}{n}, 1 - \frac{1}{n}]$. Then $\bigcup_{i \in \mathbb{N}} Q_n = (0, 1)$ is an open set.

Properties of closed sets

1. \mathbb{R} and \emptyset are closed sets.
2. If A and B are closed sets, so is $A \cup B$. (finite union)
3. If $\{F_i\}_{i \in I}$ is a collection of closed sets, then $\bigcap_{i \in I} F_i$ is closed. (arbitrary intersection)

Let $Q_n = [-1 + \frac{1}{n}, 1 - \frac{1}{n}]$. Then $\bigcup_{i \in \mathbb{N}} Q_n = (0, 1)$ is an open set.

Properties of closed sets

1. \mathbb{R} and \emptyset are closed sets.
2. If A and B are closed sets, so is $A \cup B$. (finite union)
3. If $\{F_i\}_{i \in I}$ is a collection of closed sets, then $\bigcap_{i \in I} F_i$ is closed. (arbitrary intersection)

Let $Q_n = [-1 + \frac{1}{n}, 1 - \frac{1}{n}]$. Then $\bigcup_{i \in \mathbb{N}} Q_n = (0, 1)$ is an open set.

Properties of closed sets

1. \mathbb{R} and \emptyset are closed sets.
2. If A and B are closed sets, so is $A \cup B$. (finite union)
3. If $\{F_i\}_{i \in I}$ is a collection of closed sets, then $\bigcap_{i \in I} F_i$ is closed. (arbitrary intersection)

Let $Q_n = [-1 + \frac{1}{n}, 1 - \frac{1}{n}]$. Then $\bigcup_{i \in \mathbb{N}} Q_n = (0, 1)$ is an open set.

Relative open and closed sets

Def. Let $G \subset D \subset \mathbb{R}$.

(a) G is said to be open in D if for each $x \in G$, there is an $r > 0$ such that

$$(x - r, x + r) \cap D \subset G.$$

(b) G is said to be *closed in D* if $D \setminus G$ is open in D .

Relative open and closed sets

Def. Let $G \subset D \subset \mathbb{R}$.

(a) G is said to be **open in D** if for each $x \in G$, there is an $r > 0$ such that

$$(x - r, x + r) \cap D \subset G.$$

(b) G is said to be *closed in D* if $D \setminus G$ is open in D .

Relative open and closed sets

Def. Let $G \subset D \subset \mathbb{R}$.

(a) G is said to be **open in D** if for each $x \in G$, there is an $r > 0$ such that

$$(x - r, x + r) \cap D \subset G.$$

(b) G is said to be **closed in D** if $D \setminus G$ is open in D .

E.g.

D	G	Is G open in \mathbb{R}	Is G open in D
$[0, 2]$	$[0, 1)$		
$[0, 2]$	$[0, 1]$		
\mathbb{N}	$A \subset \mathbb{N}$		

E.g.

D	G	Is G open in \mathbb{R}	Is G open in D
$[0, 2]$	$[0, 1)$	Neither open nor closed	
$[0, 2]$	$[0, 1]$		
\mathbb{N}	$A \subset \mathbb{N}$		

E.g.

D	G	Is G open in \mathbb{R}	Is G open in D
$[0, 2]$	$[0, 1)$	Neither open nor closed	open
$[0, 2]$	$[0, 1]$		
\mathbb{N}	$A \subset \mathbb{N}$		

E.g.

D	G	Is G open in \mathbb{R}	Is G open in D
$[0, 2]$	$[0, 1)$	Neither open nor closed closed	open
$[0, 2]$	$[0, 1]$		
\mathbb{N}	$A \subset \mathbb{N}$		

E.g.

D	G	Is G open in \mathbb{R}	Is G open in D
$[0, 2]$	$[0, 1)$	Neither open nor closed	open
$[0, 2]$	$[0, 1]$	closed	closed
\mathbb{N}	$A \subset \mathbb{N}$		

E.g.

D	G	Is G open in \mathbb{R}	Is G open in D
$[0, 2]$	$[0, 1)$	Neither open nor closed	open
$[0, 2]$	$[0, 1]$	closed	closed
\mathbb{N}	$A \subset \mathbb{N}$	closed	

E.g.

D	G	Is G open in \mathbb{R}	Is G open in D
$[0, 2]$	$[0, 1)$	Neither open nor closed	open
$[0, 2]$	$[0, 1]$	closed	closed
\mathbb{N}	$A \subset \mathbb{N}$	closed	open

Thm. Let $D \subset \mathbb{R}$. A set $G \subset D$ is open in D if and only if there is an open set O of \mathbb{R} such that $G = D \cap O$.

In other words, the open sets in D are precisely the open sets of \mathbb{R} intersected with D .

Thm. Let $D \subset \mathbb{R}$. A set $G \subset D$ is open in D if and only if there is an open set O of \mathbb{R} such that $G = D \cap O$.

In other words, the open sets in D are precisely the open sets of \mathbb{R} intersected with D .

HW Ex. 2.38, 2.46, 2.47, 2.49, 2.52 on p. 63 – 64.

Chapter 3. Real Number System and Calculus

§ 3.1 Real number system

§ 3.2 Sequences of real numbers

§ 3.3 Open and closed sets

§ 3.4 Real-valued functions

§ 3.5 Liminf and limsup of sets

§ 3.6 Some techniques in calculus

Def. A *real-valued function* is a function whose range is a subset of \mathbb{R} . If $f : \Omega \rightarrow \mathbb{R}$, we say that f is a *real-valued function on Ω* .

Algebraic operations

Let f, g be real-valued functions on Ω and let $\alpha \in \mathbb{R}$. Then for all $x \in \Omega$,

$$(f + g)(x) := f(x) + g(x)$$

$$(\alpha f)(x) := \alpha f(x)$$

$$(f \cdot g)(x) := f(x)g(x)$$

Def. A *real-valued function* is a function whose range is a subset of \mathbb{R} . If $f : \Omega \rightarrow \mathbb{R}$, we say that f is a *real-valued function on Ω* .

Algebraic operations

Let f, g be real-valued functions on Ω and let $\alpha \in \mathbb{R}$. Then for all $x \in \Omega$,

$$(f + g)(x) := f(x) + g(x)$$

$$(\alpha f)(x) := \alpha f(x)$$

$$(f \cdot g)(x) := f(x)g(x)$$

Def. A *real-valued function* is a function whose range is a subset of \mathbb{R} . If $f : \Omega \rightarrow \mathbb{R}$, we say that f is a *real-valued function on Ω* .

Algebraic operations

Let f, g be real-valued functions on Ω and let $\alpha \in \mathbb{R}$. Then for all $x \in \Omega$,

$$(f + g)(x) := f(x) + g(x)$$

$$(\alpha f)(x) := \alpha f(x)$$

$$(f \cdot g)(x) := f(x)g(x)$$

(Local) Continuity

Def. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a function. f is *continuous at a point c* if the limit of $f(x)$, as x approaches c , exists and is equal to $f(c)$.

Def'. (Epsilon-delta definition) The function f is *continuous at a point c* if

$$\forall \epsilon > 0 \exists \delta > 0 \forall x \in \mathbb{R} \{ |x - c| \leq \delta \rightarrow |f(x) - f(c)| \leq \epsilon \}$$

(Local) Continuity

Def. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a function. f is *continuous at a point c* if the limit of $f(x)$, as x approaches c , exists and is equal to $f(c)$.

Def'. (Epsilon-delta definition) The function f is *continuous at a point c* if

$$\forall \epsilon > 0 \exists \delta > 0 \forall x \in \mathbb{R} \{ |x - c| \leq \delta \rightarrow |f(x) - f(c)| \leq \epsilon \}$$

Here is a more abstract definition of continuous functions:

Thm let $D \subset \mathbb{R}$ and $f : D \rightarrow \mathbb{R}$. Then f is continuous on D if and only if $f^{-1}(O)$ is open in D for each open set O in \mathbb{R} .

Here is a more abstract definition of continuous functions:

Thm let $D \subset \mathbb{R}$ and $f : D \rightarrow \mathbb{R}$. Then f is continuous on D if and only if $f^{-1}(O)$ is open in D for each open set O in \mathbb{R} .

Def. f is *left-continuous at c* if


$$\lim_{x \rightarrow c+} f(x) = f(c)$$

Def. f is *right-continuous at c* if

$$\lim_{x \rightarrow c-} f(x) = f(c)$$

Def. f is *left-continuous at c* if

$$\lim_{x \rightarrow c^+} f(x) = f(c)$$




Def. f is *right-continuous at c* if

$$\lim_{x \rightarrow c^-} f(x) = f(c)$$

Def. f is *left-continuous at c* if

$$\lim_{x \rightarrow c+} f(x) = f(c)$$




Def. f is *right-continuous at c* if

$$\lim_{x \rightarrow c-} f(x) = f(c)$$


Def. f is *left-continuous at c* if

$$\lim_{x \rightarrow c^+} f(x) = f(c)$$



Def. f is *right-continuous at c* if

$$\lim_{x \rightarrow c^-} f(x) = f(c)$$



Def. f is *lower semi-continuous at x_0* if

$$\liminf_{x \rightarrow x_0} f(x) \geq f(x_0)$$

$f(x_0)$ can be all points
at or below the blue point.

f is *upper semi-continuous at x_0*
if

$$\limsup_{x \rightarrow x_0} f(x) \leq f(x_0)$$

$f(x_0)$ can be all points
at or above the blue point

Def. f is *lower semi-continuous at x_0* if

$$\liminf_{x \rightarrow x_0} f(x) \geq f(x_0)$$

$f(x_0)$ can be all points
at or below the blue point.


f is *upper semi-continuous at x_0*
if

$$\limsup_{x \rightarrow x_0} f(x) \leq f(x_0)$$

$f(x_0)$ can be all points
at or above the blue point

Def. f is *lower semi-continuous at x_0* if

$$\liminf_{x \rightarrow x_0} f(x) \geq f(x_0)$$



$f(x_0)$ can be all points
at or below the blue point.


f is *upper semi-continuous at x_0* if

$$\limsup_{x \rightarrow x_0} f(x) \leq f(x_0)$$

$f(x_0)$ can be all points
at or above the blue point

Def. f is *lower semi-continuous at x_0* if

$$\liminf_{x \rightarrow x_0} f(x) \geq f(x_0)$$



$f(x_0)$ can be all points
at or below the blue point.


f is *upper semi-continuous at x_0* if

$$\limsup_{x \rightarrow x_0} f(x) \leq f(x_0)$$

$f(x_0)$ can be all points
at or above the blue point

Def. f is *lower semi-continuous at x_0* if

$$\liminf_{x \rightarrow x_0} f(x) \geq f(x_0)$$



$f(x_0)$ can be all points
at or below the blue point.


f is *upper semi-continuous at x_0*
if

$$\limsup_{x \rightarrow x_0} f(x) \leq f(x_0)$$

$f(x_0)$ can be all points
at or above the blue point.

Def. f is *lower semi-continuous at x_0* if

$$\liminf_{x \rightarrow x_0} f(x) \geq f(x_0)$$



$f(x_0)$ can be all points
at or below the blue point.


f is *upper semi-continuous at x_0*
if

$$\limsup_{x \rightarrow x_0} f(x) \leq f(x_0)$$

$f(x_0)$ can be all points
at or above the blue point.

Def. f is *lower semi-continuous at x_0* if


$$\liminf_{x \rightarrow x_0} f(x) \geq f(x_0)$$



$f(x_0)$ can be all points
at or below the blue point.

f is *upper semi-continuous at x_0*
if

$$\limsup_{x \rightarrow x_0} f(x) \leq f(x_0)$$



$f(x_0)$ can be all points
at or above the blue point.

(Global) Uniform Continuity

Def. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a function. Let I be an interval of \mathbb{R} . Then f is *uniformly continuous over I* if for every real number $\epsilon > 0$, there exists a real number $\delta > 0$ such that for every $x, y \in I$ with $|x - y| < \delta$, then $|f(x) - f(y)| < \epsilon$.

Def.' f is *uniformly continuous over I* if

$$\forall \epsilon > 0 \exists \delta > 0 \forall x \in I \forall y \in I \{ |x - y| < \delta \rightarrow |f(x) - f(y)| < \epsilon \}$$

(Global) Uniform Continuity

Def. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a function. Let I be an interval of \mathbb{R} . Then f is *uniformly continuous over I* if for every real number $\epsilon > 0$, there exists a real number $\delta > 0$ such that for every $x, y \in I$ with $|x - y| < \delta$, then $|f(x) - f(y)| < \epsilon$.

Def.' f is *uniformly continuous over I* if

$$\forall \epsilon > 0 \exists \delta > 0 \forall x \in I \forall y \in I \{ |x - y| < \delta \rightarrow |f(x) - f(y)| < \epsilon \}$$

f is continuous at $x_0 \in I$ iff

$$\forall \epsilon > 0 \forall x \in I \exists \delta > 0 \{ |x - x_0| < \delta \rightarrow |f(x) - f(x_0)| < \epsilon \}$$

Π_2 -form

f is uniformly continuous over I iff

$$\forall \epsilon > 0 \exists \delta > 0 \forall x \in I \forall y \in I \{ |x - y| < \delta \rightarrow |f(x) - f(y)| < \epsilon \}$$

Π_3 -form

f is continuous at $x_0 \in I$ iff

$$\forall \epsilon > 0 \, \forall x \in I \, \exists \delta > 0 \, \{|x - x_0| < \delta \rightarrow |f(x) - f(x_0)| < \epsilon\}$$

Π_2 -form

f is uniformly continuous over I iff

$$\forall \epsilon > 0 \, \exists \delta > 0 \, \forall x \in I \, \forall y \in I \, \{|x - y| < \delta \rightarrow |f(x) - f(y)| < \epsilon\}$$

Π_3 -form

f is continuous at $x_0 \in I$ iff

$$\forall \epsilon > 0 \forall x \in I \exists \delta > 0 \{ |x - x_0| < \delta \rightarrow |f(x) - f(x_0)| < \epsilon \}$$

Π_2 -form

f is uniformly continuous over I iff

$$\forall \epsilon > 0 \exists \delta > 0 \forall x \in I \forall y \in I \{ |x - y| < \delta \rightarrow |f(x) - f(y)| < \epsilon \}$$

Π_3 -form

f is continuous at $x_0 \in I$ iff

$$\forall \epsilon > 0 \forall x \in I \exists \delta > 0 \{ |x - x_0| < \delta \rightarrow |f(x) - f(x_0)| < \epsilon \}$$

Π_2 -form

f is uniformly continuous over I iff

$$\forall \epsilon > 0 \exists \delta > 0 \forall x \in I \forall y \in I \{ |x - y| < \delta \rightarrow |f(x) - f(y)| < \epsilon \}$$

Π_3 -form

Properties

Prop. 1 Every uniformly continuous function is continuous, but the converse does not hold.

Ex. $f(x) = x^3$ is a continuous functions on $I = \mathbb{R}$. Show that f is not uniformly continuous on $I = \mathbb{R}$.

Sol. In order to show f is not uniformly continuous on I , we need to show

$$\begin{aligned} & \neg \left(\forall \epsilon > 0 \exists \delta > 0 \forall x \in I \forall y \in I \{ |x - y| < \delta \rightarrow |f(x) - f(y)| < \epsilon \} \right) \\ \Leftrightarrow & \exists \epsilon > 0 \forall \delta > 0 \exists x \in I \exists y \in I \neg \{ |x - y| < \delta \rightarrow |f(x) - f(y)| < \epsilon \} \\ \Leftrightarrow & \exists \epsilon > 0 \forall \delta > 0 \exists x \in I \exists y \in I \neg \{ \neg \{ |x - y| < \delta \} \vee |f(x) - f(y)| < \epsilon \} \\ \Leftrightarrow & \exists \epsilon > 0 \forall \delta > 0 \exists x \in I \exists y \in I \{ |x - y| < \delta \wedge |f(x) - f(y)| \geq \epsilon \} \end{aligned}$$

Properties

Prop. 1 Every uniformly continuous function is continuous, but the converse does not hold.

Ex. $f(x) = x^3$ is a continuous functions on $I = \mathbb{R}$. Show that f is not uniformly continuous on $I = \mathbb{R}$.

Sol. In order to show f is not uniformly continuous on I , we need to show

$$\begin{aligned} & \neg \left(\forall \epsilon > 0 \exists \delta > 0 \forall x \in I \forall y \in I \{ |x - y| < \delta \rightarrow |f(x) - f(y)| < \epsilon \} \right) \\ \Leftrightarrow & \exists \epsilon > 0 \forall \delta > 0 \exists x \in I \exists y \in I \neg \{ |x - y| < \delta \rightarrow |f(x) - f(y)| < \epsilon \} \\ \Leftrightarrow & \exists \epsilon > 0 \forall \delta > 0 \exists x \in I \exists y \in I \neg \{ \neg \{ |x - y| < \delta \} \vee |f(x) - f(y)| < \epsilon \} \\ \Leftrightarrow & \exists \epsilon > 0 \forall \delta > 0 \exists x \in I \exists y \in I \{ |x - y| < \delta \wedge |f(x) - f(y)| \geq \epsilon \} \end{aligned}$$

Properties

Prop. 1 Every uniformly continuous function is continuous, but the converse does not hold.

Ex. $f(x) = x^3$ is a continuous functions on $I = \mathbb{R}$. Show that f is not uniformly continuous on $I = \mathbb{R}$.

Sol. In order to show f is not uniformly continuous on I , we need to show

$$\begin{aligned} & \neg \left(\forall \epsilon > 0 \exists \delta > 0 \forall x \in I \forall y \in I \{ |x - y| < \delta \rightarrow |f(x) - f(y)| < \epsilon \} \right) \\ \Leftrightarrow & \exists \epsilon > 0 \forall \delta > 0 \exists x \in I \exists y \in I \neg \{ |x - y| < \delta \rightarrow |f(x) - f(y)| < \epsilon \} \\ \Leftrightarrow & \exists \epsilon > 0 \forall \delta > 0 \exists x \in I \exists y \in I \neg \{ \neg \{ |x - y| < \delta \} \vee |f(x) - f(y)| < \epsilon \} \\ \Leftrightarrow & \exists \epsilon > 0 \forall \delta > 0 \exists x \in I \exists y \in I \{ |x - y| < \delta \wedge |f(x) - f(y)| \geq \epsilon \} \end{aligned}$$

Sol. (Continued) In other words, we need to find $\epsilon > 0$ such that no matter how small $\delta > 0$ is chosen, we can always find out $x, y \in I$ with $|x - y| < \delta$ and $|f(x) - f(y)| \geq \epsilon$.

Let's choose $\epsilon = 1$. For any $\delta > 0$, we can choose

$$x = \frac{1}{\sqrt{\delta}} \quad \text{and} \quad y = \frac{1}{\sqrt{\delta}} + \frac{\delta}{3}.$$

Then we see that

$$|x - y| \leq \delta$$

and

$$\begin{aligned} |f(x) - f(y)| &= |x^3 - y^3| \\ &= |x - y| \times |x^2 + xy + y^2| \\ &\geq \frac{\delta}{3} \times \left(\frac{1}{(\sqrt{\delta})^2} + \frac{1}{\sqrt{\delta}} \times \frac{1}{\sqrt{\delta}} + \frac{1}{(\sqrt{\delta})^2} \right) \\ &= 1 = \epsilon. \end{aligned}$$

□

Sol. (Continued) In other words, we need to find $\epsilon > 0$ such that no matter how small $\delta > 0$ is chosen, we can always find out $x, y \in I$ with $|x - y| < \delta$ and $|f(x) - f(y)| \geq \epsilon$.

Let's choose $\epsilon = 1$. For any $\delta > 0$, we can choose

$$x = \frac{1}{\sqrt{\delta}} \quad \text{and} \quad y = \frac{1}{\sqrt{\delta}} + \frac{\delta}{3}.$$

Then we see that

$$|x - y| \leq \delta$$

and

$$\begin{aligned} |f(x) - f(y)| &= |x^3 - y^3| \\ &= |x - y| \times |x^2 + xy + y^2| \\ &\geq \frac{\delta}{3} \times \left(\frac{1}{(\sqrt{\delta})^2} + \frac{1}{\sqrt{\delta}} \times \frac{1}{\sqrt{\delta}} + \frac{1}{(\sqrt{\delta})^2} \right) \\ &= 1 = \epsilon. \end{aligned}$$

□

Prop. 2 If I is compact⁷ set such as $I = [a, b]$, then

f is continuous at all points in $I \iff f$ is uniformly continuous on I .

E.g. $f(x) = 1/x$ is not uniformly continuous on $(0, 1)$.

$f(x) = x^3$ is uniformly continuous on $[-1, 1]$ but neither on \mathbb{R} nor on $[0, \infty)$.

⁷namely, bounded and closed

Prop. 2 If I is compact⁷ set such as $I = [a, b]$, then

f is continuous at all points in $I \iff f$ is uniformly continuous on I .

E.g. $f(x) = 1/x$ is not uniformly continuous on $(0, 1)$.

$f(x) = x^3$ is uniformly continuous on $[-1, 1]$ but neither on \mathbb{R} nor on $[0, \infty)$.

⁷namely, bounded and closed

Prop. 2 If I is compact⁷ set such as $I = [a, b]$, then

f is continuous at all points in $I \iff f$ is uniformly continuous on I .

E.g. $f(x) = 1/x$ is not uniformly continuous on $(0, 1)$.

$f(x) = x^3$ is uniformly continuous on $[-1, 1]$ but neither on \mathbb{R} nor on $[0, \infty)$.

⁷namely, bounded and closed

Why does it matter at all?

Answer: Uniformly continuous functions map Cauchy sequences to Cauchy sequences.

Thm Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a uniformly continuous function and let $\{x_n\}_{n=1}^{\infty}$ be a Cauchy sequence. Then $\{f(x_n)\}_{n=1}^{\infty}$ is also a Cauchy sequence.

Why does it matter at all?

Answer: Uniformly continuous functions map Cauchy sequences to Cauchy sequences.

Thm Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a uniformly continuous function and let $\{x_n\}_{n=1}^{\infty}$ be a Cauchy sequence. Then $\{f(x_n)\}_{n=1}^{\infty}$ is also a Cauchy sequence.

Notation For $D \subset \mathbb{R}$, let $C(D)$ denote the set of continuous functions defined on D .

Thm (*Algebra of $C(D)$*) Let $D \subset \mathbb{R}$. Then the collection $C(D)$ of continuous functions on D is an algebra of functions, that is, for all $f, g \in C(D)$ and $\alpha \in \mathbb{R}$,

$$f + g \in C(D)$$

$$\alpha f \in C(D)$$

$$f \cdot g \in C(D)$$

Remark Can one add one more operation in this algebra: Let $\{f_n\}_{n=1}^{\infty}$ be a sequence of functions in $C(D)$, under what condition the limit $f_n \rightarrow f$ is closed?

Notation For $D \subset \mathbb{R}$, let $C(D)$ denote the set of continuous functions defined on D .

Thm (*Algebra of $C(D)$*) Let $D \subset \mathbb{R}$. Then the collection $C(D)$ of continuous functions on D is an algebra of functions, that is, for all $f, g \in C(D)$ and $\alpha \in \mathbb{R}$,

$$f + g \in C(D)$$

$$\alpha f \in C(D)$$

$$f \cdot g \in C(D)$$

Remark Can one add one more operation in this algebra: Let $\{f_n\}_{n=1}^{\infty}$ be a sequence of functions in $C(D)$, under what condition the limit $f_n \rightarrow f$ is closed?

Notation For $D \subset \mathbb{R}$, let $C(D)$ denote the set of continuous functions defined on D .

Thm (*Algebra of $C(D)$*) Let $D \subset \mathbb{R}$. Then the collection $C(D)$ of continuous functions on D is an algebra of functions, that is, for all $f, g \in C(D)$ and $\alpha \in \mathbb{R}$,

$$f + g \in C(D)$$

$$\alpha f \in C(D)$$

$$f \cdot g \in C(D)$$

Remark Can one add one more operation in this algebra: Let $\{f_n\}_{n=1}^{\infty}$ be a sequence of functions in $C(D)$, under what condition the limit $f_n \rightarrow f$ is closed?

Def. $\{f_n\}_{n=1}^{\infty}$ be a sequence of real-valued functions on Ω , namely, $f_n : \Omega \rightarrow \mathbb{R}$ for each $n \in \mathbb{N}$.

We say that $\{f_n\}_{n=1}^{\infty}$ *converges pointwise on Ω* if for each $x \in \Omega$, the sequence $\{f_n(x)\}_{n=1}^{\infty}$ of real numbers converges in \mathbb{R} .

Def. $\{f_n\}_{n=1}^{\infty}$ be a sequence of real-valued functions on Ω , namely, $f_n : \Omega \rightarrow \mathbb{R}$ for each $n \in \mathbb{N}$.

We say that $\{f_n\}_{n=1}^{\infty}$ *converges pointwise on Ω* if for each $x \in \Omega$, the sequence $\{f_n(x)\}_{n=1}^{\infty}$ of real numbers converges in \mathbb{R} .

E.g.

- (a) $f_n \in C(\mathbb{R})$ defined as $f_n = (1 + x/n)^n$. Then f_n converges pointwise on \mathbb{R} to $f(x) = e^x$. It is clear that $f \in C(\mathbb{R})$.
- (b) Let $D = [0, 1]$ and $f_n \in C(D)$ be defined as $f_n = x^n$. Then f_n converges pointwise to

$$f(x) = \begin{cases} 0 & \text{if } 0 \leq x < 1 \\ 1 & \text{if } x = 1 \end{cases}$$

It is clear that $f \notin C(D)$.

E.g.

(a) $f_n \in C(\mathbb{R})$ defined as $f_n = (1 + x/n)^n$. Then f_n converges pointwise on \mathbb{R} to $f(x) = e^x$. It is clear that $f \in C(\mathbb{R})$.

(b) Let $D = [0, 1]$ and $f_n \in C(D)$ be defined as $f_n = x^n$. Then f_n converges pointwise to

$$f(x) = \begin{cases} 0 & \text{if } 0 \leq x < 1 \\ 1 & \text{if } x = 1 \end{cases}$$

It is clear that $f \notin C(D)$.

E.g.

- (a) $f_n \in C(\mathbb{R})$ defined as $f_n = (1 + x/n)^n$. Then f_n converges pointwise on \mathbb{R} to $f(x) = e^x$. It is clear that $f \in C(\mathbb{R})$.
- (b) Let $D = [0, 1]$ and $f_n \in C(D)$ be defined as $f_n = x^n$. Then f_n converges pointwise to

$$f(x) = \begin{cases} 0 & \text{if } 0 \leq x < 1 \\ 1 & \text{if } x = 1 \end{cases}$$

It is clear that $f \notin C(D)$.

Def. Let \mathcal{F} be a collection of real-valued functions on Ω . We say that \mathcal{F} is *closed under pointwise limits* if whenever $\{f_n\}_{n=1}^{\infty} \subset \mathcal{F}$ and $f_n \rightarrow f$ pointwise on Ω , then $f \in \mathcal{F}$.

Def. Let $\{f_n\}_{n=1}^{\infty}$ be a sequence of real-valued functions on a set Ω .

We say that $\{f_n\}_{n=1}^{\infty}$ *converges uniformly* to the real-valued function f on Ω , if

$$\boxed{\forall \epsilon > 0 \exists N \in \mathbb{N} \forall n \geq N} \quad \forall x \in \Omega \quad |f_n(x) - f(x)| < \epsilon,$$

written as $f_n \rightarrow f$ *uniformly*.

Recall $\{f_n\}_{n=1}^{\infty}$ converges pointwise to f if

$$\forall x \in \Omega \quad \boxed{\forall \epsilon > 0 \exists N \in \mathbb{N} \forall n \geq N} \quad |f_n(x) - f(x)| < \epsilon,$$

Def. Let $\{f_n\}_{n=1}^{\infty}$ be a sequence of real-valued functions on a set Ω .

We say that $\{f_n\}_{n=1}^{\infty}$ *converges uniformly* to the real-valued function f on Ω , if

$$\boxed{\forall \epsilon > 0 \exists N \in \mathbb{N} \forall n \geq N} \quad \forall x \in \Omega \quad |f_n(x) - f(x)| < \epsilon,$$

written as $f_n \rightarrow f$ *uniformly*.

Recall $\{f_n\}_{n=1}^{\infty}$ converges pointwise to f if

$$\forall x \in \Omega \quad \boxed{\forall \epsilon > 0 \exists N \in \mathbb{N} \forall n \geq N} \quad |f_n(x) - f(x)| < \epsilon,$$

Def. Let $\{f_n\}_{n=1}^{\infty}$ be a sequence of real-valued functions on a set Ω .

We say that $\{f_n\}_{n=1}^{\infty}$ *converges uniformly* to the real-valued function f on Ω , if

$$\boxed{\forall \epsilon > 0 \exists N \in \mathbb{N} \forall n \geq N} \forall x \in \Omega \quad |f_n(x) - f(x)| < \epsilon,$$

written as $f_n \rightarrow f$ *uniformly*.

Recall $\{f_n\}_{n=1}^{\infty}$ converges pointwise to f if

$$\forall x \in \Omega \quad \boxed{\forall \epsilon > 0 \exists N \in \mathbb{N} \forall n \geq N} \quad |f_n(x) - f(x)| < \epsilon,$$

Def. Let $\{f_n\}_{n=1}^{\infty}$ be a sequence of real-valued functions on a set Ω .

We say that $\{f_n\}_{n=1}^{\infty}$ *converges uniformly* to the real-valued function f on Ω , if

$$\boxed{\forall \epsilon > 0 \exists N \in \mathbb{N} \forall n \geq N} \forall x \in \Omega |f_n(x) - f(x)| < \epsilon,$$

written as $f_n \rightarrow f$ *uniformly*.

Recall $\{f_n\}_{n=1}^{\infty}$ *converges pointwise* to f if

$$\forall x \in \Omega \boxed{\forall \epsilon > 0 \exists N \in \mathbb{N} \forall n \geq N} |f_n(x) - f(x)| < \epsilon,$$

Prop. Let $D \subset \mathbb{R}$. Suppose that $\{f_n\}_{n=1}^{\infty} \subset C(D)$ and that $f_n \rightarrow f$ uniformly.
Then $f \in C(D)$.

Proof.

Prop. Let $D \subset \mathbb{R}$. Suppose that $\{f_n\}_{n=1}^{\infty} \subset C(D)$ and that $f_n \rightarrow f$ uniformly.
Then $f \in C(D)$.

Proof.

Prop. Let $D \subset \mathbb{R}$. Suppose that $\{f_n\}_{n=1}^{\infty} \subset C(D)$ and that $f_n \rightarrow f$ uniformly.
Then $f \in C(D)$.

Proof.

Therefore, the collection $\mathcal{C}(D)$ of real-valued continuous functions is closed under: $+$, \cdot , scalar multiplication, and uniform convergence.

HW Ex. 3.59, 2.60, 2.61, 2.66, on p. 71 – 73.

Chapter 3. Real Number System and Calculus

§ 3.1 Real number system

§ 3.2 Sequences of real numbers

§ 3.3 Open and closed sets

§ 3.4 Real-valued functions

§ 3.5 Liminf and limsup of sets

§ 3.6 Some techniques in calculus

Some part of subsection is taken from Chapter 1 Section 4 of

*P. Billingsley, **Probability and Measure**, Wiley, 1995.*

Def. For a sequence $A_1, A_2 \cdots$ of sets, define the *limits superior and inferior* of the sequence $\{A_n\}$ as

$$\limsup_n A_n := \bigcap_{n=1}^{\infty} \bigcup_{k=n}^{\infty} A_k, \quad \text{and} \quad \liminf_n A_n := \bigcup_{n=1}^{\infty} \bigcap_{k=n}^{\infty} A_k.$$

Remark Both $\limsup_n A_n$ and $\liminf_n A_n$ are sets.

Def. For a sequence $A_1, A_2 \cdots$ of sets, define the *limits superior and inferior* of the sequence $\{A_n\}$ as

$$\limsup_n A_n := \bigcap_{n=1}^{\infty} \bigcup_{k=n}^{\infty} A_k, \quad \text{and} \quad \liminf_n A_n := \bigcup_{n=1}^{\infty} \bigcap_{k=n}^{\infty} A_k.$$

Remark Both $\limsup_n A_n$ and $\liminf_n A_n$ are sets.

Use the relation:

set	logic
\cap	\forall
\cup	\exists

$$\begin{aligned}
 \omega \in \limsup_n A_n &\iff \omega \in \bigcap_{n=1}^{\infty} \bigcup_{k=n}^{\infty} A_k \\
 &\iff (\forall n \geq 1) (\exists k \geq n) \omega \in A_k \\
 &\iff \omega \text{ lies in } \textit{infinitely many} \text{ of the } A_n
 \end{aligned}$$

Notation

$$\limsup_n A_n = [A_n \text{ i.o.}]$$

Use the relation:

set	logic
\bigcap	\forall
\bigcup	\exists

$$\begin{aligned}
 \omega \in \limsup_n A_n &\iff \omega \in \bigcap_{n=1}^{\infty} \bigcup_{k=n}^{\infty} A_k \\
 &\iff (\forall n \geq 1) (\exists k \geq n) \omega \in A_k \\
 &\iff \omega \text{ lies in } \textit{infinitely many} \text{ of the } A_n
 \end{aligned}$$

Notation

$$\limsup_n A_n = [A_n \text{ i.o.}]$$

Use the relation:

set	logic
\bigcap	\forall
\bigcup	\exists

$$\begin{aligned}
 \omega \in \limsup_n A_n &\iff \omega \in \bigcap_{n=1}^{\infty} \bigcup_{k=n}^{\infty} A_k \\
 &\iff (\forall n \geq 1) (\exists k \geq n) \omega \in A_k \\
 &\iff \omega \text{ lies in } \textit{infinitely many} \text{ of the } A_n
 \end{aligned}$$

Notation

$$\limsup_n A_n = [A_n \text{ i.o.}]$$

Use the relation:

set	logic
\cap	\forall
\cup	\exists

$$\begin{aligned}
 \omega \in \liminf_n A_n &\iff \omega \in \bigcup_{n=1}^{\infty} \bigcap_{k=n}^{\infty} A_k \\
 &\iff (\exists n \geq 1) (\forall k \geq n) \omega \in A_k \\
 &\iff \omega \text{ lies in } \textit{all but finitely many} \text{ of the } A_n
 \end{aligned}$$

Notation

$$\liminf_n A_n = [A_n \text{ all but finitely many}]$$

Use the relation:

set	logic
\bigcap	\forall
\bigcup	\exists

$$\begin{aligned}
 \omega \in \liminf_n A_n &\iff \omega \in \bigcup_{n=1}^{\infty} \bigcap_{k=n}^{\infty} A_k \\
 &\iff (\exists n \geq 1) (\forall k \geq n) \omega \in A_k \\
 &\iff \omega \text{ lies in } \textit{all but finitely many} \text{ of the } A_n
 \end{aligned}$$

Notation

$$\liminf_n A_n = [A_n \text{ all but finitely many}]$$

Use the relation:

set	logic
\bigcap	\forall
\bigcup	\exists

$$\begin{aligned}
 \omega \in \liminf_n A_n &\iff \omega \in \bigcup_{n=1}^{\infty} \bigcap_{k=n}^{\infty} A_k \\
 &\iff (\exists n \geq 1) (\forall k \geq n) \omega \in A_k \\
 &\iff \omega \text{ lies in } \textit{all but finitely many} \text{ of the } A_n
 \end{aligned}$$

Notation

$$\liminf_n A_n = [A_n \text{ all but finitely many}]$$

Def. If both $\limsup_n A_n$ and $\liminf_n A_n$ exist and are equal, then the *limit set* of the sequence $\{A_n\}$ is defined to be

$$\lim_n A_n := \limsup_n A_n = \liminf_n A_n,$$

which is also often written as $A_n \rightarrow A$.

Def. If both $\limsup_n A_n$ and $\liminf_n A_n$ exist and are equal, then the *limit set* of the sequence $\{A_n\}$ is defined to be

$$\lim_n A_n := \limsup_n A_n = \liminf_n A_n,$$

which is also often written as $A_n \rightarrow A$.

Properties

(i) By De Morgan's law,

$$\liminf_n A_n = \bigcup_{n=1}^{\infty} \bigcap_{k=n}^{\infty} A_k = \bigcup_{n=1}^{\infty} \left(\bigcup_{k=n}^{\infty} A_k^c \right)^c = \left(\bigcap_{n=1}^{\infty} \bigcup_{k=n}^{\infty} A_k^c \right)^c = \left(\limsup_n A_n^c \right)^c$$

Properties

(ii) Monotone increasing and decreasing sets:

$$\begin{array}{ccc}
 \left(\bigcap_{k=n}^{\infty} A_k \right) & \uparrow & \left(\bigcup_{n=1}^{\infty} \bigcap_{k=n}^{\infty} A_k \right) = \liminf_n A_n \implies \lim_{n \rightarrow \infty} \bigcap_{k=n}^{\infty} A_k = \liminf_n A_n \\
 \cap & & \cap \\
 A_n & & \\
 \cap & & \\
 \left(\bigcup_{k=n}^{\infty} A_k \right) & \downarrow & \left(\bigcap_{n=1}^{\infty} \bigcup_{k=n}^{\infty} A_k \right) = \limsup_n A_n \implies \lim_{n \rightarrow \infty} \bigcup_{k=n}^{\infty} A_k = \limsup_n A_n \\
 \cup & & \cup
 \end{array}$$

Interpretation Under a Probability Space

Suppose that $\{A_n\}$ are events from a probability space (Ω, \mathbb{P})

(i) The above Property (ii) can be translated to a probability statement:

$$\begin{array}{ccc}
 \mathbb{P} \left(\bigcap_{k=n}^{\infty} A_k \right) & \uparrow & \mathbb{P} \left(\bigcup_{n=1}^{\infty} \bigcap_{k=n}^{\infty} A_k \right) = \mathbb{P} \left(\liminf_n A_n \right) \\
 & & \uparrow \wedge \\
 & & \liminf_{n \rightarrow \infty} \mathbb{P}(A_n) \\
 \mathbb{P}(A_n) & & \uparrow \wedge \\
 & & \limsup_{n \rightarrow \infty} \mathbb{P}(A_n) \\
 & & \uparrow \wedge \\
 \mathbb{P} \left(\bigcup_{k=n}^{\infty} A_k \right) & \downarrow & \mathbb{P} \left(\bigcap_{n=1}^{\infty} \bigcup_{k=n}^{\infty} A_k \right) = \mathbb{P} \left(\limsup_n A_n \right)
 \end{array}$$

(ii) *Borel Cantelli lemma*

$$\sum_n \mathbb{P}(A_n) \text{ converges} \quad \Rightarrow \quad \mathbb{P}(A_n \text{ i.o.}) = 0.$$

Proof.

$$\begin{aligned} 1 &\geq \mathbb{P}(A_n^c \text{ all but finitely many}) = 1 - \mathbb{P}(\{A_n^c \text{ all but finitely many}\}^c) \\ &= 1 - \mathbb{P}(A_n \text{ i.o.}) \\ &= 1 - \lim_{n \rightarrow \infty} \mathbb{P}\left(\bigcup_{k \geq n} A_k\right) \\ &\geq 1 - \lim_{n \rightarrow \infty} \sum_{k=n}^{\infty} \mathbb{P}(A_k) \\ &= 1 - 0 = 1. \end{aligned}$$

□

(ii) *Borel Cantelli lemma*

$$\sum_n \mathbb{P}(A_n) \text{ converges} \quad \Rightarrow \quad \mathbb{P}(A_n \text{ i.o.}) = 0.$$

Proof.

$$\begin{aligned} 1 &\geq \mathbb{P}(A_n^c \text{ all but finitely many}) = 1 - \mathbb{P}(\{A_n^c \text{ all but finitely many}\}^c) \\ &= 1 - \mathbb{P}(A_n \text{ i.o.}) \\ &= 1 - \lim_{n \rightarrow \infty} \mathbb{P}\left(\bigcup_{k \geq n} A_k\right) \\ &\geq 1 - \lim_{n \rightarrow \infty} \sum_{k=n}^{\infty} \mathbb{P}(A_k) \\ &= 1 - 0 = 1. \end{aligned}$$

□

Exercise

(i) Let $A_n = \left(-\frac{1}{n}, 1 - \frac{1}{n}\right]$:

$$A_1 = (-1, 0]$$

$$A_2 = \left(-\frac{1}{2}, \frac{1}{2}\right]$$

$$A_3 = \left(-\frac{1}{3}, \frac{2}{3}\right]$$

$$A_4 = \left(-\frac{1}{4}, \frac{3}{4}\right]$$

$$\vdots \quad \vdots$$

$$A_{100} = \left(-\frac{1}{100}, \frac{99}{100}\right]$$

$$\vdots \quad \vdots$$

Show that

$$\limsup_n A_n = \liminf_n A_n = [0, 1).$$

Exercise

(i) Let $A_n = \left(-\frac{1}{n}, 1 - \frac{1}{n}\right]$:

$$A_1 = (-1, 0]$$

$$A_2 = \left(-\frac{1}{2}, \frac{1}{2}\right]$$

$$A_3 = \left(-\frac{1}{3}, \frac{2}{3}\right]$$

$$A_4 = \left(-\frac{1}{4}, \frac{3}{4}\right]$$

$$\vdots \quad \vdots$$

$$A_{100} = \left(-\frac{1}{100}, \frac{99}{100}\right]$$

$$\vdots \quad \vdots$$

Show that

$$\limsup_n A_n = \liminf_n A_n = [0, 1).$$

Sol.

$$\liminf_n A_n = \bigcup_{n=1}^{\infty} \bigcap_{k=n}^{\infty} \left(-\frac{1}{k}, \frac{k-1}{k}\right] = \bigcup_{n=1}^{\infty} \left[0, \frac{n-1}{n}\right] = [0, 1)$$

and

$$\limsup_n A_n = \bigcap_{n=1}^{\infty} \bigcup_{k=n}^{\infty} \left(-\frac{1}{k}, \frac{k-1}{k}\right] = \bigcap_{n=1}^{\infty} \left(-\frac{1}{n}, 1\right) = [0, 1).$$

Finally,

$$\limsup_n A_n = \liminf_n A_n = [0, 1).$$

□

Sol.

$$\liminf_n A_n = \bigcup_{n=1}^{\infty} \bigcap_{k=n}^{\infty} \left(-\frac{1}{k}, \frac{k-1}{k} \right] = \bigcup_{n=1}^{\infty} \left[0, \frac{n-1}{n} \right] = [0, 1)$$

and

$$\limsup_n A_n = \bigcap_{n=1}^{\infty} \bigcup_{k=n}^{\infty} \left(-\frac{1}{k}, \frac{k-1}{k} \right] = \bigcap_{n=1}^{\infty} \left(-\frac{1}{n}, 1 \right) = (-1, 1].$$

Finally,

$$\limsup_n A_n = \liminf_n A_n = [0, 1).$$

□

Sol.

$$\liminf_n A_n = \bigcup_{n=1}^{\infty} \bigcap_{k=n}^{\infty} \left(-\frac{1}{k}, \frac{k-1}{k} \right] = \bigcup_{n=1}^{\infty} \left[0, \frac{n-1}{n} \right] = [0, 1)$$

and

$$\limsup_n A_n = \bigcap_{n=1}^{\infty} \bigcup_{k=n}^{\infty} \left(-\frac{1}{k}, \frac{k-1}{k} \right] = \bigcap_{n=1}^{\infty} \left(-\frac{1}{n}, 1 \right) = (-1, 1].$$

Finally,

$$\limsup_n A_n = \liminf_n A_n = [0, 1).$$

□

Exercise

(ii) Let $A_n = \left(\frac{(-1)^n}{n}, 1 - \frac{(-1)^n}{n} \right]$:

$$A_1 = (-1, 2] \qquad A_2 = \left(\frac{1}{2}, \frac{1}{2} \right]$$

$$A_3 = \left(-\frac{1}{3}, \frac{4}{3} \right] \qquad A_4 = \left(\frac{1}{4}, \frac{3}{4} \right]$$

$$A_5 = \left(-\frac{1}{5}, \frac{6}{5} \right] \qquad A_6 = \left(\frac{1}{6}, \frac{5}{6} \right]$$

$$\vdots \qquad \vdots \qquad \qquad \qquad \vdots \qquad \vdots$$

$$A_{99} = \left(-\frac{1}{99}, \frac{100}{99} \right] \qquad A_{100} = \left(\frac{1}{100}, \frac{99}{100} \right]$$

$$\vdots \qquad \vdots \qquad \qquad \qquad \vdots \qquad \vdots$$

Show that $\lim_n A_n$ doesn't exist by demonstrating that

$$\liminf_n A_n = (0, 1) \subset [0, 1] = \limsup_n A_n.$$

Exercise

(ii) Let $A_n = \left(\frac{(-1)^n}{n}, 1 - \frac{(-1)^n}{n} \right]$:

$$A_1 = (-1, 2] \qquad A_2 = \left(\frac{1}{2}, \frac{1}{2} \right]$$

$$A_3 = \left(-\frac{1}{3}, \frac{4}{3} \right] \qquad A_4 = \left(\frac{1}{4}, \frac{3}{4} \right]$$

$$A_5 = \left(-\frac{1}{5}, \frac{6}{5} \right] \qquad A_6 = \left(\frac{1}{6}, \frac{5}{6} \right]$$

$$\vdots \qquad \vdots \qquad \qquad \qquad \vdots \qquad \vdots$$

$$A_{99} = \left(-\frac{1}{99}, \frac{100}{99} \right] \qquad A_{100} = \left(\frac{1}{100}, \frac{99}{100} \right]$$

$$\vdots \qquad \vdots \qquad \qquad \qquad \vdots \qquad \vdots$$

Show that $\lim_n A_n$ doesn't exist by demonstrating that

$$\liminf_n A_n = (0, 1) \subset [0, 1] = \limsup_n A_n.$$

Sol.

$$\begin{aligned}
 & \liminf_n A_n \\
 &= \bigcup_{n=1}^{\infty} \bigcap_{k=n}^{\infty} \left(\frac{(-1)^k}{k}, \frac{k - (-1)^k}{k} \right] \\
 &= \left\{ \bigcup_{n=1,3,5}^{\infty} \bigcap_{k=n}^{\infty} \left(\frac{(-1)^k}{k}, \frac{k - (-1)^k}{k} \right] \right\} \cup \left\{ \bigcup_{n=2,4,6}^{\infty} \bigcap_{k=n}^{\infty} \left(\frac{(-1)^k}{k}, \frac{k - (-1)^k}{k} \right] \right\} \\
 &= \left\{ \bigcup_{n=1,3,5}^{\infty} \left(\frac{1}{n+1}, \frac{n}{n+1} \right] \right\} \cup \left\{ \bigcup_{n=2,4,6}^{\infty} \left(\frac{1}{n}, \frac{n-1}{n} \right] \right\} \\
 &= (0, 1) \cup (0, 1) \\
 &= (0, 1)
 \end{aligned}$$

Sol. (continued) Similarly,

$$\begin{aligned}
 & \limsup_n A_n \\
 &= \bigcap_{n=1}^{\infty} \bigcup_{k=n}^{\infty} \left(\frac{(-1)^k}{k}, \frac{k - (-1)^k}{k} \right] \\
 &= \left\{ \bigcap_{n=1,3,5}^{\infty} \bigcup_{k=n}^{\infty} \left(\frac{(-1)^k}{k}, \frac{k - (-1)^k}{k} \right] \right\} \cap \left\{ \bigcap_{n=2,4,6}^{\infty} \bigcup_{k=n}^{\infty} \left(\frac{(-1)^k}{k}, \frac{k - (-1)^k}{k} \right] \right\} \\
 &= \left\{ \bigcap_{n=1,3,5}^{\infty} \left(-\frac{1}{n}, \frac{n+1}{n} \right] \right\} \cap \left\{ \bigcap_{n=2,4,6}^{\infty} \left(-\frac{1}{n+1}, \frac{n+2}{n+1} \right] \right\} \\
 &= [0, 1] \cap [0, 1] \\
 &= [0, 1]
 \end{aligned}$$

□

HW (Ex. 1.8 (c) on p. 11 of McDonald and Weiss) Let

$$A_n = \begin{cases} \left[0, 1 + \frac{1}{n}\right] & \text{if } n \text{ is an even integer,} \\ \left[-1 - \frac{1}{n}, 0\right] & \text{if } n \text{ is an odd integer.} \end{cases}$$

Determine $\liminf_{n \rightarrow \infty} A_n$ and $\limsup_{n \rightarrow \infty} A_n$.

Solution:

$$\liminf_{n \rightarrow \infty} A_n = \{0\} \subset [0, 1] = \limsup_{n \rightarrow \infty} A_n$$

HW (Ex. 1.8 (c) on p. 11 of McDonald and Weiss) Let

$$A_n = \begin{cases} \left[0, 1 + \frac{1}{n}\right] & \text{if } n \text{ is an even integer,} \\ \left[-1 - \frac{1}{n}, 0\right] & \text{if } n \text{ is an odd integer.} \end{cases}$$

Determine $\liminf_{n \rightarrow \infty} A_n$ and $\limsup_{n \rightarrow \infty} A_n$.

Solution:

$$\liminf_{n \rightarrow \infty} A_n = \{0\} \subset [0, 1] = \limsup_{n \rightarrow \infty} A_n$$

Chapter 3. Real Number System and Calculus

§ 3.1 Real number system

§ 3.2 Sequences of real numbers

§ 3.3 Open and closed sets

§ 3.4 Real-valued functions

§ 3.5 Liminf and limsup of sets

§ 3.6 Some techniques in calculus

Examples

1. $\int_0^1 \tan^{-1}(x) dx$
2. $\int_0^x t^2 e^t dt$
3. $\int e^x \sin(x) dx$
4. $\int_0^1 (x^2 + 1) e^{-x} dx$
5. $\int_4^9 \frac{\ln y}{y} dy$
6. $\int_1^3 r^3 \ln r dr$
7. $\int_1^2 (\ln x)^2 dx$
8. $\int_{\sqrt{\pi/2}}^{\sqrt{\pi}} \theta^3 \cos(\theta^2) d\theta$
9. $\int_0^{\pi/2} \sin^{2n+1}(x) dx = \frac{2 \cdot 4 \cdot 6 \cdots 2n}{1 \cdot 3 \cdot 5 \cdot 7 \cdots (2n+1)}$
10. $\int_0^{\pi/2} \sin^{2n}(x) dx = \frac{1 \cdot 3 \cdot 5 \cdot 7 \cdots (2n-1)}{2 \cdot 4 \cdot 6 \cdots 2n} \frac{\pi}{2}$

Examples

1. $\int_0^1 \tan^{-1}(x) dx$
2. $\int_0^x t^2 e^t dt$
3. $\int e^x \sin(x) dx$
4. $\int_0^1 (x^2 + 1) e^{-x} dx$
5. $\int_4^9 \frac{\ln y}{y} dy$
6. $\int_1^3 r^3 \ln r dr$
7. $\int_1^2 (\ln x)^2 dx$
8. $\int_{\sqrt{\pi/2}}^{\sqrt{\pi}} \theta^3 \cos(\theta^2) d\theta$
9. $\int_0^{\pi/2} \sin^{2n+1}(x) dx = \frac{2 \cdot 4 \cdot 6 \cdots 2n}{1 \cdot 3 \cdot 5 \cdot 7 \cdots (2n+1)}$
10. $\int_0^{\pi/2} \sin^{2n}(x) dx = \frac{1 \cdot 3 \cdot 5 \cdot 7 \cdots (2n-1)}{2 \cdot 4 \cdot 6 \cdots 2n} \frac{\pi}{2}$

Examples

1. $\int_0^1 \tan^{-1}(x) dx$
2. $\int_0^x t^2 e^t dt$
3. $\int e^x \sin(x) dx$
4. $\int_0^1 (x^2 + 1) e^{-x} dx$
5. $\int_4^9 \frac{\ln y}{y} dy$
6. $\int_1^3 r^3 \ln r dr$
7. $\int_1^2 (\ln x)^2 dx$
8. $\int_{\sqrt{\pi/2}}^{\sqrt{\pi}} \theta^3 \cos(\theta^2) d\theta$
9. $\int_0^{\pi/2} \sin^{2n+1}(x) dx = \frac{2 \cdot 4 \cdot 6 \cdots 2n}{1 \cdot 3 \cdot 5 \cdot 7 \cdots (2n+1)}$
10. $\int_0^{\pi/2} \sin^{2n}(x) dx = \frac{1 \cdot 3 \cdot 5 \cdot 7 \cdots (2n-1)}{2 \cdot 4 \cdot 6 \cdots 2n} \frac{\pi}{2}$

Examples

1. $\int_0^1 \tan^{-1}(x) dx$
2. $\int_0^x t^2 e^t dt$
3. $\int e^x \sin(x) dx$
4. $\int_0^1 (x^2 + 1) e^{-x} dx$
5. $\int_4^9 \frac{\ln y}{y} dy$
6. $\int_1^3 r^3 \ln r dr$
7. $\int_1^2 (\ln x)^2 dx$
8. $\int_{\sqrt{\pi/2}}^{\sqrt{\pi}} \theta^3 \cos(\theta^2) d\theta$
9. $\int_0^{\pi/2} \sin^{2n+1}(x) dx = \frac{2 \cdot 4 \cdot 6 \cdots 2n}{1 \cdot 3 \cdot 5 \cdot 7 \cdots (2n+1)}$
10. $\int_0^{\pi/2} \sin^{2n}(x) dx = \frac{1 \cdot 3 \cdot 5 \cdot 7 \cdots (2n-1)}{2 \cdot 4 \cdot 6 \cdots 2n} \frac{\pi}{2}$

Examples

1. $\int_0^1 \tan^{-1}(x) dx$
2. $\int_0^x t^2 e^t dt$
3. $\int e^x \sin(x) dx$
4. $\int_0^1 (x^2 + 1) e^{-x} dx$
5. $\int_4^9 \frac{\ln y}{y} dy$
6. $\int_1^3 r^3 \ln r dr$
7. $\int_1^2 (\ln x)^2 dx$
8. $\int_{\sqrt{\pi/2}}^{\sqrt{\pi}} \theta^3 \cos(\theta^2) d\theta$
9. $\int_0^{\pi/2} \sin^{2n+1}(x) dx = \frac{2 \cdot 4 \cdot 6 \cdots 2n}{1 \cdot 3 \cdot 5 \cdot 7 \cdots (2n+1)}$
10. $\int_0^{\pi/2} \sin^{2n}(x) dx = \frac{1 \cdot 3 \cdot 5 \cdot 7 \cdots (2n-1)}{2 \cdot 4 \cdot 6 \cdots 2n} \frac{\pi}{2}$

Examples

1. $\int_0^1 \tan^{-1}(x) dx$
2. $\int_0^x t^2 e^t dt$
3. $\int e^x \sin(x) dx$
4. $\int_0^1 (x^2 + 1) e^{-x} dx$
5. $\int_4^9 \frac{\ln y}{y} dy$
6. $\int_1^3 r^3 \ln r dr$
7. $\int_1^2 (\ln x)^2 dx$
8. $\int_{\sqrt{\pi/2}}^{\sqrt{\pi}} \theta^3 \cos(\theta^2) d\theta$
9. $\int_0^{\pi/2} \sin^{2n+1}(x) dx = \frac{2 \cdot 4 \cdot 6 \cdots 2n}{1 \cdot 3 \cdot 5 \cdot 7 \cdots (2n+1)}$
10. $\int_0^{\pi/2} \sin^{2n}(x) dx = \frac{1 \cdot 3 \cdot 5 \cdot 7 \cdots (2n-1)}{2 \cdot 4 \cdot 6 \cdots 2n} \frac{\pi}{2}$

Examples

1. $\int_0^1 \tan^{-1}(x) dx$
2. $\int_0^x t^2 e^t dt$
3. $\int e^x \sin(x) dx$
4. $\int_0^1 (x^2 + 1) e^{-x} dx$
5. $\int_4^9 \frac{\ln y}{y} dy$
6. $\int_1^3 r^3 \ln r dr$
7. $\int_1^2 (\ln x)^2 dx$
8. $\int_{\sqrt{\pi/2}}^{\sqrt{\pi}} \theta^3 \cos(\theta^2) d\theta$
9. $\int_0^{\pi/2} \sin^{2n+1}(x) dx = \frac{2 \cdot 4 \cdot 6 \cdots 2n}{1 \cdot 3 \cdot 5 \cdot 7 \cdots (2n+1)}$
10. $\int_0^{\pi/2} \sin^{2n}(x) dx = \frac{1 \cdot 3 \cdot 5 \cdot 7 \cdots (2n-1)}{2 \cdot 4 \cdot 6 \cdots 2n} \frac{\pi}{2}$

Examples

1. $\int_0^1 \tan^{-1}(x) dx$
2. $\int_0^x t^2 e^t dt$
3. $\int e^x \sin(x) dx$
4. $\int_0^1 (x^2 + 1) e^{-x} dx$
5. $\int_4^9 \frac{\ln y}{y} dy$
6. $\int_1^3 r^3 \ln r dr$
7. $\int_1^2 (\ln x)^2 dx$
8. $\int_{\sqrt{\pi/2}}^{\sqrt{\pi}} \theta^3 \cos(\theta^2) d\theta$
9. $\int_0^{\pi/2} \sin^{2n+1}(x) dx = \frac{2 \cdot 4 \cdot 6 \cdots 2n}{1 \cdot 3 \cdot 5 \cdot 7 \cdots (2n+1)}$
10. $\int_0^{\pi/2} \sin^{2n}(x) dx = \frac{1 \cdot 3 \cdot 5 \cdot 7 \cdots (2n-1)}{2 \cdot 4 \cdot 6 \cdots 2n} \frac{\pi}{2}$

Examples

1. $\int_0^1 \tan^{-1}(x) dx$
2. $\int_0^x t^2 e^t dt$
3. $\int e^x \sin(x) dx$
4. $\int_0^1 (x^2 + 1) e^{-x} dx$
5. $\int_4^9 \frac{\ln y}{y} dy$
6. $\int_1^3 r^3 \ln r dr$
7. $\int_1^2 (\ln x)^2 dx$
8. $\int_{\sqrt{\pi/2}}^{\sqrt{\pi}} \theta^3 \cos(\theta^2) d\theta$
9. $\int_0^{\pi/2} \sin^{2n+1}(x) dx = \frac{2 \cdot 4 \cdot 6 \cdots 2n}{1 \cdot 3 \cdot 5 \cdot 7 \cdots (2n+1)}$
10. $\int_0^{\pi/2} \sin^{2n}(x) dx = \frac{1 \cdot 3 \cdot 5 \cdot 7 \cdots (2n-1)}{2 \cdot 4 \cdot 6 \cdots 2n} \frac{\pi}{2}$

Examples

1. $\int_0^1 \tan^{-1}(x) dx$
2. $\int_0^x t^2 e^t dt$
3. $\int e^x \sin(x) dx$
4. $\int_0^1 (x^2 + 1) e^{-x} dx$
5. $\int_4^9 \frac{\ln y}{y} dy$
6. $\int_1^3 r^3 \ln r dr$
7. $\int_1^2 (\ln x)^2 dx$
8. $\int_{\sqrt{\pi/2}}^{\sqrt{\pi}} \theta^3 \cos(\theta^2) d\theta$
9. $\int_0^{\pi/2} \sin^{2n+1}(x) dx = \frac{2 \cdot 4 \cdot 6 \cdots 2n}{1 \cdot 3 \cdot 5 \cdot 7 \cdots (2n+1)}$
10. $\int_0^{\pi/2} \sin^{2n}(x) dx = \frac{1 \cdot 3 \cdot 5 \cdot 7 \cdots (2n-1)}{2 \cdot 4 \cdot 6 \cdots 2n} \frac{\pi}{2}$

Examples

1. e^x

2. $\sin(x)$

3. e^{x^2}

Examples

1. e^x

2. $\sin(x)$

3. e^{x^2}

Examples

1. e^x

2. $\sin(x)$

3. e^{x^2}