

An Invitation to Probability

Le Chen

lzc0090@auburn.edu

(Creative Commons Zero 1.0)

Science Center Auditorium

5:30pm – 6:30pm, June 08, 2022

Github Hash: 618c9b1

2022-06-08 13:37:16 -0500

https://github.com/chenle02/2012_Science_Summer_Institute_Auburn_Probability_by_Le

Summer Science Institute 2022
Auburn University

What is chance

How to measure chance

Birthday problem

Rolling three dices


Poker

What is CHANCE?

Tyche – The Greek goddess of chance




Fortuna – The goddess of chance in Roman religion



... et bonnes
merveilles que l'on fait
pour se promener ou
autre chose humaine
nous nous l'en
de l'autre que l'on peut faire
ce n'est pas de la
foe et des temps et des lieux. Et aussi
que vous pourrez autre chose que
rien valoir au temps que l'on fait
pour les dommages autres fera que l'on
peut être malheureux. Et cette heure que
l'on fait en n'importe ce que

bonne a la femme pour seulement une
deux ou trois fois la paix ou autre chose
que l'on fait en la fortune d'autre chose que
en la fortune d'autre chose que
par force ou force de fortune et par mou-
vement de fortune que l'on fait ne
peut pas faire autre chose que l'on fait
que l'on fait a l'entretien et bonté
d'autre chose que l'on fait en la
fortune le même mal que l'on fait
que l'on fait et c'est une heure que l'on fait
que l'on fait que l'on fait que l'on fait
que l'on fait que l'on fait que l'on fait


I Ching in China (~ 6th century B.C.)



1

¹Image is from Bennett (1998), *Randomness*, Harvard University Press.

Democritus (460 – 370 BC) — Father of modern science



Atomic theory of the universe:

A physical chance affecting all the atoms
that made up the universe.

Games of chance — using knucklebones or dice



Known to Egyptians, Babylonians, Romans, ...

There was no qualitative theory of chance in these times.

How to measure chance?

How about measure length?



The determination of a “right and lawful rood” or rod in the early sixteenth century in Germany by measuring an essentially random selection of 16 men as they leave church ².

²Stephen Stigler (1996). Statistics and the Question of Standards, *Journal of Research of the National Institute of Standards and Technology*, vol. 101.

The same for chance

To measure probability,

1. we first find or make equally probable cases,
2. then we count.

The probability of an event A , denoted by $\mathbb{P}(A)$, is then

$$\mathbb{P}(A) = \frac{\text{no. of cases in which } A \text{ occurs}}{\text{total no. of cases}}.$$

The same for chance

To measure probability,

1. we first find or make equally probable cases,
2. then we count.

The probability of an event A , denoted by $\mathbb{P}(A)$, is then

$$\mathbb{P}(A) = \frac{\text{no. of cases in which } A \text{ occurs}}{\text{total no. of cases}}.$$

The same for chance

To measure probability,

1. we first find or make equally probable cases,
2. then we count.

The probability of an event A , denoted by $\mathbb{P}(A)$, is then

$$\mathbb{P}(A) = \frac{\text{no. of cases in which } A \text{ occurs}}{\text{total no. of cases}}.$$

The same for chance

To measure probability,

1. we first find or make equally probable cases,
2. then we count.

The probability of an event A , denoted by $\mathbb{P}(A)$, is then

$$\mathbb{P}(A) = \frac{\text{no. of cases in which } A \text{ occurs}}{\text{total no. of cases}}.$$

The same for chance

To measure probability,

1. we first find or make equally probable cases,
2. then we count.

The probability of an event A , denoted by $\mathbb{P}(A)$, is then

$$\mathbb{P}(A) = \frac{\text{no. of cases in which } A \text{ occurs}}{\text{total no. of cases}}.$$

Now we see that probability has to satisfy the following properties:

1. Probability should be never negative.
2. If A occurs in all cases, then $\mathbb{P}(A) = 1$.
3. If A and B never occur in the same case, then

$$\mathbb{P}(A \text{ or } B) = \mathbb{P}(A) + \mathbb{P}(B)$$

In particular, the probability of an event not occurring is equal to

$$\mathbb{P}(\text{not } A) = 1 - \mathbb{P}(A).$$

Now we see that probability has to satisfy the following properties:

1. Probability should be never negative.
2. If A occurs in all cases, then $\mathbb{P}(A) = 1$.
3. If A and B never occur in the same case, then

$$\mathbb{P}(A \text{ or } B) = \mathbb{P}(A) + \mathbb{P}(B).$$

In particular, the probability of an event not occurring is equal to

$$\mathbb{P}(\text{not } A) = 1 - \mathbb{P}(A).$$

Now we see that probability has to satisfy the following properties:

1. Probability should be never negative.
2. If A occurs in all cases, then $\mathbb{P}(A) = 1$.
3. If A and B never occur in the same case, then

$$\mathbb{P}(A \text{ or } B) = \mathbb{P}(A) + \mathbb{P}(B).$$

In particular, the probability of an event not occurring is equal to

$$\mathbb{P}(\text{not } A) = 1 - \mathbb{P}(A).$$

Now we see that probability has to satisfy the following properties:

1. Probability should be never negative.
2. If A occurs in all cases, then $\mathbb{P}(A) = 1$.
3. If A and B never occur in the same case, then

$$\mathbb{P}(A \text{ or } B) = \mathbb{P}(A) + \mathbb{P}(B).$$

In particular, the probability of an event not occurring is equal to

$$\mathbb{P}(\text{not } A) = 1 - \mathbb{P}(A).$$

Now we see that probability has to satisfy the following properties:

1. Probability should be never negative.
2. If A occurs in all cases, then $\mathbb{P}(A) = 1$.
3. If A and B never occur in the same case, then

$$\mathbb{P}(A \text{ or } B) = \mathbb{P}(A) + \mathbb{P}(B).$$

In particular, the probability of an event not occurring is equal to

$$\mathbb{P}(\text{not } A) = 1 - \mathbb{P}(A).$$

Now we see that probability has to satisfy the following properties:

1. Probability should be never negative.
2. If A occurs in all cases, then $\mathbb{P}(A) = 1$.
3. If A and B never occur in the same case, then

$$\mathbb{P}(A \text{ or } B) = \mathbb{P}(A) + \mathbb{P}(B).$$

In particular, the probability of an event not occurring is equal to


$$\mathbb{P}(\text{not } A) = 1 - \mathbb{P}(A).$$

How to generate the equi-probable cases?

Prim sticks (variations of dice)³ and deck of poker...

³Image is from Bennett (1998), *Randomness*, Harvard University Press.

How to generate the equi-probable cases?



Prim sticks (variations of dice) ³ and deck of poker...

³Image is from Bennett (1998), *Randomness*, Harvard University Press.



Girolamo Cardano (1501 – 1576), an Italian polymath, whose interests and proficiencies ranged through those of mathematician, physician, biologist, physicist, chemist, astrologer, astronomer, philosopher, writer, and gambler. He was one of the most influential mathematicians of the Renaissance, and was one of the key figures in the foundation of probability and the earliest introducer of the binomial coefficients and the binomial theorem in the Western world.

Birthday Problem

Question: How likely do two students have the same birth day if there are

- ▶ 2
- ▶ 5
- ▶ 15
- ▶ 23
- ▶ 46
- ▶ 64
- ▶ 366

students in the class?

Assuming that

1. each year has 365 days (i.e., neglecting leap years).
2. birthdays are equi-probable.
3. birthdays are independent (no twins in the class).

Birthday Problem

Question: How likely do two students have the same birth day if there are

- ▶ 2
- ▶ 5
- ▶ 15
- ▶ 23
- ▶ 46
- ▶ 64
- ▶ 366

students in the class?

Assuming that

1. each year has 365 days (i.e., neglecting leap years).
2. birthdays are equi-probable.
3. birthdays are independent (no twins in the class).

Birthday Problem

Question: How likely do two students have the same birth day if there are

- ▶ 2
- ▶ 5
- ▶ 15
- ▶ 23
- ▶ 46
- ▶ 64
- ▶ 366

students in the class?

Assuming that

1. each year has 365 days (i.e., neglecting leap years).
2. birthdays are equi-probable.
3. birthdays are independent (no twins in the class).

Birthday Problem

Question: How likely do two students have the same birth day if there are

- ▶ 2
- ▶ 5
- ▶ 15
- ▶ 23
- ▶ 46
- ▶ 64
- ▶ 366

students in the class?

Assuming that

1. each year has 365 days (i.e., neglecting leap years).
2. birthdays are equi-probable.
3. birthdays are independent (no twins in the class).

Birthday Problem

Question: How likely do two students have the same birth day if there are

- ▶ 2
- ▶ 5
- ▶ 15
- ▶ 23
- ▶ 46
- ▶ 64
- ▶ 366

students in the class?

Assuming that

1. each year has 365 days (i.e., neglecting leap years).
2. birthdays are equi-probable.
3. birthdays are independent (no twins in the class).

Birthday Problem

Question: How likely do two students have the same birth day if there are

- ▶ 2
- ▶ 5
- ▶ 15
- ▶ 23
- ▶ 46
- ▶ 64
- ▶ 366

students in the class?

Assuming that

1. each year has 365 days (i.e., neglecting leap years).
2. birthdays are equi-probable.
3. birthdays are independent (no twins in the class).

Birthday Problem

Question: How likely do two students have the same birth day if there are

- ▶ 2
- ▶ 5
- ▶ 15
- ▶ 23
- ▶ 46
- ▶ 64
- ▶ 366

students in the class?

Assuming that

1. each year has 365 days (i.e., neglecting leap years).
2. birthdays are equi-probable.
3. birthdays are independent (no twins in the class).

Birthday Problem

Question: How likely do two students have the same birth day if there are

- ▶ 2
- ▶ 5
- ▶ 15
- ▶ 23
- ▶ 46
- ▶ 64
- ▶ 366

students in the class?

Assuming that

1. each year has 365 days (i.e., neglecting leap years).
2. birthdays are equi-probable.
3. birthdays are independent (no twins in the class).

Birthday Problem

Question: How likely do two students have the same birth day if there are

- ▶ 2
- ▶ 5
- ▶ 15
- ▶ 23
- ▶ 46
- ▶ 64
- ▶ 366

students in the class?

Assuming that

1. each year has 365 days (i.e., neglecting leap years).
2. birthdays are equi-probable.
3. birthdays are independent (no twins in the class).

Birthday Problem

Question: How likely do two students have the same birth day if there are

- ▶ 2
- ▶ 5
- ▶ 15
- ▶ 23
- ▶ 46
- ▶ 64
- ▶ 366

students in the class?

Assuming that

1. each year has 365 days (i.e., neglecting leap years),
2. birthdays are equi-probable,
3. birthdays are independent (no twins in the class).

Birthday Problem

Question: How likely do two students have the same birth day if there are

- ▶ 2
- ▶ 5
- ▶ 15
- ▶ 23
- ▶ 46
- ▶ 64
- ▶ 366

students in the class?

Assuming that

1. each year has 365 days (i.e., neglecting leap years),
2. birthdays are equi-probable,
3. birthdays are independent (no twins in the class).

Birthday Problem

Question: How likely do two students have the same birth day if there are

- ▶ 2
- ▶ 5
- ▶ 15
- ▶ 23
- ▶ 46
- ▶ 64
- ▶ 366

students in the class?

Assuming that

1. each year has 365 days (i.e., neglecting leap years),
2. birthdays are equi-probable,
3. birthdays are independent (no twins in the class).

- ▶ Suppose $n = 5$.
- ▶ Let A be the event that there is a shared birthday among these n students.
- ▶ It is not easy to compute $\mathbb{P}(A)$ directly.
- ▶ However, one can compute $\mathbb{P}(\text{not } A)$ by counting:

the probability that no two students have the same birthday, or

all students have different birthday,

is equal to

$$\mathbb{P}(\text{not } A) = \frac{365}{365} \frac{364}{365} \frac{363}{365} \frac{362}{365} \frac{361}{365}$$

- ▶ Hence,

$$\mathbb{P}(A) = 1 - \mathbb{P}(\text{not } A) = 1 - \frac{365}{365} \frac{364}{365} \frac{363}{365} \frac{362}{365} \frac{361}{365} \approx 0.027.$$

- ▶ Suppose $n = 5$.
- ▶ Let A be the event that there is a shared birthday among these n students.
- ▶ It is not easy to compute $\mathbb{P}(A)$ directly.
- ▶ However, one can compute $\mathbb{P}(\text{not } A)$ by counting:

the probability that no two students have the same birthday, or

all students have different birthday,

is equal to

$$\mathbb{P}(\text{not } A) = \frac{365}{365} \frac{364}{365} \frac{363}{365} \frac{362}{365} \frac{361}{365}$$

- ▶ Hence,

$$\mathbb{P}(A) = 1 - \mathbb{P}(\text{not } A) = 1 - \frac{365}{365} \frac{364}{365} \frac{363}{365} \frac{362}{365} \frac{361}{365} \approx 0.027.$$

- ▶ Suppose $n = 5$.
- ▶ Let A be the event that there is a shared birthday among these n students.
- ▶ It is not easy to compute $\mathbb{P}(A)$ directly.
- ▶ However, one can compute $\mathbb{P}(\text{not } A)$ by counting:

the probability that no two students have the same birthday, or

all students have different birthday,

is equal to

$$\mathbb{P}(\text{not } A) = \frac{365}{365} \frac{364}{365} \frac{363}{365} \frac{362}{365} \frac{361}{365}$$

- ▶ Hence,

$$\mathbb{P}(A) = 1 - \mathbb{P}(\text{not } A) = 1 - \frac{365}{365} \frac{364}{365} \frac{363}{365} \frac{362}{365} \frac{361}{365} \approx 0.027.$$

- ▶ Suppose $n = 5$.
- ▶ Let A be the event that there is a shared birthday among these n students.
- ▶ It is not easy to compute $\mathbb{P}(A)$ directly.
- ▶ However, one can compute $\mathbb{P}(\text{not } A)$ by counting:

the probability that no two students have the same birthday, or

all students have different birthday,

is equal to

$$\mathbb{P}(\text{not } A) = \frac{365}{365} \frac{364}{365} \frac{363}{365} \frac{362}{365} \frac{361}{365}$$

- ▶ Hence,

$$\mathbb{P}(A) = 1 - \mathbb{P}(\text{not } A) = 1 - \frac{365}{365} \frac{364}{365} \frac{363}{365} \frac{362}{365} \frac{361}{365} \approx 0.027.$$

- ▶ Suppose $n = 5$.
- ▶ Let A be the event that there is a shared birthday among these n students.
- ▶ It is not easy to compute $\mathbb{P}(A)$ directly.
- ▶ However, one can compute $\mathbb{P}(\text{not } A)$ by counting:

the probability that no two students have the same birthday, or

all students have different birthday,

is equal to

$$\mathbb{P}(\text{not } A) = \frac{365}{365} \frac{364}{365} \frac{363}{365} \frac{362}{365} \frac{361}{365}$$

- ▶ Hence,

$$\mathbb{P}(A) = 1 - \mathbb{P}(\text{not } A) = 1 - \frac{365}{365} \frac{364}{365} \frac{363}{365} \frac{362}{365} \frac{361}{365} \approx 0.027.$$

- ▶ Suppose $n = 5$.
- ▶ Let A be the event that there is a shared birthday among these n students.
- ▶ It is not easy to compute $\mathbb{P}(A)$ directly.
- ▶ However, one can compute $\mathbb{P}(\text{not } A)$ by counting:

the probability that no two students have the same birthday, or

all students have different birthday,

is equal to

$$\mathbb{P}(\text{not } A) = \frac{365}{365} \frac{364}{365} \frac{363}{365} \frac{362}{365} \frac{361}{365}$$

- ▶ Hence,

$$\mathbb{P}(A) = 1 - \mathbb{P}(\text{not } A) = 1 - \frac{365}{365} \frac{364}{365} \frac{363}{365} \frac{362}{365} \frac{361}{365} \approx 0.027.$$

$$\mathbb{P}(A) = 1 - \frac{365 \times 364 \times \cdots \times (365 - n + 1)}{365^n}$$

n	2	5	15	23	46	64	366
$\mathbb{P}(A)$							1.0

$$\mathbb{P}(A) = 1 - \frac{365 \times 364 \times \cdots \times (365 - n + 1)}{365^n}$$

n	2	5	15	23	46	64	366
$\mathbb{P}(A)$	0.0027						1.0

$$\mathbb{P}(A) = 1 - \frac{365 \times 364 \times \cdots \times (365 - n + 1)}{365^n}$$

n	2	5	15	23	46	64	366
$\mathbb{P}(A)$	0.0027	0.0271					1.0

$$\mathbb{P}(A) = 1 - \frac{365 \times 364 \times \cdots \times (365 - n + 1)}{365^n}$$

n	2	5	15	23	46	64	366
$\mathbb{P}(A)$	0.0027	0.0271	0.2529				1.0

$$\mathbb{P}(A) = 1 - \frac{365 \times 364 \times \cdots \times (365 - n + 1)}{365^n}$$

n	2	5	15	23	46	64	366
$\mathbb{P}(A)$	0.0027	0.0271	0.2529	0.5073			1.0

$$\mathbb{P}(A) = 1 - \frac{365 \times 364 \times \cdots \times (365 - n + 1)}{365^n}$$


n	2	5	15	23	46	64	366
$\mathbb{P}(A)$	0.0027	0.0271	0.2529	0.5073	0.9483		1.0

$$\mathbb{P}(A) = 1 - \frac{365 \times 364 \times \cdots \times (365 - n + 1)}{365^n}$$

n	2	5	15	23	46	64	366
$\mathbb{P}(A)$	0.0027	0.0271	0.2529	0.5073	0.9483	0.9972	1.0

$$\mathbb{P}(A) = 1 - \frac{365 \times 364 \times \cdots \times (365 - n + 1)}{365^n}$$

n	2	5	15	23	46	64	366
$\mathbb{P}(A)$	0.0027	0.0271	0.2529	0.5073	0.9483	0.9972	1.0



When time permits, we will try some simulations !

Source codes are here:

https://github.com/chenle02/2012_Science_Summer_Institute_Auburn_Probability_by_Le

More surprises of this type

1. Let C be the number of categories (i.e., $C = 365$) and n be the number of students. Here is one useful approximation:

when $n = 1.2 \sqrt{C}$, the chance of shared something is close to $1/2$.

E.g., $1.2 \sqrt{365} = 22.9$ (Birthday problem), $1.2 \sqrt{60} = 9.3$ (*Watch problem*).

2. How large should n be to have approximately even probability of a triple birthday match?

Answer: 81.

3. Would that be a surprise if you find out with someone, not only you share the same birthday, but also the same father's birthday and the same grandfather's birthday?

More surprises of this type

1. Let C be the number of categories (i.e., $C = 365$) and n be the number of students. Here is one useful approximation:

when $n = 1.2 \sqrt{C}$, the chance of shared something is close to $1/2$.

E.g., $1.2 \sqrt{365} = 22.9$ (Birthday problem), $1.2 \sqrt{60} = 9.3$ (*Watch problem*).

2. How large should n be to have approximately even probability of a triple birthday match?

Answer: 81.

3. Would that be a surprise if you find out with someone, not only you share the same birthday, but also the same father's birthday and the same grandfather's birthday?

More surprises of this type

1. Let C be the number of categories (i.e., $C = 365$) and n be the number of students. Here is one useful approximation:

when $n = 1.2 \sqrt{C}$, the chance of shared something is close to $1/2$.

E.g., $1.2 \sqrt{365} = 22.9$ (Birthday problem), $1.2 \sqrt{60} = 9.3$ (*Watch problem*).

2. How large should n be to have approximately even probability of a triple birthday match?

Answer: 81.

3. Would that be a surprise if you find out with someone, not only you share the same birthday, but also the same father's birthday and the same grandfather's birthday?

More surprises of this type

1. Let C be the number of categories (i.e., $C = 365$) and n be the number of students. Here is one useful approximation:

when $n = 1.2 \sqrt{C}$, the chance of shared something is close to $1/2$.

E.g., $1.2 \sqrt{365} = 22.9$ (Birthday problem), $1.2 \sqrt{60} = 9.3$ (*Watch problem*).

2. How large should n be to have approximately even probability of a triple birthday match?

Answer: 81.

3. Would that be a surprise if you find out with someone, not only you share the same birthday, but also the same father's birthday and the same grandfather's birthday?

More surprises of this type

1. Let C be the number of categories (i.e., $C = 365$) and n be the number of students. Here is one useful approximation:

when $n = 1.2 \sqrt{C}$, the chance of shared something is close to $1/2$.

E.g., $1.2 \sqrt{365} = 22.9$ (Birthday problem), $1.2 \sqrt{60} = 9.3$ (*Watch problem*).

2. How large should n be to have approximately even probability of a triple birthday match?

Answer: 81.

3. Would that be a surprise if you find out with someone, not only you share the same birthday, but also the same father's birthday and the same grandfather's birthday?

More surprises of this type

1. Let C be the number of categories (i.e., $C = 365$) and n be the number of students. Here is one useful approximation:

when $n = 1.2 \sqrt{C}$, the chance of shared something is close to $1/2$.

E.g., $1.2 \sqrt{365} = 22.9$ (Birthday problem), $1.2 \sqrt{60} = 9.3$ (*Watch problem*).

2. How large should n be to have approximately even probability of a triple birthday match?

Answer: 81.

3. Would that be a surprise if you find out with someone, not only you share the same birthday, but also the same father's birthday and the same grandfather's birthday?

Number of generations	1	2	3	4	5
n to have about even odds	23				

Number of generations	1	2	3	4	5
n to have about even odds	23	438			

Number of generations	1	2	3	4	5
n to have about even odds	23	438	8,368		

Number of generations	1	2	3	4	5
n to have about even odds	23	438	8,368	159,870	

Number of generations	1	2	3	4	5
n to have about even odds	23	438	8,368	159,870	3,054,312

A question asked by *Grand Duke of Tuscany* to *Galileo* in early seventh century

Three dice are thrown, such as

9 : 


10 : 

11 : 


12 : 

Counting combinations of numbers, 10 and 11 can be made in 6 ways, as can 9 and 12. Yet it is known that long observation has made dice-players consider 10 and 11 to be more advantageous than 9 and 12. How can this be?


9




10



11



12



$$3 = \square + \square + \square$$

$$4 = \square + \square + \square$$

$$5 = \square + \square + \square = \square + \square + \square$$

$$6 = \square + \bullet + \blacksquare = \square + \circlearrowleft + \blacksquare = \blacksquare + \circlearrowright + \blacksquare$$

$$7 = \square + \square + \blacksquare = \square + \blacksquare + \blacksquare = \blacksquare + \blacksquare + \blacksquare = \blacksquare + \square + \square$$

$$8 = \square + \square + \blacksquare + \blacksquare = \square + \square + \blacksquare + \blacksquare = \square + \square + \blacksquare + \blacksquare = \square + \square + \blacksquare + \blacksquare$$

$$9 = \square + \square + \square = \square + \square + \square$$

$$10 = \square + \square + \blacksquare = \square + \blacksquare + \blacksquare = \square + \square + \blacksquare = \square + \square + \blacksquare = \square + \blacksquare + \blacksquare = \square + \square + \blacksquare = \square + \square + \blacksquare$$

$$11 = \square + \square + \square = \square + \square + \square$$

$$12 = \square + \square + \square = \square + \square + \square$$

$$13 = \square + \square + \square = \square + \square + \square$$

$$14 = \square + \square + \square = \square + \square + \square = \square + \square + \square = \square + \square + \square$$

$$15 = \square + \square + \square - \square + \square + \square - \square + \square + \square$$

16 + + +

17 8+8+8

10 11 12

$$3 = 1 + 1 + 1$$

$$4 = 1 + 1 + 2$$

$$5 = 1 + 1 + 3 = 1 + 2 + 2$$

$$6 = 1 + 1 + 4 = 1 + 2 + 3 = 2 + 2 + 2$$

$$7 = 1 + 1 + 5 = 1 + 2 + 4 = 2 + 2 + 3 = 3 + 3 + 1$$

$$8 = 1 + 1 + 6 = 1 + 2 + 5 = 1 + 3 + 4 = 2 + 2 + 4 = 2 + 3 + 3$$

$$9 = 1 + 2 + 6 = 1 + 3 + 5 = 1 + 4 + 4 = 2 + 2 + 5 = 2 + 3 + 4 = 3 + 3 + 3$$

$$10 = 1 + 3 + 6 = 1 + 4 + 5 = 2 + 2 + 6 = 2 + 3 + 5 = 2 + 4 + 4 = 3 + 3 + 4$$

$$11 = 1 + 4 + 6 = 1 + 5 + 5 = 2 + 3 + 6 = 2 + 4 + 5 = 3 + 3 + 5 = 3 + 4 + 4$$

$$12 = 1 + 5 + 6 = 2 + 4 + 6 = 2 + 5 + 5 = 3 + 3 + 6 = 3 + 4 + 5 = 4 + 4 + 4$$

$$13 = 1 + 6 + 6 = 2 + 5 + 6 = 3 + 4 + 6 = 3 + 5 + 5 = 4 + 4 + 5$$

$$14 = 2 + 6 + 6 = 3 + 5 + 6 = 4 + 4 + 6 = 4 + 5 + 5$$

$$15 = 3 + 6 + 6 = 4 + 5 + 6 = 5 + 5 + 5$$

$$16 = 4 + 6 + 6 = 5 + 5 + 6$$

$$17 = 5 + 6 + 6$$

$$18 = 6 + 6 + 6$$

k	Probability of a sum of k	\approx
3	1/216	0.5%
4	3/216	1.4%
5	6/216	2.8%
6	10/216	4.6%
7	15/216	7.0%
8	21/216	9.7%
9	25/216	11.6%
10	27/216	12.5%
11	27/216	12.5%
12	25/216	11.6%
13	21/216	9.7%
14	15/216	7.0%
15	10/216	4.6%
16	6/216	2.8%
17	3/216	1.4%
18	1/216	0.5%

Game of rolling three dices dated back to Roman Empire.

3	18	Punctatura	1	Cadentia	1
4	17	Punctatura	1	Cadentia	3
5	16	Punctatura	2	Cadentia	6
6	15	Punctatura	3	Cadentia	10
7	14	Punctatura	4	Cadentia	15
8	13	Punctatura	5	Cadentia	21
9	12	Punctatura	6	Cadentia	25
10	11	Punctatura	6	Cadentia	27

Richard de Fournival discovered in 13th century the summary of
216 possible sequences ⁴
in his poem, *De Vetula*, written between 1220 to 1250.

⁴Image is from Bennett (1998), *Randomness*, Harvard University Press.



POKER

HAND RANKINGS



#1 ROYAL FLUSH



#6 STRAIGHT



#2 STRAIGHT FLUSH



#7 THREE OF A KIND



#3 FOUR OF A KIND



#8 TWO PAIR



#4 FULL HOUSE



#9 ONE PAIR



#5 FLUSH



#10 HIGH CARD

Table 10-1. Ways to Deal Five-Card Poker Hands

Hand	Number of ways	Probability that opponent has a better hand ^a when the number of opposing hands is:					
		1	2	3	4	5	6
Straight flush	40	0	0	0	0	0	0
Four of a kind	624	.00002	.00003	.00005	.00006	.00008	.00009
Full house	3,744	.00026	.00051	.00078	.00102	.00128	.00153
Flush	5,108	.00170	.00339	.00509	.00677	.00847	.01015
Straight	10,200	.00366	.00731	.01094	.01457	.01817	.02177
Three of a kind	54,912	.00759	.01511	.02259	.03000	.03736	.04466
Two pairs:							
Aces high	19,008	.02871	.05660	.08369	.11001	.13556	.16038
Kings	17,424	.03603	.07076	.10424	.13651	.16762	.19761
Queens	15,840	.04273	.08364	.12280	.16028	.19617	.23052
Jacks	14,256	.04883	.09527	.13945	.18146	.22143	.25945
Tens	12,672	.05431	.10568	.15425	.20019	.24362	.28470
Nines	11,088	.05919	.11487	.16726	.21655	.26292	.30655
Eights	9,504	.06345	.12288	.17854	.23067	.27948	.32520
Sevens	7,920	.06711	.12972	.18812	.24261	.29344	.34086
Sixes	6,336	.07016	.13540	.19606	.25246	.30491	.35367
Fives	4,752	.07260	.13992	.20236	.26027	.31397	.36378
Fours	3,168	.07443	.14331	.20707	.26608	.32071	.37126
Threes	1,584	.07564	.14556	.21019	.26993	.32515	.37620

Table 10-1. continued

Hand	Number of ways	Probability that opponent has a better hand ^a when the number of opposing hands is:					
		1	2	3	4	5	6
One pair:							
Aces	84,480	.07625	.14669	.21176	.27187	.32739	.37868
Kings	"	.10876	.20569	.29208	.36907	.43769	.49885
Queens	"	.14126	.26257	.36674	.45620	.53302	.59899
Jacks	"	.17377	.31734	.43597	.53398	.61496	.68187
Tens	"	.20627	.37000	.49995	.60310	.68497	.74995
Nines	"	.23878	.42054	.55891	.66423	.74441	.80544
Eights	"	.27129	.46898	.61303	.71801	.79451	.85026
Sevens	"	.30379	.51529	.66254	.76506	.83643	.88612
Sixes	"	.33630	.55950	.70764	.80596	.87121	.91452
Fives	"	.36880	.60159	.74852	.84127	.89981	.93676
Fours	"	.40131	.64157	.78541	.87153	.92309	.95395
Threes	"	.43381	.67943	.81850	.89724	.94182	.96706
Deuces	"	.46632	.71518	.84800	.91888	.95671	.97690
No pair	1,302,540	.49882	.74882	.87411	.93691	.96838	.98415
All hands	2,598,960						

a. Assuming that you have the best hand of its type—for example, ace-high if you have no pair.

Thank you for your
listening and participating !

References:

- ▶ Persi Diaconis and Brian Skyrms (2017). *The great ideas about chance*. Princeton University Press.
- ▶ Deborah J. Bennett (1998). *Randomness*. Harvard University Press.
- ▶ John D. McGervey (1986). *Probabilities in everyday life*. Nelson Hall Publishers.