

Analysis of Tetris Ballistic Deposition and the Robustness of the KPZ Universality Class

Le Chen
Auburn University

Acknwolegement

NSF 2246850, NSF 2443823, & Simons Foundation Travel Grant (2022-2027)

Talk available at: github.com/chenle02

Emerging Synergies between Stochastic Analysis and Statistical Mechanics
Banff, Alberta, Canada
October 28, 2025

Integration of Research, Education, and Outreach

Outreach

- ▶ Auburn Summer Science Institute (AU-SSI): **2024, 2025**
Selected talented high school students
- ▶ Destination STEM: **2023, 2024**
Junior middle school students

Education

- ▶ Graduate Student Seminars (Mathematics), Auburn: **2022–2025**
- ▶ Math 7820/7830: Applied Stochastic Processes Course project, **2023/24**

Research

- ▶ Simulation and modeling packages (open source)
- ▶ Forming conjectures and validating results

Most materials are available at

github.com/chenle02

Math 7820/30: Applied Stochastic Processes (2023/24):



Mauricio Montes and Ian Ruau

Simulation package:

https://github.com/chenle02/Simulations_on_Some_Surface_Growth_Models

```
pip install tetris-ballistic
```



Image is generated by OpenAI's *DALL-E*

Random deposition



Ballistic deposition

Image is generated by OpenAI's *DALL-E*

Plan

Introduction to growth model and SPDE

Family-Vicsek scaling and experiments

More examples

Tetromino Pieces

Plan

Introduction to growth model and SPDE


Family-Vicsek scaling and experiments

More examples


Tetromino Pieces

How do surfaces grow?


Three models




Random deposition




Random deposition




Random deposition




Random deposition




Random deposition with surface relaxation




Random deposition with surface relaxation




Random deposition with surface relaxation




Random deposition with surface relaxation




Random deposition with surface relaxation




Ballistic deposition




Ballistic deposition




Ballistic deposition




Ballistic deposition



Ballistic deposition



Average height and fluctuation




Average height and fluctuation

$$\bar{h}(t) = \frac{1}{L} \sum_{x=1}^L h(t, x)$$

$$h(8, x) = \begin{array}{ccccc} 0 & 3 & 3 & 3 & 0 \end{array}$$

$$\text{Fluctuation } W(L, t) = \sqrt{\frac{1}{L} \sum_{x=1}^L [h(t, x) - \bar{h}(t)]^2}$$




Average height and fluctuation

$$\bar{h}(t) = \frac{1}{L} \sum_{x=1}^L h(t, x)$$

$$h(8, x) = \begin{array}{ccccccc} & 0 & 3 & 3 & 3 & 0 & 0 \\ \end{array}$$

$$\text{Fluctuation } W(L, t) = \sqrt{\frac{1}{L} \sum_{x=1}^L [h(t, x) - \bar{h}(t)]^2}$$



Random Deposition (independent columns, nonsticky)

Model. L independent columns. At each integer time $t = 1, 2, \dots$, drop *one* particle on a uniformly random column. Heights $h(t, x)$, mean $\bar{h}(t) = \frac{1}{L} \sum_{x=1}^L h(t, x) = \frac{t}{L}$, width

$$W^2(L, t) = \frac{1}{L} \sum_{x=1}^L (h(t, x) - \bar{h}(t))^2.$$

Single-column law: After t drops total,

$$h(t, x) \sim \text{Binomial}\left(t, \frac{1}{L}\right), \quad \mathbb{E}[h(t, x)] = \frac{t}{L}, \quad \text{Var}(h(t, x)) = t \frac{1}{L} \left(1 - \frac{1}{L}\right).$$

Fluctuation: By i.i.d. columns,

$$\mathbb{E}[W^2(L, t)] = \frac{1}{L} \sum_{x=1}^L \mathbb{E}[h(t, x)^2] - \mathbb{E}[\bar{h}^2(t)] = \mathbb{E}[h(t, 1)^2] - \left(\frac{t}{L}\right)^2 = \left(1 - \frac{1}{L}\right) \text{Var}(h(t, 1)).$$

Hence

$$\boxed{\mathbb{E}[W^2(L, t)] = \left(1 - \frac{1}{L}\right) t \frac{1}{L} \left(1 - \frac{1}{L}\right) = \frac{t}{L} \left(1 - \frac{1}{L}\right)^2}$$

and

$$\boxed{W(L, t) \simeq \left(1 - \frac{1}{L}\right) \left(\frac{t}{L}\right)^{1/2}}$$

Scaling. Growth exponent $\beta = \frac{1}{2}$.

Random Deposition (independent columns, nonsticky)

Model. L independent columns. At each integer time $t = 1, 2, \dots$, drop *one* particle on a uniformly random column. Heights $h(t, x)$, mean $\bar{h}(t) = \frac{1}{L} \sum_{x=1}^L h(t, x) = \frac{t}{L}$, width

$$W^2(L, t) = \frac{1}{L} \sum_{x=1}^L (h(t, x) - \bar{h}(t))^2.$$

Single-column law: After t drops total,

$$h(t, x) \sim \text{Binomial}\left(t, \frac{1}{L}\right), \quad \mathbb{E}[h(t, x)] = \frac{t}{L}, \quad \text{Var}(h(t, x)) = t \frac{1}{L} \left(1 - \frac{1}{L}\right).$$

Fluctuation: By i.i.d. columns,

$$\mathbb{E}[W^2(L, t)] = \frac{1}{L} \sum_{x=1}^L \mathbb{E}[h(t, x)^2] - \mathbb{E}[\bar{h}^2(t)] = \mathbb{E}[h(t, 1)^2] - \left(\frac{t}{L}\right)^2 = \left(1 - \frac{1}{L}\right) \text{Var}(h(t, 1)).$$

Hence

$$\boxed{\mathbb{E}[W^2(L, t)] = \left(1 - \frac{1}{L}\right) t \frac{1}{L} \left(1 - \frac{1}{L}\right) = \frac{t}{L} \left(1 - \frac{1}{L}\right)^2}$$

and

$$\boxed{W(L, t) \simeq \left(1 - \frac{1}{L}\right) \left(\frac{t}{L}\right)^{1/2}}$$

Scaling. Growth exponent $\beta = \frac{1}{2}$.

Random Deposition (independent columns, nonsticky)

Model. L independent columns. At each integer time $t = 1, 2, \dots$, drop *one* particle on a uniformly random column. Heights $h(t, x)$, mean $\bar{h}(t) = \frac{1}{L} \sum_{x=1}^L h(t, x) = \frac{t}{L}$, width

$$W^2(L, t) = \frac{1}{L} \sum_{x=1}^L (h(t, x) - \bar{h}(t))^2.$$

Single-column law: After t drops total,

$$h(t, x) \sim \text{Binomial}\left(t, \frac{1}{L}\right), \quad \mathbb{E}[h(t, x)] = \frac{t}{L}, \quad \text{Var}(h(t, x)) = t \frac{1}{L} \left(1 - \frac{1}{L}\right).$$

Fluctuation: By i.i.d. columns,

$$\mathbb{E}[W^2(L, t)] = \frac{1}{L} \sum_{x=1}^L \mathbb{E}[h(t, x)^2] - \mathbb{E}[\bar{h}^2(t)] = \mathbb{E}[h(t, 1)^2] - \left(\frac{t}{L}\right)^2 = \left(1 - \frac{1}{L}\right) \text{Var}(h(t, 1)).$$

Hence


$$\boxed{\mathbb{E}[W^2(L, t)] = \left(1 - \frac{1}{L}\right) t \frac{1}{L} \left(1 - \frac{1}{L}\right) = \frac{t}{L} \left(1 - \frac{1}{L}\right)^2}$$

and

$$\boxed{W(L, t) \simeq \left(1 - \frac{1}{L}\right) \left(\frac{t}{L}\right)^{1/2}}$$

Scaling. Growth exponent $\beta = \frac{1}{2}$.

Questions




$$W(t, x) \sim t^{1/2}$$


$$W(t, x) \sim t^{??}$$

$$W(t, x) \sim t^{??}$$

Log-Log plots for nonsticky piece 19



Log-Log plots for sticky piece 19



Plan

Introduction to growth model and SPDE

Family-Vicsek scaling and experiments

More examples

Tetromino Pieces

Family–Vicsek scaling theory

Recall height function $h(x, t)$, $x = 1, \dots, L$. Mean $\bar{h}(t) = \frac{1}{L} \sum_{x=1}^L h(x, t)$.

$$w(L, t) := \sqrt{\frac{1}{L} \sum_{x=1}^L (h(x, t) - \bar{h}(t))^2} \quad (\text{Fluctuation}).$$

Empirical scaling (log–log)

Early time $w(L, t) \sim t^\beta$

β : growth exponent

Late time $w(L, t) \rightarrow w_{\text{sat}}(L) \sim L^\alpha$

α : roughness exponent


Crossover time $t_x(L) \sim L^z$

z : dynamic exponent

Family, F., Vicsek, T., *Journal of Physics A: Mathematical and General*, 1985

Family-Vicsek scaling theory

α : Roughness exp.; β : Growth exp.; z : Dynamic exp.



(Image from Barabási-Stanley's book 95)

Family–Vicsek scaling theory

α : Roughness exp.; β : Growth exp.; z : Dynamic exp.

$$w(L, t) \sim L^\alpha f\left(\frac{t}{L^z}\right)$$


with

$$f(u) \sim \begin{cases} u^\beta, & u \ll 1, \\ \text{const}, & u \gg 1. \end{cases} \Rightarrow \beta = \frac{\alpha}{z}$$

Immediate consequences:

$$w_{\text{sat}}(L) \sim L^\alpha, \quad t_x(L) \sim L^z.$$

Interpretation: dynamic renormalization $x \rightarrow bx$, $t \rightarrow b^z t$, $h \rightarrow b^\alpha h$ leaves $w(L, t)/L^\alpha$ invariant as a function of t/L^z .



(Image from Barabási-Stanley's book 95)

Family and Vicsek reported from their simulation that

$$\beta = 0.30 \pm 0.02 \approx \frac{1}{3}.$$

Difficulty: Robust way to compute the slope in order to obtain an estimate of β .

Family, F., Vicsek, T., *Journal of Physics A: Mathematical and General*, 1985

Continuum viewpoint: EW and KPZ



Edwards–Wilkinson (EW)

Linear diffusion + noise

$$\partial_t h = \nu \nabla^2 h + \eta, \quad \langle \eta \eta \rangle \propto \delta(x) \delta(t).$$

Scale invariance (1D): $z = 2$, $\alpha = \frac{1}{2}$
 $\Rightarrow \beta = \frac{1}{4}$.

$$\alpha = \frac{1}{2}, z = 2, \beta = \frac{1}{4}$$

Kardar–Parisi–Zhang (KPZ)

Nonlinear growth

$$\partial_t h = \nu \nabla^2 h + \frac{\lambda}{2} (\nabla h)^2 + \eta.$$

Galilean/tilt invariance: $\alpha + z = 2$. In 1D
(exact):

$$\alpha = \frac{1}{2}, z = \frac{3}{2}, \beta = \frac{1}{3}$$

BD, RSOS, Eden \Rightarrow KPZ universality in 1+1D.

Continuum viewpoint: EW and KPZ



Edwards–Wilkinson (EW)

Linear diffusion + noise

$$\partial_t h = \nu \nabla^2 h + \eta, \quad \langle \eta \eta \rangle \propto \delta(x) \delta(t).$$

Scale invariance (1D): $z = 2$, $\alpha = \frac{1}{2}$
 $\Rightarrow \beta = \frac{1}{4}$.

$$\boxed{\alpha = \frac{1}{2}, z = 2, \beta = \frac{1}{4}}$$

Kardar–Parisi–Zhang (KPZ)

Nonlinear growth

$$\partial_t h = \nu \nabla^2 h + \frac{\lambda}{2} (\nabla h)^2 + \eta.$$

Galilean/tilt invariance: $\alpha + z = 2$. In 1D (exact):

$$\boxed{\alpha = \frac{1}{2}, z = \frac{3}{2}, \beta = \frac{1}{3}}$$

BD, RSOS, Eden \Rightarrow KPZ universality in 1+1D.

Continuum viewpoint: EW and KPZ



Edwards–Wilkinson (EW)

Linear diffusion + noise

$$\partial_t h = \nu \nabla^2 h + \eta, \quad \langle \eta \eta \rangle \propto \delta(x) \delta(t).$$

Scale invariance (1D): $z = 2$, $\alpha = \frac{1}{2}$

$$\Rightarrow \beta = \frac{1}{4}.$$

$$\boxed{\alpha = \frac{1}{2}, z = 2, \beta = \frac{1}{4}}$$

Barber–Parisi–Zhang (BPZ)

Nonlinear growth

$$\partial_t h = \nu \nabla^2 h + \frac{\lambda}{2} (\nabla h)^2 + \eta.$$

Galilean/tilt invariance: $\alpha + z = 2$. In 1D (exact):

$$\boxed{\alpha = \frac{1}{2}, z = \frac{3}{2}, \beta = \frac{1}{3}}$$

BD, RSOS, Eden \Rightarrow KPZ universality in 1+1D.

Continuum viewpoint: EW and KPZ



Edwards–Wilkinson (EW)

Linear diffusion + noise

$$\partial_t h = \nu \nabla^2 h + \eta, \quad \langle \eta \eta \rangle \propto \delta(x) \delta(t).$$

Scale invariance (1D): $z = 2$, $\alpha = \frac{1}{2}$

$$\Rightarrow \beta = \frac{1}{4}.$$

$$\boxed{\alpha = \frac{1}{2}, z = 2, \beta = \frac{1}{4}}$$

Kardar–Parisi–Zhang (KPZ)

Nonlinear growth

$$\partial_t h = \nu \nabla^2 h + \frac{\lambda}{2} (\nabla h)^2 + \eta.$$

Galilean/tilt invariance: $\alpha + z = 2$. In 1D (exact):

$$\boxed{\alpha = \frac{1}{2}, z = \frac{3}{2}, \beta = \frac{1}{3}}$$

BD, RSOS, Eden \Rightarrow KPZ universality in 1+1D.

Continuum viewpoint: EW and KPZ



Edwards–Wilkinson (EW)

Linear diffusion + noise

$$\partial_t h = \nu \nabla^2 h + \eta, \quad \langle \eta \eta \rangle \propto \delta(x) \delta(t).$$

Scale invariance (1D): $z = 2$, $\alpha = \frac{1}{2}$

$$\Rightarrow \beta = \frac{1}{4}.$$

$$\boxed{\alpha = \frac{1}{2}, z = 2, \beta = \frac{1}{4}}$$

Kardar–Parisi–Zhang (KPZ)

Nonlinear growth

$$\partial_t h = \nu \nabla^2 h + \frac{\lambda}{2} (\nabla h)^2 + \eta.$$

Galilean/tilt invariance: $\alpha + z = 2$. In 1D (exact):

$$\boxed{\alpha = \frac{1}{2}, z = \frac{3}{2}, \beta = \frac{1}{3}}$$

BD, RSOS, Eden \Rightarrow KPZ universality in 1+1D.

Continuum viewpoint: EW and KPZ



Edwards–Wilkinson (EW)

Linear diffusion + noise

$$\partial_t h = \nu \nabla^2 h + \eta, \quad \langle \eta \eta \rangle \propto \delta(x) \delta(t).$$

Scale invariance (1D): $z = 2$, $\alpha = \frac{1}{2}$
 $\Rightarrow \beta = \frac{1}{4}$.

$$\boxed{\alpha = \frac{1}{2}, z = 2, \beta = \frac{1}{4}}$$

Kardar–Parisi–Zhang (KPZ)

Nonlinear growth

$$\partial_t h = \nu \nabla^2 h + \frac{\lambda}{2} (\nabla h)^2 + \eta.$$

Galilean/tilt invariance: $\alpha + z = 2$. In 1D (exact):

$$\boxed{\alpha = \frac{1}{2}, z = \frac{3}{2}, \beta = \frac{1}{3}}$$

BD, RSOS, Eden \Rightarrow KPZ universality in 1+1D.

Reading plots (as in Barabási–Stanley)

- ▶ On w vs. t (log–log) for several L : early slopes $\approx \beta$; plateaus $\propto L^\alpha$; horizontal shift of crossovers $\propto L^z$.
- ▶ After rescaling: plot w/L^α vs. t/L^z ; all curves collapse to $f(u)$ with $f(u) \sim u^\beta$ ($u \ll 1$) and $f(u) \rightarrow \text{const}$ ($u \gg 1$).

Rule of thumb (1D):

$$\text{EW: } (\alpha, \beta, z) = \left(\frac{1}{2}, \frac{1}{4}, 2\right), \quad \text{KPZ/BD: } (\alpha, \beta, z) = \left(\frac{1}{2}, \frac{1}{3}, \frac{3}{2}\right).$$

Reading plots (as in Barabási–Stanley)

- ▶ On w vs. t (log–log) for several L : early slopes $\approx \beta$; plateaus $\propto L^\alpha$; horizontal shift of crossovers $\propto L^z$.
- ▶ After rescaling: plot w/L^α vs. t/L^z ; all curves collapse to $f(u)$ with $f(u) \sim u^\beta$ ($u \ll 1$) and $f(u) \rightarrow \text{const}$ ($u \gg 1$).

Rule of thumb (1D):

$$\text{EW: } (\alpha, \beta, z) = \left(\frac{1}{2}, \frac{1}{4}, 2\right), \quad \text{KPZ/BD: } (\alpha, \beta, z) = \left(\frac{1}{2}, \frac{1}{3}, \frac{3}{2}\right).$$

Exactly solvable baseline: Random Deposition (RD)

One particle per step, random column; after t steps heights are multinomial:

$$\mathbb{E}[w^2(L, t)] = \frac{t}{L} \left(1 - \frac{1}{L}\right) \sim \frac{t}{L}.$$

Hence $w \sim (t/L)^{1/2}$ ($\beta = \frac{1}{2}$), *no lateral correlations*, no finite z and no true saturation. RD serves as a growth-only sanity check; FV scaling requires correlated dynamics (EW/KPZ).

Structure factor (optional but handy)

Fourier modes $h_k(t)$, structure factor $S(k, t) = \langle |h_k(t)|^2 \rangle$ obey

$$S(k, t) \sim k^{-(2\alpha+1)} g(k^z t), \quad g(y) \sim \begin{cases} y^{2\beta+1} & y \ll 1, \\ \text{const} & y \gg 1. \end{cases}$$

Useful when $w(L, t)$ is noisy; fit (α, z) from spectral slopes and collapse $S(k, t)$.

Cheat sheet / takeaways

$$w(L, t) \sim L^\alpha f\left(\frac{t}{L^z}\right), \quad f(u) \sim u^\beta \text{ (} u \ll 1 \text{), } f(u) \rightarrow \text{const (} u \gg 1 \text{), } \beta = \alpha/z$$

1D benchmarks: EW (α, β, z) = ($\frac{1}{2}, \frac{1}{4}, 2$); KPZ/BD ($\frac{1}{2}, \frac{1}{3}, \frac{3}{2}$).

Workflow: measure $w(L, t)$ \Rightarrow fit α and β \Rightarrow infer/fit z \Rightarrow collapse w/L^α vs. t/L^z .

Simulations on
Random deposition vs. Ballistic decomposition

Study of growing interfaces in a thin film


— Convection of nematic liquid crystal*

Show movies !

Takeuchi, K. A., Sano, M., Sasamoto, T., Spohn, H., *Sci. Rep.*, 2011

Study of growing interfaces in a thin film

— Convection of nematic liquid crystal*



Prediction from KPZ equation:


$$h \asymp v_\infty t + (\Gamma t)^{1/3} \xi$$

Takeuchi, K. A., Sano, M., Sasamoto, T., Spohn, H., *Sci. Rep.*, 2011

Study of growing interfaces in a thin film

— Convection of nematic liquid crystal*

$$h \asymp v_\infty t + (\Gamma t)^{1/3} \xi$$



KPZ Equation '86

$$\frac{\partial}{\partial t} h(t, x) = \frac{1}{2} \Delta h(t, x) + \frac{\lambda}{2} (\nabla h)^2 + \dot{W}(t, x) \quad (\text{KPZ})$$



Mehran Kardar (1957 –) Giorgio Parisi (1948 –)



Yicheng Zhang

Kardar, M., Parisi, G., Zhang, Y.-C., *Phys. Rev. Lett.*, 1986

Plan


Introduction to growth model and SPDE

Family-Vicsek scaling and experiments


More examples

Tetromino Pieces

More models? Even more simpler?




Paper – a random environment




Zhang, J., Zhang, Y.-C., Alstrøm, P., Levinsen, M., *Phys. A: Stat. Mech. Appl.*, 1992

Paper wetting experiment




Barabási, A.-L., Stanley, H. E., 1995

Paper wetting experiment





Barabási, A.-L., Stanley, H. E., 1995

Paper burning experiment



Zhang, J., Zhang, Y.-C., Alstrøm, P., Levinsen, M., *Phys. A: Stat. Mech. Appl.*, 1992

Paper rupture experiment



Kertész, J., Horváth, V. k., Weber, F., *Fractals*, 1993

Plan


Introduction to growth model and SPDE

Family-Vicsek scaling and experiments

More examples

Tetromino Pieces


Tetrominoes



- ▶ “1x1”: Single (extra single-site particle)
- ▶ “I”: Horizontal, Vertical
- ▶ “J, L, T”: Up, Right, Down, Left
- ▶ “S, Z”: Horizontal, Vertical
- ▶ “O”: Single (2x2 square)
- ▶ Sticky
- ▶ Nonsticky

$$(1 + 1 \times 2 + 3 \times 4 + 2 \times 2 + 1) \times 2 = 20 \times 2 = 40 \text{ types of pieces}$$


Tetrominoes



- ▶ “1x1”: Single (extra single-site particle)
- ▶ “I”: Horizontal, Vertical
- ▶ “J, L, T”: Up, Right, Down, Left
- ▶ “S, Z”: Horizontal, Vertical
- ▶ “O”: Single (2x2 square)
- ▶ Sticky
- ▶ Nonsticky

$$(1 + 1 \times 2 + 3 \times 4 + 2 \times 2 + 1) \times 2 = 20 \times 2 = 40 \text{ types of pieces}$$

Tetrominoes



- ▶ “1x1”: Single (extra single-site particle)
- ▶ “I”: Horizontal, Vertical
- ▶ “J, L, T”: Up, Right, Down, Left
- ▶ “S, Z”: Horizontal, Vertical
- ▶ “O”: Single (2x2 square)
- ▶ Sticky
- ▶ Nonsticky

$$(1 + 1 \times 2 + 3 \times 4 + 2 \times 2 + 1) \times 2 = 20 \times 2 = 40 \text{ types of pieces}$$

Configure files

```
steps: 12000  
width: 100  
height: 300  
seed: 12  
Piece-00: [20, 0]  
Piece-01: [20, 0]  
Piece-02: [20, 0]  
Piece-03: [20, 0]  
Piece-04: [20, 0]  
Piece-05: [20, 0]  
Piece-06: [20, 0]  
Piece-07: [20, 0]  
Piece-08: [20, 0]  
Piece-09: [20, 0]  
Piece-10: [20, 0]  
Piece-11: [20, 0]  
Piece-12: [20, 0]  
Piece-13: [20, 0]  
Piece-14: [20, 0]  
Piece-15: [20, 0]  
Piece-16: [20, 0]  
Piece-17: [20, 0]  
Piece-18: [20, 0]  
Piece-19: [20, 0]
```

All nonsticky pieces
with equal prob.

```
steps: 12000  
width: 100  
height: 300  
seed: 12  
Piece-00: [0, 20]  
Piece-01: [0, 20]  
Piece-02: [0, 20]  
Piece-03: [0, 20]  
Piece-04: [0, 20]  
Piece-05: [0, 20]  
Piece-06: [0, 20]  
Piece-07: [0, 20]  
Piece-08: [0, 20]  
Piece-09: [0, 20]  
Piece-10: [0, 20]  
Piece-11: [0, 20]  
Piece-12: [0, 20]  
Piece-13: [0, 20]  
Piece-14: [0, 20]  
Piece-15: [0, 20]  
Piece-16: [0, 20]  
Piece-17: [0, 20]  
Piece-18: [0, 20]  
Piece-19: [0, 20]
```

All sticky pieces
with equal prob.

```
steps: 12000  
width: 100  
height: 300  
seed: 12  
Piece-00: [0, 0]  
Piece-01: [0, 0]  
Piece-02: [0, 0]  
Piece-03: [0, 0]  
Piece-04: [0, 0]  
Piece-05: [0, 0]  
Piece-06: [0, 0]  
Piece-07: [0, 0]  
Piece-08: [0, 0]  
Piece-09: [0, 0]  
Piece-10: [0, 0]  
Piece-11: [0, 0]  
Piece-12: [0, 0]  
Piece-13: [0, 0]  
Piece-14: [0, 0]  
Piece-15: [0, 0]  
Piece-16: [0, 0]  
Piece-17: [0, 0]  
Piece-18: [0, 0]  
Piece-19: [20, 80]
```

20% nonsticky
+ 80% sticky
of 1x1 piece

Question

For various configurations of Tetromino pieces, do the resulting surface robustly exhibit Family–Vicsek scaling?

Will the scaling exponent β be always close to $\frac{1}{3}$?

Setup

Consider a mixture of $\ell\%$ ($\ell \in [0, 100]$) nonsticky pieces and $(100-\alpha)\%$ sticky pieces, where only the 1×1 piece is sticky.

Question

How does α influence the scaling exponents, in particular, $\beta \stackrel{?}{\approx} \frac{1}{3}$?

Simulations and log-log plots:

https://chenle02.github.io/2025-10-28_Emerging_Synergies_Banff_Le/exp13/videos_and_images_display.html

Setup

Consider a mixture of $\ell\%$ ($\ell \in [0, 100]$) nonsticky pieces and $(100-\alpha)\%$ sticky pieces, where only the 1×1 piece is sticky.

Question

How does α influence the scaling exponents, in particular, $\beta \approx \frac{1}{3}$?

Simulations and log-log plots:

https://chenle02.github.io/2025-10-28_Emerging_Synergies_Banff_Le/exp13/videos_and_images_display.html

Outreach Highlights



AU-SSI 2023



AU-SSI 2024



Destination STEM 2023

Main References:

- Barabási, A.-L., & Stanley, H. E. (1995). *Fractal concepts in surface growth*. Cambridge University Press, Cambridge.
- Family, F., & Vicsek, T. (1985). Scaling of the active zone in the eden process on percolation networks and the ballistic deposition model. *Journal of Physics A: Mathematical and General*, 18(2), L75.
- Kardar, M., Parisi, G., & Zhang, Y.-C. (1986). Dynamic scaling of growing interfaces. *Phys. Rev. Lett.*, 56(9), 889.
- Kertész, J., Horváth, V. k., & Weber, F. (1993). Self-affine rupture lines in paper sheets. *Fractals*, 01(01), 67–74.
- Takeuchi, K. A., Sano, M., Sasamoto, T., & Spohn, H. (2011). Growing interfaces uncover universal fluctuations behind scale invariance. *Sci. Rep.*, 1(1), 1–5.
- Zhang, J., Zhang, Y.-C., Alstrøm, P., & Levinsen, M. (1992). Modeling forest fire by a paper-burning experiment, a realization of the interface growth mechanism. *Phys. A: Stat. Mech. Appl.*, 189(3), 383–389.

Thank you!

Questions?