Numerical approximations for stochastic rough integrations

Yanghui Liu

Baruch College, CUNY

AMS Fall Southeastern Virtual Sectional Meeting, November 20-21, 2021.

A joint work with Z. Selk and S. Tindel.



Outline

1 Elements of rough paths

2 Numerical methods for rough integrals

3 Numerical methods for stochastic rough integrals

Presentation Outline

1 Elements of rough paths

2 Numerical methods for rough integrals

3 Numerical methods for stochastic rough integrals

• We recall some elements of rough paths.

- We recall some elements of rough paths.
- Consider a \mathbb{R}^d -valued smooth path

$$x=(x^1,\ldots,x^d).$$

- We recall some elements of rough paths.
- Consider a \mathbb{R}^d -valued smooth path

$$X=(X^1,\ldots,X^d).$$

Consider multiple integrals of x:

$$\mathbf{x}_{st}^{1} := \int_{s}^{t} dx_{u} = x_{t} - x_{s} \qquad \mathbf{x}_{st}^{2} := \int_{s}^{t} \int_{s}^{u} dx_{v} \otimes dx_{u}$$
$$\mathbf{x}_{st}^{3} := \int_{s}^{t} \int_{s}^{u} \int_{s}^{v} dx_{w} \otimes dx_{v} \otimes dx_{u}$$

and similar definition for m-th order multiple integral \mathbf{x}^m .

- We recall some elements of rough paths.
- Consider a \mathbb{R}^d -valued smooth path

$$x=(x^1,\ldots,x^d).$$

Consider multiple integrals of x:

$$\mathbf{x}_{st}^{1} := \int_{s}^{t} dx_{u} = x_{t} - x_{s} \qquad \mathbf{x}_{st}^{2} := \int_{s}^{t} \int_{s}^{u} dx_{v} \otimes dx_{u}$$
$$\mathbf{x}_{st}^{3} := \int_{s}^{t} \int_{s}^{u} \int_{s}^{v} dx_{w} \otimes dx_{v} \otimes dx_{u}$$

and similar definition for m-th order multiple integral \mathbf{x}^m .

The *n*-th order signature of x:

$$S_n(x)_{st} = (\mathbf{x}_{st}^1, \mathbf{x}_{st}^2, \dots, \mathbf{x}_{st}^n)$$

It is known that for fixed s and t, the $S_n(x)_{st}$ contains all information of x for $n = \infty$.



• Consider *p*-variation continuous path *x* in \mathbb{R}^d . We denote $\delta x_{st} = x_t - x_s$.

- Consider *p*-variation continuous path *x* in \mathbb{R}^d . We denote $\delta x_{st} = x_t x_s$.
- Recall that the p-variation norm is defined as

$$||x||_{\rho\text{-var}} = \left(\sup_{(t_k)\in\mathcal{P}}\sum_{k}|\delta x_{t_kt_{k+1}}|^{\rho}\right)^{1/\rho},$$

where \mathcal{P} is the set of finite partitions of the time interval [0, T].

- Consider *p*-variation continuous path *x* in \mathbb{R}^d . We denote $\delta x_{st} = x_t x_s$.
- Recall that the p-variation norm is defined as

$$||x||_{\rho\text{-var}} = \left(\sup_{(t_k)\in\mathcal{P}}\sum_k |\delta x_{t_kt_{k+1}}|^{\rho}\right)^{1/\rho},$$

where P is the set of finite partitions of the time interval [0, T].

• Suppose that there exists a smooth path x^{ϵ} such that for $n = \lfloor p \rfloor$ the nth order signature $S_n(x^{\epsilon})$ of x^{ϵ} converges under the p-variation norm.

- Consider *p*-variation continuous path *x* in \mathbb{R}^d . We denote $\delta x_{st} = x_t x_s$.
- Recall that the p-variation norm is defined as

$$\|\mathbf{x}\|_{p\text{-var}} = \left(\sup_{(t_k)\in\mathcal{P}}\sum_{k}|\delta x_{t_kt_{k+1}}|^p\right)^{1/p},$$

where P is the set of finite partitions of the time interval [0, T].

- Suppose that there exists a smooth path x^{ϵ} such that for $n = \lfloor p \rfloor$ the nth order signature $S_n(x^{\epsilon})$ of x^{ϵ} converges under the p-variation norm.
- Denote the limit by $S_n(x) := \mathbf{x}$. \mathbf{x} is called a p-rough path.

$$y_t = y_0 + \int_0^t V(y_u) dx_u, \quad t \in [0, T].$$

Consider differential equation for a smooth function x:

$$y_t = y_0 + \int_0^t V(y_u) dx_u, \quad t \in [0, T].$$

• $S_n(x)$: signature of x.

$$y_t = y_0 + \int_0^t V(y_u) dx_u, \quad t \in [0, T].$$

- $S_n(x)$: signature of x.
- $S_n(y)$: signature of solution y.

$$y_t = y_0 + \int_0^t V(y_u) dx_u, \quad t \in [0, T].$$

- $S_n(x)$: signature of x.
- $S_n(y)$: signature of solution y.
- Consider the Itô map $I: S_n(x) \to S_n(y)$.

$$y_t = y_0 + \int_0^t V(y_u) dx_u, \quad t \in [0, T].$$

- $S_n(x)$: signature of x.
- $S_n(y)$: signature of solution y.
- Consider the Itô map $I: S_n(x) \to S_n(y)$.
- I is continuous under p-var norm with p < n + 1. (Lyons '98)

$$y_t = y_0 + \int_0^t V(y_u) dx_u, \quad t \in [0, T].$$

- $S_n(x)$: signature of x.
- S_n(y): signature of solution y.
- Consider the Itô map $I: S_n(x) \to S_n(y)$.
- I is continuous under p-var norm with p < n + 1. (Lyons '98)
- We extend the Itô map I to p-variation rough paths. We define $\mathbf{y} = I(\mathbf{x})$ as the solution of the differential equation

$$d\mathbf{y}_t = V(y_t)d\mathbf{x}_t.$$



• The rough paths theory provides a framework for differential equations driven by an arbitrary irregular noise.

- The rough paths theory provides a framework for differential equations driven by an arbitrary irregular noise.
- Such solutions are path-wise solutions. No probability structure is required (e.g. Markovian or martingale).

- The rough paths theory provides a framework for differential equations driven by an arbitrary irregular noise.
- Such solutions are path-wise solutions. No probability structure is required (e.g. Markovian or martingale).
- The rough paths framework provides the stability of the Itô map.

- The rough paths theory provides a framework for differential equations driven by an arbitrary irregular noise.
- Such solutions are path-wise solutions. No probability structure is required (e.g. Markovian or martingale).
- The rough paths framework provides the stability of the Itô map.
- The signature has been applied to model complex data streams.

Presentation Outline

Elements of rough paths

2 Numerical methods for rough integrals

3 Numerical methods for stochastic rough integrals

• Consider a *p*-rough path **x**. Let \mathbf{x}^{ϵ} be such that $\mathbf{x}^{\epsilon} \to \mathbf{x}$.

- Consider a *p*-rough path **x**. Let \mathbf{x}^{ϵ} be such that $\mathbf{x}^{\epsilon} \to \mathbf{x}$.
- For $V \in C^{\infty}$ we define the rough integral:

$$\int_0^T V(x)d\mathbf{x} = \lim_{\epsilon \to 0} \int_0^T V(x^{\epsilon})dx^{\epsilon}.$$

- Consider a *p*-rough path **x**. Let \mathbf{x}^{ϵ} be such that $\mathbf{x}^{\epsilon} \to \mathbf{x}$.
- For $V \in C^{\infty}$ we define the rough integral:

$$\int_0^T V(x)d\mathbf{x} = \lim_{\epsilon \to 0} \int_0^T V(x^{\epsilon})dx^{\epsilon}.$$

• Let (t_k) be a partition of [0, T]. For each k we consider the approximation

$$\int_{t_k}^{t_{k+1}} V(x_t) d\mathbf{x}_t \approx \int_{t_k}^{t_{k+1}} V(x_{t_k}) dx_t = V(x_{t_k}) \delta x_{t_k t_{k+1}}.$$

- Consider a *p*-rough path **x**. Let \mathbf{x}^{ϵ} be such that $\mathbf{x}^{\epsilon} \to \mathbf{x}$.
- For $V \in C^{\infty}$ we define the rough integral:

$$\int_0^T V(x)d\mathbf{x} = \lim_{\epsilon \to 0} \int_0^T V(x^{\epsilon})dx^{\epsilon}.$$

• Let (t_k) be a partition of [0, T]. For each k we consider the approximation

$$\int_{t_k}^{t_{k+1}} V(x_t) d\mathbf{x}_t \approx \int_{t_k}^{t_{k+1}} V(x_{t_k}) dx_t = V(x_{t_k}) \delta x_{t_k t_{k+1}}.$$

This leads to the Riemann sum approximation

$$\int_0^T V(x_t) d\mathbf{x}_t = \sum_k \int_{t_k}^{t_{k+1}} V(x_t) d\mathbf{x}_t \approx \sum_k V(x_{t_k}) \delta x_{t_k t_{k+1}}.$$



- Consider a *p*-rough path **x**. Let \mathbf{x}^{ϵ} be such that $\mathbf{x}^{\epsilon} \to \mathbf{x}$.
- For $V \in C^{\infty}$ we define the rough integral:

$$\int_0^T V(x)d\mathbf{x} = \lim_{\epsilon \to 0} \int_0^T V(x^{\epsilon})dx^{\epsilon}.$$

• Let (t_k) be a partition of [0, T]. For each k we consider the approximation

$$\int_{t_k}^{t_{k+1}} V(x_t) d\mathbf{x}_t \approx \int_{t_k}^{t_{k+1}} V(x_{t_k}) dx_t = V(x_{t_k}) \delta x_{t_k t_{k+1}}.$$

This leads to the Riemann sum approximation

$$\int_0^T V(x_t) d\mathbf{x}_t = \sum_k \int_{t_k}^{t_{k+1}} V(x_t) d\mathbf{x}_t \approx \sum_k V(x_{t_k}) \delta x_{t_k t_{k+1}}.$$

• Note that if $p \ge 2$ the Riemann sum $\sum_k V(x_{t_k}) \delta x_{t_k t_{k+1}}$ diverges.



• Denote $\mathcal{L}V = \partial VV$ and $\mathcal{L}^n = \mathcal{L} \circ \cdots \circ \mathcal{L}$.

- Denote $\mathcal{L}V = \partial VV$ and $\mathcal{L}^n = \mathcal{L} \circ \cdots \circ \mathcal{L}$.
- We consider an improved approximation

$$\int_{t_k}^{t_{k+1}} V(x_t) d\mathbf{x}_t \approx$$

- Denote $\mathcal{L}V = \partial VV$ and $\mathcal{L}^n = \mathcal{L} \circ \cdots \circ \mathcal{L}$.
- We consider an improved approximation

$$\int_{t_k}^{t_{k+1}} V(x_t) d\mathbf{x}_t \approx \int_{t_k}^{t_{k+1}} \underbrace{\left(V(x_{t_k}) + (\mathcal{L}V)(x_{t_k})\mathbf{x}_{t_kt}^1 + \dots + (\mathcal{L}^{n-1}V)(x_{t_k})\mathbf{x}_{t_kt}^{n-1}\right)}_{\text{The first } n \text{ terms of Taylor expansion of } V(y_t) \text{ at } t} d\mathbf{x}_t$$

- Denote $\mathcal{L}V = \partial VV$ and $\mathcal{L}^n = \mathcal{L} \circ \cdots \circ \mathcal{L}$.
- We consider an improved approximation

$$\begin{split} \int_{t_k}^{t_{k+1}} V(x_t) d\textbf{x}_t &\approx \int_{t_k}^{t_{k+1}} \underbrace{\left(V(x_{t_k}) + (\mathcal{L}V)(x_{t_k})\textbf{x}_{t_kt}^1 + \dots + (\mathcal{L}^{n-1}V)(x_{t_k})\textbf{x}_{t_kt}^{n-1}\right)}_{\text{The first } n \text{ terms of Taylor expansion of } V(y_t) \text{ at } t} \\ &= V(x_{t_k}) \delta x_{t_kt_{k+1}} + (\mathcal{L}V)(x_{t_k})\textbf{x}_{t_kt_{k+1}}^2 + \dots + (\mathcal{L}^{n-1}V)(x_{t_k})\textbf{x}_{t_kt_{k+1}}^n \end{split}$$

- Denote $\mathcal{L}V = \partial VV$ and $\mathcal{L}^n = \mathcal{L} \circ \cdots \circ \mathcal{L}$.
- We consider an improved approximation

$$\int_{t_k}^{t_{k+1}} V(x_t) d\mathbf{x}_t \approx \int_{t_k}^{t_{k+1}} \underbrace{\left(V(x_{t_k}) + (\mathcal{L}V)(x_{t_k})\mathbf{x}_{t_kt}^1 + \dots + (\mathcal{L}^{n-1}V)(x_{t_k})\mathbf{x}_{t_kt}^{n-1}\right)}_{\text{The first } n \text{ terms of Taylor expansion of } V(y_t) \text{ at } t}$$

$$= V(x_{t_k}) \delta x_{t_kt_{k+1}} + (\mathcal{L}V)(x_{t_k})\mathbf{x}_{t_kt_{k+1}}^2 + \dots + (\mathcal{L}^{n-1}V)(x_{t_k})\mathbf{x}_{t_kt_{k+1}}^n$$

We obtain the compensated Riemann sum:

$$\int_0^T V(x_t) d\mathbf{x}_t \approx \sum_k V(x_{t_k}) \delta x_{t_k t_{k+1}} + (\mathcal{L}V)(x_{t_k}) \mathbf{x}_{t_k t_{k+1}}^2 + \dots + (\mathcal{L}^{n-1}V)(x_{t_k}) \mathbf{x}_{t_k t_{k+1}}^n.$$

- Denote $\mathcal{L}V = \partial VV$ and $\mathcal{L}^n = \mathcal{L} \circ \cdots \circ \mathcal{L}$.
- We consider an improved approximation

$$\int_{t_k}^{t_{k+1}} V(x_t) d\mathbf{x}_t \approx \int_{t_k}^{t_{k+1}} \underbrace{\left(V(x_{t_k}) + (\mathcal{L}V)(x_{t_k})\mathbf{x}_{t_kt}^1 + \dots + (\mathcal{L}^{n-1}V)(x_{t_k})\mathbf{x}_{t_kt}^{n-1}\right)}_{\text{The first } n \text{ terms of Taylor expansion of } V(y_t) \text{ at } t}$$

$$= V(x_{t_k}) \delta x_{t_kt_{k+1}} + (\mathcal{L}V)(x_{t_k})\mathbf{x}_{t_kt_{k+1}}^2 + \dots + (\mathcal{L}^{n-1}V)(x_{t_k})\mathbf{x}_{t_kt_{k+1}}^n$$

We obtain the compensated Riemann sum:

$$\int_0^T V(x_t) d\mathbf{x}_t \approx \sum_k V(x_{t_k}) \delta x_{t_k t_{k+1}} + (\mathcal{L}V)(x_{t_k}) \mathbf{x}_{t_k t_{k+1}}^2 + \cdots + (\mathcal{L}^{n-1}V)(x_{t_k}) \mathbf{x}_{t_k t_{k+1}}^n.$$

• The compensated Riemann sum converges to the rough integral $\int_0^T V(x) d\mathbf{x}$ when n > p-1.

- Denote $\mathcal{L}V = \partial VV$ and $\mathcal{L}^n = \mathcal{L} \circ \cdots \circ \mathcal{L}$.
- We consider an improved approximation

$$\int_{t_k}^{t_{k+1}} V(x_t) d\mathbf{x}_t \approx \int_{t_k}^{t_{k+1}} \underbrace{\left(V(x_{t_k}) + (\mathcal{L}V)(x_{t_k})\mathbf{x}_{t_kt}^1 + \dots + (\mathcal{L}^{n-1}V)(x_{t_k})\mathbf{x}_{t_kt}^{n-1}\right)}_{\text{The first } n \text{ terms of Taylor expansion of } V(y_t) \text{ at } t}$$

$$= V(x_{t_k}) \delta x_{t_kt_{k+1}} + (\mathcal{L}V)(x_{t_k})\mathbf{x}_{t_kt_{k+1}}^2 + \dots + (\mathcal{L}^{n-1}V)(x_{t_k})\mathbf{x}_{t_kt_{k+1}}^n$$

We obtain the compensated Riemann sum:

$$\int_0^T V(x_t) d\mathbf{x}_t \approx \sum_k V(x_{t_k}) \delta x_{t_k t_{k+1}} + (\mathcal{L}V)(x_{t_k}) \mathbf{x}_{t_k t_{k+1}}^2 + \cdots + (\mathcal{L}^{n-1}V)(x_{t_k}) \mathbf{x}_{t_k t_{k+1}}^n.$$

- The compensated Riemann sum converges to the rough integral $\int_0^T V(x) d\mathbf{x}$ when n > p 1.
- Note that the compensated Riemann sum requires the computations of signatures of ${\bf x}$.



• Consider a p-rough path \mathbf{x} with p < 4.

- Consider a p-rough path x with p < 4.
- Consider continuous paths y, y' and y" such that

$$\delta y_{st} = y_s' \mathbf{x}_{st}^1 + y_s'' \mathbf{x}_{st}^2 + r_{st}^0, \qquad \delta y_{st}' = y_s'' \mathbf{x}_{st}^1 + r_{st}^1,$$

- Consider a p-rough path x with p < 4.
- Consider continuous paths y, y' and y" such that

$$\delta y_{st} = y_s' \mathbf{x}_{st}^1 + y_s'' \mathbf{x}_{st}^2 + r_{st}^0, \qquad \delta y_{st}' = y_s'' \mathbf{x}_{st}^1 + r_{st}^1,$$

Then the corresponding compensated Riemann sum converges:

$$\sum_{k} y_{t_{k}} \delta \mathbf{x}_{t_{k}t_{k+1}}^{1} + y_{t_{k}}' \delta \mathbf{x}_{t_{k}t_{k+1}}^{2} + y_{t_{k}}'' \delta \mathbf{x}_{t_{k}t_{k+1}}^{3} \to \int_{0}^{T} y_{t} d\mathbf{x}_{t}.$$

- Consider a p-rough path x with p < 4.
- Consider continuous paths y, y' and y'' such that

$$\delta y_{st} = y_s' \mathbf{x}_{st}^1 + y_s'' \mathbf{x}_{st}^2 + r_{st}^0, \qquad \delta y_{st}' = y_s'' \mathbf{x}_{st}^1 + r_{st}^1,$$

Then the corresponding compensated Riemann sum converges:

$$\sum_{k} y_{t_{k}} \delta \mathbf{x}_{t_{k}t_{k+1}}^{1} + y_{t_{k}}' \delta \mathbf{x}_{t_{k}t_{k+1}}^{2} + y_{t_{k}}'' \delta \mathbf{x}_{t_{k}t_{k+1}}^{3} \to \int_{0}^{T} y_{t} d\mathbf{x}_{t}.$$

• (y, y', y'') is called a controlled paths of **x** of order 2.

- Consider a p-rough path x with p < 4.
- Consider continuous paths y, y' and y'' such that

$$\delta y_{st} = y_s' \mathbf{x}_{st}^1 + y_s'' \mathbf{x}_{st}^2 + r_{st}^0, \qquad \delta y_{st}' = y_s'' \mathbf{x}_{st}^1 + r_{st}^1,$$

Then the corresponding compensated Riemann sum converges:

$$\sum_{k} y_{t_{k}} \delta \mathbf{x}_{t_{k}t_{k+1}}^{1} + y_{t_{k}}' \delta \mathbf{x}_{t_{k}t_{k+1}}^{2} + y_{t_{k}}'' \delta \mathbf{x}_{t_{k}t_{k+1}}^{3} \to \int_{0}^{T} y_{t} d\mathbf{x}_{t}.$$

- (y, y', y'') is called a controlled paths of **x** of order 2.
- Such path contains most of the interesting examples. e.g. y = V(x) or y is the solution of a RDE.

Presentation Outline

1 Elements of rough paths

2 Numerical methods for rough integrals

3 Numerical methods for stochastic rough integrals

• Let $X = (X^1, \dots, X^d)$ be a continuous centered Gaussian process with i.i.d. components.

- Let $X = (X^1, \dots, X^d)$ be a continuous centered Gaussian process with i.i.d. components.
- The covariance function of X is defined as follows

$$R(s,t) = \mathbb{E}[X_s^j X_t^j].$$

- Let $X = (X^1, \dots, X^d)$ be a continuous centered Gaussian process with i.i.d. components.
- The covariance function of X is defined as follows

$$R(s,t) = \mathbb{E}[X_s^j X_t^j].$$

• The information concerning *X* is mostly encoded in the rectangular increments of *R*:

$$R_{uv}^{st} := R(t,v) - R(t,u) - R(s,v) + R(s,u) = \mathbb{E}[\delta X_{st}^j \delta X_{uv}^j].$$

- Let $X = (X^1, ..., X^d)$ be a continuous centered Gaussian process with i.i.d. components.
- The covariance function of X is defined as follows

$$R(s,t) = \mathbb{E}[X_s^j X_t^j].$$

• The information concerning *X* is mostly encoded in the rectangular increments of *R*:

$$R_{uv}^{st} := R(t,v) - R(t,u) - R(s,v) + R(s,u) = \mathbb{E}[\delta X_{st}^j \delta X_{uv}^j].$$

• For $\rho \ge 1$ we define the ρ -variation of R as

$$\|R\|_{
ho ext{-var}} = \sup_{(t_i),(t_j')} \left(\sum_{i,j} \left| R_{t_i t_{i+1}}^{t_j' t_{j+1}'} \right|^
ho \right)^{1/
ho},$$

where (t_j) and (t'_j) are partitions on [0, T].

- Let $X = (X^1, ..., X^d)$ be a continuous centered Gaussian process with i.i.d. components.
- The covariance function of X is defined as follows

$$R(s,t) = \mathbb{E}[X_s^j X_t^j].$$

• The information concerning *X* is mostly encoded in the rectangular increments of *R*:

$$R_{uv}^{st} := R(t, v) - R(t, u) - R(s, v) + R(s, u) = \mathbb{E}[\delta X_{st}^j \delta X_{uv}^j].$$

• For $\rho \geq 1$ we define the ρ -variation of R as

$$\|R\|_{
ho ext{-var}} = \sup_{(t_i),(t_j')} \left(\sum_{i,j} \left| R_{t_i t_{i+1}}^{t_j' t_{j+1}'} \right|^
ho \right)^{1/
ho},$$

where (t_i) and (t'_i) are partitions on [0, T].

• If R has finite ρ -variation for $\rho \in [1,2)$, then X gives raise to a p-rough path, provided $p > 2\rho$. (Friz-Victoir '11).

$$\text{tr-}\mathcal{J}_0^T(y,X) = \sum_k \frac{y_{t_k} + y_{t_{k+1}}}{2} \cdot \delta X_{t_k t_{k+1}}.$$

$$\operatorname{tr-}\mathcal{J}_0^T(y,X) = \sum_k \frac{y_{t_k} + y_{t_{k+1}}}{2} \cdot \delta X_{t_k t_{k+1}}.$$

Theorem (Liu-Selk-Tindel '21)

$$\operatorname{tr-}\mathcal{J}_0^T(y,X) = \sum_{k} \frac{y_{t_k} + y_{t_{k+1}}}{2} \cdot \delta X_{t_k t_{k+1}}.$$

Theorem (Liu-Selk-Tindel '21)

• Suppose that $\|R\|_{\rho\text{-var}}<\infty$. Then as the mesh size of the partition (t_k) goes to 0 we have

$$\operatorname{tr-}\mathcal{J}_0^T(y,X) o \int_0^T y_t d\mathbf{X}_t$$
 in probability.

$$\operatorname{tr-}\mathcal{J}_0^T(y,X) = \sum_{k} \frac{y_{t_k} + y_{t_{k+1}}}{2} \cdot \delta X_{t_k t_{k+1}}.$$

Theorem (Liu-Selk-Tindel '21)

• Suppose that $\|R\|_{\rho\text{-var}} < \infty$. Then as the mesh size of the partition (t_k) goes to 0 we have

$$\operatorname{tr-}\mathcal{J}_0^T(y,X) o \int_0^T y_t d\mathbf{X}_t$$
 in probability.

• Suppose that there exists a constant C>0 such that $\|R\|_{\rho\text{-var},[s,t]\times[0,T]}\leq C|t-s|$ for all $[s,t]\subset[0,T]$. Then we have

$$\operatorname{tr-}\mathcal{J}_0^T(y,X) \to \int_0^T y_t d\mathbf{X}_t$$
 almost surely.



With a careful rearrangement of the trapezoid rule we can recast it as

$$\text{tr-}\,\mathcal{J}_0^{\,T}(y,X) = \sum_k \, \mathit{I}_1 + \mathit{I}_2 + \mathit{I}_3 + \mathit{I}_4,$$

With a careful rearrangement of the trapezoid rule we can recast it as

$$\text{tr-}\,\mathcal{J}_0^{\,\mathsf{T}}(y,X) = \sum_k \, \mathit{I}_1 \,+\, \mathit{I}_2 \,+\, \mathit{I}_3 \,+\, \mathit{I}_4,$$

where

$$\begin{split} I_1 &= y_{t_k} \cdot \mathbf{X}_{t_k t_{k+1}}^1 + y_{t_k}' \cdot \mathbf{X}_{t_k t_{k+1}}^2 + y_{t_k}'' \cdot \mathbf{X}_{t_k t_{k+1}}^3 \\ I_2 &= \frac{1}{2} y_{t_k}' \mathbf{X}_{t_k t_{k+1}}^1 \cdot \mathbf{X}_{t_k t_{k+1}}^1 - y_{t_k}' \cdot \mathbf{X}_{t_k t_{k+1}}^2 \\ I_3 &= \frac{1}{2} y_{t_k}'' \mathbf{X}_{t_k t_{k+1}}^2 \cdot \mathbf{X}_{t_k t_{k+1}}^1 - y_{t_k}'' \cdot \mathbf{X}_{t_k t_{k+1}}^3 \\ I_4 &= \frac{1}{2} r_{t_k t_{k+1}}^0 \cdot \mathbf{X}_{t_k t_{k+1}}^1. \end{split}$$

With a careful rearrangement of the trapezoid rule we can recast it as

$$\text{tr-}\,\mathcal{J}_0^{\,\mathsf{T}}(y,X) = \sum_k \, \mathit{I}_1 \,+\, \mathit{I}_2 \,+\, \mathit{I}_3 \,+\, \mathit{I}_4,$$

where

$$\begin{split} I_1 &= y_{t_k} \cdot \mathbf{X}_{t_k t_{k+1}}^1 + y_{t_k}' \cdot \mathbf{X}_{t_k t_{k+1}}^2 + y_{t_k}'' \cdot \mathbf{X}_{t_k t_{k+1}}^3 \\ I_2 &= \frac{1}{2} y_{t_k}' \mathbf{X}_{t_k t_{k+1}}^1 \cdot \mathbf{X}_{t_k t_{k+1}}^1 - y_{t_k}' \cdot \mathbf{X}_{t_k t_{k+1}}^2 \\ I_3 &= \frac{1}{2} y_{t_k}'' \mathbf{X}_{t_k t_{k+1}}^2 \cdot \mathbf{X}_{t_k t_{k+1}}^1 - y_{t_k}'' \cdot \mathbf{X}_{t_k t_{k+1}}^3 \\ I_4 &= \frac{1}{2} r_{t_k t_{k+1}}^0 \cdot \mathbf{X}_{t_k t_{k+1}}^1. \end{split}$$

• I_1 is the compensated Riemann sum of $\int_0^T y d\mathbf{X}$.

With a careful rearrangement of the trapezoid rule we can recast it as

$$\operatorname{tr-} \mathcal{J}_0^T(y,X) = \sum_k I_1 + I_2 + I_3 + I_4,$$

where

$$\begin{split} I_1 &= y_{t_k} \cdot \mathbf{X}_{t_k t_{k+1}}^1 + y_{t_k}' \cdot \mathbf{X}_{t_k t_{k+1}}^2 + y_{t_k}'' \cdot \mathbf{X}_{t_k t_{k+1}}^3 \\ I_2 &= \frac{1}{2} y_{t_k}' \mathbf{X}_{t_k t_{k+1}}^1 \cdot \mathbf{X}_{t_k t_{k+1}}^1 - y_{t_k}' \cdot \mathbf{X}_{t_k t_{k+1}}^2 \\ I_3 &= \frac{1}{2} y_{t_k}'' \mathbf{X}_{t_k t_{k+1}}^2 \cdot \mathbf{X}_{t_k t_{k+1}}^1 - y_{t_k}'' \cdot \mathbf{X}_{t_k t_{k+1}}^3 \\ I_4 &= \frac{1}{2} r_{t_k}^0 t_{k+1} \cdot \mathbf{X}_{t_k t_{k+1}}^1 . \end{split}$$

- I_1 is the compensated Riemann sum of $\int_0^T y d\mathbf{X}$.
- I_2 and I_3 are weighted random sums of the forms: $\sum_k y'_{t_k} h^n_{t_k t_{k+1}}$ and $\sum_k y''_{t_k} \tilde{h}^n_{t_k t_{k+1}}$.

With a careful rearrangement of the trapezoid rule we can recast it as

$$\operatorname{tr-} \mathcal{J}_0^T(y,X) = \sum_k I_1 + I_2 + I_3 + I_4,$$

where

$$\begin{split} I_1 &= y_{t_k} \cdot \mathbf{X}_{t_k t_{k+1}}^1 + y_{t_k}' \cdot \mathbf{X}_{t_k t_{k+1}}^2 + y_{t_k}'' \cdot \mathbf{X}_{t_k t_{k+1}}^3 \\ I_2 &= \frac{1}{2} y_{t_k}' \mathbf{X}_{t_k t_{k+1}}^1 \cdot \mathbf{X}_{t_k t_{k+1}}^1 - y_{t_k}' \cdot \mathbf{X}_{t_k t_{k+1}}^2 \\ I_3 &= \frac{1}{2} y_{t_k}'' \mathbf{X}_{t_k t_{k+1}}^2 \cdot \mathbf{X}_{t_k t_{k+1}}^1 - y_{t_k}'' \cdot \mathbf{X}_{t_k t_{k+1}}^3 \\ I_4 &= \frac{1}{2} r_{t_k t_{k+1}}^0 \cdot \mathbf{X}_{t_k t_{k+1}}^1. \end{split}$$

- I_1 is the compensated Riemann sum of $\int_0^T y d\mathbf{X}$.
- l_2 and l_3 are weighted random sums of the forms: $\sum_k y'_{t_k} h^n_{t_k t_{k+1}}$ and $\sum_k y''_{t_k} \tilde{h}^n_{t_k t_{k+1}}$.
- The convergences of I_i , i = 2, 3, 4 can be shown based on a transfer principle combined with some 2d young-type estimates.



• In order to bound a weighted sum $\sum_{k=0}^{n-1} y_{t_k} h_{t_k t_{k+1}}^n$ it suffices to consider the following elementary weighted sums:

$$\sum_{s \leq t_k < t} h_{t_k t_{k+1}}^n, \quad \sum_{s \leq t_k < t} \mathbf{X}_{st_k}^1 h_{t_k t_{k+1}}^n, \cdots \sum_{s \leq t_k < t} \mathbf{X}_{st_k}^\ell h_{t_k t_{k+1}}^n,$$

where ℓ is an integer depending on X and h^n .

• In order to bound a weighted sum $\sum_{k=0}^{n-1} y_{t_k} h_{t_k t_{k+1}}^n$ it suffices to consider the following elementary weighted sums:

$$\sum_{s \leq t_k < t} h^n_{t_k t_{k+1}}, \quad \sum_{s \leq t_k < t} \mathbf{X}^1_{s t_k} h^n_{t_k t_{k+1}}, \cdots \sum_{s \leq t_k < t} \mathbf{X}^\ell_{s t_k} h^n_{t_k t_{k+1}},$$

where ℓ is an integer depending on X and h^n .

 These special weighted sums belong to finite Wiener chaos and are easier to handle.

• In order to bound a weighted sum $\sum_{k=0}^{n-1} y_{t_k} h_{t_k t_{k+1}}^n$ it suffices to consider the following elementary weighted sums:

$$\sum_{s \leq t_k < t} h^n_{t_k t_{k+1}}, \quad \sum_{s \leq t_k < t} \mathbf{X}^1_{s t_k} h^n_{t_k t_{k+1}}, \cdots \sum_{s \leq t_k < t} \mathbf{X}^\ell_{s t_k} h^n_{t_k t_{k+1}},$$

where ℓ is an integer depending on X and h^n .

- These special weighted sums belong to finite Wiener chaos and are easier to handle.
- For example, in order to estimate $I_2 = \frac{1}{2} \sum_k y'_{t_k} (\mathbf{X}^1_{t_k t_{k+1}} \cdot \mathbf{X}^1_{t_k t_{k+1}} \cdot \mathbf{X}^2_{t_k t_{k+1}})$ it suffices to bound

$$\sum_{k} (\boldsymbol{X}_{t_k t_{k+1}}^1 \cdot \boldsymbol{X}_{t_k t_{k+1}}^1 - \boldsymbol{X}_{t_k t_{k+1}}^2) \quad \text{and} \quad \sum_{k} \boldsymbol{X}_{t_k}^1 (\boldsymbol{X}_{t_k t_{k+1}}^1 \cdot \boldsymbol{X}_{t_k t_{k+1}}^1 - \boldsymbol{X}_{t_k t_{k+1}}^2)$$



• In order to bound a weighted sum $\sum_{k=0}^{n-1} y_{t_k} h_{t_k t_{k+1}}^n$ it suffices to consider the following elementary weighted sums:

$$\sum_{s \leq t_k < t} h^n_{t_k t_{k+1}}, \quad \sum_{s \leq t_k < t} \mathbf{X}^1_{s t_k} h^n_{t_k t_{k+1}}, \cdots \sum_{s \leq t_k < t} \mathbf{X}^\ell_{s t_k} h^n_{t_k t_{k+1}},$$

where ℓ is an integer depending on X and h^n .

- These special weighted sums belong to finite Wiener chaos and are easier to handle.
- For example, in order to estimate $I_2 = \frac{1}{2} \sum_k y'_{t_k} (\mathbf{X}^1_{t_k t_{k+1}} \cdot \mathbf{X}^1_{t_k t_{k+1}} \cdot \mathbf{X}^2_{t_k t_{k+1}})$ it suffices to bound

$$\sum_{k} (\boldsymbol{X}_{t_{k}t_{k+1}}^{1} \cdot \boldsymbol{X}_{t_{k}t_{k+1}}^{1} - \boldsymbol{X}_{t_{k}t_{k+1}}^{2}) \quad \text{and} \quad \sum_{k} \boldsymbol{X}_{t_{k}}^{1} (\boldsymbol{X}_{t_{k}t_{k+1}}^{1} \cdot \boldsymbol{X}_{t_{k}t_{k+1}}^{1} - \boldsymbol{X}_{t_{k}t_{k+1}}^{2})$$

 Such transfer principle for limit theorems of weighted sums are obtained and has been applied to very general weighted sum.



$$\mathsf{m-}\mathcal{J}_0^T(f(X),X) = \sum_{k=0}^{n-1} f\Big(\frac{X_{t_k} + X_{t_{k+1}}}{2}\Big) \cdot \delta X_{t_k t_{k+1}}.$$

$$\mathsf{m-}\mathcal{J}_0^T(f(X),X) = \sum_{k=0}^{n-1} f\Big(\frac{X_{t_k} + X_{t_{k+1}}}{2}\Big) \cdot \delta X_{t_k t_{k+1}}.$$

Corollary

$$\mathsf{m-}\mathcal{J}_0^T(f(X),X) = \sum_{k=0}^{n-1} f\Big(\frac{X_{t_k} + X_{t_{k+1}}}{2}\Big) \cdot \delta X_{t_k t_{k+1}}.$$

Corollary

• Suppose that $\|R\|_{\rho\text{-var}}<\infty$. Then as the mesh size of the partition (t_k) goes to 0 we have

$$\text{m-}\mathcal{J}_0^T(f(X),X) o \int_0^T f(X_t) d\mathbf{X}_t$$
 in probability.

$$\mathsf{m}\text{-}\mathcal{J}_0^T(f(X),X) = \sum_{k=0}^{n-1} f\Big(\frac{X_{t_k} + X_{t_{k+1}}}{2}\Big) \cdot \delta X_{t_k t_{k+1}}.$$

Corollary

• Suppose that $\|R\|_{\rho\text{-var}}<\infty.$ Then as the mesh size of the partition (t_k) goes to 0 we have

$$\text{m-}\mathcal{J}_0^T(f(X),X) \to \int_0^T f(X_t) d\mathbf{X}_t$$
 in probability.

• If we assume further that $\|R\|_{\rho\text{-var};[s,t]\times[0,T]}\leq C|t-s|$. Then the convergence holds almost surely.

For $a, b \in \mathbb{R}^d$ we consider the following mean value identity

$$\frac{f(a)+f(b)}{2}-f\left(\frac{a+b}{2}\right)=\frac{1}{2}\partial^2 f(c)\left(\frac{b-a}{2}\right)^{\otimes 2},$$

where $c \in \mathbb{R}^3$ satisfies $c = a + \theta(b - a)$ for some $\theta \in [0, 1]$.

For $a, b \in \mathbb{R}^d$ we consider the following mean value identity

$$\frac{f(a)+f(b)}{2}-f\left(\frac{a+b}{2}\right)=\frac{1}{2}\partial^2 f(c)\left(\frac{b-a}{2}\right)^{\otimes 2},$$

where $c \in \mathbb{R}^3$ satisfies $c = a + \theta(b - a)$ for some $\theta \in [0, 1]$.

• Apply the mean value identity with $a = X_{t_k}$ and $b = X_{t_{k+1}}$ to the difference

$$\operatorname{tr-} \mathcal{J}_0^T(f(X),X) - \operatorname{m-} \mathcal{J}_0^T(f(X),X).$$

We will obtain some weighted sums similar to I_3 and I_4 in the previous proof.

For $a, b \in \mathbb{R}^d$ we consider the following mean value identity

$$\frac{f(a)+f(b)}{2}-f\left(\frac{a+b}{2}\right)=\frac{1}{2}\partial^2 f(c)\left(\frac{b-a}{2}\right)^{\otimes 2},$$

where $c \in \mathbb{R}^3$ satisfies $c = a + \theta(b - a)$ for some $\theta \in [0, 1]$.

• Apply the mean value identity with $a = X_{t_k}$ and $b = X_{t_{k+1}}$ to the difference

$$\operatorname{tr-} \mathcal{J}_0^T(f(X),X) - \operatorname{m-} \mathcal{J}_0^T(f(X),X).$$

We will obtain some weighted sums similar to I_3 and I_4 in the previous proof.

 We conclude that the two numerical integral methods converge to the same limit.



Thank you very much for your attention!