

# Math 362: Mathematical Statistics II

Le Chen

le.chen@emory.edu  
chenle02@gmail.com

Emory University  
Atlanta, GA

Last updated on Spring 2021  
Last compiled on January 15, 2023

2021 Spring

Creative Commons License  
(CC By-NC-SA)

# Chapter 10. Goodness-of-fit Tests

## § 10.1 Introduction

## § 10.2 The Multinomial Distribution

## § 10.3 Goodness-of-Fit Tests: All Parameters Known

## § 10.4 Goodness-of-Fit Tests: Parameters Unknown

## § 10.5 Contingency Tables

# Plan

§ 10.1 Introduction

§ 10.2 The Multinomial Distribution

§ 10.3 Goodness-of-Fit Tests: All Parameters Known

§ 10.4 Goodness-of-Fit Tests: Parameters Unknown

§ 10.5 Contingency Tables

# Chapter 10. Goodness-of-fit Tests

§ 10.1 Introduction

§ 10.2 The Multinomial Distribution

§ 10.3 Goodness-of-Fit Tests: All Parameters Known

§ 10.4 Goodness-of-Fit Tests: Parameters Unknown

§ 10.5 Contingency Tables

$p_i$ are known	$p_i$ are unknown
$D = \sum_{i=1}^t \frac{(X_i - np_i)^2}{np_i}$	$D_1 = \sum_{i=1}^t \frac{(X_i - n\hat{p}_i)^2}{n\hat{p}_i}$
$\chi^2$ with f.d. $t - 1$	$\chi^2$ with f.d. $t - 1 - s$
$d = \sum_{i=1}^t \frac{(k_i - np_{i0})^2}{np_{i0}}$	$d_1 = \sum_{i=1}^t \frac{(k_i - n\hat{p}_{i0})^2}{n\hat{p}_{i0}}$
$np_{i0} \geq 5$	$n\hat{p}_{i0} \geq 5$
$d > \chi_{1-\alpha, t-1}^2$	$d_1 > \chi_{1-\alpha, t-1-s}^2$

†  $s$  is the number of unknown parameters.

df = number of classes - 1 - number of unknown parameters.

E.g. 1 Binomial data: 4096 students, each shots basketball 4 times. Let  $X_i$  be the number of hits for the  $i$ th student.

People believe that  $X_i$  should following  $\text{binomial}(4, p)$ , that is, shotting basketball should be something like trying to get red chocolate beans from a jar of beans of two colors.

Find the MLE for  $p$ . Use the data to make a conclusion.

Sol. 1)  $H_0 : X_i \sim \text{binomal}(4, p)$ .

2) Under  $H_0$ , the MLE for  $p$  is  $p_e = \dots = 0.251$

E.g. 1 Binomial data: 4096 students, each shots basketball 4 times. Let  $X_i$  be the number of hits for the  $i$ th student.



Number of Hits, $i$	Obs. Freq., $k_i$
0	1280
1	1717
2	915
3	167
4	17

People believe that  $X_i$  should following binomial(4,  $p$ ), that is, shotting basketball should be something like trying to get red chocolate beans from a jar of beans of two colors.

Find the MLE for  $p$ . Use the data to make a conclusion.

Sol. 1)  $H_0 : X_i \sim \text{binomal}(4, p)$ .

2) Under  $H_0$ , the MLE for  $p$  is  $p_e = \dots = 0.251$

E.g. 1 Binomial data: 4096 students, each shots basketball 4 times. Let  $X_i$  be the number of hits for the  $i$ th student.



Number of Hits, $i$	Obs. Freq., $k_i$
0	1280
1	1717
2	915
3	167
4	17

People believe that  $X_i$  should following binomial(4,  $p$ ), that is, shotting basketball should be something like trying to get red chocolate beans from a jar of beans of two colors.

Find the MLE for  $p$ . Use the data to make a conclusion.

Sol. 1)  $H_0 : X_i \sim \text{binomal}(4, p)$ .

2) Under  $H_0$ , the MLE for  $p$  is  $p_e = \dots = 0.251$



E.g. 1 Binomial data: 4096 students, each shots basketball 4 times. Let  $X_i$  be the number of hits for the  $i$ th student.



Number of Hits, $i$	Obs. Freq., $k_i$
0	1280
1	1717
2	915
3	167
4	17

People believe that  $X_i$  should following binomial(4,  $p$ ), that is, shotting basketball should be something like trying to get red chocolate beans from a jar of beans of two colors.

Find the MLE for  $p$ . Use the data to make a conclusion.

Sol. 1)  $H_0 : X_i \sim \text{binomal}(4, p)$ .

2) Under  $H_0$ , the MLE for  $p$  is  $p_e = \dots = 0.251$

E.g. 1 Binomial data: 4096 students, each shots basketball 4 times. Let  $X_i$  be the number of hits for the  $i$ th student.



Number of Hits, $i$	Obs. Freq., $k_i$
0	1280
1	1717
2	915
3	167
4	17

People believe that  $X_i$  should following binomial(4,  $p$ ), that is, shotting basketball should be something like trying to get red chocolate beans from a jar of beans of two colors.

Find the MLE for  $p$ . Use the data to make a conclusion.

Sol. 1)  $H_0 : X_i \sim \text{binomal}(4, p)$ .

2) Under  $H_0$ , the MLE for  $p$  is  $p_e = \dots = 0.251$

3) Compute the expected frequencies:

$$\implies d_1 = \dots = 6.401.$$

4) Critical region:  $(\chi^2_{.95, 5-1-1}, +\infty) = (7.815, +\infty)$

5) Conclusion: Fail to reject.

6) Alternatively,  $P\text{-value} = \mathbb{P}(\chi^2_3 \geq 6.401) = 0.094, \dots$  discuss...



3) Compute the expected frequencies:

<b>Table 10.4.1</b>		
Number of Hits, $i$	Obs. Freq., $k_i$	Estimated Exp. Freq., $n \hat{p}_{i_o}$
$r'_i s \left\{ \begin{array}{l} 0 \\ 1 \\ 2 \\ 3 \\ 4 \end{array} \right.$	1280	1289.1
	1717	1728.0
	915	868.6
	167	194.0
	17	16.3

$$\implies d_1 = \dots = 6.401.$$

4) Critical region:  $(\chi^2_{.95, 5-1-1}, +\infty) = (7.815, +\infty)$

5) Conclusion: Fail to reject.

6) Alternatively,  $P\text{-value} = \mathbb{P}(\chi^2_3 \geq 6.401) = 0.094, \dots$  discuss...



3) Compute the expected frequencies:

<b>Table 10.4.1</b>		
Number of Hits, $i$	Obs. Freq., $k_i$	Estimated Exp. Freq., $n \hat{p}_{i_o}$
$r'_i s \left\{ \begin{array}{l} 0 \\ 1 \\ 2 \\ 3 \\ 4 \end{array} \right.$	1280	1289.1
	1717	1728.0
	915	868.6
	167	194.0
	17	16.3

$$\implies d_1 = \dots = 6.401.$$

4) Critical region:  $(\chi^2_{.95, 5-1-1}, +\infty) = (7.815, +\infty)$

5) Conclusion: Fail to reject.

6) Alternatively,  $P\text{-value} = \mathbb{P}(\chi^2_3 \geq 6.401) = 0.094, \dots$  discuss...



3) Compute the expected frequencies:

<b>Table 10.4.1</b>		
Number of Hits, $i$	Obs. Freq., $k_i$	Estimated Exp. Freq., $n \hat{p}_{i_0}$
$r'_i s \left\{ \begin{array}{l} 0 \\ 1 \\ 2 \\ 3 \\ 4 \end{array} \right.$	1280	1289.1
	1717	1728.0
	915	868.6
	167	194.0
	17	16.3

$$\implies d_1 = \dots = 6.401.$$

4) Critical region:  $(\chi^2_{.95, 5-1-1}, +\infty) = (7.815, +\infty)$

5) Conclusion: Fail to reject.

6) Alternatively,  $P\text{-value} = \mathbb{P}(\chi^2_3 \geq 6.401) = 0.094, \dots$  discuss...

□

E.g. 2 Does the number of death per day follow the Poisson distribution?

E.g. 2 Does the number of death per day follow the Poisson distribution?

Number of Deaths, $i$	Obs. Freq., $k_i$
0	162
1	267
2	271
3	185
4	111
5	61
6	27
7	8
8	3
9	1
10+	0
	<hr/> 1096





Sol. 1) Let  $X_i$  be the number of death in  $i$ th day,  $1 \leq i \leq 1096$ .

2)  $H_0 : X_i$  follow  $\text{Poisson}(\lambda)$ .

3) The MLE for  $\lambda$  is:  $\lambda_e = \dots = 2.157$ .

4) Compute the expected frequencies:

$$E_{ij} = n \cdot \hat{p}_{ij} = 1096 \cdot 0.0002 = 0.218$$

5) P-value =  $1 - \chi^2_{(1096-1)}(0.05) = 0.0002$  - Reject!





Sol. 1) Let  $X_i$  be the number of death in  $i$ th day,  $1 \leq i \leq 1096$ .

2)  $H_0 : X_i$  follow  $\text{Poisson}(\lambda)$ .

3) The MLE for  $\lambda$  is:  $\lambda_e = \dots = 2.157$ .

4) Compute the expected frequencies:

Table 10.4.2

Number of Deaths, $i$	Obs. Freq., $k_i$	Est. Exp. Freq., $n \hat{p}_{i\omega}$
0	162	126.8
1	267	273.5
2	271	294.9
3	185	212.1
4	111	114.3
5	61	49.3
6	27	17.8
7	8	5.5
8	3	1.4
9	1	0.3
10+	0	0.1
	<u>1096</u>	<u>1096</u>

**Table 10.4.3**

Number of Deaths, $i$	Obs. Freq., $k_i$	Est. Exp. Freq., $n \hat{p}_{i_0}$
$r_1, r_2, \dots, r_8$	0	126.8
	1	273.5
	2	294.9
	3	212.1
	4	114.3
	5	49.3
	6	17.8
	7+	7.3
	1096	1096

Sol. 1) Let  $X_i$  be the number of death in  $i$ th day,  $1 \leq i \leq 1096$ .

2)  $H_0 : X_i$  follow  $\text{Poisson}(\lambda)$ .

3) The MLE for  $\lambda$  is:  $\lambda_e = \dots = 2.157$ .

4) Compute the expected frequencies:

Table 10.4.2

Number of Deaths, $i$	Obs. Freq., $k_i$	Est. Exp. Freq., $n \hat{p}_{i\omega}$
0	162	126.8
1	267	273.5
2	271	294.9
3	185	212.1
4	111	114.3
5	61	49.3
6	27	17.8
7	8	5.5
8	3	1.4
9	1	0.3
10+	0	0.1
	<u>1096</u>	<u>1096</u>

**Table 10.4.3**

Number of Deaths, $i$	Obs. Freq., $k_i$	Est. Exp. Freq., $n \hat{p}_{i_0}$	
$r_1, r_2, \dots, r_8$	0	162	126.8
	1	267	273.5
	2	271	294.9
	3	185	212.1
	4	111	114.3
	5	61	49.3
	6	27	17.8
	7+	12	7.3
	1096	1096	

$$\implies d_1 = \dots = 25.98.$$

Sol. 1) Let  $X_i$  be the number of death in  $i$ th day,  $1 \leq i \leq 1096$ .

2)  $H_0 : X_i$  follow  $\text{Poisson}(\lambda)$ .

3) The MLE for  $\lambda$  is:  $\lambda_e = \dots = 2.157$ .

4) Compute the expected frequencies:

Table 10.4.2		
Number of Deaths, $i$	Obs. Freq., $k_i$	Est. Exp. Freq., $n \hat{p}_{i_0}$
0	162	126.8
1	267	273.5
2	271	294.9
3	185	212.1
4	111	114.3
5	61	49.3
6	27	17.8
7	8	5.5
8	3	1.4
9	1	0.3
10+	0	0.1
	1096	1096

Table 10.4.3		
Number of Deaths, $i$	Obs. Freq., $k_i$	Est. Exp. Freq., $n \hat{p}_{i_0}$
$r_1, r_2, \dots, r_8$	0	126.8
	1	273.5
	2	294.9
	3	212.1
	4	114.3
	5	49.3
	6	17.8
	7+	7.3
		1096

$$\implies d_1 = \dots = 25.98.$$

5)  $P\text{-value} = \mathbb{P}(\chi_{1,8-1-1}^2 \geq 25.98) = 0.00022$ . Reject!

□