

Math 362: Mathematical Statistics II

Le Chen

le.chen@emory.edu

Emory University

Atlanta, GA

Last updated on December 20, 2020

2021 Spring

Chapter 5: Estimation

Chapter 5. Estimation

§ 5.1 Introduction

§ 5.2 Estimating parameters: MLE and MME

§ 5.1 Introduction

Motivating example: Given an unfair coin, or p -coin, such that

$$X = \begin{cases} 1 & \text{head with probability } p, \\ 0 & \text{tail with probability } 1 - p, \end{cases}$$

how would you determine the value p ?


Solutions:

1. You need to try the coin several times, say, three times. What you obtain is “HHT”.
2. Draw a conclusion from the experiment you just made.

Rationale: The choice of the parameter p should be the value that maximizes the probability of the sample.

$$\begin{aligned}\mathbb{P}(X_1 = 1, X_2 = 1, X_3 = 0) &= P(X_1 = 1)P(X_2 = 1)P(X_3 = 0) \\ &= p^2(1 - p).\end{aligned}$$

```
1 # Hello, R.  
2 p <- seq(0,1,0.01)  
3 plot(p,p^2*(1-p),  
4     type="l",  
5     col="red")  
6 title("Likelihood")  
7 # add a vertical dotted (4) blue  
  line  
8 abline(v=0.67, col="blue", lty=4)  
9 # add some text  
10 text(0.67,0.01, "2/3")
```



Maximize $f(p) = p^2(1 - p) \dots$

A random sample of size n from the population – Bernoulli(p):

- ▶ X_1, \dots, X_n are i.i.d.¹ random variables, each following Bernoulli(p).
- ▶ Suppose the outcomes of the random sample are: $X_1 = k_1, \dots, X_n = k_n$.
- ▶ What is your choice of p based on the above random sample?

$$p = \frac{1}{n} \sum_{i=1}^n k_i =: \bar{k}.$$

¹independent and identically distributed

A random sample of size n from the population with given pdf:

- ▶ X_1, \dots, X_n are i.i.d. random variables, each following the same given pdf.
- ▶ a **statistic** or an **estimator** is a function of the random sample.

Statistic/Estimator is a random variable!

e.g.,

$$\hat{p} = \frac{1}{n} \sum_{i=1}^n X_i.$$

- ▶ The outcome of a statistic/estimator is called an **estimate**. e.g.,

$$p_e = \frac{1}{n} \sum_{i=1}^n k_i.$$

Chapter 5. Estimation

§ 5.1 Introduction

§ 5.2 Estimating parameters: MLE and MME

§ 5.2 Estimating parameters: MLE and MME

Two methods:

Corresponding estimator

1. Method of maximum likelihood.

MLE

2. Method of moments.

MME

Maximum Likelihood Estimation

Definition 5.2.1. For a random sample of size n from the discrete (resp. continuous) population/pdf $p_X(k; \theta)$ (resp. $f_Y(y; \theta)$), the **likelihood function**, $L(\theta)$, is the product of the pdf evaluated at $X_i = k_i$ (resp. $Y_i = y_i$), i.e.,

$$L(\theta) = \prod_{i=1}^n p_X(k_i; \theta) \quad \left(\text{resp. } L(\theta) = \prod_{i=1}^n f_Y(y_i; \theta) \right).$$

Definition 5.2.2. Let $L(\theta)$ be as defined in Definition 5.2.1. If θ_e is a value of the parameter such that $L(\theta_e) \geq L(\theta)$ for all possible values of θ , then we call θ_e the **maximum likelihood estimate** for θ .

Examples for MLE

Often but not always MLE can be obtained by setting the first derivative equal to zero:

E.g. 1. Poisson distribution: $p_X(k) = e^{-\lambda} \frac{\lambda^k}{k!}$, $k = 0, 1, \dots$.

$$L(\lambda) = \prod_{i=1}^n e^{-\lambda} \frac{\lambda^{k_i}}{k_i!} = e^{-n\lambda} \lambda^{\sum_{i=1}^n k_i} \left(\prod_{i=1}^n k_i! \right)^{-1}.$$

$$\ln L(\lambda) = -n\lambda + \left(\sum_{i=1}^n k_i \right) \ln \lambda - \ln \left(\prod_{i=1}^n k_i! \right).$$

$$\frac{d}{d\lambda} \ln L(\lambda) = -n + \frac{1}{\lambda} \sum_{i=1}^n k_i.$$

$$\frac{d}{d\lambda} \ln L(\lambda) = 0 \quad \Longrightarrow \quad \boxed{\lambda_e = \frac{1}{n} \sum_{i=1}^n k_i =: \bar{k}}.$$

Comment: The critical point is indeed global maximum because

$$\frac{d^2}{d\lambda^2} \ln L(\lambda) = -\frac{1}{\lambda^2} \sum_{i=1}^n k_i < 0.$$

The following two cases are related to waiting time:

E.g. 2. Exponential distribution: $f_Y(y) = \lambda e^{-\lambda y}$ for $y \geq 0$.

$$L(\lambda) = \prod_{i=1}^n \lambda e^{-\lambda y_i} = \lambda^n \exp \left(-\lambda \sum_{i=1}^n y_i \right)$$

$$\ln L(\lambda) = n \ln \lambda - \lambda \sum_{i=1}^n y_i.$$

$$\frac{d}{d\lambda} \ln L(\lambda) = \frac{n}{\lambda} - \sum_{i=1}^n y_i.$$

$$\frac{d}{d\lambda} \ln L(\lambda) = 0 \quad \Longrightarrow \quad \boxed{\lambda_e = \frac{n}{\sum_{i=1}^n y_i} =: \frac{1}{\bar{y}}}.$$

A random sample of size n from the following population:

E.g. 3. Gamma distribution: $f_Y(y; \lambda) = \frac{\lambda^r}{\Gamma(r)} y^{r-1} e^{-\lambda y}$ for $y \geq 0$ with $r > 1$ known.

$$L(\lambda) = \prod_{i=1}^n \frac{\lambda^r}{\Gamma(r)} y_i^{r-1} e^{-\lambda y_i} = \lambda^{r n} \Gamma(r)^{-n} \left(\prod_{i=1}^n y_i^{r-1} \right) \exp \left(-\lambda \sum_{i=1}^n y_i \right)$$

$$\ln L(\lambda) = r n \ln \lambda - n \ln \Gamma(r) + \ln \left(\prod_{i=1}^n y_i^{r-1} \right) - \lambda \sum_{i=1}^n y_i.$$

$$\frac{d}{d\lambda} \ln L(\lambda) = \frac{r n}{\lambda} - \sum_{i=1}^n y_i.$$

$$\frac{d}{d\lambda} \ln L(\lambda) = 0 \quad \Longrightarrow \quad \boxed{\lambda_e = \frac{r n}{\sum_{i=1}^n y_i} = \frac{r}{\bar{y}}}.$$

Comment:

- When $r = 1$, this reduces to the exponential distribution case.
- If r is also unknown, it will be much more complicated.
No closed-form solution. One needs numerical solver².
Try MME instead.

²[DW, Example 7.2.25]

A detailed study with data:

E.g. 4. Geometric distribution: $p_X(k; p) = (1 - p)^{k-1}p$, $k = 1, 2, \dots$.

$$L(p) = \prod_{i=1}^n (1 - p)^{k_i-1} p = (1 - p)^{-n + \sum_{i=1}^n k_i} p^n.$$

$$\ln L(p) = \left(-n + \sum_{i=1}^n k_i \right) \ln(1 - p) + n \ln p.$$

$$\frac{d}{dp} \ln L(p) = -\frac{-n + \sum_{i=1}^n k_i}{1 - p} + \frac{n}{p}.$$

$$\frac{d}{dp} \ln L(p) = 0 \quad \Longrightarrow \quad \boxed{p_e = \frac{n}{\sum_{i=1}^n k_i} = \frac{1}{\bar{k}}}.$$

Comment: Its cousin distribution, the negative binomial distribution can be worked out similarly (See Ex 5.2.14).

MLE for Geom. Distr. of sample size $n = 128$

k	Observed frequency	Predicted frequency
1	72	74.14
2	35	31.2
3	11	13.13
4	6	5.52
5	2	2.32
6	2	0.98

Real p = unknown and MLE for $p = 0.5792$

```

1 # The example from the book.
2 library(pracma) # Load the library "Practical Numerical Math Functions"
3 k<-c(72, 35, 11, 6, 2, 2) # observed freq.
4 a=1:6
5 pe=sum(k)/dot(k,a) # MLE for p.
6 f=a
7 for (i in 1:6) {
8   f[i] = round((1-pe)^(i-1) * pe * sum(k),2)
9 }
10 # Initialize the table
11 d <-matrix(1:18, nrow = 6, ncol = 3)
12 # Now adding the column names
13 colnames(d) <- c("k",
14                  "Observed freq.",
15                  "Predicted freq.")
16 d[1:6,1]<-a
17 d[1:6,2]<-k
18 d[1:6,3]<-f
19 grid.table(d) # Show the table
20 PlotResults("unknown", pe, d, "Geometric.pdf") # Output the results using a user
    defined function

```


MLE for Geom. Distr. of sample size $n = 128$

k	Observed frequency	Predicted frequency
1	42	40.96
2	31	27.85
3	15	18.94
4	11	12.88
5	9	8.76
6	5	5.96
7	7	4.05
8	2	2.75
9	1	1.87
10	2	1.27
11	1	0.87
13	1	0.59
14	1	0.4

Real $p = 0.3333$ and MLE for $p = 0.32$

```

1 # Now let's generate random samples from a Geometric distribution with  $p=1/3$  with
   the same size of the sample.
2 p = 1/3
3 n = 128
4 gdata<-rgeom(n, p)+1 # Generate random samples
5 g<- table(gdata) # Count frequency of your data.
6 g<- t(rbind(as.numeric(rownames(g)), g)) # Transpose and combine two columns.
7 pe=n/dot(g[,1],g[,2]) # MLE for p.
8 f <- g[,1] # Initialize f
9 for (i in 1:nrow(g)) {
10   f[i] = round((1-pe)^(i-1) * pe * n,2)
11 } # Compute the expected frequency
12 g<-cbind(g,f) # Add one columns to your matrix.
13 colnames(g) <- c("k",
14                 "Observed freq.",
15                 "Predicted freq.") # Specify the column names.
16 d_df <- as.data.frame(d) # One can use data frame to store data
17 d_df # Show data on your terminal
18 PlotResults(p, pe, g, "Geometric2.pdf") # Output the results using a user defined
   function

```

MLE for Geom. Distr. of sample size $n = 300$

k	Observed frequency	Predicted frequency
1	99	105.88
2	69	68.51
3	47	44.33
4	28	28.69
5	27	18.56
6	9	12.01
7	8	7.77
8	5	5.03
9	5	3.25
10	3	2.11

Real $p = 0.3333$ and MLE for $p = 0.3529$

In case we have several parameters:

E.g. 5. Normal distribution: $f_Y(y; \mu, \sigma^2) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(y-\mu)^2}{2\sigma^2}}$, $y \in \mathbb{R}$.

$$L(\mu, \sigma^2) = \prod_{i=1}^n \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(y_i-\mu)^2}{2\sigma^2}} = (2\pi\sigma^2)^{-n/2} \exp\left(-\frac{1}{2\sigma^2} \sum_{i=1}^n (y_i - \mu)^2\right)$$

$$\ln L(\mu, \sigma^2) = -\frac{n}{2} \ln(2\pi\sigma^2) - \frac{1}{2\sigma^2} \sum_{i=1}^n (y_i - \mu)^2.$$

$$\begin{cases} \frac{\partial}{\partial \mu} \ln L(\mu, \sigma^2) = \frac{1}{\sigma^2} \sum_{i=1}^n (y_i - \mu) \\ \frac{\partial}{\partial \sigma^2} \ln L(\mu, \sigma^2) = -\frac{n}{2\sigma^2} + \frac{1}{2\sigma^4} \sum_{i=1}^n (y_i - \mu)^2 \end{cases}$$

$$\begin{cases} \frac{\partial}{\partial \mu} \ln L(\mu, \sigma^2) = 0 \\ \frac{\partial}{\partial \sigma^2} \ln L(\mu, \sigma^2) = 0 \end{cases}$$

\Rightarrow

$$\begin{cases} \mu_e = \bar{y} \\ \sigma_e^2 = \frac{1}{n} \sum_{i=1}^n (y_i - \bar{y})^2 \end{cases}$$

In case when the parameters determine the support of the density:
(Non regular case)

E.g. 6. Uniform distribution on $[a, b]$ with $a < b$: $f_Y(y; a, b) = \frac{1}{b-a}$ if $y \in [a, b]$.

$$L(a, b) = \begin{cases} \prod_{i=1}^n \frac{1}{b-a} = \frac{1}{(b-a)^n} & \text{if } a \leq y_1, \dots, y_n \leq b, \\ 0 & \text{otherwise.} \end{cases}$$

$L(a, b)$ is monotone increasing in a and decreasing in b . Hence, in order to maximize $L(a, b)$, one needs to choose

$$a_e = y_{\min} \quad \text{and} \quad b_e = y_{\max}.$$

E.g. 7. $f_Y(y; \theta) = \frac{2y}{\theta^2}$ for $y \in [0, \theta]$.

$$L(\theta) = \begin{cases} \prod_{i=1}^n \frac{2y_i}{\theta^2} = 2^n \theta^{-2n} \prod_{i=1}^n y_i & \text{if } 0 \leq y_1, \dots, y_n \leq \theta, \\ 0 & \text{otherwise.} \end{cases}$$

\Downarrow

$$\theta_e = y_{\max}.$$

In case of discrete parameter:

E.g. 8. Wildlife sampling. Capture-tag-recapture.... In the history, a tags have been put. In order to estimate the population size N , one randomly captures n animals, and there are k tagged. Find the MLE for N .


Sol. The population follows hypergeometric distr.:

$$p_X(k; N) = \frac{\binom{a}{k} \binom{N-a}{n-k}}{\binom{N}{n}}.$$

$$L(N) = \frac{\binom{a}{k} \binom{N-a}{n-k}}{\binom{N}{n}}$$

How to maximize $L(N)$?

```
1 > a=10
2 > k=5
3 > n=20
4 > N=seq(a,a+100)
5 > p=choose(a,k)*choose(N-a,n-k
6 > plot(N,p,type = "p")
7 > print(paste("The MLE is", n*a/
8 [1] "The MLE is 40"
```



The graph suggests to study the following quantity:

$$r(N) := \frac{L(N)}{L(N-1)} = \frac{N-n}{N} \times \frac{N-a}{N-a-n+k}$$

$$r(N) < 1 \quad \Longleftrightarrow \quad na < Nk \quad \text{i.e., } N > \frac{na}{k}$$

$$N_e = \arg \max \left\{ L(N) : N = \left\lfloor \frac{na}{k} \right\rfloor, \left\lceil \frac{na}{k} \right\rceil \right\}.$$

□

Method of Moments Estimation

Rationale: The population moments should be close to the sample moments, i.e.,

$$E(Y^k) \approx \frac{1}{n} \sum_{i=1}^n y_i^k, \quad k = 1, 2, 3, \dots$$

Definition 5.2.3. For a random sample of size n from the discrete (resp. continuous) population/pdf $p_X(k; \theta_1, \dots, \theta_s)$ (resp. $f_Y(y; \theta_1, \dots, \theta_s)$), solutions to

$$\begin{cases} E(Y) = \frac{1}{n} \sum_{i=1}^n y_i \\ \vdots \\ E(Y^s) = \frac{1}{n} \sum_{i=1}^n y_i^s \end{cases}$$

which are denoted by $\theta_{1e}, \dots, \theta_{se}$, are called the **method of moments estimates** of $\theta_1, \dots, \theta_s$.

Examples for MME

MME is often the same as MLE:

E.g. 1. Normal distribution: $f_Y(y; \mu, \sigma^2) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(y-\mu)^2}{2\sigma^2}}$, $y \in \mathbb{R}$.

$$\left\{ \begin{array}{l} \mu = \mathbb{E}(Y) = \frac{1}{n} \sum_{i=1}^n y_i = \bar{y} \\ \sigma^2 + \mu^2 = \mathbb{E}(Y^2) = \frac{1}{n} \sum_{i=1}^n y_i^2 \end{array} \right. \Rightarrow \left\{ \begin{array}{l} \mu_e = \bar{y} \\ \sigma_e^2 = \frac{1}{n} \sum_{i=1}^n y_i^2 - \mu_e^2 \\ \quad = \frac{1}{n} \sum_{i=1}^n (y_i - \bar{y})^2 \end{array} \right.$$

More examples when MLE coincides with MME: Poisson, Exponential, Geometric.

MME is often much more tractable than MLE:

E.g. 2. Gamma distribution³: $f_Y(y; r, \lambda) = \frac{\lambda^r}{\Gamma(r)} y^{r-1} e^{-\lambda y}$ for $y \geq 0$.

$$\begin{cases} \frac{r}{\lambda} = \mathbb{E}(Y) = \frac{1}{n} \sum_{i=1}^n y_i = \bar{y} \\ \frac{r}{\lambda^2} + \frac{r^2}{\lambda^2} = \mathbb{E}(Y^2) = \frac{1}{n} \sum_{i=1}^n y_i^2 \end{cases} \Rightarrow \begin{cases} r_e = \frac{\bar{y}^2}{\hat{\sigma}^2} \\ \lambda_e = \frac{\bar{y}}{\hat{\sigma}^2} = \frac{r_e}{\bar{y}} \end{cases}$$

where \bar{y} is the sample mean and $\hat{\sigma}^2$ is the sample variance:
 $\hat{\sigma}^2 := \frac{1}{n} \sum_{i=1}^n (y_i - \bar{y})^2$.

Comments: MME for λ is consistent with MLE when r is known.

³Check Theorem 4.6.3 on p. 269 for mean and variance

Another tractable example for MME, while less tractable for MLE:

E.g. 3. Neg. binomial distribution: $p_X(k; p, r) = \binom{k+r-1}{k} (1-p)^k p^r$,
 $k = 0, 1, \dots$.


$$\begin{cases} \frac{r(1-p)}{p} = \mathbb{E}(X) = \bar{k} \\ \frac{r(1-p)}{p^2} = \text{Var}(X) = \hat{\sigma}^2 \end{cases} \Rightarrow \begin{cases} p_e = \frac{\bar{k}}{\hat{\sigma}^2} \\ r_e = \frac{\bar{k}^2}{\hat{\sigma}^2 - \bar{k}} \end{cases}$$

(Case Study 5.2.2 continued)

Table 5.2.4

Number	Observed Frequency	Expected Frequency
0-5	0	0
6-10	10	7.7
11-15	20	21.4
16-20	23	28.4
21-25	27	22.4
26-30	12	12.3
31-35	5	5.3
36-40	2	1.8
> 40	1	0.7

Data from: <http://www.seattlecentral.edu/qelp/sets/039/039.html>



$$r_e = 12.74 \text{ and } p_e = 0.391.$$

E.g. 4. $f_Y(y; \theta) = \frac{2y}{\theta^2}$ for $y \in [0, \theta]$.

$$\bar{y} = \mathbb{E}[Y] = \int_0^\theta \frac{2y^2}{\theta^2} dy = \frac{2}{3} \frac{y^3}{\theta^2} \Big|_{y=0}^{y=\theta} = \frac{2}{3} \theta.$$

\Downarrow

$$\boxed{\theta_e = \frac{3}{2} \bar{y}.$$