

# Math 362: Mathematical Statistics II

Le Chen

le.chen@emory.edu  
chenle02@gmail.com

Emory University  
Atlanta, GA

Last updated on Spring 2021  
Last compiled on January 15, 2023

2021 Spring

Creative Commons License  
(CC By-NC-SA)

# Chapter 5. Estimation

§ 5.1 Introduction

§ 5.2 Estimating parameters: MLE and MME

§ 5.3 Interval Estimation

§ 5.4 Properties of Estimators

§ 5.5 Minimum-Variance Estimators: The Cramér-Rao Lower Bound

§ 5.6 Sufficient Estimators

§ 5.7 Consistency

§ 5.8 Bayesian Estimation

# Plan

§ 5.1 Introduction

§ 5.2 Estimating parameters: MLE and MME

§ 5.3 Interval Estimation

**§ 5.4 Properties of Estimators**

§ 5.5 Minimum-Variance Estimators: The Cramér-Rao Lower Bound

§ 5.6 Sufficient Estimators

§ 5.7 Consistency

§ 5.8 Bayesian Estimation

# Chapter 5. Estimation

§ 5.1 Introduction

§ 5.2 Estimating parameters: MLE and MME

§ 5.3 Interval Estimation

**§ 5.4 Properties of Estimators**

§ 5.5 Minimum-Variance Estimators: The Cramér-Rao Lower Bound

§ 5.6 Sufficient Estimators

§ 5.7 Consistency

§ 5.8 Bayesian Estimation

## § 5.4 Properties of Estimators

**Question:** Estimators are not in general unique (MLE or MME ...). How to select one estimator?

Recall: For a random sample of size  $n$  from the population with given pdf, we have  $X_1, \dots, X_n$ , which are i.i.d. r.v.'s. The estimator  $\hat{\theta}$  is a function of  $X_i$ 's:

$$\hat{\theta} = \hat{\theta}(X_1, \dots, X_n).$$

### Criteria:

1. Unbiased. (Mean)
2. Efficiency, the minimum-variance estimator. (Variance)
3. Sufficiency.
4. Consistency. (Asymptotic behavior)

## § 5.4 Properties of Estimators

**Question:** Estimators are not in general unique (MLE or MME ...). How to select one estimator?

Recall: For a random sample of size  $n$  from the population with given pdf, we have  $X_1, \dots, X_n$ , which are i.i.d. r.v.'s. The estimator  $\hat{\theta}$  is a function of  $X_i$ 's:

$$\hat{\theta} = \hat{\theta}(X_1, \dots, X_n).$$

### Criteria:

1. Unbiased. (Mean)
2. Efficiency, the minimum-variance estimator. (Variance)
3. Sufficiency.
4. Consistency. (Asymptotic behavior)

## § 5.4 Properties of Estimators

**Question:** Estimators are not in general unique (MLE or MME ...). How to select one estimator?

Recall: For a random sample of size  $n$  from the population with given pdf, we have  $X_1, \dots, X_n$ , which are i.i.d. r.v.'s. The estimator  $\hat{\theta}$  is a function of  $X_i$ 's:

$$\hat{\theta} = \hat{\theta}(X_1, \dots, X_n).$$

### Criteria:

1. Unbiased. (Mean)
2. Efficiency, the minimum-variance estimator. (Variance)
3. Sufficiency.
4. Consistency. (Asymptotic behavior)

## § 5.4 Properties of Estimators

**Question:** Estimators are not in general unique (MLE or MME ...). How to select one estimator?

Recall: For a random sample of size  $n$  from the population with given pdf, we have  $X_1, \dots, X_n$ , which are i.i.d. r.v.'s. The estimator  $\hat{\theta}$  is a function of  $X_i$ 's:



$$\hat{\theta} = \hat{\theta}(X_1, \dots, X_n).$$

### Criteria:

1. Unbiased. (Mean)
2. Efficiency, the minimum-variance estimator. (Variance)
3. Sufficiency.
4. Consistency. (Asymptotic behavior)



# Unbiasedness



**Definition 5.4.1.** Given a random sample of size  $n$  whose population distribution depends on an unknown parameter  $\theta$ , let  $\hat{\theta}$  be an estimator of  $\theta$ .

Then  $\hat{\theta}$  is called **unbiased** if  $\mathbb{E}(\hat{\theta}) = \theta$ ;

and  $\hat{\theta}$  is called **asymptotically unbiased** if  $\lim_{n \rightarrow \infty} \mathbb{E}(\hat{\theta}) = \theta$ .

E.g. 1.  $f_Y(y; \theta) = \frac{2y}{\theta^2}$  if  $y \in [0, \theta]$ .

$$- \hat{\theta}_1 = \frac{3}{2} \bar{Y}$$

$$- \hat{\theta}_2 = Y_{\max}.$$

$$- \hat{\theta}_3 = \frac{2n+1}{2n} Y_{\max}.$$

E.g. 2. Let  $X_1, \dots, X_n$  be a random sample of size  $n$  with the unknown parameter  $\theta = \mathbb{E}(X)$ . Show that for any constants  $a_i$ 's,

$$\hat{\theta} = \sum_{i=1}^n a_i X_i \quad \text{is unbiased} \quad \Longleftrightarrow \quad \sum_{i=1}^n a_i = 1.$$

E.g. 1.  $f_Y(y; \theta) = \frac{2y}{\theta^2}$  if  $y \in [0, \theta]$ .

$$- \hat{\theta}_1 = \frac{3}{2} \bar{Y}$$

$$- \hat{\theta}_2 = Y_{\max}.$$

$$- \hat{\theta}_3 = \frac{2n+1}{2n} Y_{\max}.$$

E.g. 2. Let  $X_1, \dots, X_n$  be a random sample of size  $n$  with the unknown parameter  $\theta = \mathbb{E}(X)$ . Show that for any constants  $a_i$ 's,

$$\hat{\theta} = \sum_{i=1}^n a_i X_i \quad \text{is unbiased} \quad \Longleftrightarrow \quad \sum_{i=1}^n a_i = 1.$$

E.g. 1.  $f_Y(y; \theta) = \frac{2y}{\theta^2}$  if  $y \in [0, \theta]$ .

$$- \hat{\theta}_1 = \frac{3}{2} \bar{Y}$$

$$- \hat{\theta}_2 = Y_{\max}.$$

$$- \hat{\theta}_3 = \frac{2n+1}{2n} Y_{\max}.$$

E.g. 2. Let  $X_1, \dots, X_n$  be a random sample of size  $n$  with the unknown parameter  $\theta = \mathbb{E}(X)$ . Show that for any constants  $a_i$ 's,

$$\hat{\theta} = \sum_{i=1}^n a_i X_i \quad \text{is unbiased} \quad \Longleftrightarrow \quad \sum_{i=1}^n a_i = 1.$$

E.g. 1.  $f_Y(y; \theta) = \frac{2y}{\theta^2}$  if  $y \in [0, \theta]$ .

$$- \hat{\theta}_1 = \frac{3}{2} \bar{Y}$$

$$- \hat{\theta}_2 = Y_{\max}.$$

$$- \hat{\theta}_3 = \frac{2n+1}{2n} Y_{\max}.$$

E.g. 2. Let  $X_1, \dots, X_n$  be a random sample of size  $n$  with the unknown parameter  $\theta = \mathbb{E}(X)$ . Show that for any constants  $a_i$ 's,

$$\hat{\theta} = \sum_{i=1}^n a_i X_i \text{ is unbiased} \iff \sum_{i=1}^n a_i = 1.$$

E.g. 1.  $f_Y(y; \theta) = \frac{2y}{\theta^2}$  if  $y \in [0, \theta]$ .

$$- \hat{\theta}_1 = \frac{3}{2} \bar{Y}$$

$$- \hat{\theta}_2 = Y_{\max}.$$

$$- \hat{\theta}_3 = \frac{2n+1}{2n} Y_{\max}.$$

E.g. 2. Let  $X_1, \dots, X_n$  be a random sample of size  $n$  with the unknown parameter  $\theta = \mathbb{E}(X)$ . Show that for any constants  $a_i$ 's,

$$\hat{\theta} = \sum_{i=1}^n a_i X_i \quad \text{is unbiased} \quad \Longleftrightarrow \quad \sum_{i=1}^n a_i = 1.$$

E.g. 3. Let  $X_1, \dots, X_n$  be a random sample of size  $n$  with the unknown parameter  $\sigma^2 = \text{Var}(X)$ .

$$- \hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2$$

$$- S^2 = \text{Sample Variance} = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2$$

$$- S = \text{Sample Standard Deviation} = \sqrt{\frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2}. \quad (\text{Biased for } \sigma!)$$

E.g. 3. Let  $X_1, \dots, X_n$  be a random sample of size  $n$  with the unknown parameter  $\sigma^2 = \text{Var}(X)$ .

$$- \hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n (x_i - \bar{X})^2$$

$$- S^2 = \text{Sample Variance} = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2$$

$$- S = \text{Sample Standard Deviation} = \sqrt{\frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2}. \quad (\text{Biased for } \sigma!)$$



E.g. 3. Let  $X_1, \dots, X_n$  be a random sample of size  $n$  with the unknown parameter  $\sigma^2 = \text{Var}(X)$ .

$$- \hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2$$

$$- S^2 = \text{Sample Variance} = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2$$

$$- S = \text{Sample Standard Deviation} = \sqrt{\frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2}. \quad (\text{Biased for } \sigma!)$$

E.g. 3. Let  $X_1, \dots, X_n$  be a random sample of size  $n$  with the unknown parameter  $\sigma^2 = \text{Var}(X)$ .

$$- \hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2$$

$$- S^2 = \text{Sample Variance} = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2$$

$$- S = \text{Sample Standard Deviation} = \sqrt{\frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2}. \quad (\text{Biased for } \sigma!)$$

E.g. 4. Exponential distr.:  $f_Y(y; \lambda) = \lambda e^{-\lambda y}$  for  $y \geq 0$ .  $\hat{\lambda} = 1/\bar{Y}$  is biased.

$n\bar{Y} = \sum_{i=1}^n Y_i \sim \text{Gamma distribution}(n, \lambda)$ . Hence,

$$\begin{aligned}\mathbb{E}(\hat{\lambda}) &= \mathbb{E}(1/\bar{Y}) = n \int_0^\infty \frac{1}{y} \frac{\lambda^n}{\Gamma(n)} y^{n-1} e^{-\lambda y} dy \\ &= \frac{n\lambda}{n-1} \int_0^\infty \underbrace{\frac{\lambda^{n-1}}{\Gamma(n-1)} y^{(n-1)-1} e^{-\lambda y}}_{\text{pdf for Gamma distr. } (n-1, \lambda)} dy \\ &= \frac{n}{n-1} \lambda.\end{aligned}$$

Biased! But  $\mathbb{E}(\hat{\lambda}) = \frac{n}{n-1} \lambda \rightarrow \lambda$  as  $n \rightarrow \infty$ . (Asymptotically unbiased.)

Note:  $\hat{\lambda}^* = \frac{n-1}{n\bar{Y}}$  is unbiased.

E.g. 4'. Exponential distr.:  $f_Y(y; \theta) = \frac{1}{\theta} e^{-y/\theta}$  for  $y \geq 0$ .  $\hat{\theta} = \bar{Y}$  is unbiased.

$$\mathbb{E}(\hat{\theta}) = \frac{1}{n} \sum_{i=1}^n \mathbb{E}(Y_i) = \frac{1}{n} \sum_{i=1}^n \theta = \theta$$

E.g. 4. Exponential distr.:  $f_Y(y; \lambda) = \lambda e^{-\lambda y}$  for  $y \geq 0$ .  $\hat{\lambda} = 1/\bar{Y}$  is biased.

$n\bar{Y} = \sum_{i=1}^n Y_i \sim \text{Gamma distribution}(n, \lambda)$ . Hence,

$$\begin{aligned}\mathbb{E}(\hat{\lambda}) &= \mathbb{E}(1/\bar{Y}) = n \int_0^\infty \frac{1}{y} \frac{\lambda^n}{\Gamma(n)} y^{n-1} e^{-\lambda y} dy \\ &= \frac{n\lambda}{n-1} \int_0^\infty \underbrace{\frac{\lambda^{n-1}}{\Gamma(n-1)} y^{(n-1)-1} e^{-\lambda y}}_{\text{pdf for Gamma distr. } (n-1, \lambda)} dy \\ &= \frac{n}{n-1} \lambda.\end{aligned}$$

Biased! But  $\mathbb{E}(\hat{\lambda}) = \frac{n}{n-1} \lambda \rightarrow \lambda$  as  $n \rightarrow \infty$ . (Asymptotically unbiased.)

Note:  $\hat{\lambda}^* = \frac{n-1}{n\bar{Y}}$  is unbiased.

E.g. 4'. Exponential distr.:  $f_Y(y; \theta) = \frac{1}{\theta} e^{-y/\theta}$  for  $y \geq 0$ .  $\hat{\theta} = \bar{Y}$  is unbiased.

$$\mathbb{E}(\hat{\theta}) = \frac{1}{n} \sum_{i=1}^n \mathbb{E}(Y_i) = \frac{1}{n} \sum_{i=1}^n \theta = \theta$$

E.g. 4. Exponential distr.:  $f_Y(y; \lambda) = \lambda e^{-\lambda y}$  for  $y \geq 0$ .  $\hat{\lambda} = 1/\bar{Y}$  is biased.

$n\bar{Y} = \sum_{i=1}^n Y_i \sim \text{Gamma distribution}(n, \lambda)$ . Hence,

$$\begin{aligned}\mathbb{E}(\hat{\lambda}) &= \mathbb{E}(1/\bar{Y}) = n \int_0^\infty \frac{1}{y} \frac{\lambda^n}{\Gamma(n)} y^{n-1} e^{-\lambda y} dy \\ &= \frac{n\lambda}{n-1} \int_0^\infty \underbrace{\frac{\lambda^{n-1}}{\Gamma(n-1)} y^{(n-1)-1} e^{-\lambda y}}_{\text{pdf for Gamma distr. } (n-1, \lambda)} dy \\ &= \frac{n}{n-1} \lambda.\end{aligned}$$

Biased! But  $\mathbb{E}(\hat{\lambda}) = \frac{n}{n-1} \lambda \rightarrow \lambda$  as  $n \rightarrow \infty$ . (Asymptotically unbiased.)

Note:  $\hat{\lambda}^* = \frac{n-1}{n\bar{Y}}$  is unbiased.

E.g. 4'. Exponential distr.:  $f_Y(y; \theta) = \frac{1}{\theta} e^{-y/\theta}$  for  $y \geq 0$ .  $\hat{\theta} = \bar{Y}$  is unbiased.

$$\mathbb{E}(\hat{\theta}) = \frac{1}{n} \sum_{i=1}^n \mathbb{E}(Y_i) = \frac{1}{n} \sum_{i=1}^n \theta = \theta$$

E.g. 4. Exponential distr.:  $f_Y(y; \lambda) = \lambda e^{-\lambda y}$  for  $y \geq 0$ .  $\hat{\lambda} = 1/\bar{Y}$  is biased.

$n\bar{Y} = \sum_{i=1}^n Y_i \sim \text{Gamma distribution}(n, \lambda)$ . Hence,

$$\begin{aligned}\mathbb{E}(\hat{\lambda}) &= \mathbb{E}(1/\bar{Y}) = n \int_0^\infty \frac{1}{y} \frac{\lambda^n}{\Gamma(n)} y^{n-1} e^{-\lambda y} dy \\ &= \frac{n\lambda}{n-1} \int_0^\infty \underbrace{\frac{\lambda^{n-1}}{\Gamma(n-1)} y^{(n-1)-1} e^{-\lambda y}}_{\text{pdf for Gamma distr. } (n-1, \lambda)} dy \\ &= \frac{n}{n-1} \lambda.\end{aligned}$$

Biased! But  $\mathbb{E}(\hat{\lambda}) = \frac{n}{n-1} \lambda \rightarrow \lambda$  as  $n \rightarrow \infty$ . (Asymptotically unbiased.)

Note:  $\hat{\lambda}^* = \frac{n-1}{n\bar{Y}}$  is unbiased.

E.g. 4'. Exponential distr.:  $f_Y(y; \theta) = \frac{1}{\theta} e^{-y/\theta}$  for  $y \geq 0$ .  $\hat{\theta} = \bar{Y}$  is unbiased.

$$\mathbb{E}(\bar{Y}) = \frac{1}{n} \sum_{i=1}^n \mathbb{E}(Y_i) = \frac{1}{n} \sum_{i=1}^n \theta = \theta$$

E.g. 4. Exponential distr.:  $f_Y(y; \lambda) = \lambda e^{-\lambda y}$  for  $y \geq 0$ .  $\hat{\lambda} = 1/\bar{Y}$  is biased.

$n\bar{Y} = \sum_{i=1}^n Y_i \sim \text{Gamma distribution}(n, \lambda)$ . Hence,

$$\begin{aligned}\mathbb{E}(\hat{\lambda}) &= \mathbb{E}\left(1/\bar{Y}\right) = n \int_0^\infty \frac{1}{y} \frac{\lambda^n}{\Gamma(n)} y^{n-1} e^{-\lambda y} dy \\ &= \frac{n\lambda}{n-1} \int_0^\infty \underbrace{\frac{\lambda^{n-1}}{\Gamma(n-1)} y^{(n-1)-1} e^{-\lambda y}}_{\text{pdf for Gamma distr. } (n-1, \lambda)} dy \\ &= \frac{n}{n-1} \lambda.\end{aligned}$$

Biased! But  $\mathbb{E}(\hat{\lambda}) = \frac{n}{n-1} \lambda \rightarrow \lambda$  as  $n \rightarrow \infty$ . (Asymptotically unbiased.)

Note:  $\hat{\lambda}^* = \frac{n-1}{n\bar{Y}}$  is unbiased.

E.g. 4'. Exponential distr.:  $f_Y(y; \theta) = \frac{1}{\theta} e^{-y/\theta}$  for  $y \geq 0$ .  $\hat{\theta} = \bar{Y}$  is unbiased.

$$\mathbb{E}(\bar{Y}) = \frac{1}{n} \sum_{i=1}^n \mathbb{E}(Y_i) = \frac{1}{n} \sum_{i=1}^n \theta = \theta$$

E.g. 4. Exponential distr.:  $f_Y(y; \lambda) = \lambda e^{-\lambda y}$  for  $y \geq 0$ .  $\hat{\lambda} = 1/\bar{Y}$  is biased.

$n\bar{Y} = \sum_{i=1}^n Y_i \sim \text{Gamma distribution}(n, \lambda)$ . Hence,

$$\begin{aligned}\mathbb{E}(\hat{\lambda}) &= \mathbb{E}\left(1/\bar{Y}\right) = n \int_0^\infty \frac{1}{y} \frac{\lambda^n}{\Gamma(n)} y^{n-1} e^{-\lambda y} dy \\ &= \frac{n\lambda}{n-1} \int_0^\infty \underbrace{\frac{\lambda^{n-1}}{\Gamma(n-1)} y^{(n-1)-1} e^{-\lambda y}}_{\text{pdf for Gamma distr. } (n-1, \lambda)} dy \\ &= \frac{n}{n-1} \lambda.\end{aligned}$$

Biased! But  $\mathbb{E}(\hat{\lambda}) = \frac{n}{n-1} \lambda \rightarrow \lambda$  as  $n \rightarrow \infty$ . (Asymptotically unbiased.)

Note:  $\hat{\lambda}^* = \frac{n-1}{n\bar{Y}}$  is unbiased.

E.g. 4'. Exponential distr.:  $f_Y(y; \lambda) = \lambda e^{-\lambda y}$  for  $y \geq 0$ .  $\hat{\lambda} = \lambda$  is unbiased.

$$\hat{\lambda} = \frac{1}{n} \sum_{i=1}^n \lambda = \frac{1}{n} \sum_{i=1}^n \lambda = \lambda$$



E.g. 4. Exponential distr.:  $f_Y(y; \lambda) = \lambda e^{-\lambda y}$  for  $y \geq 0$ .  $\hat{\lambda} = 1/\bar{Y}$  is biased.

$n\bar{Y} = \sum_{i=1}^n Y_i \sim \text{Gamma distribution}(n, \lambda)$ . Hence,

$$\begin{aligned}\mathbb{E}(\hat{\lambda}) &= \mathbb{E}\left(1/\bar{Y}\right) = n \int_0^\infty \frac{1}{y} \frac{\lambda^n}{\Gamma(n)} y^{n-1} e^{-\lambda y} dy \\ &= \frac{n\lambda}{n-1} \int_0^\infty \underbrace{\frac{\lambda^{n-1}}{\Gamma(n-1)} y^{(n-1)-1} e^{-\lambda y}}_{\text{pdf for Gamma distr. } (n-1, \lambda)} dy \\ &= \frac{n}{n-1} \lambda.\end{aligned}$$

Biased! But  $\mathbb{E}(\hat{\lambda}) = \frac{n}{n-1} \lambda \rightarrow \lambda$  as  $n \rightarrow \infty$ . (Asymptotically unbiased.)

Note:  $\hat{\lambda}^* = \frac{n-1}{n\bar{Y}}$  is unbiased.

E.g. 4'. Exponential distr.:  $f_Y(y; \lambda) = \lambda e^{-\lambda y}$  for  $y \geq 0$ .  $\hat{\lambda} = 1/\bar{Y}$  is unbiased.

$$\hat{\lambda} = \frac{1}{\bar{Y}} = \frac{1}{\frac{1}{n} \sum_{i=1}^n Y_i} = \frac{n}{\sum_{i=1}^n Y_i} = \frac{n}{n\bar{Y}} = \frac{1}{\bar{Y}}$$

E.g. 4. Exponential distr.:  $f_Y(y; \lambda) = \lambda e^{-\lambda y}$  for  $y \geq 0$ .  $\hat{\lambda} = 1/\bar{Y}$  is biased.

$n\bar{Y} = \sum_{i=1}^n Y_i \sim \text{Gamma distribution}(n, \lambda)$ . Hence,

$$\begin{aligned}\mathbb{E}(\hat{\lambda}) &= \mathbb{E}(1/\bar{Y}) = n \int_0^\infty \frac{1}{y} \frac{\lambda^n}{\Gamma(n)} y^{n-1} e^{-\lambda y} dy \\ &= \frac{n\lambda}{n-1} \int_0^\infty \underbrace{\frac{\lambda^{n-1}}{\Gamma(n-1)} y^{(n-1)-1} e^{-\lambda y}}_{\text{pdf for Gamma distr. } (n-1, \lambda)} dy \\ &= \frac{n}{n-1} \lambda.\end{aligned}$$

Biased! But  $\mathbb{E}(\hat{\lambda}) = \frac{n}{n-1} \lambda \rightarrow \lambda$  as  $n \rightarrow \infty$ . (Asymptotically unbiased.)

Note:  $\hat{\lambda}^* = \frac{n-1}{n\bar{Y}}$  is unbiased.

E.g. 4'. Exponential distr.:  $f_Y(y; \theta) = \frac{1}{\theta} e^{-y/\theta}$  for  $y \geq 0$ .  $\hat{\theta} = \bar{Y}$  is unbiased.

$$\mathbb{E}(\hat{\theta}) = \frac{1}{n} \sum_{i=1}^n \mathbb{E}(Y_i) = \frac{1}{n} \sum_{i=1}^n \theta = \theta.$$

E.g. 4. Exponential distr.:  $f_Y(y; \lambda) = \lambda e^{-\lambda y}$  for  $y \geq 0$ .  $\hat{\lambda} = 1/\bar{Y}$  is biased.

$n\bar{Y} = \sum_{i=1}^n Y_i \sim \text{Gamma distribution}(n, \lambda)$ . Hence,

$$\begin{aligned}\mathbb{E}(\hat{\lambda}) &= \mathbb{E}(1/\bar{Y}) = n \int_0^\infty \frac{1}{y} \frac{\lambda^n}{\Gamma(n)} y^{n-1} e^{-\lambda y} dy \\ &= \frac{n\lambda}{n-1} \int_0^\infty \underbrace{\frac{\lambda^{n-1}}{\Gamma(n-1)} y^{(n-1)-1} e^{-\lambda y}}_{\text{pdf for Gamma distr. } (n-1, \lambda)} dy \\ &= \frac{n}{n-1} \lambda.\end{aligned}$$

Biased! But  $\mathbb{E}(\hat{\lambda}) = \frac{n}{n-1} \lambda \rightarrow \lambda$  as  $n \rightarrow \infty$ . (Asymptotically unbiased.)

Note:  $\hat{\lambda}^* = \frac{n-1}{n\bar{Y}}$  is unbiased.

E.g. 4'. Exponential distr.:  $f_Y(y; \theta) = \frac{1}{\theta} e^{-y/\theta}$  for  $y \geq 0$ .  $\hat{\theta} = \bar{Y}$  is unbiased.

$$\mathbb{E}(\hat{\theta}) = \frac{1}{n} \sum_{i=1}^n \mathbb{E}(Y_i) = \frac{1}{n} \sum_{i=1}^n \theta = \theta.$$

E.g. 4. Exponential distr.:  $f_Y(y; \lambda) = \lambda e^{-\lambda y}$  for  $y \geq 0$ .  $\hat{\lambda} = 1/\bar{Y}$  is biased.

$n\bar{Y} = \sum_{i=1}^n Y_i \sim \text{Gamma distribution}(n, \lambda)$ . Hence,

$$\begin{aligned}\mathbb{E}(\hat{\lambda}) &= \mathbb{E}(1/\bar{Y}) = n \int_0^{\infty} \frac{1}{y} \frac{\lambda^n}{\Gamma(n)} y^{n-1} e^{-\lambda y} dy \\ &= \frac{n\lambda}{n-1} \int_0^{\infty} \underbrace{\frac{\lambda^{n-1}}{\Gamma(n-1)} y^{(n-1)-1} e^{-\lambda y}}_{\text{pdf for Gamma distr. } (n-1, \lambda)} dy \\ &= \frac{n}{n-1} \lambda.\end{aligned}$$


Biased! But  $\mathbb{E}(\hat{\lambda}) = \frac{n}{n-1} \lambda \rightarrow \lambda$  as  $n \rightarrow \infty$ . (Asymptotically unbiased.)

Note:  $\hat{\lambda}^* = \frac{n-1}{n\bar{Y}}$  is unbiased.

E.g. 4'. Exponential distr.:  $f_Y(y; \theta) = \frac{1}{\theta} e^{-y/\theta}$  for  $y \geq 0$ .  $\hat{\theta} = \bar{Y}$  is unbiased.

$$\mathbb{E}(\hat{\theta}) = \frac{1}{n} \sum_{i=1}^n \mathbb{E}(Y_i) = \frac{1}{n} \sum_{i=1}^n \theta = \theta.$$

# Efficiency



**Definition 5.4.2.** Let  $\hat{\theta}_1$  and  $\hat{\theta}_2$  be two unbiased estimators for a parameter  $\theta$ . If  $\text{Var}(\hat{\theta}_1) < \text{Var}(\hat{\theta}_2)$ , then we say that  $\hat{\theta}_1$  is **more efficient** than  $\hat{\theta}_2$ . The **relative efficiency** of  $\hat{\theta}_1$  w.r.t.  $\hat{\theta}_2$  is the ratio  $\text{Var}(\hat{\theta}_2)/\text{Var}(\hat{\theta}_1)$ .

E.g. 1.  $f_Y(y; \theta) = \frac{2y}{\theta^2}$  if  $y \in [0, \theta]$ . Which is more efficient? Find the relative efficiency of  $\hat{\theta}_1$  w.r.t.  $\hat{\theta}_3$ .

$$- \hat{\theta}_1 = \frac{3}{2} \bar{Y}$$

$$- \hat{\theta}_3 = \frac{2n+1}{2n} Y_{\max}.$$

E.g. 2. Let  $X_1, \dots, X_n$  be a random sample of size  $n$  with the unknown parameter  $\theta = \mathbb{E}(X)$  (suppose  $\sigma^2 = \text{Var}(X) < \infty$ ).

Among all possible unbiased estimators  $\theta = \sum_{i=1}^n a_i X_i$  with  $\sum_{i=1}^n a_i = 1$ , Find the most efficient one

Sol:

$$\text{Var}(\theta) = \sum_{i=1}^n a_i^2 \text{Var}(X_i) = \sigma^2 \sum_{i=1}^n a_i^2 \geq \sigma^2 \frac{1}{n} \left( \sum_{i=1}^n a_i \right)^2 = \frac{1}{n} \sigma^2$$

with equality iff  $a_1 = \dots = a_n = 1/n$ .

Hence, the most efficient one is the sample mean  $\bar{X} = X$ .

E.g. 1.  $f_Y(y; \theta) = \frac{2y}{\theta^2}$  if  $y \in [0, \theta]$ . Which is more efficient? Find the relative efficiency of  $\hat{\theta}_1$  w.r.t.  $\hat{\theta}_3$ .

$$- \hat{\theta}_1 = \frac{3}{2} \bar{Y}$$

$$- \hat{\theta}_3 = \frac{2n+1}{2n} Y_{\max}.$$

E.g. 2. Let  $X_1, \dots, X_n$  be a random sample of size  $n$  with the unknown parameter  $\theta = \mathbb{E}(X)$  (suppose  $\sigma^2 = \text{Var}(X) < \infty$ ).

Among all possible unbiased estimators  $\theta = \sum_{i=1}^n a_i X_i$  with  $\sum_{i=1}^n a_i = 1$ , Find the most efficient one

Sol:

$$\text{Var}(\theta) = \sum_{i=1}^n a_i^2 \text{Var}(X_i) = \sigma^2 \sum_{i=1}^n a_i^2 \geq \sigma^2 \frac{1}{n} \left( \sum_{i=1}^n a_i \right)^2 = \frac{1}{n} \sigma^2$$

with equality if  $a_1 = \dots = a_n = 1/n$ .

Hence, the most efficient one is the sample mean  $\bar{X} = \bar{X}_n$ .

E.g. 1.  $f_Y(y; \theta) = \frac{2y}{\theta^2}$  if  $y \in [0, \theta]$ . Which is more efficient? Find the relative efficiency of  $\hat{\theta}_1$  w.r.t.  $\hat{\theta}_3$ .

$$- \hat{\theta}_1 = \frac{3}{2} \bar{Y}$$

$$- \hat{\theta}_3 = \frac{2n+1}{2n} Y_{\max}.$$

E.g. 2. Let  $X_1, \dots, X_n$  be a random sample of size  $n$  with the unknown parameter  $\theta = \mathbb{E}(X)$  (suppose  $\sigma^2 = \text{Var}(X) < \infty$ ).

Among all possible unbiased estimators  $\theta = \sum_{i=1}^n a_i X_i$  with  $\sum_{i=1}^n a_i = 1$ , Find the most efficient one.

Sol:

$$\text{Var}(\theta) = \sum_{i=1}^n a_i^2 \text{Var}(X_i) = \sigma^2 \sum_{i=1}^n a_i^2 = \sigma^2 \frac{1}{n} \left( \sum_{i=1}^n a_i \right)^2 = \frac{1}{n} \sigma^2$$

with equality iff  $a_1 = \dots = a_n = 1/n$ .

Hence, the most efficient one is the sample mean  $\bar{X} = \bar{X}_n$ .



E.g. 1.  $f_Y(y; \theta) = \frac{2y}{\theta^2}$  if  $y \in [0, \theta]$ . Which is more efficient? Find the relative efficiency of  $\hat{\theta}_1$  w.r.t.  $\hat{\theta}_3$ .

$$- \hat{\theta}_1 = \frac{3}{2} \bar{Y}$$

$$- \hat{\theta}_3 = \frac{2n+1}{2n} Y_{\max}.$$

E.g. 2. Let  $X_1, \dots, X_n$  be a random sample of size  $n$  with the unknown parameter  $\theta = \mathbb{E}(X)$  (suppose  $\sigma^2 = \text{Var}(X) < \infty$ ).

Among all possible unbiased estimators  $\hat{\theta} = \sum_{i=1}^n a_i X_i$  with  $\sum_{i=1}^n a_i = 1$ . Find the most efficient one.

Sol:

$$\text{Var}(\hat{\theta}) = \sum_{i=1}^n a_i^2 \text{Var}(X) = \sigma^2 \sum_{i=1}^n a_i^2 \geq \sigma^2 \frac{1}{n} \left( \sum_{i=1}^n a_i \right)^2 = \frac{1}{n} \sigma^2,$$

with equality iff  $a_1 = \dots = a_n = 1/n$ .

Hence, the most efficient one is the sample mean  $\hat{\theta} = \bar{X}$ . □

E.g. 1.  $f_Y(y; \theta) = \frac{2y}{\theta^2}$  if  $y \in [0, \theta]$ . Which is more efficient? Find the relative efficiency of  $\hat{\theta}_1$  w.r.t.  $\hat{\theta}_3$ .

$$- \hat{\theta}_1 = \frac{3}{2} \bar{Y}$$

$$- \hat{\theta}_3 = \frac{2n+1}{2n} Y_{\max}.$$

E.g. 2. Let  $X_1, \dots, X_n$  be a random sample of size  $n$  with the unknown parameter  $\theta = \mathbb{E}(X)$  (suppose  $\sigma^2 = \text{Var}(X) < \infty$ ).

Among all possible unbiased estimators  $\hat{\theta} = \sum_{i=1}^n a_i X_i$  with  $\sum_{i=1}^n a_i = 1$ . Find the most efficient one.

Sol:

$$\text{Var}(\hat{\theta}) = \sum_{i=1}^n a_i^2 \text{Var}(X) = \sigma^2 \sum_{i=1}^n a_i^2 \geq \sigma^2 \frac{1}{n} \left( \sum_{i=1}^n a_i \right)^2 = \frac{1}{n} \sigma^2,$$

with equality iff  $a_1 = \dots = a_n = 1/n$ .

Hence, the most efficient one is the sample mean  $\hat{\theta} = \bar{X}$ . □

E.g. 1.  $f_Y(y; \theta) = \frac{2y}{\theta^2}$  if  $y \in [0, \theta]$ . Which is more efficient? Find the relative efficiency of  $\hat{\theta}_1$  w.r.t.  $\hat{\theta}_3$ .

$$- \hat{\theta}_1 = \frac{3}{2} \bar{Y}$$

$$- \hat{\theta}_3 = \frac{2n+1}{2n} Y_{\max}.$$

E.g. 2. Let  $X_1, \dots, X_n$  be a random sample of size  $n$  with the unknown parameter  $\theta = \mathbb{E}(X)$  (suppose  $\sigma^2 = \text{Var}(X) < \infty$ ).

Among all possible unbiased estimators  $\hat{\theta} = \sum_{i=1}^n a_i X_i$  with  $\sum_{i=1}^n a_i = 1$ . Find the most efficient one.

Sol:

$$\text{Var}(\hat{\theta}) = \sum_{i=1}^n a_i^2 \text{Var}(X) = \sigma^2 \sum_{i=1}^n a_i^2 \geq \sigma^2 \frac{1}{n} \left( \sum_{i=1}^n a_i \right)^2 = \frac{1}{n} \sigma^2,$$

with equality iff  $a_1 = \dots = a_n = 1/n$ .

Hence, the most efficient one is the sample mean  $\hat{\theta} = \bar{X}$ . □

E.g. 1.  $f_Y(y; \theta) = \frac{2y}{\theta^2}$  if  $y \in [0, \theta]$ . Which is more efficient? Find the relative efficiency of  $\hat{\theta}_1$  w.r.t.  $\hat{\theta}_3$ .

$$- \hat{\theta}_1 = \frac{3}{2} \bar{Y}$$

$$- \hat{\theta}_3 = \frac{2n+1}{2n} Y_{\max}.$$

E.g. 2. Let  $X_1, \dots, X_n$  be a random sample of size  $n$  with the unknown parameter  $\theta = \mathbb{E}(X)$  (suppose  $\sigma^2 = \text{Var}(X) < \infty$ ).

Among all possible unbiased estimators  $\hat{\theta} = \sum_{i=1}^n a_i X_i$  with  $\sum_{i=1}^n a_i = 1$ . Find the most efficient one.

Sol:

$$\text{Var}(\hat{\theta}) = \sum_{i=1}^n a_i^2 \text{Var}(X) = \sigma^2 \sum_{i=1}^n a_i^2 \geq \sigma^2 \frac{1}{n} \left( \sum_{i=1}^n a_i \right)^2 = \frac{1}{n} \sigma^2,$$

with equality iff  $a_1 = \dots = a_n = 1/n$ .

Hence, the most efficient one is the sample mean  $\hat{\theta} = \bar{X}$ . □