

Math 221: LINEAR ALGEBRA

Chapter 2. Matrix Algebra

§2-1. Matrix Addition, Scalar Multiplication and Transposition

Le Chen¹

Emory University, 2021 Spring

(last updated on 01/12/2023)



Creative Commons License
(CC BY-NC-SA)

¹ Slides are adapted from those by Karen Seyffarth from University of Calgary.

Linear Algebra with Applications

Lecture Notes

Current Lecture Notes Revision: Version 2018 — Revision B

These lecture notes were originally developed by Karen Seyffarth of the University of Calgary. Edits, additions, and revisions have been made to these notes by the editorial team at Lyryx Learning to accompany their text [Linear Algebra with Applications](#) based on W. K. Nicholson's original text.

In addition we recognize the following contributors. All new content contributed is released under the same license as noted below.

- Ilijas Farah, York University

BE A CHAMPION OF OER!

Contribute suggestions for improvements, new content, or errata:

A new topic

A new example or problem

A new or better proof to an existing theorem

Any other suggestions to improve the material

Contact Lyryx at info@lyryx.com with your ideas.

License



Attribution-NonCommercial-ShareAlike (CC BY-NC-SA)

This license lets others remix, tweak, and build upon your work non-commercially, as long as they credit you and license their new creations under the identical terms.

Copyright

Matrices – Definitions and Basic Properties

Matrix Addition

Scalar Multiplication

The Transpose

Copyright

Matrices – Definitions and Basic Properties

Matrix Addition

Scalar Multiplication

The Transpose

Matrices – Definitions and Basic Properties

Matrices – Definitions and Basic Properties

Definition

Let m and n be positive integers.

- An $m \times n$ matrix is a rectangular array of numbers having m rows and n columns. Such a matrix is said to have size $m \times n$.

Matrices – Definitions and Basic Properties

Definition

Let m and n be positive integers.

- ▶ An $m \times n$ matrix is a rectangular array of numbers having m rows and n columns. Such a matrix is said to have size $m \times n$.
- ▶ A row matrix (or row) is a $1 \times n$ matrix, and a column matrix (or column) is an $m \times 1$ matrix.

Matrices – Definitions and Basic Properties

Definition

Let m and n be positive integers.

- ▶ An $m \times n$ matrix is a rectangular array of numbers having m rows and n columns. Such a matrix is said to have size $m \times n$.
- ▶ A row matrix (or row) is a $1 \times n$ matrix, and a column matrix (or column) is an $m \times 1$ matrix.
- ▶ A square matrix is an $n \times n$ matrix.

Matrices – Definitions and Basic Properties

Definition

Let m and n be positive integers.

- ▶ An $m \times n$ matrix is a rectangular array of numbers having m rows and n columns. Such a matrix is said to have size $m \times n$.
- ▶ A row matrix (or row) is a $1 \times n$ matrix, and a column matrix (or column) is an $m \times 1$ matrix.
- ▶ A square matrix is an $n \times n$ matrix.
- ▶ The (i,j) -entry of a matrix is the entry in row i and column j . For a matrix A , the (i,j) -entry of A is often written as a_{ij} .

Matrices – Definitions and Basic Properties

Definition

Let m and n be positive integers.

- ▶ An $m \times n$ matrix is a rectangular array of numbers having m rows and n columns. Such a matrix is said to have size $m \times n$.
- ▶ A row matrix (or row) is a $1 \times n$ matrix, and a column matrix (or column) is an $m \times 1$ matrix.
- ▶ A square matrix is an $n \times n$ matrix.
- ▶ The (i,j) -entry of a matrix is the entry in row i and column j . For a matrix A , the (i,j) -entry of A is often written as a_{ij} .

General notation for an $m \times n$ matrix, A :

$$A = \begin{bmatrix} a_{11} & a_{12} & a_{13} & \dots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \dots & a_{2n} \\ a_{31} & a_{32} & a_{33} & \dots & a_{3n} \\ \vdots & \vdots & \vdots & & \vdots \\ a_{m1} & a_{m2} & a_{m3} & \dots & a_{mn} \end{bmatrix} = [a_{ij}]$$

Remark (Basic Properties)

1. Equality: two matrices are equal if and only if they have the same size and the corresponding entries are equal.

Remark (Basic Properties)

1. Equality: two matrices are equal if and only if they have the same size and the corresponding entries are equal.
2. Zero Matrix: an $m \times n$ matrix with all entries equal to zero.

Remark (Basic Properties)

1. Equality: two matrices are equal if and only if they have the same size and the corresponding entries are equal.
2. Zero Matrix: an $m \times n$ matrix with all entries equal to zero.
3. Addition: matrices must have the same size; add corresponding entries.

Remark (Basic Properties)

1. Equality: two matrices are equal if and only if they have the same size and the corresponding entries are equal.
2. Zero Matrix: an $m \times n$ matrix with all entries equal to zero.
3. Addition: matrices must have the same size; add corresponding entries.
4. Scalar Multiplication: multiply each entry of the matrix by the scalar.

Remark (Basic Properties)

1. Equality: two matrices are equal if and only if they have the same size and the corresponding entries are equal.
2. Zero Matrix: an $m \times n$ matrix with all entries equal to zero.
3. Addition: matrices must have the same size; add corresponding entries.
4. Scalar Multiplication: multiply each entry of the matrix by the scalar.
5. Negative of a Matrix: for an $m \times n$ matrix A , its negative is denoted $-A$ and $-A = (-1)A$.

Remark (Basic Properties)

1. Equality: two matrices are equal if and only if they have the same size and the corresponding entries are equal.
2. Zero Matrix: an $m \times n$ matrix with all entries equal to zero.
3. Addition: matrices must have the same size; add corresponding entries.
4. Scalar Multiplication: multiply each entry of the matrix by the scalar.
5. Negative of a Matrix: for an $m \times n$ matrix A , its negative is denoted $-A$ and $-A = (-1)A$.
6. Subtraction: for $m \times n$ matrices A and B , $A - B = A + (-1)B$.

Copyright

Matrices – Definitions and Basic Properties

Matrix Addition

Scalar Multiplication

The Transpose

Matrix Addition

Matrix Addition

Definition

Let $A = [a_{ij}]$ and $B = [b_{ij}]$ be two $m \times n$ matrices. Then $A + B = C$ where C is the $m \times n$ matrix $C = [c_{ij}]$ defined by

$$c_{ij} = a_{ij} + b_{ij}$$

Matrix Addition

Definition

Let $A = [a_{ij}]$ and $B = [b_{ij}]$ be two $m \times n$ matrices. Then $A + B = C$ where C is the $m \times n$ matrix $C = [c_{ij}]$ defined by

$$c_{ij} = a_{ij} + b_{ij}$$

Example

Let $A = \begin{bmatrix} 1 & 3 \\ 2 & 5 \end{bmatrix}$, $B = \begin{bmatrix} 0 & -2 \\ 6 & 1 \end{bmatrix}$. Then,

$$\begin{aligned} A + B &= \begin{bmatrix} 1+0 & 3+(-2) \\ 2+6 & 5+1 \end{bmatrix} \\ &= \begin{bmatrix} 1 & 1 \\ 8 & 6 \end{bmatrix} \end{aligned}$$

Theorem (Properties of Matrix Addition)

Let A, B and C be $m \times n$ matrices. Then the following properties hold.

Theorem (Properties of Matrix Addition)

Let A, B and C be $m \times n$ matrices. Then the following properties hold.

1. $A + B = B + A$ (matrix addition is commutative).

Theorem (Properties of Matrix Addition)

Let A, B and C be $m \times n$ matrices. Then the following properties hold.

1. $A + B = B + A$ (matrix addition is commutative).
2. $(A + B) + C = A + (B + C)$ (matrix addition is associative).

Theorem (Properties of Matrix Addition)

Let A, B and C be $m \times n$ matrices. Then the following properties hold.

1. $A + B = B + A$ (matrix addition is commutative).
2. $(A + B) + C = A + (B + C)$ (matrix addition is associative).
3. There exists an $m \times n$ zero matrix, $\mathbf{0}$, such that $A + \mathbf{0} = A$.
(existence of an additive identity).

Theorem (Properties of Matrix Addition)

Let A, B and C be $m \times n$ matrices. Then the following properties hold.

1. $A + B = B + A$ (matrix addition is commutative).
2. $(A + B) + C = A + (B + C)$ (matrix addition is associative).
3. There exists an $m \times n$ zero matrix, $\mathbf{0}$, such that $A + \mathbf{0} = A$.
(existence of an additive identity).
4. There exists an $m \times n$ matrix $-\mathbf{A}$ such that $A + (-A) = \mathbf{0}$.
(existence of an additive inverse).

Copyright

Matrices – Definitions and Basic Properties

Matrix Addition

Scalar Multiplication

The Transpose

Scalar Multiplication

Definition

Let $A = [a_{ij}]$ be an $m \times n$ matrix and let k be a scalar. Then $kA = [ka_{ij}]$.

Scalar Multiplication

Definition

Let $A = [a_{ij}]$ be an $m \times n$ matrix and let k be a scalar. Then $kA = [ka_{ij}]$.

Example

$$\text{Let } A = \begin{bmatrix} 2 & 0 & -1 \\ 3 & 1 & -2 \\ 0 & 4 & 5 \end{bmatrix}.$$

Scalar Multiplication

Definition

Let $A = [a_{ij}]$ be an $m \times n$ matrix and let k be a scalar. Then $kA = [ka_{ij}]$.

Example

$$\text{Let } A = \begin{bmatrix} 2 & 0 & -1 \\ 3 & 1 & -2 \\ 0 & 4 & 5 \end{bmatrix}.$$

Then

$$\begin{aligned} 3A &= \begin{bmatrix} 3(2) & 3(0) & 3(-1) \\ 3(3) & 3(1) & 3(-2) \\ 3(0) & 3(4) & 3(5) \end{bmatrix} \\ &= \begin{bmatrix} 6 & 0 & -3 \\ 9 & 3 & -6 \\ 0 & 12 & 15 \end{bmatrix} \end{aligned}$$

Theorem (Properties of Scalar Multiplication)

Let A, B be $m \times n$ matrices and let $k, p \in \mathbb{R}$ (scalars). Then the following properties hold.

Theorem (Properties of Scalar Multiplication)

Let A, B be $m \times n$ matrices and let $k, p \in \mathbb{R}$ (scalars). Then the following properties hold.

1. $k(A + B) = kA + kB$.
(scalar multiplication distributes over matrix addition).

Theorem (Properties of Scalar Multiplication)

Let A, B be $m \times n$ matrices and let $k, p \in \mathbb{R}$ (scalars). Then the following properties hold.

1. $k(A + B) = kA + kB$.
(scalar multiplication distributes over matrix addition).
2. $(k + p)A = kA + pA$.
(addition distributes over scalar multiplication).

Theorem (Properties of Scalar Multiplication)

Let A, B be $m \times n$ matrices and let $k, p \in \mathbb{R}$ (scalars). Then the following properties hold.

1. $k(A + B) = kA + kB$.
(scalar multiplication distributes over matrix addition).
2. $(k + p)A = kA + pA$.
(addition distributes over scalar multiplication).
3. $k(pA) = (kp)A$. (scalar multiplication is associative).

Theorem (Properties of Scalar Multiplication)

Let A, B be $m \times n$ matrices and let $k, p \in \mathbb{R}$ (scalars). Then the following properties hold.

1. $k(A + B) = kA + kB$.
(scalar multiplication distributes over matrix addition).
2. $(k + p)A = kA + pA$.
(addition distributes over scalar multiplication).
3. $k(pA) = (kp)A$. (scalar multiplication is associative).
4. $1A = A$. (existence of a multiplicative identity).

Example

$$2 \begin{bmatrix} -1 & 0 \\ 1 & 1 \end{bmatrix} + 4 \begin{bmatrix} -2 & 1 \\ 3 & 0 \end{bmatrix} - \begin{bmatrix} 6 & 8 \\ 1 & -1 \end{bmatrix} =$$

Example

$$2 \begin{bmatrix} -1 & 0 \\ 1 & 1 \end{bmatrix} + 4 \begin{bmatrix} -2 & 1 \\ 3 & 0 \end{bmatrix} - \begin{bmatrix} 6 & 8 \\ 1 & -1 \end{bmatrix} = \begin{bmatrix} -16 & -4 \\ 13 & 3 \end{bmatrix}$$

Example

$$2 \begin{bmatrix} -1 & 0 \\ 1 & 1 \end{bmatrix} + 4 \begin{bmatrix} -2 & 1 \\ 3 & 0 \end{bmatrix} - \begin{bmatrix} 6 & 8 \\ 1 & -1 \end{bmatrix} = \begin{bmatrix} -16 & -4 \\ 13 & 3 \end{bmatrix}$$

Problem

Let A and B be $m \times n$ matrices. Simplify the expression

$$2[9(A - B) + 7(2B - A)] - 2[3(2B + A) - 2(A + 3B) - 5(A + B)]$$

Example

$$2 \begin{bmatrix} -1 & 0 \\ 1 & 1 \end{bmatrix} + 4 \begin{bmatrix} -2 & 1 \\ 3 & 0 \end{bmatrix} - \begin{bmatrix} 6 & 8 \\ 1 & -1 \end{bmatrix} = \begin{bmatrix} -16 & -4 \\ 13 & 3 \end{bmatrix}$$

Problem

Let A and B be $m \times n$ matrices. Simplify the expression

$$2[9(A - B) + 7(2B - A)] - 2[3(2B + A) - 2(A + 3B) - 5(A + B)]$$

Solution

$$\begin{aligned} & 2[9(A - B) + 7(2B - A)] - 2[3(2B + A) - 2(A + 3B) - 5(A + B)] \\ = & 2(9A - 9B + 14B - 7A) - 2(6B + 3A - 2A - 6B - 5A - 5B) \\ = & 2(2A + 5B) - 2(-4A - 5B) \\ = & 12A + 20B \end{aligned}$$

Copyright

Matrices – Definitions and Basic Properties

Matrix Addition

Scalar Multiplication

The Transpose

Matrix Transpose

Definition

If A is an $m \times n$ matrix, then its **transpose**, denoted A^T , is the $n \times m$ whose i^{th} row is the i^{th} column of A , $1 \leq i \leq n$; i.e., if $A = [a_{ij}]$, then

$$A^T = [a_{ij}]^T = [a_{ji}]$$

i.e., the (i,j) -entry of A^T is the (j,i) -entry of A .

Matrix Transpose

Definition

If A is an $m \times n$ matrix, then its **transpose**, denoted A^T , is the $n \times m$ whose i^{th} row is the i^{th} column of A , $1 \leq i \leq n$; i.e., if $A = [a_{ij}]$, then

$$A^T = [a_{ij}]^T = [a_{ji}]$$

i.e., the (i, j) -entry of A^T is the (j, i) -entry of A .

Theorem (Properties of the Transpose of a Matrix)

Let A and B be $m \times n$ matrices, C be a $n \times p$ matrix, and $r \in \mathbb{R}$ a scalar.
Then

Matrix Transpose

Definition

If A is an $m \times n$ matrix, then its **transpose**, denoted A^T , is the $n \times m$ whose i^{th} row is the i^{th} column of A , $1 \leq i \leq n$; i.e., if $A = [a_{ij}]$, then

$$A^T = [a_{ij}]^T = [a_{ji}]$$

i.e., the (i, j) -entry of A^T is the (j, i) -entry of A .

Theorem (Properties of the Transpose of a Matrix)

Let A and B be $m \times n$ matrices, C be a $n \times p$ matrix, and $r \in \mathbb{R}$ a scalar. Then

1. $(A^T)^T = A$

Matrix Transpose

Definition

If A is an $m \times n$ matrix, then its **transpose**, denoted A^T , is the $n \times m$ whose i^{th} row is the i^{th} column of A , $1 \leq i \leq n$; i.e., if $A = [a_{ij}]$, then

$$A^T = [a_{ij}]^T = [a_{ji}]$$

i.e., the (i, j) -entry of A^T is the (j, i) -entry of A .

Theorem (Properties of the Transpose of a Matrix)

Let A and B be $m \times n$ matrices, C be a $n \times p$ matrix, and $r \in \mathbb{R}$ a scalar. Then

1. $(A^T)^T = A$
2. $(rA)^T = rA^T$

Matrix Transpose

Definition

If A is an $m \times n$ matrix, then its **transpose**, denoted A^T , is the $n \times m$ whose i^{th} row is the i^{th} column of A , $1 \leq i \leq n$; i.e., if $A = [a_{ij}]$, then

$$A^T = [a_{ij}]^T = [a_{ji}]$$

i.e., the (i, j) -entry of A^T is the (j, i) -entry of A .

Theorem (Properties of the Transpose of a Matrix)

Let A and B be $m \times n$ matrices, C be a $n \times p$ matrix, and $r \in \mathbb{R}$ a scalar. Then

1. $(A^T)^T = A$

2. $(rA)^T = rA^T$

3. $(A + B)^T = A^T + B^T$

Matrix Transpose

Definition

If A is an $m \times n$ matrix, then its **transpose**, denoted A^T , is the $n \times m$ whose i^{th} row is the i^{th} column of A , $1 \leq i \leq n$; i.e., if $A = [a_{ij}]$, then

$$A^T = [a_{ij}]^T = [a_{ji}]$$

i.e., the (i, j) -entry of A^T is the (j, i) -entry of A .

Theorem (Properties of the Transpose of a Matrix)

Let A and B be $m \times n$ matrices, C be a $n \times p$ matrix, and $r \in \mathbb{R}$ a scalar. Then

$$1. (A^T)^T = A$$

$$2. (rA)^T = rA^T$$

$$3. (A + B)^T = A^T + B^T$$

$$4. (AC)^T = C^T A^T$$

Matrix Transpose

Definition

If A is an $m \times n$ matrix, then its **transpose**, denoted A^T , is the $n \times m$ whose i^{th} row is the i^{th} column of A , $1 \leq i \leq n$; i.e., if $A = [a_{ij}]$, then

$$A^T = [a_{ij}]^T = [a_{ji}]$$

i.e., the (i,j) -entry of A^T is the (j,i) -entry of A .

Theorem (Properties of the Transpose of a Matrix)

Let A and B be $m \times n$ matrices, C be a $n \times p$ matrix, and $r \in \mathbb{R}$ a scalar. Then

$$1. (A^T)^T = A$$

$$2. (rA)^T = rA^T$$

$$3. (A + B)^T = A^T + B^T$$

$$4. (AC)^T = C^T A^T$$

To prove each these properties, you only need to compute the (i,j) -entries of the matrices on the left-hand side and the right-hand side.

Matrix Transpose

Definition

If A is an $m \times n$ matrix, then its **transpose**, denoted A^T , is the $n \times m$ whose i^{th} row is the i^{th} column of A , $1 \leq i \leq n$; i.e., if $A = [a_{ij}]$, then

$$A^T = [a_{ij}]^T = [a_{ji}]$$

i.e., the (i,j) -entry of A^T is the (j,i) -entry of A .

Theorem (Properties of the Transpose of a Matrix)

Let A and B be $m \times n$ matrices, C be a $n \times p$ matrix, and $r \in \mathbb{R}$ a scalar. Then

1. $(A^T)^T = A$

2. $(rA)^T = rA^T$

3. $(A + B)^T = A^T + B^T$

4. $(AC)^T = C^T A^T$

To prove each these properties, you only need to compute the (i,j) -entries of the matrices on the left-hand side and the right-hand side. **And you can do it!**

Problem

Find the matrix A if $\left(A + 3 \begin{bmatrix} 1 & -1 & 0 \\ 1 & 2 & 4 \end{bmatrix}\right)^T = \begin{bmatrix} 2 & 1 \\ 0 & 5 \\ 3 & 8 \end{bmatrix}$.

Problem

Find the matrix A if $\left(A + 3 \begin{bmatrix} 1 & -1 & 0 \\ 1 & 2 & 4 \end{bmatrix}\right)^T = \begin{bmatrix} 2 & 1 \\ 0 & 5 \\ 3 & 8 \end{bmatrix}$.

Solution

$$\left[\left(A + 3 \begin{bmatrix} 1 & -1 & 0 \\ 1 & 2 & 4 \end{bmatrix}\right)^T\right]^T = \begin{bmatrix} 2 & 1 \\ 0 & 5 \\ 3 & 8 \end{bmatrix}^T$$

Problem

Find the matrix A if $\left(A + 3 \begin{bmatrix} 1 & -1 & 0 \\ 1 & 2 & 4 \end{bmatrix}\right)^T = \begin{bmatrix} 2 & 1 \\ 0 & 5 \\ 3 & 8 \end{bmatrix}$.

Solution

$$\begin{aligned} \left[\left(A + 3 \begin{bmatrix} 1 & -1 & 0 \\ 1 & 2 & 4 \end{bmatrix} \right)^T \right]^T &= \begin{bmatrix} 2 & 1 \\ 0 & 5 \\ 3 & 8 \end{bmatrix}^T \\ A + 3 \begin{bmatrix} 1 & -1 & 0 \\ 1 & 2 & 4 \end{bmatrix} &= \begin{bmatrix} 2 & 0 & 3 \\ 1 & 5 & 8 \end{bmatrix} \end{aligned}$$

Problem

Find the matrix A if $\left(A + 3 \begin{bmatrix} 1 & -1 & 0 \\ 1 & 2 & 4 \end{bmatrix}\right)^T = \begin{bmatrix} 2 & 1 \\ 0 & 5 \\ 3 & 8 \end{bmatrix}$.

Solution

$$\begin{aligned} \left[\left(A + 3 \begin{bmatrix} 1 & -1 & 0 \\ 1 & 2 & 4 \end{bmatrix} \right)^T \right]^T &= \begin{bmatrix} 2 & 1 \\ 0 & 5 \\ 3 & 8 \end{bmatrix}^T \\ A + 3 \begin{bmatrix} 1 & -1 & 0 \\ 1 & 2 & 4 \end{bmatrix} &= \begin{bmatrix} 2 & 0 & 3 \\ 1 & 5 & 8 \end{bmatrix} \\ A &= \begin{bmatrix} 2 & 0 & 3 \\ 1 & 5 & 8 \end{bmatrix} - 3 \begin{bmatrix} 1 & -1 & 0 \\ 1 & 2 & 4 \end{bmatrix} \end{aligned}$$

Problem

Find the matrix A if $\left(A + 3 \begin{bmatrix} 1 & -1 & 0 \\ 1 & 2 & 4 \end{bmatrix}\right)^T = \begin{bmatrix} 2 & 1 \\ 0 & 5 \\ 3 & 8 \end{bmatrix}$.

Solution

$$\begin{aligned} \left[\left(A + 3 \begin{bmatrix} 1 & -1 & 0 \\ 1 & 2 & 4 \end{bmatrix} \right)^T \right]^T &= \begin{bmatrix} 2 & 1 \\ 0 & 5 \\ 3 & 8 \end{bmatrix}^T \\ A + 3 \begin{bmatrix} 1 & -1 & 0 \\ 1 & 2 & 4 \end{bmatrix} &= \begin{bmatrix} 2 & 0 & 3 \\ 1 & 5 & 8 \end{bmatrix} \\ A &= \begin{bmatrix} 2 & 0 & 3 \\ 1 & 5 & 8 \end{bmatrix} - 3 \begin{bmatrix} 1 & -1 & 0 \\ 1 & 2 & 4 \end{bmatrix} \\ A &= \begin{bmatrix} -1 & 3 & 3 \\ -2 & -1 & -4 \end{bmatrix} \end{aligned}$$

Definition

Let $A = [a_{ij}]$ be an $m \times n$ matrix. The entries $a_{11}, a_{22}, a_{33}, \dots$ are called the **main diagonal** of A .

Definition

Let $A = [a_{ij}]$ be an $m \times n$ matrix. The entries $a_{11}, a_{22}, a_{33}, \dots$ are called the **main diagonal** of A .

Definition (Symmetric Matrices)

The matrix A is called **symmetric** if and only if $A^T = A$. Note that this immediately implies that A is a square matrix.

Definition

Let $A = [a_{ij}]$ be an $m \times n$ matrix. The entries $a_{11}, a_{22}, a_{33}, \dots$ are called the **main diagonal** of A .

Definition (Symmetric Matrices)

The matrix A is called **symmetric** if and only if $A^T = A$. Note that this immediately implies that A is a square matrix.

Examples

$$\begin{bmatrix} 2 & -3 \\ -3 & 17 \end{bmatrix}, \begin{bmatrix} -1 & 0 & 5 \\ 0 & 2 & 11 \\ 5 & 11 & -3 \end{bmatrix}, \begin{bmatrix} 0 & 2 & 5 & -1 \\ 2 & 1 & -3 & 0 \\ 5 & -3 & 2 & -7 \\ -1 & 0 & -7 & 4 \end{bmatrix}$$

are symmetric matrices, and each is symmetric about its main diagonal.

Definition

An $n \times n$ matrix A is said to be **skew symmetric** if $A^T = -A$.

Definition

An $n \times n$ matrix A is said to be **skew symmetric** if $A^T = -A$.

Example (Skew Symmetric Matrices)

$$\begin{bmatrix} 0 & 2 \\ -2 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 9 & 4 \\ -9 & 0 & -3 \\ -4 & 3 & 0 \end{bmatrix}$$

Definition

An $n \times n$ matrix A is said to be **skew symmetric** if $A^T = -A$.

Example (Skew Symmetric Matrices)

$$\begin{bmatrix} 0 & 2 \\ -2 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 9 & 4 \\ -9 & 0 & -3 \\ -4 & 3 & 0 \end{bmatrix}$$

Problem

Show that if A is a square matrix, then $A - A^T$ is skew-symmetric.

Definition

An $n \times n$ matrix A is said to be **skew symmetric** if $A^T = -A$.

Example (Skew Symmetric Matrices)

$$\begin{bmatrix} 0 & 2 \\ -2 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 9 & 4 \\ -9 & 0 & -3 \\ -4 & 3 & 0 \end{bmatrix}$$

Problem

Show that if A is a square matrix, then $A - A^T$ is skew-symmetric.

Solution

We must show that $(A - A^T)^T = -(A - A^T)$.

Definition

An $n \times n$ matrix A is said to be **skew symmetric** if $A^T = -A$.

Example (Skew Symmetric Matrices)

$$\begin{bmatrix} 0 & 2 \\ -2 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 9 & 4 \\ -9 & 0 & -3 \\ -4 & 3 & 0 \end{bmatrix}$$

Problem

Show that if A is a square matrix, then $A - A^T$ is skew-symmetric.

Solution

We must show that $(A - A^T)^T = -(A - A^T)$. Using the properties of matrix addition, scalar multiplication, and transposition

$$(A - A^T)^T = A^T - (A^T)^T = A^T - A = -(A - A^T).$$