

# Math 221: LINEAR ALGEBRA

## Chapter 5. Vector Space $\mathbb{R}^n$

### §5-3. Orthogonality

Le Chen<sup>1</sup>

Emory University, 2021 Spring

(last updated on 01/12/2023)



Creative Commons License  
(CC BY-NC-SA)

<sup>1</sup>Slides are adapted from those by Karen Seyffarth from University of Calgary.



# Linear Algebra with Applications

## Lecture Notes

### Current Lecture Notes Revision: Version 2018 — Revision B

These lecture notes were originally developed by Karen Seyffarth of the University of Calgary. Edits, additions, and revisions have been made to these notes by the editorial team at Lyryx Learning to accompany their text [Linear Algebra with Applications](#) based on W. K. Nicholson's original text.

In addition we recognize the following contributors. All new content contributed is released under the same license as noted below.

- Ilijas Farah, York University

### BE A CHAMPION OF OER!

Contribute suggestions for improvements, new content, or errata:

A new topic

A new example or problem

A new or better proof to an existing theorem

Any other suggestions to improve the material

Contact Lyryx at [info@lyryx.com](mailto:info@lyryx.com) with your ideas.

### License



Attribution-NonCommercial-ShareAlike (CC BY-NC-SA)

This license lets others remix, tweak, and build upon your work non-commercially, as long as they credit you and license their new creations under the identical terms.

Copyright

The Dot Product

The Cauchy Inequality

Orthogonality

Orthogonality and Independence

Fourier Expansion

Copyright

## The Dot Product

The Cauchy Inequality

Orthogonality

Orthogonality and Independence

Fourier Expansion

## Dot Product

# Dot Product

## Definitions

Let  $\vec{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}$  and  $\vec{y} = \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{bmatrix}$  be vectors in  $\mathbb{R}^n$ .

# Dot Product

## Definitions

Let  $\vec{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}$  and  $\vec{y} = \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{bmatrix}$  be vectors in  $\mathbb{R}^n$ .

1. The **dot product** of  $\vec{x}$  and  $\vec{y}$  is

$$\vec{x} \cdot \vec{y} = x_1 y_1 + x_2 y_2 + \cdots x_n y_n = \vec{x}^T \vec{y}.$$



# Dot Product

## Definitions

Let  $\vec{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}$  and  $\vec{y} = \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{bmatrix}$  be vectors in  $\mathbb{R}^n$ .

1. The **dot product** of  $\vec{x}$  and  $\vec{y}$  is

$$\vec{x} \cdot \vec{y} = x_1 y_1 + x_2 y_2 + \cdots + x_n y_n = \vec{x}^T \vec{y}.$$

2. The **length** or **norm** of  $\vec{x}$ , denoted  $||\vec{x}||$  is

$$||\vec{x}|| = \sqrt{x_1^2 + x_2^2 + \cdots + x_n^2} = \sqrt{\vec{x} \cdot \vec{x}} = \sqrt{\vec{x}^T \vec{x}}.$$

# Dot Product

## Definitions

Let  $\vec{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}$  and  $\vec{y} = \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{bmatrix}$  be vectors in  $\mathbb{R}^n$ .

1. The **dot product** of  $\vec{x}$  and  $\vec{y}$  is

$$\vec{x} \cdot \vec{y} = x_1 y_1 + x_2 y_2 + \cdots x_n y_n = \vec{x}^T \vec{y}.$$

2. The **length** or **norm** of  $\vec{x}$ , denoted  $||\vec{x}||$  is

$$||\vec{x}|| = \sqrt{x_1^2 + x_2^2 + \cdots + x_n^2} = \sqrt{\vec{x} \cdot \vec{x}} = \sqrt{\vec{x}^T \vec{x}}.$$

3.  $\vec{x}$  is called a **unit vector** if  $||\vec{x}|| = 1$ .

## Theorem (Properties of length and the dot product)

Let  $\vec{x}, \vec{y}, \vec{z} \in \mathbb{R}^n$ , and let  $a \in \mathbb{R}$ . Then

## Theorem (Properties of length and the dot product)

Let  $\vec{x}, \vec{y}, \vec{z} \in \mathbb{R}^n$ , and let  $a \in \mathbb{R}$ . Then

1.  $\vec{x} \cdot \vec{y} = \vec{y} \cdot \vec{x}$  (the dot product is commutative)

## Theorem (Properties of length and the dot product)

Let  $\vec{x}, \vec{y}, \vec{z} \in \mathbb{R}^n$ , and let  $a \in \mathbb{R}$ . Then

1.  $\vec{x} \cdot \vec{y} = \vec{y} \cdot \vec{x}$  (the dot product is commutative)
2.  $\vec{x} \cdot (\vec{y} + \vec{z}) = \vec{x} \cdot \vec{y} + \vec{x} \cdot \vec{z}$  (the dot product distributes over addition)

## Theorem (Properties of length and the dot product)

Let  $\vec{x}, \vec{y}, \vec{z} \in \mathbb{R}^n$ , and let  $a \in \mathbb{R}$ . Then

1.  $\vec{x} \cdot \vec{y} = \vec{y} \cdot \vec{x}$  (the dot product is commutative)
2.  $\vec{x} \cdot (\vec{y} + \vec{z}) = \vec{x} \cdot \vec{y} + \vec{x} \cdot \vec{z}$  (the dot product distributes over addition)
3.  $(a\vec{x}) \cdot \vec{y} = a(\vec{x} \cdot \vec{y}) = \vec{x} \cdot (a\vec{y})$

## Theorem (Properties of length and the dot product)

Let  $\vec{x}, \vec{y}, \vec{z} \in \mathbb{R}^n$ , and let  $a \in \mathbb{R}$ . Then

1.  $\vec{x} \cdot \vec{y} = \vec{y} \cdot \vec{x}$  (the dot product is commutative)
2.  $\vec{x} \cdot (\vec{y} + \vec{z}) = \vec{x} \cdot \vec{y} + \vec{x} \cdot \vec{z}$  (the dot product distributes over addition)
3.  $(a\vec{x}) \cdot \vec{y} = a(\vec{x} \cdot \vec{y}) = \vec{x} \cdot (a\vec{y})$
4.  $\|\vec{x}\|^2 = \vec{x} \cdot \vec{x}$ .

## Theorem (Properties of length and the dot product)

Let  $\vec{x}, \vec{y}, \vec{z} \in \mathbb{R}^n$ , and let  $a \in \mathbb{R}$ . Then

1.  $\vec{x} \cdot \vec{y} = \vec{y} \cdot \vec{x}$  (the dot product is commutative)
2.  $\vec{x} \cdot (\vec{y} + \vec{z}) = \vec{x} \cdot \vec{y} + \vec{x} \cdot \vec{z}$  (the dot product distributes over addition)
3.  $(a\vec{x}) \cdot \vec{y} = a(\vec{x} \cdot \vec{y}) = \vec{x} \cdot (a\vec{y})$
4.  $\|\vec{x}\|^2 = \vec{x} \cdot \vec{x}$ .
5.  $\|\vec{x}\| \geq 0$  with equality if and only if  $\vec{x} = \vec{0}_n$ .



## Theorem (Properties of length and the dot product)

Let  $\vec{x}, \vec{y}, \vec{z} \in \mathbb{R}^n$ , and let  $a \in \mathbb{R}$ . Then

1.  $\vec{x} \cdot \vec{y} = \vec{y} \cdot \vec{x}$  (the dot product is commutative)
2.  $\vec{x} \cdot (\vec{y} + \vec{z}) = \vec{x} \cdot \vec{y} + \vec{x} \cdot \vec{z}$  (the dot product distributes over addition)
3.  $(a\vec{x}) \cdot \vec{y} = a(\vec{x} \cdot \vec{y}) = \vec{x} \cdot (a\vec{y})$
4.  $\|\vec{x}\|^2 = \vec{x} \cdot \vec{x}$ .
5.  $\|\vec{x}\| \geq 0$  with equality if and only if  $\vec{x} = \vec{0}_n$ .
6.  $\|a\vec{x}\| = |a| \|\vec{x}\|$ .

## Example

Let  $\vec{x}, \vec{y} \in \mathbb{R}^n$ . Then

$$\begin{aligned} \|\vec{x} + \vec{y}\|^2 &= (\vec{x} + \vec{y}) \cdot (\vec{x} + \vec{y}) \\ &= \vec{x} \cdot \vec{x} + \vec{x} \cdot \vec{y} + \vec{y} \cdot \vec{x} + \vec{y} \cdot \vec{y} \\ &= \vec{x} \cdot \vec{x} + 2(\vec{x} \cdot \vec{y}) + \vec{y} \cdot \vec{y} \\ &= \|\vec{x}\|^2 + 2(\vec{x} \cdot \vec{y}) + \|\vec{y}\|^2. \end{aligned}$$

## Problem

Let  $\{\vec{f}_1, \vec{f}_2, \dots, \vec{f}_k\} \in \mathbb{R}^n$  and suppose  $\mathbb{R}^n = \text{span}\{\vec{f}_1, \vec{f}_2, \dots, \vec{f}_k\}$ . Furthermore, suppose that there exists a vector  $\vec{x} \in \mathbb{R}^n$  for which  $\vec{x} \cdot \vec{f}_j = 0$  for all  $j$ ,  $1 \leq j \leq k$ . Show that  $\vec{x} = \vec{0}_n$ .

## Problem

Let  $\{\vec{f}_1, \vec{f}_2, \dots, \vec{f}_k\} \in \mathbb{R}^n$  and suppose  $\mathbb{R}^n = \text{span}\{\vec{f}_1, \vec{f}_2, \dots, \vec{f}_k\}$ . Furthermore, suppose that there exists a vector  $\vec{x} \in \mathbb{R}^n$  for which  $\vec{x} \cdot \vec{f}_j = 0$  for all  $j$ ,  $1 \leq j \leq k$ . Show that  $\vec{x} = \vec{0}_n$ .

## Proof.

Write  $\vec{x} = t_1 \vec{f}_1 + t_2 \vec{f}_2 + \dots + t_k \vec{f}_k$  for some  $t_1, t_2, \dots, t_k \in \mathbb{R}$  (this is possible because  $\vec{f}_1, \vec{f}_2, \dots, \vec{f}_k$  span  $\mathbb{R}^n$ , is this representation unique?).

## Problem

Let  $\{\vec{f}_1, \vec{f}_2, \dots, \vec{f}_k\} \in \mathbb{R}^n$  and suppose  $\mathbb{R}^n = \text{span}\{\vec{f}_1, \vec{f}_2, \dots, \vec{f}_k\}$ . Furthermore, suppose that there exists a vector  $\vec{x} \in \mathbb{R}^n$  for which  $\vec{x} \cdot \vec{f}_j = 0$  for all  $j$ ,  $1 \leq j \leq k$ . Show that  $\vec{x} = \vec{0}_n$ .

## Proof.

Write  $\vec{x} = t_1 \vec{f}_1 + t_2 \vec{f}_2 + \dots + t_k \vec{f}_k$  for some  $t_1, t_2, \dots, t_k \in \mathbb{R}$  (this is possible because  $\vec{f}_1, \vec{f}_2, \dots, \vec{f}_k$  span  $\mathbb{R}^n$ , is this representation unique?). Then

$$\|\vec{x}\|^2 = \vec{x} \cdot \vec{x}$$

## Problem

Let  $\{\vec{f}_1, \vec{f}_2, \dots, \vec{f}_k\} \in \mathbb{R}^n$  and suppose  $\mathbb{R}^n = \text{span}\{\vec{f}_1, \vec{f}_2, \dots, \vec{f}_k\}$ . Furthermore, suppose that there exists a vector  $\vec{x} \in \mathbb{R}^n$  for which  $\vec{x} \cdot \vec{f}_j = 0$  for all  $j$ ,  $1 \leq j \leq k$ . Show that  $\vec{x} = \vec{0}_n$ .

## Proof.

Write  $\vec{x} = t_1 \vec{f}_1 + t_2 \vec{f}_2 + \dots + t_k \vec{f}_k$  for some  $t_1, t_2, \dots, t_k \in \mathbb{R}$  (this is possible because  $\vec{f}_1, \vec{f}_2, \dots, \vec{f}_k$  span  $\mathbb{R}^n$ , is this representation unique?). Then

$$\begin{aligned} \|\vec{x}\|^2 &= \vec{x} \cdot \vec{x} \\ &= \vec{x} \cdot (t_1 \vec{f}_1 + t_2 \vec{f}_2 + \dots + t_k \vec{f}_k) \end{aligned}$$

## Problem

Let  $\{\vec{f}_1, \vec{f}_2, \dots, \vec{f}_k\} \in \mathbb{R}^n$  and suppose  $\mathbb{R}^n = \text{span}\{\vec{f}_1, \vec{f}_2, \dots, \vec{f}_k\}$ . Furthermore, suppose that there exists a vector  $\vec{x} \in \mathbb{R}^n$  for which  $\vec{x} \cdot \vec{f}_j = 0$  for all  $j$ ,  $1 \leq j \leq k$ . Show that  $\vec{x} = \vec{0}_n$ .

## Proof.

Write  $\vec{x} = t_1 \vec{f}_1 + t_2 \vec{f}_2 + \dots + t_k \vec{f}_k$  for some  $t_1, t_2, \dots, t_k \in \mathbb{R}$  (this is possible because  $\vec{f}_1, \vec{f}_2, \dots, \vec{f}_k$  span  $\mathbb{R}^n$ , is this representation unique?). Then

$$\begin{aligned} \|\vec{x}\|^2 &= \vec{x} \cdot \vec{x} \\ &= \vec{x} \cdot (t_1 \vec{f}_1 + t_2 \vec{f}_2 + \dots + t_k \vec{f}_k) \\ &= \vec{x} \cdot (t_1 \vec{f}_1) + \vec{x} \cdot (t_2 \vec{f}_2) + \dots + \vec{x} \cdot (t_k \vec{f}_k) \end{aligned}$$

## Problem

Let  $\{\vec{f}_1, \vec{f}_2, \dots, \vec{f}_k\} \in \mathbb{R}^n$  and suppose  $\mathbb{R}^n = \text{span}\{\vec{f}_1, \vec{f}_2, \dots, \vec{f}_k\}$ . Furthermore, suppose that there exists a vector  $\vec{x} \in \mathbb{R}^n$  for which  $\vec{x} \cdot \vec{f}_j = 0$  for all  $j$ ,  $1 \leq j \leq k$ . Show that  $\vec{x} = \vec{0}_n$ .

## Proof.

Write  $\vec{x} = t_1 \vec{f}_1 + t_2 \vec{f}_2 + \dots + t_k \vec{f}_k$  for some  $t_1, t_2, \dots, t_k \in \mathbb{R}$  (this is possible because  $\vec{f}_1, \vec{f}_2, \dots, \vec{f}_k$  span  $\mathbb{R}^n$ , is this representation unique?). Then

$$\begin{aligned} \|\vec{x}\|^2 &= \vec{x} \cdot \vec{x} \\ &= \vec{x} \cdot (t_1 \vec{f}_1 + t_2 \vec{f}_2 + \dots + t_k \vec{f}_k) \\ &= \vec{x} \cdot (t_1 \vec{f}_1) + \vec{x} \cdot (t_2 \vec{f}_2) + \dots + \vec{x} \cdot (t_k \vec{f}_k) \\ &= t_1 (\vec{x} \cdot \vec{f}_1) + t_2 (\vec{x} \cdot \vec{f}_2) + \dots + t_k (\vec{x} \cdot \vec{f}_k) \end{aligned}$$



## Problem

Let  $\{\vec{f}_1, \vec{f}_2, \dots, \vec{f}_k\} \in \mathbb{R}^n$  and suppose  $\mathbb{R}^n = \text{span}\{\vec{f}_1, \vec{f}_2, \dots, \vec{f}_k\}$ . Furthermore, suppose that there exists a vector  $\vec{x} \in \mathbb{R}^n$  for which  $\vec{x} \cdot \vec{f}_j = 0$  for all  $j$ ,  $1 \leq j \leq k$ . Show that  $\vec{x} = \vec{0}_n$ .

## Proof.

Write  $\vec{x} = t_1 \vec{f}_1 + t_2 \vec{f}_2 + \dots + t_k \vec{f}_k$  for some  $t_1, t_2, \dots, t_k \in \mathbb{R}$  (this is possible because  $\vec{f}_1, \vec{f}_2, \dots, \vec{f}_k$  span  $\mathbb{R}^n$ , is this representation unique?). Then

$$\begin{aligned} ||\vec{x}||^2 &= \vec{x} \cdot \vec{x} \\ &= \vec{x} \cdot (t_1 \vec{f}_1 + t_2 \vec{f}_2 + \dots + t_k \vec{f}_k) \\ &= \vec{x} \cdot (t_1 \vec{f}_1) + \vec{x} \cdot (t_2 \vec{f}_2) + \dots + \vec{x} \cdot (t_k \vec{f}_k) \\ &= t_1 (\vec{x} \cdot \vec{f}_1) + t_2 (\vec{x} \cdot \vec{f}_2) + \dots + t_k (\vec{x} \cdot \vec{f}_k) \\ &= t_1(0) + t_2(0) + \dots + t_k(0) = 0. \end{aligned}$$

## Problem

Let  $\{\vec{f}_1, \vec{f}_2, \dots, \vec{f}_k\} \in \mathbb{R}^n$  and suppose  $\mathbb{R}^n = \text{span}\{\vec{f}_1, \vec{f}_2, \dots, \vec{f}_k\}$ . Furthermore, suppose that there exists a vector  $\vec{x} \in \mathbb{R}^n$  for which  $\vec{x} \cdot \vec{f}_j = 0$  for all  $j$ ,  $1 \leq j \leq k$ . Show that  $\vec{x} = \vec{0}_n$ .

## Proof.

Write  $\vec{x} = t_1\vec{f}_1 + t_2\vec{f}_2 + \dots + t_k\vec{f}_k$  for some  $t_1, t_2, \dots, t_k \in \mathbb{R}$  (this is possible because  $\vec{f}_1, \vec{f}_2, \dots, \vec{f}_k$  span  $\mathbb{R}^n$ , is this representation unique?). Then

$$\begin{aligned} \|\vec{x}\|^2 &= \vec{x} \cdot \vec{x} \\ &= \vec{x} \cdot (t_1\vec{f}_1 + t_2\vec{f}_2 + \dots + t_k\vec{f}_k) \\ &= \vec{x} \cdot (t_1\vec{f}_1) + \vec{x} \cdot (t_2\vec{f}_2) + \dots + \vec{x} \cdot (t_k\vec{f}_k) \\ &= t_1(\vec{x} \cdot \vec{f}_1) + t_2(\vec{x} \cdot \vec{f}_2) + \dots + t_k(\vec{x} \cdot \vec{f}_k) \\ &= t_1(0) + t_2(0) + \dots + t_k(0) = 0. \end{aligned}$$

Since  $\|\vec{x}\|^2 = 0$ ,  $\|\vec{x}\| = 0$ . By the previous theorem,  $\|\vec{x}\| = 0$  if and only if  $\vec{x} = \vec{0}_n$ . Therefore,  $\vec{x} = \vec{0}_n$ . ■

Copyright

The Dot Product

**The Cauchy Inequality**

Orthogonality

Orthogonality and Independence


Fourier Expansion



# Cauchy-Schwartz Inequality

## Theorem (Cauchy-Schwartz Inequality)

If  $\vec{x}, \vec{y} \in \mathbb{R}^n$ , then  $|\vec{x} \cdot \vec{y}| \leq \|\vec{x}\| \|\vec{y}\|$  with equality if and only if  $\{\vec{x}, \vec{y}\}$  is linearly dependent.



$$\left| \frac{\vec{x}}{\|\vec{x}\|} \cdot \frac{\vec{y}}{\|\vec{y}\|} \right| \leq 1$$

$\{\vec{x}, \vec{y}\}$  is linearly dependent  $\Leftrightarrow \vec{x} = t\vec{y}$ , for some  $t \in \mathbb{R}$ .

**Proof.**

Let  $\vec{x}, \vec{y} \in \mathbb{R}^n$  and  $t \in \mathbb{R}$ . Then

$$\begin{aligned} 0 \leq \|t\vec{x} + \vec{y}\|^2 &= (t\vec{x} + \vec{y}) \cdot (t\vec{x} + \vec{y}) \\ &= t^2\vec{x} \cdot \vec{x} + 2t\vec{x} \cdot \vec{y} + \vec{y} \cdot \vec{y} \\ &= t^2\|\vec{x}\|^2 + 2t(\vec{x} \cdot \vec{y}) + \|\vec{y}\|^2. \end{aligned}$$

The quadratic  $t^2\|\vec{x}\|^2 + 2t(\vec{x} \cdot \vec{y}) + \|\vec{y}\|^2$  in  $t$  is always nonnegative, so it does not have distinct real roots. Thus, if we use the quadratic formula to solve for  $t$ , the discriminant must be non-positive, i.e.,

$$\Delta = (2\vec{x} \cdot \vec{y})^2 - 4\|\vec{x}\|^2\|\vec{y}\|^2 \leq 0$$

Therefore,  $(2\vec{x} \cdot \vec{y})^2 \leq 4\|\vec{x}\|^2\|\vec{y}\|^2$ . Since both sides of the inequality are nonnegative, we can take (positive) square roots of both sides:

$$|2\vec{x} \cdot \vec{y}| \leq 2\|\vec{x}\| \|\vec{y}\|$$

Therefore,  $|\vec{x} \cdot \vec{y}| \leq \|\vec{x}\| \|\vec{y}\|$ . What remains is to show that  $|\vec{x} \cdot \vec{y}| = \|\vec{x}\| \|\vec{y}\|$  if and only if  $\{\vec{x}, \vec{y}\}$  is linearly dependent. ■

Proof. (continued)

First suppose that  $\{\vec{x}, \vec{y}\}$  is dependent. Then by symmetry (of  $\vec{x}$  and  $\vec{y}$ ),  $\vec{x} = k\vec{y}$  for some  $k \in \mathbb{R}$ . Hence

$$|\vec{x} \cdot \vec{y}| = |(k\vec{y}) \cdot \vec{y}| = |k| |\vec{y} \cdot \vec{y}| = |k| \|\vec{y}\|^2, \quad \text{and} \quad \|\vec{x}\| \|\vec{y}\| = \|k\vec{y}\| \|\vec{y}\| = |k| \|\vec{y}\|^2,$$

$$\text{so } |\vec{x} \cdot \vec{y}| = \|\vec{x}\| \|\vec{y}\|.$$

Conversely, suppose  $\{\vec{x}, \vec{y}\}$  is independent; then  $t\vec{x} + \vec{y} \neq \vec{0}_n$  for all  $t \in \mathbb{R}$ , so  $\|t\vec{x} + \vec{y}\|^2 > 0$  for all  $t \in \mathbb{R}$ . Thus the quadratic

$$t^2 \|\vec{x}\|^2 + 2t(\vec{x} \cdot \vec{y}) + \|\vec{y}\|^2 > 0$$

so has no real roots. It follows that the the discriminant is negative, i.e.,

$$(2\vec{x} \cdot \vec{y})^2 - 4\|\vec{x}\|^2 \|\vec{y}\|^2 < 0.$$

Therefore,  $(2\vec{x} \cdot \vec{y})^2 < 4\|\vec{x}\|^2 \|\vec{y}\|^2$ ; taking square roots of both sides (they are both nonnegative) and dividing by two gives us

$$|\vec{x} \cdot \vec{y}| < \|\vec{x}\| \|\vec{y}\|,$$

showing that equality is impossible. ■

### Corollary (Triangle Inequality I )

If  $\vec{x}, \vec{y} \in \mathbb{R}^n$ , then  $\|\vec{x} + \vec{y}\| \leq \|\vec{x}\| + \|\vec{y}\|$ .



### Corollary (Triangle Inequality I )

If  $\vec{x}, \vec{y} \in \mathbb{R}^n$ , then  $\|\vec{x} + \vec{y}\| \leq \|\vec{x}\| + \|\vec{y}\|$ .

Proof.

$$\|\vec{x} + \vec{y}\|^2 = (\vec{x} + \vec{y}) \cdot (\vec{x} + \vec{y})$$

### Corollary (Triangle Inequality I )

If  $\vec{x}, \vec{y} \in \mathbb{R}^n$ , then  $\|\vec{x} + \vec{y}\| \leq \|\vec{x}\| + \|\vec{y}\|$ .

Proof.

$$\begin{aligned}\|\vec{x} + \vec{y}\|^2 &= (\vec{x} + \vec{y}) \cdot (\vec{x} + \vec{y}) \\ &= \vec{x} \cdot \vec{x} + 2\vec{x} \cdot \vec{y} + \vec{y} \cdot \vec{y}\end{aligned}$$

### Corollary (Triangle Inequality I )

If  $\vec{x}, \vec{y} \in \mathbb{R}^n$ , then  $||\vec{x} + \vec{y}|| \leq ||\vec{x}|| + ||\vec{y}||$ .

Proof.

$$\begin{aligned} ||\vec{x} + \vec{y}||^2 &= (\vec{x} + \vec{y}) \cdot (\vec{x} + \vec{y}) \\ &= \vec{x} \cdot \vec{x} + 2\vec{x} \cdot \vec{y} + \vec{y} \cdot \vec{y} \\ &= ||\vec{x}||^2 + 2\vec{x} \cdot \vec{y} + ||\vec{y}||^2 \end{aligned}$$

### Corollary (Triangle Inequality I )

If  $\vec{x}, \vec{y} \in \mathbb{R}^n$ , then  $||\vec{x} + \vec{y}|| \leq ||\vec{x}|| + ||\vec{y}||$ .

Proof.

$$\begin{aligned} ||\vec{x} + \vec{y}||^2 &= (\vec{x} + \vec{y}) \cdot (\vec{x} + \vec{y}) \\ &= \vec{x} \cdot \vec{x} + 2\vec{x} \cdot \vec{y} + \vec{y} \cdot \vec{y} \\ &= ||\vec{x}||^2 + 2\vec{x} \cdot \vec{y} + ||\vec{y}||^2 \\ &\leq ||\vec{x}||^2 + 2||\vec{x}|| ||\vec{y}|| + ||\vec{y}||^2 \text{ by the Cauchy Inequality} \end{aligned}$$

### Corollary (Triangle Inequality I )

If  $\vec{x}, \vec{y} \in \mathbb{R}^n$ , then  $\|\vec{x} + \vec{y}\| \leq \|\vec{x}\| + \|\vec{y}\|$ .

Proof.

$$\begin{aligned}\|\vec{x} + \vec{y}\|^2 &= (\vec{x} + \vec{y}) \cdot (\vec{x} + \vec{y}) \\ &= \vec{x} \cdot \vec{x} + 2\vec{x} \cdot \vec{y} + \vec{y} \cdot \vec{y} \\ &= \|\vec{x}\|^2 + 2\vec{x} \cdot \vec{y} + \|\vec{y}\|^2 \\ &\leq \|\vec{x}\|^2 + 2\|\vec{x}\| \|\vec{y}\| + \|\vec{y}\|^2 \text{ by the Cauchy Inequality} \\ &= (\|\vec{x}\| + \|\vec{y}\|)^2.\end{aligned}$$

### Corollary (Triangle Inequality I )

If  $\vec{x}, \vec{y} \in \mathbb{R}^n$ , then  $\|\vec{x} + \vec{y}\| \leq \|\vec{x}\| + \|\vec{y}\|$ .

Proof.

$$\begin{aligned}\|\vec{x} + \vec{y}\|^2 &= (\vec{x} + \vec{y}) \cdot (\vec{x} + \vec{y}) \\ &= \vec{x} \cdot \vec{x} + 2\vec{x} \cdot \vec{y} + \vec{y} \cdot \vec{y} \\ &= \|\vec{x}\|^2 + 2\vec{x} \cdot \vec{y} + \|\vec{y}\|^2 \\ &\leq \|\vec{x}\|^2 + 2\|\vec{x}\| \|\vec{y}\| + \|\vec{y}\|^2 \text{ by the Cauchy Inequality} \\ &= (\|\vec{x}\| + \|\vec{y}\|)^2.\end{aligned}$$

Since both sides of the inequality are nonnegative, we take (positive) square roots of both sides:

$$\|\vec{x} + \vec{y}\| \leq \|\vec{x}\| + \|\vec{y}\|.$$



## Definition

If  $\vec{x}, \vec{y} \in \mathbb{R}^n$ , then the **distance** between  $\vec{x}$  and  $\vec{y}$  is defined as

$$d(\vec{x}, \vec{y}) = \|\vec{x} - \vec{y}\|.$$

## Definition

If  $\vec{x}, \vec{y} \in \mathbb{R}^n$ , then the **distance** between  $\vec{x}$  and  $\vec{y}$  is defined as

$$d(\vec{x}, \vec{y}) = \|\vec{x} - \vec{y}\|.$$

## Theorem (Properties of the distance function)

Let  $\vec{x}, \vec{y}, \vec{z} \in \mathbb{R}^n$ . Then



## Definition

If  $\vec{x}, \vec{y} \in \mathbb{R}^n$ , then the **distance** between  $\vec{x}$  and  $\vec{y}$  is defined as

$$d(\vec{x}, \vec{y}) = \|\vec{x} - \vec{y}\|.$$

## Theorem (Properties of the distance function)

Let  $\vec{x}, \vec{y}, \vec{z} \in \mathbb{R}^n$ . Then

1.  $d(\vec{x}, \vec{y}) \geq 0$ .

## Definition

If  $\vec{x}, \vec{y} \in \mathbb{R}^n$ , then the **distance** between  $\vec{x}$  and  $\vec{y}$  is defined as

$$d(\vec{x}, \vec{y}) = \|\vec{x} - \vec{y}\|.$$

## Theorem (Properties of the distance function)

Let  $\vec{x}, \vec{y}, \vec{z} \in \mathbb{R}^n$ . Then

1.  $d(\vec{x}, \vec{y}) \geq 0$ .
2.  $d(\vec{x}, \vec{y}) = 0$  if and only if  $\vec{x} = \vec{y}$ .

## Definition

If  $\vec{x}, \vec{y} \in \mathbb{R}^n$ , then the **distance** between  $\vec{x}$  and  $\vec{y}$  is defined as

$$d(\vec{x}, \vec{y}) = \|\vec{x} - \vec{y}\|.$$

## Theorem (Properties of the distance function)

Let  $\vec{x}, \vec{y}, \vec{z} \in \mathbb{R}^n$ . Then

1.  $d(\vec{x}, \vec{y}) \geq 0$ .
2.  $d(\vec{x}, \vec{y}) = 0$  if and only if  $\vec{x} = \vec{y}$ .
3.  $d(\vec{x}, \vec{y}) = d(\vec{y}, \vec{x})$ .

## Definition

If  $\vec{x}, \vec{y} \in \mathbb{R}^n$ , then the **distance** between  $\vec{x}$  and  $\vec{y}$  is defined as

$$d(\vec{x}, \vec{y}) = \|\vec{x} - \vec{y}\|.$$

## Theorem (Properties of the distance function)

Let  $\vec{x}, \vec{y}, \vec{z} \in \mathbb{R}^n$ . Then

1.  $d(\vec{x}, \vec{y}) \geq 0$ .
2.  $d(\vec{x}, \vec{y}) = 0$  if and only if  $\vec{x} = \vec{y}$ .
3.  $d(\vec{x}, \vec{y}) = d(\vec{y}, \vec{x})$ .
4.  $d(\vec{x}, \vec{z}) \leq d(\vec{x}, \vec{y}) + d(\vec{y}, \vec{z})$  (Triangle Inequality II).

## Definition

If  $\vec{x}, \vec{y} \in \mathbb{R}^n$ , then the **distance** between  $\vec{x}$  and  $\vec{y}$  is defined as

$$d(\vec{x}, \vec{y}) = \|\vec{x} - \vec{y}\|.$$

## Theorem (Properties of the distance function)

Let  $\vec{x}, \vec{y}, \vec{z} \in \mathbb{R}^n$ . Then

1.  $d(\vec{x}, \vec{y}) \geq 0$ .
2.  $d(\vec{x}, \vec{y}) = 0$  if and only if  $\vec{x} = \vec{y}$ .
3.  $d(\vec{x}, \vec{y}) = d(\vec{y}, \vec{x})$ .
4.  $d(\vec{x}, \vec{z}) \leq d(\vec{x}, \vec{y}) + d(\vec{y}, \vec{z})$  (Triangle Inequality II).

## Proof. (Proof of the Triangle Inequality II)

$$d(\vec{x}, \vec{z}) = \|\vec{x} - \vec{z}\| = \|(\vec{x} - \vec{y}) + (\vec{y} - \vec{z})\|$$

## Definition

If  $\vec{x}, \vec{y} \in \mathbb{R}^n$ , then the **distance** between  $\vec{x}$  and  $\vec{y}$  is defined as

$$d(\vec{x}, \vec{y}) = \|\vec{x} - \vec{y}\|.$$

## Theorem (Properties of the distance function)

Let  $\vec{x}, \vec{y}, \vec{z} \in \mathbb{R}^n$ . Then

1.  $d(\vec{x}, \vec{y}) \geq 0$ .
2.  $d(\vec{x}, \vec{y}) = 0$  if and only if  $\vec{x} = \vec{y}$ .
3.  $d(\vec{x}, \vec{y}) = d(\vec{y}, \vec{x})$ .
4.  $d(\vec{x}, \vec{z}) \leq d(\vec{x}, \vec{y}) + d(\vec{y}, \vec{z})$  (Triangle Inequality II).

## Proof. (Proof of the Triangle Inequality II)

$$\begin{aligned} d(\vec{x}, \vec{z}) = \|\vec{x} - \vec{z}\| &= \|(\vec{x} - \vec{y}) + (\vec{y} - \vec{z})\| \\ &\leq \|\vec{x} - \vec{y}\| + \|\vec{y} - \vec{z}\| \text{ by Triangle Inequality I} \end{aligned}$$

## Definition

If  $\vec{x}, \vec{y} \in \mathbb{R}^n$ , then the **distance** between  $\vec{x}$  and  $\vec{y}$  is defined as

$$d(\vec{x}, \vec{y}) = \|\vec{x} - \vec{y}\|.$$

## Theorem (Properties of the distance function)

Let  $\vec{x}, \vec{y}, \vec{z} \in \mathbb{R}^n$ . Then

1.  $d(\vec{x}, \vec{y}) \geq 0$ .
2.  $d(\vec{x}, \vec{y}) = 0$  if and only if  $\vec{x} = \vec{y}$ .
3.  $d(\vec{x}, \vec{y}) = d(\vec{y}, \vec{x})$ .
4.  $d(\vec{x}, \vec{z}) \leq d(\vec{x}, \vec{y}) + d(\vec{y}, \vec{z})$  (Triangle Inequality II).

## Proof. (Proof of the Triangle Inequality II)

$$\begin{aligned} d(\vec{x}, \vec{z}) = \|\vec{x} - \vec{z}\| &= \|(\vec{x} - \vec{y}) + (\vec{y} - \vec{z})\| \\ &\leq \|\vec{x} - \vec{y}\| + \|\vec{y} - \vec{z}\| \text{ by Triangle Inequality I} \\ &= d(\vec{x}, \vec{y}) + d(\vec{y}, \vec{z}). \end{aligned}$$



Copyright

The Dot Product

The Cauchy Inequality

Orthogonality

Orthogonality and Independence

Fourier Expansion





# Orthogonality

## Definitions

- Let  $\vec{x}, \vec{y} \in \mathbb{R}^n$ . We say that two vectors  $\vec{x}$  and  $\vec{y}$  are **orthogonal** if  $\vec{x} \cdot \vec{y} = 0$ .

# Orthogonality

## Definitions

- ▶ Let  $\vec{x}, \vec{y} \in \mathbb{R}^n$ . We say that two vectors  $\vec{x}$  and  $\vec{y}$  are **orthogonal** if  $\vec{x} \cdot \vec{y} = 0$ .
- ▶ More generally,  $X = \{\vec{x}_1, \vec{x}_2, \dots, \vec{x}_k\} \subseteq \mathbb{R}^n$  is an **orthogonal set** if each  $\vec{x}_i$  is nonzero, and every pair of **distinct** vectors of  $X$  is orthogonal, i.e.,  $\vec{x}_i \cdot \vec{x}_j = 0$  for all  $i \neq j$ ,  $1 \leq i, j \leq k$ .

# Orthogonality

## Definitions

- ▶ Let  $\vec{x}, \vec{y} \in \mathbb{R}^n$ . We say that two vectors  $\vec{x}$  and  $\vec{y}$  are **orthogonal** if  $\vec{x} \cdot \vec{y} = 0$ .
- ▶ More generally,  $X = \{\vec{x}_1, \vec{x}_2, \dots, \vec{x}_k\} \subseteq \mathbb{R}^n$  is an **orthogonal set** if each  $\vec{x}_i$  is nonzero, and every pair of **distinct** vectors of  $X$  is orthogonal, i.e.,  $\vec{x}_i \cdot \vec{x}_j = 0$  for all  $i \neq j$ ,  $1 \leq i, j \leq k$ .
- ▶ A set  $X = \{\vec{x}_1, \vec{x}_2, \dots, \vec{x}_k\} \subseteq \mathbb{R}^n$  is an **orthonormal** set if  $X$  is an orthogonal set of **unit vectors**, i.e.,  $\|\vec{x}_i\| = 1$  for all  $i$ ,  $1 \leq i \leq k$ .

## Examples

1. The standard basis  $\{\vec{e}_1, \dots, \vec{e}_n\}$  of  $\mathbb{R}^n$  is an orthonormal set (and hence an orthogonal set).

## Examples

1. The standard basis  $\{\vec{e}_1, \dots, \vec{e}_n\}$  of  $\mathbb{R}^n$  is an orthonormal set (and hence an orthogonal set).

2.

$$\left\{ \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ -1 \\ -1 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 1 \\ -1 \end{bmatrix} \right\}$$

is an orthogonal (but not orthonormal) subset of  $\mathbb{R}^4$ .

## Examples

1. The standard basis  $\{\vec{e}_1, \dots, \vec{e}_n\}$  of  $\mathbb{R}^n$  is an orthonormal set (and hence an orthogonal set).

2.

$$\left\{ \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ -1 \\ -1 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 1 \\ -1 \end{bmatrix} \right\}$$

is an orthogonal (but not orthonormal) subset of  $\mathbb{R}^4$ .

3. If  $\{\vec{x}_1, \vec{x}_2, \dots, \vec{x}_k\}$  is an orthogonal subset of  $\mathbb{R}^n$  and  $p \neq 0$ , then  $\{p\vec{x}_1, p\vec{x}_2, \dots, p\vec{x}_k\}$  is an orthogonal subset of  $\mathbb{R}^n$ .

## Examples

1. The standard basis  $\{\vec{e}_1, \dots, \vec{e}_n\}$  of  $\mathbb{R}^n$  is an orthonormal set (and hence an orthogonal set).

2.

$$\left\{ \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ -1 \\ -1 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 1 \\ -1 \end{bmatrix} \right\}$$

is an orthogonal (but not orthonormal) subset of  $\mathbb{R}^4$ .

3. If  $\{\vec{x}_1, \vec{x}_2, \dots, \vec{x}_k\}$  is an orthogonal subset of  $\mathbb{R}^n$  and  $p \neq 0$ , then  $\{p\vec{x}_1, p\vec{x}_2, \dots, p\vec{x}_k\}$  is an orthogonal subset of  $\mathbb{R}^n$ .

4.

$$\left\{ \frac{1}{2} \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}, \frac{1}{2} \begin{bmatrix} 1 \\ 1 \\ -1 \\ -1 \end{bmatrix}, \frac{1}{2} \begin{bmatrix} 1 \\ -1 \\ 1 \\ -1 \end{bmatrix} \right\}$$

is an orthonormal subset of  $\mathbb{R}^4$ .





## Definition

**Normalizing an orthogonal set** is the process of turning an orthogonal (but not orthonormal) set into an orthonormal set. If  $\{\vec{x}_1, \vec{x}_2, \dots, \vec{x}_k\}$  is an orthogonal subset of  $\mathbb{R}^n$ , then

$$\left\{ \frac{1}{\|\vec{x}_1\|} \vec{x}_1, \frac{1}{\|\vec{x}_2\|} \vec{x}_2, \dots, \frac{1}{\|\vec{x}_k\|} \vec{x}_k \right\}$$

is an orthonormal set.



### Problem

Verify that

$$\left\{ \begin{bmatrix} 1 \\ -1 \\ 2 \end{bmatrix}, \begin{bmatrix} 0 \\ 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 5 \\ 1 \\ -2 \end{bmatrix} \right\}$$

is an orthogonal set, and normalize this set.

## Solution

$$\begin{bmatrix} 1 \\ -1 \\ 2 \end{bmatrix} \cdot \begin{bmatrix} 0 \\ 2 \\ 1 \end{bmatrix} = 0 - 2 + 2 = 0,$$

$$\begin{bmatrix} 0 \\ 2 \\ 1 \end{bmatrix} \cdot \begin{bmatrix} 5 \\ 1 \\ -2 \end{bmatrix} = 0 + 2 - 2 = 0,$$

$$\begin{bmatrix} 1 \\ -1 \\ 2 \end{bmatrix} \cdot \begin{bmatrix} 5 \\ 1 \\ -2 \end{bmatrix} = 5 - 1 - 4 = 0,$$

proving that the set is orthogonal. Normalizing gives us the orthonormal set

$$\left\{ \frac{1}{\sqrt{6}} \begin{bmatrix} 1 \\ -1 \\ 2 \end{bmatrix}, \frac{1}{\sqrt{5}} \begin{bmatrix} 0 \\ 2 \\ 1 \end{bmatrix}, \frac{1}{\sqrt{30}} \begin{bmatrix} 5 \\ 1 \\ -2 \end{bmatrix} \right\}.$$



## Theorem (Pythagoras' Theorem)

If  $\{\vec{x}_1, \vec{x}_2, \dots, \vec{x}_k\} \subseteq \mathbb{R}^n$  is orthogonal, then

$$\|\vec{x}_1 + \vec{x}_2 + \dots + \vec{x}_k\|^2 = \|\vec{x}_1\|^2 + \|\vec{x}_2\|^2 + \dots + \|\vec{x}_k\|^2.$$

## Theorem (Pythagoras' Theorem)

If  $\{\vec{x}_1, \vec{x}_2, \dots, \vec{x}_k\} \subseteq \mathbb{R}^n$  is orthogonal, then

$$||\vec{x}_1 + \vec{x}_2 + \dots + \vec{x}_k||^2 = ||\vec{x}_1||^2 + ||\vec{x}_2||^2 + \dots + ||\vec{x}_k||^2.$$

Proof.

Start with

$$||\vec{x}_1 + \vec{x}_2 + \dots + \vec{x}_k||^2 = (\vec{x}_1 + \vec{x}_2 + \dots + \vec{x}_k) \cdot (\vec{x}_1 + \vec{x}_2 + \dots + \vec{x}_k)$$

## Theorem (Pythagoras' Theorem)

If  $\{\vec{x}_1, \vec{x}_2, \dots, \vec{x}_k\} \subseteq \mathbb{R}^n$  is orthogonal, then

$$\|\vec{x}_1 + \vec{x}_2 + \dots + \vec{x}_k\|^2 = \|\vec{x}_1\|^2 + \|\vec{x}_2\|^2 + \dots + \|\vec{x}_k\|^2.$$

Proof.

Start with

$$\begin{aligned} \|\vec{x}_1 + \vec{x}_2 + \dots + \vec{x}_k\|^2 &= (\vec{x}_1 + \vec{x}_2 + \dots + \vec{x}_k) \cdot (\vec{x}_1 + \vec{x}_2 + \dots + \vec{x}_k) \\ &= (\vec{x}_1 \cdot \vec{x}_1 + \vec{x}_1 \cdot \vec{x}_2 + \dots + \vec{x}_1 \cdot \vec{x}_k) \\ &\quad + (\vec{x}_2 \cdot \vec{x}_1 + \vec{x}_2 \cdot \vec{x}_2 + \dots + \vec{x}_2 \cdot \vec{x}_k) \\ &\quad \vdots \qquad \qquad \qquad \vdots \qquad \qquad \qquad \vdots \\ &\quad + (\vec{x}_k \cdot \vec{x}_1 + \vec{x}_k \cdot \vec{x}_2 + \dots + \vec{x}_k \cdot \vec{x}_k) \\ &= \vec{x}_1 \cdot \vec{x}_1 + \vec{x}_2 \cdot \vec{x}_2 + \dots + \vec{x}_k \cdot \vec{x}_k \end{aligned}$$

## Theorem (Pythagoras' Theorem)

If  $\{\vec{x}_1, \vec{x}_2, \dots, \vec{x}_k\} \subseteq \mathbb{R}^n$  is orthogonal, then

$$\|\vec{x}_1 + \vec{x}_2 + \dots + \vec{x}_k\|^2 = \|\vec{x}_1\|^2 + \|\vec{x}_2\|^2 + \dots + \|\vec{x}_k\|^2.$$

Proof.

Start with

$$\begin{aligned} \|\vec{x}_1 + \vec{x}_2 + \dots + \vec{x}_k\|^2 &= (\vec{x}_1 + \vec{x}_2 + \dots + \vec{x}_k) \cdot (\vec{x}_1 + \vec{x}_2 + \dots + \vec{x}_k) \\ &= (\vec{x}_1 \cdot \vec{x}_1 + \vec{x}_1 \cdot \vec{x}_2 + \dots + \vec{x}_1 \cdot \vec{x}_k) \\ &\quad + (\vec{x}_2 \cdot \vec{x}_1 + \vec{x}_2 \cdot \vec{x}_2 + \dots + \vec{x}_2 \cdot \vec{x}_k) \\ &\quad \vdots \qquad \qquad \qquad \vdots \qquad \qquad \qquad \vdots \\ &\quad + (\vec{x}_k \cdot \vec{x}_1 + \vec{x}_k \cdot \vec{x}_2 + \dots + \vec{x}_k \cdot \vec{x}_k) \\ &= \vec{x}_1 \cdot \vec{x}_1 + \vec{x}_2 \cdot \vec{x}_2 + \dots + \vec{x}_k \cdot \vec{x}_k \\ &= \|\vec{x}_1\|^2 + \|\vec{x}_2\|^2 + \dots + \|\vec{x}_k\|^2. \end{aligned}$$



## Theorem (Pythagoras' Theorem)

If  $\{\vec{x}_1, \vec{x}_2, \dots, \vec{x}_k\} \subseteq \mathbb{R}^n$  is orthogonal, then

$$\|\vec{x}_1 + \vec{x}_2 + \dots + \vec{x}_k\|^2 = \|\vec{x}_1\|^2 + \|\vec{x}_2\|^2 + \dots + \|\vec{x}_k\|^2.$$

Proof.

Start with

$$\begin{aligned} \|\vec{x}_1 + \vec{x}_2 + \dots + \vec{x}_k\|^2 &= (\vec{x}_1 + \vec{x}_2 + \dots + \vec{x}_k) \cdot (\vec{x}_1 + \vec{x}_2 + \dots + \vec{x}_k) \\ &= (\vec{x}_1 \cdot \vec{x}_1 + \vec{x}_1 \cdot \vec{x}_2 + \dots + \vec{x}_1 \cdot \vec{x}_k) \\ &\quad + (\vec{x}_2 \cdot \vec{x}_1 + \vec{x}_2 \cdot \vec{x}_2 + \dots + \vec{x}_2 \cdot \vec{x}_k) \\ &\quad \vdots \qquad \qquad \qquad \vdots \qquad \qquad \qquad \vdots \\ &\quad + (\vec{x}_k \cdot \vec{x}_1 + \vec{x}_k \cdot \vec{x}_2 + \dots + \vec{x}_k \cdot \vec{x}_k) \\ &= \vec{x}_1 \cdot \vec{x}_1 + \vec{x}_2 \cdot \vec{x}_2 + \dots + \vec{x}_k \cdot \vec{x}_k \\ &= \|\vec{x}_1\|^2 + \|\vec{x}_2\|^2 + \dots + \|\vec{x}_k\|^2. \end{aligned}$$

The second last equality follows from the fact that the set is orthogonal, so for all  $i$  and  $j$ ,  $i \neq j$  and  $1 \leq i, j \leq k$ ,  $\vec{x}_i \cdot \vec{x}_j = 0$ . Thus, the only nonzero terms are the ones of the form  $\vec{x}_i \cdot \vec{x}_i$ ,  $1 \leq i \leq k$ . ■

Copyright

The Dot Product

The Cauchy Inequality

Orthogonality

**Orthogonality and Independence**

Fourier Expansion

# Orthogonality and Independence

# Orthogonality and Independence

## Theorem

If  $S = \{\vec{x}_1, \vec{x}_2, \dots, \vec{x}_k\} \subseteq \mathbb{R}^n$  is an orthogonal set, then  $S$  is independent.

# Orthogonality and Independence

## Theorem

If  $S = \{\vec{x}_1, \vec{x}_2, \dots, \vec{x}_k\} \subseteq \mathbb{R}^n$  is an orthogonal set, then  $S$  is independent.

## Proof.

Form the linear equation:  $t_1\vec{x}_1 + t_2\vec{x}_2 + \dots + t_k\vec{x}_k = \vec{0}$ . We need to check whether there is only trivial solution.

# Orthogonality and Independence

## Theorem

If  $S = \{\vec{x}_1, \vec{x}_2, \dots, \vec{x}_k\} \subseteq \mathbb{R}^n$  is an orthogonal set, then  $S$  is independent.

## Proof.

Form the linear equation:  $t_1\vec{x}_1 + t_2\vec{x}_2 + \dots + t_k\vec{x}_k = \vec{0}$ . We need to check whether there is only trivial solution. Notice that for all  $i$ ,  $1 \leq i \leq k$ ,

$$0 = (t_1\vec{x}_1 + t_2\vec{x}_2 + \dots + t_k\vec{x}_k) \cdot \vec{x}_i = t_i\vec{x}_i \cdot \vec{x}_i = t_i\|\vec{x}_i\|^2,$$

since  $t_j\vec{x}_j \cdot \vec{x}_i = 0$  for all  $j$ ,  $1 \leq j \leq k$  where  $j \neq i$ .

# Orthogonality and Independence


## Theorem

If  $S = \{\vec{x}_1, \vec{x}_2, \dots, \vec{x}_k\} \subseteq \mathbb{R}^n$  is an orthogonal set, then  $S$  is independent.

## Proof.

Form the linear equation:  $t_1\vec{x}_1 + t_2\vec{x}_2 + \dots + t_k\vec{x}_k = \vec{0}$ . We need to check whether there is only trivial solution. Notice that for all  $i$ ,  $1 \leq i \leq k$ ,

$$0 = (t_1\vec{x}_1 + t_2\vec{x}_2 + \dots + t_k\vec{x}_k) \cdot \vec{x}_i = t_i\vec{x}_i \cdot \vec{x}_i = t_i\|\vec{x}_i\|^2,$$

since  $t_j\vec{x}_j \cdot \vec{x}_i = 0$  for all  $j$ ,  $1 \leq j \leq k$  where  $j \neq i$ . Since  $\vec{x}_i \neq \vec{0}_n$  and  $t_i\|\vec{x}_i\|^2 = 0$ , it follows that  $t_i = 0$  for all  $i$ ,  $1 \leq i \leq k$ . Therefore,  $S$  is linearly independent. 

## Example

Given an arbitrary vector

$$\vec{x} = \begin{bmatrix} a_1 \\ a_2 \\ \vdots \\ a_n \end{bmatrix} \in \mathbb{R}^n,$$


it is trivial to express  $\vec{x}$  as a linear combination of the standard basis vectors of  $\mathbb{R}^n$ ,  $\{\vec{e}_1, \vec{e}_2, \dots, \vec{e}_n\}$ :

$$\vec{x} = a_1 \vec{e}_1 + a_2 \vec{e}_2 + \dots + a_n \vec{e}_n.$$



## Problem

Given any orthogonal basis  $B$  of  $\mathbb{R}^n$  (so not necessarily the standard basis), and an arbitrary vector  $\vec{x} \in \mathbb{R}^n$ , how do we express  $\vec{x}$  as a linear combination of the vectors in  $B$ ?



Copyright

The Dot Product

The Cauchy Inequality

Orthogonality

Orthogonality and Independence

**Fourier Expansion**



# Fourier Expansion

## Theorem (Fourier Expansion)

Let  $\{\vec{f}_1, \vec{f}_2, \dots, \vec{f}_m\}$  be an orthogonal basis of a subspace  $U$  of  $\mathbb{R}^n$ . Then for any  $\vec{x} \in U$ ,

$$\vec{x} = \left( \frac{\vec{x} \cdot \vec{f}_1}{\|\vec{f}_1\|^2} \right) \vec{f}_1 + \left( \frac{\vec{x} \cdot \vec{f}_2}{\|\vec{f}_2\|^2} \right) \vec{f}_2 + \dots + \left( \frac{\vec{x} \cdot \vec{f}_m}{\|\vec{f}_m\|^2} \right) \vec{f}_m.$$

This expression is called the **Fourier expansion** of  $\vec{x}$ , and

$$\frac{\vec{x} \cdot \vec{f}_j}{\|\vec{f}_j\|^2}, \quad j = 1, 2, \dots, m$$

are called the **Fourier coefficients**.

### Example

Let  $\vec{f}_1 = \begin{bmatrix} 1 \\ -1 \\ 2 \end{bmatrix}$ ,  $\vec{f}_2 = \begin{bmatrix} 0 \\ 2 \\ 1 \end{bmatrix}$ , and  $\vec{f}_3 = \begin{bmatrix} 5 \\ 1 \\ -2 \end{bmatrix}$ , and let  $\vec{x} = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$ .

We have seen that  $B = \{\vec{f}_1, \vec{f}_2, \vec{f}_3\}$  is an orthogonal subset of  $\mathbb{R}^3$ .

### Example

Let  $\vec{f}_1 = \begin{bmatrix} 1 \\ -1 \\ 2 \end{bmatrix}$ ,  $\vec{f}_2 = \begin{bmatrix} 0 \\ 2 \\ 1 \end{bmatrix}$ , and  $\vec{f}_3 = \begin{bmatrix} 5 \\ 1 \\ -2 \end{bmatrix}$ , and let  $\vec{x} = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$ .

We have seen that  $B = \{\vec{f}_1, \vec{f}_2, \vec{f}_3\}$  is an orthogonal subset of  $\mathbb{R}^3$ .

It follows that  $B$  is an orthogonal basis of  $\mathbb{R}^3$ . (Why?)

### Example

$$\text{Let } \vec{f}_1 = \begin{bmatrix} 1 \\ -1 \\ 2 \end{bmatrix}, \vec{f}_2 = \begin{bmatrix} 0 \\ 2 \\ 1 \end{bmatrix}, \text{ and } \vec{f}_3 = \begin{bmatrix} 5 \\ 1 \\ -2 \end{bmatrix}, \text{ and let } \vec{x} = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}.$$

We have seen that  $B = \{\vec{f}_1, \vec{f}_2, \vec{f}_3\}$  is an orthogonal subset of  $\mathbb{R}^3$ .

It follows that  $B$  is an orthogonal basis of  $\mathbb{R}^3$ . (Why?)

To express  $\vec{x}$  as a linear combination of the vectors of  $B$ , apply the Fourier Expansion Theorem. Assume  $\vec{x} = t_1\vec{f}_1 + t_2\vec{f}_2 + t_3\vec{f}_3$ .

### Example

$$\text{Let } \vec{f}_1 = \begin{bmatrix} 1 \\ -1 \\ 2 \end{bmatrix}, \vec{f}_2 = \begin{bmatrix} 0 \\ 2 \\ 1 \end{bmatrix}, \text{ and } \vec{f}_3 = \begin{bmatrix} 5 \\ 1 \\ -2 \end{bmatrix}, \text{ and let } \vec{x} = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}.$$

We have seen that  $B = \{\vec{f}_1, \vec{f}_2, \vec{f}_3\}$  is an orthogonal subset of  $\mathbb{R}^3$ .

It follows that  $B$  is an orthogonal basis of  $\mathbb{R}^3$ . (Why?)

To express  $\vec{x}$  as a linear combination of the vectors of  $B$ , apply the Fourier Expansion Theorem. Assume  $\vec{x} = t_1\vec{f}_1 + t_2\vec{f}_2 + t_3\vec{f}_3$ . Then

$$t_1 = \frac{\vec{x} \cdot \vec{f}_1}{\|\vec{f}_1\|^2} = \frac{2}{6}, \quad t_2 = \frac{\vec{x} \cdot \vec{f}_2}{\|\vec{f}_2\|^2} = \frac{3}{5}, \quad \text{and} \quad t_3 = \frac{\vec{x} \cdot \vec{f}_3}{\|\vec{f}_3\|^2} = \frac{4}{30}.$$



### Example

$$\text{Let } \vec{f}_1 = \begin{bmatrix} 1 \\ -1 \\ 2 \end{bmatrix}, \vec{f}_2 = \begin{bmatrix} 0 \\ 2 \\ 1 \end{bmatrix}, \text{ and } \vec{f}_3 = \begin{bmatrix} 5 \\ 1 \\ -2 \end{bmatrix}, \text{ and let } \vec{x} = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}.$$

We have seen that  $B = \{\vec{f}_1, \vec{f}_2, \vec{f}_3\}$  is an orthogonal subset of  $\mathbb{R}^3$ .

It follows that  $B$  is an orthogonal basis of  $\mathbb{R}^3$ . (Why?)

To express  $\vec{x}$  as a linear combination of the vectors of  $B$ , apply the Fourier Expansion Theorem. Assume  $\vec{x} = t_1\vec{f}_1 + t_2\vec{f}_2 + t_3\vec{f}_3$ . Then

$$t_1 = \frac{\vec{x} \cdot \vec{f}_1}{\|\vec{f}_1\|^2} = \frac{2}{6}, \quad t_2 = \frac{\vec{x} \cdot \vec{f}_2}{\|\vec{f}_2\|^2} = \frac{3}{5}, \quad \text{and} \quad t_3 = \frac{\vec{x} \cdot \vec{f}_3}{\|\vec{f}_3\|^2} = \frac{4}{30}.$$

Therefore,

$$\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} = \frac{1}{3} \begin{bmatrix} 1 \\ -1 \\ 2 \end{bmatrix} + \frac{3}{5} \begin{bmatrix} 0 \\ 2 \\ 1 \end{bmatrix} + \frac{2}{15} \begin{bmatrix} 5 \\ 1 \\ -2 \end{bmatrix}.$$

Proof. (Fourier Expansion)

Let  $\vec{x} \in U$ . Since  $\{\vec{f}_1, \vec{f}_2, \dots, \vec{f}_m\}$  is a basis of  $U$ ,  $\vec{x} = t_1\vec{f}_1 + t_2\vec{f}_2 + \dots + t_m\vec{f}_m$  for some  $t_1, t_2, \dots, t_m \in \mathbb{R}$ .

### Proof. (Fourier Expansion)

Let  $\vec{x} \in U$ . Since  $\{\vec{f}_1, \vec{f}_2, \dots, \vec{f}_m\}$  is a basis of  $U$ ,  $\vec{x} = t_1\vec{f}_1 + t_2\vec{f}_2 + \dots + t_m\vec{f}_m$  for some  $t_1, t_2, \dots, t_m \in \mathbb{R}$ . Notice that for any  $i$ ,  $1 \leq i \leq m$ ,

$$\vec{x} \cdot \vec{f}_i = (t_1\vec{f}_1 + t_2\vec{f}_2 + \dots + t_m\vec{f}_m) \cdot \vec{f}_i$$

### Proof. (Fourier Expansion)

Let  $\vec{x} \in U$ . Since  $\{\vec{f}_1, \vec{f}_2, \dots, \vec{f}_m\}$  is a basis of  $U$ ,  $\vec{x} = t_1\vec{f}_1 + t_2\vec{f}_2 + \dots + t_m\vec{f}_m$  for some  $t_1, t_2, \dots, t_m \in \mathbb{R}$ . Notice that for any  $i$ ,  $1 \leq i \leq m$ ,

$$\begin{aligned}\vec{x} \cdot \vec{f}_i &= (t_1\vec{f}_1 + t_2\vec{f}_2 + \dots + t_m\vec{f}_m) \cdot \vec{f}_i \\ &= t_i\vec{f}_i \cdot \vec{f}_i \quad \text{since } \{\vec{f}_1, \vec{f}_2, \dots, \vec{f}_m\} \text{ is orthogonal} \\ &= t_i \|\vec{f}_i\|^2.\end{aligned}$$

### Proof. (Fourier Expansion)

Let  $\vec{x} \in U$ . Since  $\{\vec{f}_1, \vec{f}_2, \dots, \vec{f}_m\}$  is a basis of  $U$ ,  $\vec{x} = t_1\vec{f}_1 + t_2\vec{f}_2 + \dots + t_m\vec{f}_m$  for some  $t_1, t_2, \dots, t_m \in \mathbb{R}$ . Notice that for any  $i$ ,  $1 \leq i \leq m$ ,

$$\begin{aligned}\vec{x} \cdot \vec{f}_i &= (t_1\vec{f}_1 + t_2\vec{f}_2 + \dots + t_m\vec{f}_m) \cdot \vec{f}_i \\ &= t_i\vec{f}_i \cdot \vec{f}_i \quad \text{since } \{\vec{f}_1, \vec{f}_2, \dots, \vec{f}_m\} \text{ is orthogonal} \\ &= t_i \|\vec{f}_i\|^2.\end{aligned}$$

Since  $\vec{f}_i$  is nonzero, we obtain

$$t_i = \frac{\vec{x} \cdot \vec{f}_i}{\|\vec{f}_i\|^2}.$$

The result now follows. ■

## Proof. (Fourier Expansion)

Let  $\vec{x} \in U$ . Since  $\{\vec{f}_1, \vec{f}_2, \dots, \vec{f}_m\}$  is a basis of  $U$ ,  $\vec{x} = t_1\vec{f}_1 + t_2\vec{f}_2 + \dots + t_m\vec{f}_m$  for some  $t_1, t_2, \dots, t_m \in \mathbb{R}$ . Notice that for any  $i$ ,  $1 \leq i \leq m$ ,

$$\begin{aligned}\vec{x} \cdot \vec{f}_i &= (t_1\vec{f}_1 + t_2\vec{f}_2 + \dots + t_m\vec{f}_m) \cdot \vec{f}_i \\ &= t_i\vec{f}_i \cdot \vec{f}_i \quad \text{since } \{\vec{f}_1, \vec{f}_2, \dots, \vec{f}_m\} \text{ is orthogonal} \\ &= t_i \|\vec{f}_i\|^2.\end{aligned}$$

Since  $\vec{f}_i$  is nonzero, we obtain

$$t_i = \frac{\vec{x} \cdot \vec{f}_i}{\|\vec{f}_i\|^2}.$$

The result now follows. ■

## Remark

If  $\{\vec{f}_1, \vec{f}_2, \dots, \vec{f}_m\}$  is an orthonormal basis, then the Fourier coefficients are simply  $t_j = \vec{x} \cdot \vec{f}_j$ ,  $j = 1, 2, \dots, m$ .

### Problem

$$\text{Let } \vec{f}_1 = \begin{bmatrix} 1 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \quad \vec{f}_2 = \begin{bmatrix} 1 \\ -1 \\ 0 \\ 0 \end{bmatrix}, \quad \vec{f}_3 = \begin{bmatrix} 0 \\ 0 \\ 1 \\ 1 \end{bmatrix}, \quad \vec{f}_4 = \begin{bmatrix} 0 \\ 0 \\ 1 \\ -1 \end{bmatrix}.$$

Show that  $B = \{\vec{f}_1, \vec{f}_2, \vec{f}_3, \vec{f}_4\}$  is an orthogonal basis of  $\mathbb{R}^4$ , and express  $\vec{x} = \begin{bmatrix} a & b & c & d \end{bmatrix}^T$  as a linear combination of  $\vec{f}_1, \vec{f}_2, \vec{f}_3$  and  $\vec{f}_4$ .

### Problem

$$\text{Let } \vec{f}_1 = \begin{bmatrix} 1 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \quad \vec{f}_2 = \begin{bmatrix} 1 \\ -1 \\ 0 \\ 0 \end{bmatrix}, \quad \vec{f}_3 = \begin{bmatrix} 0 \\ 0 \\ 1 \\ 1 \end{bmatrix}, \quad \vec{f}_4 = \begin{bmatrix} 0 \\ 0 \\ 1 \\ -1 \end{bmatrix}.$$

Show that  $B = \{\vec{f}_1, \vec{f}_2, \vec{f}_3, \vec{f}_4\}$  is an orthogonal basis of  $\mathbb{R}^4$ , and express  $\vec{x} = \begin{bmatrix} a & b & c & d \end{bmatrix}^T$  as a linear combination of  $\vec{f}_1, \vec{f}_2, \vec{f}_3$  and  $\vec{f}_4$ .

### Solution

Computing  $\vec{f}_i \cdot \vec{f}_j$  for  $1 \leq i < j \leq 4$  gives us

$$\begin{aligned} \vec{f}_1 \cdot \vec{f}_2 &= 0, & \vec{f}_1 \cdot \vec{f}_3 &= 0, & \vec{f}_1 \cdot \vec{f}_4 &= 0, \\ \vec{f}_2 \cdot \vec{f}_3 &= 0, & \vec{f}_2 \cdot \vec{f}_4 &= 0, & \vec{f}_3 \cdot \vec{f}_4 &= 0. \end{aligned}$$

Hence,  $B$  is an orthogonal set.



### Problem

$$\text{Let } \vec{f}_1 = \begin{bmatrix} 1 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \quad \vec{f}_2 = \begin{bmatrix} 1 \\ -1 \\ 0 \\ 0 \end{bmatrix}, \quad \vec{f}_3 = \begin{bmatrix} 0 \\ 0 \\ 1 \\ 1 \end{bmatrix}, \quad \vec{f}_4 = \begin{bmatrix} 0 \\ 0 \\ 1 \\ -1 \end{bmatrix}.$$

Show that  $B = \{\vec{f}_1, \vec{f}_2, \vec{f}_3, \vec{f}_4\}$  is an orthogonal basis of  $\mathbb{R}^4$ , and express  $\vec{x} = \begin{bmatrix} a & b & c & d \end{bmatrix}^T$  as a linear combination of  $\vec{f}_1, \vec{f}_2, \vec{f}_3$  and  $\vec{f}_4$ .

### Solution

Computing  $\vec{f}_i \cdot \vec{f}_j$  for  $1 \leq i < j \leq 4$  gives us

$$\begin{aligned} \vec{f}_1 \cdot \vec{f}_2 &= 0, & \vec{f}_1 \cdot \vec{f}_3 &= 0, & \vec{f}_1 \cdot \vec{f}_4 &= 0, \\ \vec{f}_2 \cdot \vec{f}_3 &= 0, & \vec{f}_2 \cdot \vec{f}_4 &= 0, & \vec{f}_3 \cdot \vec{f}_4 &= 0. \end{aligned}$$

Hence,  $B$  is an orthogonal set. It follows that  $B$  is independent, and since  $|B| = 4 = \dim(\mathbb{R}^4)$ ,  $B$  also spans  $\mathbb{R}^4$ . Therefore,  $B$  is an orthogonal basis of  $\mathbb{R}^4$ .

### Problem

$$\text{Let } \vec{f}_1 = \begin{bmatrix} 1 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \quad \vec{f}_2 = \begin{bmatrix} 1 \\ -1 \\ 0 \\ 0 \end{bmatrix}, \quad \vec{f}_3 = \begin{bmatrix} 0 \\ 0 \\ 1 \\ 1 \end{bmatrix}, \quad \vec{f}_4 = \begin{bmatrix} 0 \\ 0 \\ 1 \\ -1 \end{bmatrix}.$$

Show that  $B = \{\vec{f}_1, \vec{f}_2, \vec{f}_3, \vec{f}_4\}$  is an orthogonal basis of  $\mathbb{R}^4$ , and express  $\vec{x} = \begin{bmatrix} a & b & c & d \end{bmatrix}^T$  as a linear combination of  $\vec{f}_1, \vec{f}_2, \vec{f}_3$  and  $\vec{f}_4$ .

### Solution

Computing  $\vec{f}_i \cdot \vec{f}_j$  for  $1 \leq i < j \leq 4$  gives us

$$\begin{aligned} \vec{f}_1 \cdot \vec{f}_2 &= 0, & \vec{f}_1 \cdot \vec{f}_3 &= 0, & \vec{f}_1 \cdot \vec{f}_4 &= 0, \\ \vec{f}_2 \cdot \vec{f}_3 &= 0, & \vec{f}_2 \cdot \vec{f}_4 &= 0, & \vec{f}_3 \cdot \vec{f}_4 &= 0. \end{aligned}$$

Hence,  $B$  is an orthogonal set. It follows that  $B$  is independent, and since  $|B| = 4 = \dim(\mathbb{R}^4)$ ,  $B$  also spans  $\mathbb{R}^4$ . Therefore,  $B$  is an orthogonal basis of  $\mathbb{R}^4$ . By the Fourier Expansion Theorem,

$$\vec{x} = \left(\frac{a+b}{2}\right) \vec{f}_1 + \left(\frac{a-b}{2}\right) \vec{f}_2 + \left(\frac{c+d}{2}\right) \vec{f}_3 + \left(\frac{c-d}{2}\right) \vec{f}_4.$$

