

# Math 221: LINEAR ALGEBRA

## Chapter 3. Determinants and Diagonalization

### §3-2. Determinants and Matrix Inverses

Le Chen<sup>1</sup>

Emory University, 2021 Spring

(last updated on 01/12/2023)



Creative Commons License  
(CC BY-NC-SA)

<sup>1</sup>Slides are adapted from those by Karen Seyffarth from University of Calgary.



# Linear Algebra with Applications

## Lecture Notes

### Current Lecture Notes Revision: Version 2018 — Revision B

These lecture notes were originally developed by Karen Seyffarth of the University of Calgary. Edits, additions, and revisions have been made to these notes by the editorial team at Lyryx Learning to accompany their text [Linear Algebra with Applications](#) based on W. K. Nicholson's original text.

In addition we recognize the following contributors. All new content contributed is released under the same license as noted below.

- Ilijas Farah, York University

### BE A CHAMPION OF OER!

Contribute suggestions for improvements, new content, or errata:

A new topic

A new example or problem

A new or better proof to an existing theorem

Any other suggestions to improve the material

Contact Lyryx at [info@lyryx.com](mailto:info@lyryx.com) with your ideas.

### License



Attribution-NonCommercial-ShareAlike (CC BY-NC-SA)

This license lets others remix, tweak, and build upon your work non-commercially, as long as they credit you and license their new creations under the identical terms.

Copyright

Determinants and Matrix Inverses

Adjugates

Cramer's Rule

Polynomial Interpolation and Vandermonde Determinant

Copyright

## Determinants and Matrix Inverses

Adjugates

Cramer's Rule

Polynomial Interpolation and Vandermonde Determinant

# Determinants and Matrix Inverses

## Theorem (Product Theorem)

If  $A$  and  $B$  are  $n \times n$  matrices, then

$$\det(AB) = \det A \det B.$$

**Proof.**

If either  $A$  or  $B$  is singular, then both sides are equal to zero.

Now assume that both  $A$  and  $B$  are nonsingular, i.e.,  $\text{rank}(A) = \text{rank}(B) = n$ . Then

$$\text{rref}(A) = \text{rref}(B) = I$$

and

$$A = E_1 E_2 \cdots E_p \quad \text{and} \quad B = F_1 F_2 \cdots F_q.$$

where  $E_i$  and  $F_j$  are elementary matrices. Then by the relation of elementary row operations with determinants (Theorem 3.1.2), we see that

$$\begin{aligned} |AB| &= |E_1 \cdots E_p F_1 \cdots F_q| \\ &= |E_1| \cdots |E_p| |F_1| \cdots |F_q| \\ &= |E_1 \cdots E_p| |F_1 \cdots F_q| \\ &= |A| |B|. \end{aligned}$$



## Theorem (Determinant of Matrix Inverse)

An  $n \times n$  matrix  $A$  is invertible if and only if  $\det A \neq 0$ . In this case,

$$\det(A^{-1}) = (\det A)^{-1} = \frac{1}{\det A}.$$

Proof.

" $\Rightarrow$ ":

$$1 = |I| = |AA^{-1}| = |A||A^{-1}| \quad \Rightarrow \quad \begin{cases} |A| \neq 0 \\ |A^{-1}| = \frac{1}{|A|}. \end{cases}$$

" $\Leftarrow$ ": If  $|A| \neq 0$ , then  $\text{rref}(A) = I$  because otherwise one obtains contradiction by Theorem 3.1.2. This is another way to say that  $A$  is invertible: (recall the matrix inverse algorithm)

$$[A|I] \rightarrow \left[ \underbrace{\text{rref}(A)}_{= I} \middle| A^{-1} \right].$$



### Example

Find all values of  $c$  for which  $A = \begin{bmatrix} c & 1 & 0 \\ 0 & 2 & c \\ -1 & c & 5 \end{bmatrix}$  is invertible.

$$\begin{aligned}\det A &= \begin{vmatrix} c & 1 & 0 \\ 0 & 2 & c \\ -1 & c & 5 \end{vmatrix} = c \begin{vmatrix} 2 & c \\ c & 5 \end{vmatrix} + (-1) \begin{vmatrix} 1 & 0 \\ 2 & c \end{vmatrix} \\ &= c(10 - c^2) - c = c(9 - c^2) = c(3 - c)(3 + c).\end{aligned}$$

Therefore,  $A$  is invertible for all  $c \neq 0, 3, -3$ .



## Theorem (Determinant of Matrix Transpose)

If  $A$  is an  $n \times n$  matrix, then  $\det(A^T) = \det A$ .

Proof.

1. This is trivially true for all elementary matrices.
2. If  $A$  is not invertible, then neither is  $A^T$ . Hence,  $\det A = 0 = \det A^T$ .
3. If  $A$  is invertible, then  $A = E_k E_{k-1} \cdots E_2 E_1$ . Hence, by Case 1,

$$\begin{aligned} |A^T| &= |(E_k E_{k-1} \cdots E_2 E_1)^T| \\ &= |E_1^T E_2^T \cdots E_{k-1}^T E_k^T| \\ &= |E_1^T| |E_2^T| \cdots |E_{k-1}^T| |E_k^T| \\ &= |E_1| |E_2| \cdots |E_{k-1}| |E_k| \\ &= |E_k| |E_{k-1}| \cdots |E_2| |E_1| \\ &= |E_k E_{k-1} \cdots E_2 E_1| \\ &= |A|. \end{aligned}$$



## Problem

Suppose  $A$  is a  $3 \times 3$  matrix. Find  $\det A$  and  $\det B$  if

$$\det(2A^{-1}) = -4 = \det(A^3(B^{-1})^T).$$

## Solution

First,

$$\begin{aligned}\det(2A^{-1}) &= -4 \\ 2^3 \det(A^{-1}) &= -4 \\ \frac{1}{\det A} &= \frac{-4}{8} = -\frac{1}{2}\end{aligned}$$

Therefore,  $\det A = -2$ .

## Solution (continued)

Now,

$$\begin{aligned}\det(\mathbf{A}^3(\mathbf{B}^{-1})^T) &= -4 \\ (\det \mathbf{A})^3 \det(\mathbf{B}^{-1}) &= -4 \\ (-2)^3 \det(\mathbf{B}^{-1}) &= -4 \\ (-8) \det(\mathbf{B}^{-1}) &= -4 \\ \frac{1}{\det \mathbf{B}} &= \frac{-4}{-8} = \frac{1}{2}\end{aligned}$$

Therefore,  $\det \mathbf{B} = 2$ .



## Problem

Suppose  $A$ ,  $B$  and  $C$  are  $4 \times 4$  matrices with

$$\det A = -1, \det B = 2, \quad \text{and} \quad \det C = 1.$$

Find  $\det(2A^2(B^{-1})(C^T)^3B(A^{-1}))$ .

## Solution

$$\begin{aligned}\det(2A^2(B^{-1})(C^T)^3B(A^{-1})) &= 2^4(\det A)^2 \frac{1}{\det B} (\det C)^3 (\det B) \frac{1}{\det A} \\ &= 16(\det A)(\det C)^3 \\ &= 16 \times (-1) \times 1^3 \\ &= -16.\end{aligned}$$




## Problem

A square matrix  $A$  is **orthogonal** if and only if  $A^T = A^{-1}$ . What are the possible values of  $\det A$  if  $A$  is orthogonal?

## Solution

Since  $A^T = A^{-1}$ ,

$$\begin{aligned}\det A^T &= \det(A^{-1}) \\ \det A &= \frac{1}{\det A} \\ (\det A)^2 &= 1\end{aligned}$$

Assuming  $A$  is a **real** matrix, this implies that  $\det A = \pm 1$ , i.e.,  $\det A = 1$  or  $\det A = -1$ . 

Copyright

Determinants and Matrix Inverses

**Adjugates**

Cramer's Rule

Polynomial Interpolation and Vandermonde Determinant

# Adjugates

For a  $2 \times 2$  matrix  $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ , we have already seen the **adjugate** of  $A$  defined as

$$\text{adj}(A) = \begin{bmatrix} d & -b \\ -c & a \end{bmatrix},$$

and observed that

$$\begin{aligned} A \text{adj}(A) &= \begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix} \\ &= \begin{bmatrix} ad - bc & 0 \\ 0 & ad - bc \end{bmatrix} \\ &= (\det A)I_2 \end{aligned}$$

Furthermore, if  $\det A \neq 0$ , then  $A$  is invertible and

$$A^{-1} = \frac{1}{\det A} \text{adj}(A).$$

### Definition (Adjugate Matrix)

If  $A$  is an  $n \times n$  matrix, then the **adjugate matrix of  $A$**  is defined to be

$$\text{adj}(A) \stackrel{\text{def}}{=} [c_{ij}(A)]^T = [(-1)^{i+j} \det(A_{ij})]^T,$$

where  $c_{ij}(A)$  is the  $(i,j)$ -cofactor of  $A$ , i.e.,  $\text{adj}(A)$  is the transpose of the **cofactor matrix** (matrix of cofactors).



## Problem

Find  $\text{adj}(A)$  when  $A = \begin{bmatrix} 2 & 1 & 3 \\ 5 & -7 & 1 \\ 3 & 0 & -6 \end{bmatrix}$  and compute  $A \text{adj}(A)$ .

## Solution

$$\text{adj}(A) = \begin{bmatrix} 42 & 6 & 22 \\ 33 & -21 & 13 \\ 21 & 3 & -19 \end{bmatrix}.$$

Notice that

$$A \text{adj}(A) = \begin{bmatrix} 2 & 1 & 3 \\ 5 & -7 & 1 \\ 3 & 0 & -6 \end{bmatrix} \begin{bmatrix} 42 & 6 & 22 \\ 33 & -21 & 13 \\ 21 & 3 & -19 \end{bmatrix} = \begin{bmatrix} 180 & 0 & 0 \\ 0 & 180 & 0 \\ 0 & 0 & 180 \end{bmatrix}$$

$$\det A = \begin{vmatrix} 2 & 1 & 3 \\ 5 & -7 & 1 \\ 3 & 0 & -6 \end{vmatrix} = \begin{vmatrix} 2 & 1 & 3 \\ 19 & 0 & 22 \\ 3 & 0 & -6 \end{vmatrix} = (-1) \begin{vmatrix} 19 & 22 \\ 3 & -6 \end{vmatrix} = 180,$$

Therefore,

$$A \text{adj}(A) = (\det A)I.$$



## Theorem (The Adjugate Formula)

If  $A$  is an  $n \times n$  matrix, then

$$A \operatorname{adj}(A) = (\det A)I = \operatorname{adj}(A)A.$$

Furthermore, if  $\det A \neq 0$ , then

$$A^{-1} = \frac{1}{\det A} \operatorname{adj}(A).$$

### Proof.

We only prove the case when  $n = 3$ .

$$A \operatorname{adj}(A) = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix} \begin{bmatrix} c_{11} & c_{21} & c_{31} \\ c_{12} & c_{22} & c_{32} \\ c_{13} & c_{23} & c_{33} \end{bmatrix} = \begin{bmatrix} |A| & 0 & 0 \\ 0 & |A| & 0 \\ 0 & 0 & |A| \end{bmatrix}$$

where, for example,

$$\begin{aligned} (3,2)\text{-th entry} &= a_{31}c_{21} + a_{32}c_{22} + a_{33}c_{23} \\ &= \det \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{31} & a_{32} & a_{33} \\ a_{31} & a_{32} & a_{33} \end{bmatrix} = 0. \end{aligned}$$



### Example

For an  $n \times n$  matrix  $A$ , show that  $\det \operatorname{adj}(A) = (\det A)^{n-1}$ .

Using the adjugate formula,

$$\begin{aligned} A \operatorname{adj}(A) &= (\det A)I \\ \det(A \operatorname{adj}(A)) &= \det((\det A)I) \\ (\det A) \times \det \operatorname{adj}(A) &= (\det A)^n (\det I) \\ (\det A) \times \det \operatorname{adj}(A) &= (\det A)^n \end{aligned}$$

If  $\det A \neq 0$ , then divide both sides of the last equation by  $\det A$ :

$$\det \operatorname{adj}(A) = (\det A)^{n-1}.$$

### Example (continued)

For the case  $\det A = 0$ , we claim that

$$\det A = 0 \quad \Rightarrow \quad \det \operatorname{adj}(A) = 0, \quad (\star)$$


which implies that

$$\det \operatorname{adj}(A) = 0 = 0^{n-1} = (\det A)^{n-1}.$$

### Proof. (of $(\star)$ )

We will prove  $(\star)$  by contradiction. Indeed, if  $\det A = 0$ , then

$$A \operatorname{adj}(A) = (\det A)I = (0)I = O,$$

i.e.,  $A \operatorname{adj}(A)$  is the zero matrix. If  $\det \operatorname{adj}(A) \neq 0$ , then  $\operatorname{adj}(A)$  would be invertible, and  $A \operatorname{adj}(A) = O$  would imply  $A = O$ . However, if  $A = O$ , then  $\operatorname{adj}(A) = O$  and is not invertible, and thus has determinant equal to zero, i.e.,  $\det \operatorname{adj}(A) = 0$ , (a contradiction!) Therefore,  $\det \operatorname{adj}(A) = 0$ , i.e.,  $(\star)$  is true. 

## Problem

Let  $A$  and  $B$  be  $n \times n$  matrices. Show that  $\det(A + B^T) = \det(A^T + B)$ .

## Solution

Notice that

$$(A + B^T)^T = A^T + (B^T)^T = A^T + B.$$

Since a matrix and its transpose have the same determinant

$$\det(A + B^T) = \det((A + B^T)^T) = \det(A^T + B).$$



## Problem

For each of the following statements, determine if it is true or false, and supply a proof or a counterexample.

1. If  $\text{adj}(A)$  exists, then  $A$  is invertible.
2. If  $A$  and  $B$  are  $n \times n$  matrices, then  $\det(AB) = \det(B^T A)$ .

## Problem

Prove or give a counterexample to the following statement:

$$\text{If } \det A = 1, \text{ then } \text{adj}(A) = A.$$

Copyright

Determinants and Matrix Inverses

Adjugates

**Cramer's Rule**

Polynomial Interpolation and Vandermonde Determinant

## Cramer's Rule

If  $A$  is an  $n \times n$  **invertible** matrix, then the solution to  $A\vec{x} = \vec{b}$  can be given in terms of determinants of matrices.

### Theorem (Cramer's Rule)

Let  $A$  be an  $n \times n$  invertible matrix, the solution to the system  $A\vec{x} = \vec{b}$  of  $n$  equations in  $n$  variables  $x_1, x_2, \dots, x_n$  is given by

$$x_1 = \frac{\det(A_1(\vec{b}))}{\det A}, \quad x_2 = \frac{\det(A_2(\vec{b}))}{\det A}, \quad \dots, \quad x_n = \frac{\det(A_n(\vec{b}))}{\det A}$$

where, for each  $j$ , the matrix  $A_j(\vec{b})$  is obtained from  $A$  by replacing column  $j$  with  $\vec{b}$ :

$$A_j(\vec{b}) = \begin{bmatrix} \vec{a}_1 & \cdots & \vec{a}_{j-1} & \vec{b} & \vec{a}_{j+1} & \cdots & \vec{a}_n \end{bmatrix}$$



Proof.

► Notice that

$$\begin{aligned} A_j(\vec{b}) &= \begin{bmatrix} \vec{a}_1 & \cdots & \vec{a}_{j-1} & \vec{b} & \vec{a}_{j+1} & \cdots & \vec{a}_n \end{bmatrix} \\ &= \begin{bmatrix} A\vec{e}_1 & \cdots & A\vec{e}_{j-1} & A\vec{x} & A\vec{e}_{j+1} & \cdots & A\vec{e}_n \end{bmatrix} \\ &= A \begin{bmatrix} \vec{e}_1 & \cdots & \vec{e}_{j-1} & \vec{x} & \vec{e}_{j+1} & \cdots & \vec{e}_n \end{bmatrix} \\ &= A I_j(\vec{x}) \end{aligned}$$

where

$$\begin{aligned} I_j(\vec{x}) &= \begin{bmatrix} \vec{e}_1 & \cdots & \vec{e}_{j-1} & \vec{x} & \vec{e}_{j+1} & \cdots & \vec{e}_n \end{bmatrix} \\ &= \begin{bmatrix} 1 & & & x_1 & & & \\ & \ddots & & \vdots & & & \\ & & 1 & x_{j-1} & & & \\ & & & x_j & & & \\ & & & x_{j+1} & 1 & & \\ & & & \vdots & & \ddots & \\ & & & x_n & & & 1 \end{bmatrix} \end{aligned}$$

Proof. (continued)

- Hence, by taking the determinants on both sides, we have

$$\begin{aligned}\det(A_j(\vec{b})) &= \det(A \ I_j(\vec{x})) \\ &= \det(A) \det(I_j(\vec{x}))\end{aligned}$$

- And because  $\det(A) \neq 0$ , we can then write:

$$\det(I_j(\vec{x})) = \frac{\det(A_j(\vec{b}))}{\det(A)}$$

- Finally, notice that  $\det(I_j(\vec{x})) = \cdots = x_j$ .



## Problem

Find  $x_3$  such that

$$\begin{array}{ccccccccc} 3x_1 & + & x_2 & - & x_3 & = & -1 \\ 5x_1 & + & 2x_2 & & & = & 2 \\ x_1 & + & x_2 & - & x_3 & = & 1 \end{array}$$

## Solution

By Cramer's rule,  $x_3 = \frac{\det A_3}{\det A}$ , where

$$A = \begin{bmatrix} 3 & 1 & -1 \\ 5 & 2 & 0 \\ 1 & 1 & -1 \end{bmatrix} \quad \text{and} \quad A_3 = \begin{bmatrix} 3 & 1 & -1 \\ 5 & 2 & 2 \\ 1 & 1 & 1 \end{bmatrix}.$$

Computing the determinants of these two matrices,

$$\det A = -4 \quad \text{and} \quad \det A_3 = -6.$$

Therefore,  $x_3 = \frac{-6}{-4} = \frac{3}{2}$ .



### Remark

For practice, you should compute  $\det A_1$  and  $\det A_2$ , where

$$A_1 = \begin{bmatrix} -1 & 1 & -1 \\ 2 & 2 & 0 \\ 1 & 1 & -1 \end{bmatrix} \quad \text{and} \quad A_2 = \begin{bmatrix} 3 & -1 & -1 \\ 5 & 2 & 0 \\ 1 & 1 & -1 \end{bmatrix},$$

and then solve for  $x_1$  and  $x_2$ .

Solution.  $x_1 = -1$ ,  $x_2 = 7/2$ .



Copyright

Determinants and Matrix Inverses

Adjugates


Cramer's Rule

Polynomial Interpolation and Vandermonde Determinant

# Polynomial Interpolation and Vandermonde Determinant

## Problem

Given data points  $(0, 1)$ ,  $(1, 2)$ ,  $(2, 5)$  and  $(3, 10)$ , find an interpolating polynomial  $p(x)$  of degree at most three, and then estimate the value of  $y$  corresponding to  $x = 3/2$ .



## Solution

We want to find the coefficients  $r_0$ ,  $r_1$ ,  $r_2$  and  $r_3$  of

$$p(x) = r_0 + r_1x + r_2x^2 + r_3x^3$$

so that  $p(0) = 1$ ,  $p(1) = 2$ ,  $p(2) = 5$ , and  $p(3) = 10$ .

$$p(0) = r_0 = 1$$

$$p(1) = r_0 + r_1 + r_2 + r_3 = 2$$

$$p(2) = r_0 + 2r_1 + 4r_2 + 8r_3 = 5$$

$$p(3) = r_0 + 3r_1 + 9r_2 + 27r_3 = 10$$

### Solution (continued)

Solve this system of four equations in the four variables  $r_0$ ,  $r_1$ ,  $r_2$  and  $r_3$ .

$$\left[ \begin{array}{cccc|c} 1 & 0 & 0 & 0 & 1 \\ 1 & 1 & 1 & 1 & 2 \\ 1 & 2 & 4 & 8 & 5 \\ 1 & 3 & 9 & 27 & 10 \end{array} \right] \rightarrow \cdots \rightarrow \left[ \begin{array}{cccc|c} 1 & 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 \end{array} \right]$$

Therefore,  $r_0 = 1$ ,  $r_1 = 0$ ,  $r_2 = 1$ ,  $r_3 = 0$ , and so


$$p(x) = 1 + x^2.$$

Finally, the estimate is

$$y = p\left(\frac{3}{2}\right) = 1 + \left(\frac{3}{2}\right)^2 = \frac{13}{4}.$$







## Theorem (Polynomial Interpolation)

Given  $n$  data points  $(x_1, y_1), (x_2, y_2), \dots, (x_n, y_n)$  with the  $x_i$  **distinct**, there is a unique polynomial

$$p(x) = r_0 + r_1x + r_2x^2 + \cdots + r_{n-1}x^{n-1}$$

such that  $p(x_i) = y_i$  for  $i = 1, 2, \dots, n$ .

The polynomial  $p(x)$  is called the **interpolating polynomial** for the data.

To find  $p(x) = r_0 + r_1x + r_2x^2 + \cdots + r_{n-1}x^{n-1}$ , set up a system of  $n$  linear equations in the  $n$  variables  $r_0, r_1, r_2, \dots, r_{n-1}$ .

$$\begin{array}{rcl}
 r_0 + r_1x_1 + r_2x_1^2 + \cdots + r_{n-1}x_1^{n-1} & = & y_1 \\
 r_0 + r_1x_2 + r_2x_2^2 + \cdots + r_{n-1}x_2^{n-1} & = & y_2 \\
 r_0 + r_1x_3 + r_2x_3^2 + \cdots + r_{n-1}x_3^{n-1} & = & y_3 \\
 \vdots & & \vdots \\
 r_0 + r_1x_n + r_2x_n^2 + \cdots + r_{n-1}x_n^{n-1} & = & y_n
 \end{array}$$

The coefficient matrix for this system is


$$\begin{bmatrix}
 1 & x_1 & x_1^2 & \cdots & x_1^{n-1} \\
 1 & x_2 & x_2^2 & \cdots & x_2^{n-1} \\
 \vdots & \vdots & \vdots & \vdots & \vdots \\
 1 & x_n & x_n^2 & \cdots & x_n^{n-1}
 \end{bmatrix}$$

- Such matrix is called **Vandermonde matrix**.
- Its determinant is called **Vandermonde determinant**.

## Theorem (Vandermonde Determinant )

Let  $a_1, a_2, \dots, a_n$  be real numbers,  $n \geq 2$ . The corresponding Vandermonde determinant is

$$\det \begin{bmatrix} 1 & a_1 & a_1^2 & \cdots & a_1^{n-1} \\ 1 & a_2 & a_2^2 & \cdots & a_2^{n-1} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & a_n & a_n^2 & \cdots & a_n^{n-1} \end{bmatrix} = \prod_{1 \leq j < i \leq n} (a_i - a_j).$$



Proof.

We will prove this by induction. It is clear that when  $n = 2$ ,

$$\det \begin{pmatrix} 1 & a_1 \\ 1 & a_2 \end{pmatrix} = a_2 - a_1 = \prod_{1 \leq j < i \leq 2} (a_i - a_j).$$

Assume that it is true for  $n - 1$ . Now let's consider the case  $n$ . Denote

$$p(\mathbf{x}) := \det \begin{bmatrix} 1 & a_1 & a_1^2 & \cdots & a_1^{n-1} \\ 1 & a_2 & a_2^2 & \cdots & a_2^{n-1} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 1 & a_{n-1} & a_{n-1}^2 & \cdots & a_{n-1}^{n-1} \\ \mathbf{1} & \mathbf{x} & \mathbf{x}^2 & \cdots & \mathbf{x}^{n-1} \end{bmatrix}.$$

Proof. (continued)

Because  $p(a_1) = \cdots = p(a_{n-1}) = 0$  (why?),  $p(x)$  has to take the following form:

$$p(x) = c(x - a_1)(x - a_2) \cdots (x - a_{n-1}).$$


To identify the constant  $c$ , notice that  $c$  is the coefficient for  $x^{n-1}$ . By cofactor expansion of the determinant along the last row,

$$\begin{aligned} c &= (-1)^{n+n} \det \begin{bmatrix} 1 & a_1 & a_1^2 & \cdots & a_1^{n-1} \\ 1 & a_2 & a_2^2 & \cdots & a_2^{n-1} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 1 & a_{n-1} & a_{n-1}^2 & \cdots & a_{n-1}^{n-1} \end{bmatrix} \\ &= \prod_{1 \leq j < i \leq n-1} (a_i - a_j). \end{aligned}$$

Proof. (continued)

Hence,

$$p(a_n) = \left( \prod_{1 \leq j < i \leq n-1} (a_i - a_j) \right) \times (a_n - a_1)(a_n - a_2) \cdots (a_n - a_{n-1})$$



$$p(a_n) = \prod_{1 \leq j < i \leq n} (a_i - a_j).$$



## Example

In our earlier example with the data points  $(0, 1)$ ,  $(1, 2)$ ,  $(2, 5)$  and  $(3, 10)$ , we have

$$a_1 = 0, \quad a_2 = 1, \quad a_3 = 2, \quad a_4 = 3$$

giving us the Vandermonde determinant

$$\begin{vmatrix} 1 & 0 & 0 & 0 \\ 1 & 1 & 1 & 1 \\ 1 & 2 & 4 & 8 \\ 1 & 3 & 9 & 27 \end{vmatrix}$$

According to the previous theorem, this determinant is equal to

$$\begin{aligned} & (a_2 - a_1)(a_3 - a_1)(a_3 - a_2)(a_4 - a_1)(a_4 - a_2)(a_4 - a_3) \\ &= (1 - 0)(2 - 0)(2 - 1)(3 - 0)(3 - 1)(3 - 2) \\ &= 2 \times 3 \times 2 \\ &= 12. \end{aligned}$$



## Corollary

The Vandermonde determinant is nonzero if  $a_1, a_2, \dots, a_n$  are distinct.

This means that given  $n$  data points  $(x_1, y_1), (x_2, y_2), \dots, (x_n, y_n)$  with **distinct**  $x_i$ , then there is a unique interpolating polynomial

$$p(x) = r_0 + r_1x + r_2x^2 + \cdots + r_{n-1}x^{n-1}.$$