

Math 221: LINEAR ALGEBRA

Chapter 3. Determinants and Diagonalization

§3-1. The Cofactor Expansion

Le Chen¹

Emory University, 2021 Spring

(last updated on 01/12/2023)



Creative Commons License
(CC BY-NC-SA)

¹Slides are adapted from those by Karen Seyffarth from University of Calgary.

Linear Algebra with Applications

Lecture Notes

Current Lecture Notes Revision: Version 2018 — Revision B

These lecture notes were originally developed by Karen Seyffarth of the University of Calgary. Edits, additions, and revisions have been made to these notes by the editorial team at Lyryx Learning to accompany their text [Linear Algebra with Applications](#) based on W. K. Nicholson's original text.

In addition we recognize the following contributors. All new content contributed is released under the same license as noted below.

- Ilijas Farah, York University

BE A CHAMPION OF OER!

Contribute suggestions for improvements, new content, or errata:

A new topic

A new example or problem

A new or better proof to an existing theorem

Any other suggestions to improve the material

Contact Lyryx at info@lyryx.com with your ideas.

License



Attribution-NonCommercial-ShareAlike (CC BY-NC-SA)

This license lets others remix, tweak, and build upon your work non-commercially, as long as they credit you and license their new creations under the identical terms.

Copyright

Determinant of Small Matrices

The Cofactor Expansion

Elementary Row Operations and Determinants

Determinant and Scalar Multiple

Determinant of Triangular Matrices

Determinant of Block Matrices

Some More Exercises

Copyright

Determinant of Small Matrices

The Cofactor Expansion

Elementary Row Operations and Determinants

Determinant and Scalar Multiple

Determinant of Triangular Matrices

Determinant of Block Matrices

Some More Exercises

Determinant of Small Matrices

Determinant of Small Matrices

Recall that if $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$, then the **determinant** of A is defined as

$$\det A = ad - bc,$$

and that A is invertible if and only if $\det A \neq 0$.


Determinant of Small Matrices

Recall that if $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$, then the **determinant** of A is defined as

$$\det A = ad - bc,$$

and that A is invertible if and only if $\det A \neq 0$.


Notation: For $\det \begin{bmatrix} a & b \\ c & d \end{bmatrix}$, we often write $\begin{vmatrix} a & b \\ c & d \end{vmatrix}$, i.e., use **vertical bars** instead of **square brackets**.



$$\det \begin{pmatrix} a & c \\ b & d \end{pmatrix} = \text{signed area of parallelogram}$$


Problem

How to define determinant for a general $n \times n$ matrix?




$\det \begin{pmatrix} \vec{r}_1 & \vec{r}_2 & \vec{r}_3 \end{pmatrix} = \text{signed volume of the parallelepiped}$

4×4




Only partially right... still missing many terms...


4×4 part I:




4×4 part II:



4×4 part III:



4×4 part IV:



$5 \times 5?$...

The determinant of an $n \times n$ matrix is more effectively defined through recursion...

Copyright

Determinant of Small Matrices

The Cofactor Expansion

Elementary Row Operations and Determinants

Determinant and Scalar Multiple

Determinant of Triangular Matrices

Determinant of Block Matrices

Some More Exercises

Cofactor and cofactor expansion

Definition

Let $A = [a_{ij}]$ be an $n \times n$ matrix.

- The **sign** of the (i,j) position is $(-1)^{i+j}$. (Thus the sign is 1 if $(i+j)$ is even, and -1 if $(i+j)$ is odd.)

$$\begin{bmatrix} a_{11} & a_{12} & a_{13} & \cdots \\ a_{21} & a_{22} & a_{23} & \cdots \\ a_{31} & a_{32} & a_{33} & \cdots \\ \vdots & \vdots & \vdots & \end{bmatrix} \Rightarrow \begin{bmatrix} + & - & + & \cdots \\ - & + & - & \cdots \\ + & - & + & \cdots \\ \vdots & \vdots & \vdots & \end{bmatrix}$$

Definition (continued)

- Let A_{ij} denote the $(n-1) \times (n-1)$ matrix obtained from A by deleting row i and column j . The (i,j) -cofactor of A is

$$c_{ij}(A) = (-1)^{i+j} \det(A_{ij}).$$

- The determinant of A is defined as

$$\det A = a_{11}c_{11}(A) + a_{12}c_{12}(A) + a_{13}c_{13}(A) + \cdots + a_{1n}c_{1n}(A)$$

and is called the cofactor expansion of $\det A$ along row 1.

Example

Let $A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix}$. Find $\det A$.

Using cofactor expansion along row 1,

$$\begin{aligned}\det A &= 1c_{11}(A) + 2c_{12}(A) + 3c_{13}(A) \\&= 1(-1)^2 \begin{vmatrix} 5 & 6 \\ 8 & 9 \end{vmatrix} + 2(-1)^3 \begin{vmatrix} 4 & 6 \\ 7 & 9 \end{vmatrix} + 3(-1)^4 \begin{vmatrix} 4 & 5 \\ 7 & 8 \end{vmatrix} \\&= (45 - 48) - 2(36 - 42) + 3(32 - 35) \\&= -3 - 2(-6) + 3(-3) \\&= -3 + 12 - 9 \\&= 0\end{aligned}$$

Example (continued)

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix}$$

Now try cofactor expansion along column 2.

Example (continued)

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix}$$

Now try cofactor expansion along column 2.

$$\begin{aligned} \det A &= 2c_{12}(A) + 5c_{22}(A) + 8c_{32}(A) \\ &= 2(-1)^3 \begin{vmatrix} 4 & 6 \\ 7 & 9 \end{vmatrix} + 5(-1)^4 \begin{vmatrix} 1 & 3 \\ 7 & 9 \end{vmatrix} + 8(-1)^5 \begin{vmatrix} 1 & 3 \\ 4 & 6 \end{vmatrix} \\ &= -2(36 - 42) + 5(9 - 21) - 8(6 - 12) \\ &= -2(-6) + 5(-12) - 8(-6) \\ &= 12 - 60 + 48 \\ &= 0. \end{aligned}$$

Example (continued)

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix}$$

Now try cofactor expansion along column 2.

$$\begin{aligned} \det A &= 2c_{12}(A) + 5c_{22}(A) + 8c_{32}(A) \\ &= 2(-1)^3 \begin{vmatrix} 4 & 6 \\ 7 & 9 \end{vmatrix} + 5(-1)^4 \begin{vmatrix} 1 & 3 \\ 7 & 9 \end{vmatrix} + 8(-1)^5 \begin{vmatrix} 1 & 3 \\ 4 & 6 \end{vmatrix} \\ &= -2(36 - 42) + 5(9 - 21) - 8(6 - 12) \\ &= -2(-6) + 5(-12) - 8(-6) \\ &= 12 - 60 + 48 \\ &= 0. \end{aligned}$$

We get the same answer!

Theorem (Cofactor Expansion Theorem)

The determinant of an $n \times n$ matrix A can be computed using the cofactor expansion along **any row or column** of A .

Theorem (Cofactor Expansion Theorem)

The determinant of an $n \times n$ matrix A can be computed using the cofactor expansion along **any row or column** of A .

Example

$$\text{Let } A = \begin{bmatrix} 0 & 1 & 2 & 1 \\ 5 & 0 & 0 & 7 \\ 0 & 1 & -1 & 0 \\ 3 & 0 & 0 & 2 \end{bmatrix}. \text{ Cofactor expansion along row 1 yields}$$

$$\begin{aligned} \det A &= 0c_{11}(A) + 1c_{12}(A) + 2c_{13}(A) + 1c_{14}(A) \\ &= 1c_{12}(A) + 2c_{13}(A) + c_{14}(A), \end{aligned}$$

whereas cofactor expansion along, row 3 yields

$$\begin{aligned} \det A &= 0c_{31}(A) + 1c_{32}(A) + (-1)c_{33}(A) + 0c_{34}(A) \\ &= 1c_{32}(A) + (-1)c_{33}(A), \end{aligned}$$

i.e., in the first case we have to compute three cofactors, but in the second we only have to compute two.


Example (continued)

We can save ourselves some work by using cofactor expansion along row 3 rather than row 1.

$$A = \begin{bmatrix} 0 & 1 & 2 & 1 \\ 5 & 0 & 0 & 7 \\ 0 & \textcolor{red}{1} & \textcolor{red}{-1} & 0 \\ 3 & 0 & 0 & 2 \end{bmatrix}$$

$$\begin{aligned} \det A &= \textcolor{red}{1}c_{32}(A) + (\textcolor{red}{-1})c_{33}(A) \\ &= 1(-1)^5 \begin{vmatrix} 0 & 2 & 1 \\ 5 & 0 & 7 \\ 3 & 0 & 2 \end{vmatrix} + (-1)(-1)^6 \begin{vmatrix} 0 & 1 & 1 \\ 5 & 0 & 7 \\ 3 & 0 & 2 \end{vmatrix} \\ &= (-1)2(-1)^3 \begin{vmatrix} 5 & 7 \\ 3 & 2 \end{vmatrix} + (-1)1(-1)^3 \begin{vmatrix} 5 & 7 \\ 3 & 2 \end{vmatrix} \\ &= 2(10 - 21) + 1(10 - 21) \\ &= 2(-11) + (-11) \\ &= -33. \end{aligned}$$

Example (continued)

Try computing $\det \begin{bmatrix} 0 & 1 & 2 & 1 \\ 5 & 0 & 0 & 7 \\ 0 & 1 & -1 & 0 \\ 3 & 0 & 0 & 2 \end{bmatrix}$ using cofactor expansion along other rows and columns, for instance column 2 or row 4. You will still get $\det A = -33$. 

Problem

Find $\det A$ for $A = \begin{bmatrix} -8 & 1 & 0 & -4 \\ 5 & 7 & 0 & -7 \\ 12 & -3 & 0 & 8 \\ -3 & 11 & 0 & 2 \end{bmatrix}$.

Problem

Find $\det A$ for $A = \begin{bmatrix} -8 & 1 & 0 & -4 \\ 5 & 7 & 0 & -7 \\ 12 & -3 & 0 & 8 \\ -3 & 11 & 0 & 2 \end{bmatrix}$.

Solution

Using cofactor expansion along column 3, $\det A = 0$.



Problem

Find $\det A$ for $A = \begin{bmatrix} -8 & 1 & 0 & -4 \\ 5 & 7 & 0 & -7 \\ 12 & -3 & 0 & 8 \\ -3 & 11 & 0 & 2 \end{bmatrix}$.

Solution

Using cofactor expansion along column 3, $\det A = 0$. ■

Remark

If A is an $n \times n$ matrix with a row or column of zeros, then $\det A = 0$.

Copyright

Determinant of Small Matrices

The Cofactor Expansion

Elementary Row Operations and Determinants

Determinant and Scalar Multiple

Determinant of Triangular Matrices

Determinant of Block Matrices

Some More Exercises

Elementary Row Operations and Determinants

Elementary Row Operations and Determinants

Example

Let $A = \begin{bmatrix} 2 & 0 & -3 \\ 0 & 4 & 0 \\ 1 & 0 & -2 \end{bmatrix}$. Then

$$\det A = 4(-1)^4 \begin{vmatrix} 2 & -3 \\ 1 & -2 \end{vmatrix} = 4(-1) = -4.$$

Elementary Row Operations and Determinants

Example

$$\text{Let } A = \begin{bmatrix} 2 & 0 & -3 \\ 0 & 4 & 0 \\ 1 & 0 & -2 \end{bmatrix}. \text{ Then}$$

$$\det A = 4(-1)^4 \begin{vmatrix} 2 & -3 \\ 1 & -2 \end{vmatrix} = 4(-1) = -4.$$

Let B_1, B_2 , and B_3 be obtained from A by performing a type 1, 2 and 3 elementary row operation, respectively, i.e.,

$$B_1 = \begin{matrix} r_2 \leftrightarrow r_3 \\ \left[\begin{array}{ccc} 2 & 0 & -3 \\ 1 & 0 & -2 \\ 0 & 4 & 0 \end{array} \right] \end{matrix} \quad \left| \quad \begin{matrix} -3 \times r_3 \rightarrow r_3 \\ \left[\begin{array}{ccc} 2 & 0 & -3 \\ 0 & 4 & 0 \\ -3 & 0 & 6 \end{array} \right] \end{matrix} \quad \left| \quad \begin{matrix} 2 \times r_1 + r_3 \rightarrow r_3 \\ \left[\begin{array}{ccc} 2 & 0 & -3 \\ 0 & 4 & 0 \\ 5 & 0 & -8 \end{array} \right] \end{matrix} \right.$$

Example (continued)

$$\det B_1 = 4(-1)^5 \begin{vmatrix} 2 & -3 \\ 1 & -2 \end{vmatrix} = (-4)(-1) = 4 = (-1) \det A.$$

Example (continued)

$$\det B_1 = 4(-1)^5 \begin{vmatrix} 2 & -3 \\ 1 & -2 \end{vmatrix} = (-4)(-1) = 4 = (-1) \det A.$$

$$\det B_2 = 4(-1)^4 \begin{vmatrix} 2 & -3 \\ -3 & 6 \end{vmatrix} = 4(12 - 9) = 4 \times 3 = 12 = -3 \det A.$$

Example (continued)

$$\det B_1 = 4(-1)^5 \begin{vmatrix} 2 & -3 \\ 1 & -2 \end{vmatrix} = (-4)(-1) = 4 = (-1) \det A.$$

$$\det B_2 = 4(-1)^4 \begin{vmatrix} 2 & -3 \\ -3 & 6 \end{vmatrix} = 4(12 - 9) = 4 \times 3 = 12 = -3 \det A.$$

$$\det B_3 = 4(-1)^4 \begin{vmatrix} 2 & -3 \\ 5 & -8 \end{vmatrix} = 4(-16 + 15) = 4(-1) = -4 = \det A.$$

Theorem (Determinant and Elementary Row Operations)

Let A be an $n \times n$ matrix.

1. If B is obtained from A by exchanging two different rows (or columns) of A , then $\det B = -\det A$.

Theorem (Determinant and Elementary Row Operations)

Let A be an $n \times n$ matrix.

1. If B is obtained from A by exchanging two different rows (or columns) of A , then $\det B = -\det A$.
2. If B is obtained from A by multiplying a row (or column) of A by a scalar $k \in \mathbb{R}$, then $\det B = k \det A$.

Theorem (Determinant and Elementary Row Operations)

Let A be an $n \times n$ matrix.

1. If B is obtained from A by exchanging two different rows (or columns) of A , then $\det B = -\det A$.
2. If B is obtained from A by multiplying a row (or column) of A by a scalar $k \in \mathbb{R}$, then $\det B = k \det A$.
3. If B is obtained from A by adding k times one row of A to a different row of A (or adding k times one column of A to a different column of A) then $\det B = \det A$.

Theorem (Determinant and Elementary Row Operations)

Let A be an $n \times n$ matrix.

1. If B is obtained from A by exchanging two different rows (or columns) of A , then $\det B = -\det A$.
2. If B is obtained from A by multiplying a row (or column) of A by a scalar $k \in \mathbb{R}$, then $\det B = k \det A$.
3. If B is obtained from A by adding k times one row of A to a different row of A (or adding k times one column of A to a different column of A) then $\det B = \det A$.
4. If A has a row or column of zeros, then $\det A = 0$.

Theorem (Determinant and Elementary Row Operations)

Let A be an $n \times n$ matrix.

1. If B is obtained from A by exchanging two different rows (or columns) of A , then $\det B = -\det A$.
2. If B is obtained from A by multiplying a row (or column) of A by a scalar $k \in \mathbb{R}$, then $\det B = k \det A$.
3. If B is obtained from A by adding k times one row of A to a different row of A (or adding k times one column of A to a different column of A) then $\det B = \det A$.
4. If A has a row or column of zeros, then $\det A = 0$.
5. If two different rows (or columns) of A are identical, then $\det A = 0$.

Example

$$\det \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix} = \begin{vmatrix} 1 & 2 & 3 \\ 0 & -3 & -6 \\ 0 & -6 & -12 \end{vmatrix} = \begin{vmatrix} -3 & -6 \\ -6 & -12 \end{vmatrix} = 36 - 36 = 0.$$



Example

$$\begin{aligned}\det \begin{bmatrix} 3 & 1 & 2 & 4 \\ -1 & -3 & 8 & 0 \\ 1 & -1 & 5 & 5 \\ 1 & 1 & 2 & -1 \end{bmatrix} &= \begin{vmatrix} 0 & -8 & 26 & 4 \\ -1 & -3 & 8 & 0 \\ 0 & -4 & 13 & 5 \\ 0 & -2 & 10 & -1 \end{vmatrix} \\ &= (-1)(-1)^3 \begin{vmatrix} -8 & 26 & 4 \\ -4 & 13 & 5 \\ -2 & 10 & -1 \end{vmatrix} \\ &= \begin{vmatrix} 0 & -14 & 8 \\ 0 & -7 & 7 \\ -2 & 10 & -1 \end{vmatrix} \\ &= (-2)(-1)^4 \begin{vmatrix} -14 & 8 \\ -7 & 7 \end{vmatrix} \\ &= -2 \begin{vmatrix} 0 & -6 \\ -7 & 7 \end{vmatrix} \\ &= (-2)(-42) = 84.\end{aligned}$$



Problem

$$\text{If } \det \begin{bmatrix} a & b & c \\ p & q & r \\ x & y & z \end{bmatrix} = -1, \text{ find } \det \begin{bmatrix} -x & -y & -z \\ 3p + a & 3q + b & 3r + c \\ 2p & 2q & 2r \end{bmatrix}.$$

Problem

$$\text{If } \det \begin{bmatrix} a & b & c \\ p & q & r \\ x & y & z \end{bmatrix} = -1, \text{ find } \det \begin{bmatrix} -x & -y & -z \\ 3p + a & 3q + b & 3r + c \\ 2p & 2q & 2r \end{bmatrix}.$$

Problem

$$\text{If } \det \begin{bmatrix} a & b & c \\ p & q & r \\ x & y & z \end{bmatrix} = -1, \text{ find } \det \begin{bmatrix} -x & -y & -z \\ 3p + a & 3q + b & 3r + c \\ 2p & 2q & 2r \end{bmatrix}.$$

Solution

$$\begin{vmatrix} -x & -y & -z \\ 3p + a & 3q + b & 3r + c \\ 2p & 2q & 2r \end{vmatrix} = (-1)(2) \begin{vmatrix} x & y & z \\ 3p + a & 3q + b & 3r + c \\ p & q & r \end{vmatrix}$$

Problem

$$\text{If } \det \begin{bmatrix} a & b & c \\ p & q & r \\ x & y & z \end{bmatrix} = -1, \text{ find } \det \begin{bmatrix} -x & -y & -z \\ 3p + a & 3q + b & 3r + c \\ 2p & 2q & 2r \end{bmatrix}.$$

Solution

$$\begin{aligned} & \begin{vmatrix} -x & -y & -z \\ 3p + a & 3q + b & 3r + c \\ 2p & 2q & 2r \end{vmatrix} = (-1)(2) \begin{vmatrix} x & y & z \\ 3p + a & 3q + b & 3r + c \\ p & q & r \end{vmatrix} \\ & = (-2) \begin{vmatrix} x & y & z \\ a & b & c \\ p & q & r \end{vmatrix} = (-2)(-1) \begin{vmatrix} a & b & c \\ x & y & z \\ p & q & r \end{vmatrix} = 2(-1) \begin{vmatrix} a & b & c \\ p & q & r \\ x & y & z \end{vmatrix} \end{aligned}$$

Problem

$$\text{If } \det \begin{bmatrix} a & b & c \\ p & q & r \\ x & y & z \end{bmatrix} = -1, \text{ find } \det \begin{bmatrix} -x & -y & -z \\ 3p + a & 3q + b & 3r + c \\ 2p & 2q & 2r \end{bmatrix}.$$

Solution

$$\begin{aligned} & \begin{vmatrix} -x & -y & -z \\ 3p + a & 3q + b & 3r + c \\ 2p & 2q & 2r \end{vmatrix} = (-1)(2) \begin{vmatrix} x & y & z \\ 3p + a & 3q + b & 3r + c \\ p & q & r \end{vmatrix} \\ &= (-2) \begin{vmatrix} x & y & z \\ a & b & c \\ p & q & r \end{vmatrix} = (-2)(-1) \begin{vmatrix} a & b & c \\ x & y & z \\ p & q & r \end{vmatrix} = 2(-1) \begin{vmatrix} a & b & c \\ p & q & r \\ x & y & z \end{vmatrix} \\ &= (-2)(-1) = 2. \end{aligned}$$



Example

$$\begin{aligned}\det \begin{bmatrix} 1 & 2 & 3 \\ 0 & 5 & 6 \\ 0 & 0 & 9 \end{bmatrix} &= 1 \det \begin{bmatrix} 5 & 6 \\ 0 & 9 \end{bmatrix} \\ &= (1)(5) \det \begin{bmatrix} 9 \end{bmatrix} \\ &= (1)(5)(9) \\ &= 45.\end{aligned}$$



Copyright

Determinant of Small Matrices

The Cofactor Expansion

Elementary Row Operations and Determinants

Determinant and Scalar Multiple

Determinant of Triangular Matrices

Determinant of Block Matrices

Some More Exercises

Determinant and Scalar Multiple

Problem

Suppose A is a 3×3 matrix with $\det A = 7$. What is $\det(-3A)$?

Determinant and Scalar Multiple

Problem

Suppose A is a 3×3 matrix with $\det A = 7$. What is $\det(-3A)$?

Solution

Write $A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$. Then $-3A = \begin{bmatrix} -3a_{11} & -3a_{12} & -3a_{13} \\ -3a_{21} & -3a_{22} & -3a_{23} \\ -3a_{31} & -3a_{32} & -3a_{33} \end{bmatrix}$.

$$\det(-3A) = \begin{vmatrix} -3a_{11} & -3a_{12} & -3a_{13} \\ -3a_{21} & -3a_{22} & -3a_{23} \\ -3a_{31} & -3a_{32} & -3a_{33} \end{vmatrix} = (-3) \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ -3a_{21} & -3a_{22} & -3a_{23} \\ -3a_{31} & -3a_{32} & -3a_{33} \end{vmatrix}$$

Determinant and Scalar Multiple

Problem

Suppose A is a 3×3 matrix with $\det A = 7$. What is $\det(-3A)$?

Solution

$$\text{Write } A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}. \text{ Then } -3A = \begin{bmatrix} -3a_{11} & -3a_{12} & -3a_{13} \\ -3a_{21} & -3a_{22} & -3a_{23} \\ -3a_{31} & -3a_{32} & -3a_{33} \end{bmatrix}.$$

$$\det(-3A) = \begin{vmatrix} -3a_{11} & -3a_{12} & -3a_{13} \\ -3a_{21} & -3a_{22} & -3a_{23} \\ -3a_{31} & -3a_{32} & -3a_{33} \end{vmatrix} = (-3) \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ -3a_{21} & -3a_{22} & -3a_{23} \\ -3a_{31} & -3a_{32} & -3a_{33} \end{vmatrix}$$

$$= (-3)(-3) \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ -3a_{31} & -3a_{32} & -3a_{33} \end{vmatrix} = (-3)(-3)(-3) \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix}$$

Determinant and Scalar Multiple

Problem

Suppose A is a 3×3 matrix with $\det A = 7$. What is $\det(-3A)$?

Solution

$$\text{Write } A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}. \text{ Then } -3A = \begin{bmatrix} -3a_{11} & -3a_{12} & -3a_{13} \\ -3a_{21} & -3a_{22} & -3a_{23} \\ -3a_{31} & -3a_{32} & -3a_{33} \end{bmatrix}.$$

$$\det(-3A) = \begin{vmatrix} -3a_{11} & -3a_{12} & -3a_{13} \\ -3a_{21} & -3a_{22} & -3a_{23} \\ -3a_{31} & -3a_{32} & -3a_{33} \end{vmatrix} = (-3) \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ -3a_{21} & -3a_{22} & -3a_{23} \\ -3a_{31} & -3a_{32} & -3a_{33} \end{vmatrix}$$

$$= (-3)(-3) \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ -3a_{31} & -3a_{32} & -3a_{33} \end{vmatrix} = (-3)(-3)(-3) \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix}$$

$$= (-3)^3 \det A = (-27) \times 7 = -189.$$



Theorem (Determinant of Scalar Multiple of Matrices)

If A is an $n \times n$ matrix and $k \in \mathbb{R}$ is a scalar, then

$$\det(kA) = k^n \det A.$$

Problem

Let

$$A = \begin{bmatrix} a & b & c \\ p & q & r \\ x & y & z \end{bmatrix} \quad \text{and} \quad B = \begin{bmatrix} 2a + p & 2b + q & 2c + r \\ 2p + x & 2q + y & 2r + z \\ 2x + a & 2y + b & 2z + c \end{bmatrix}$$

Show that $\det B = 9 \det A$.

Solution

$$\det B = \begin{vmatrix} 2a + p & 2b + q & 2c + r \\ 2p + x & 2q + y & 2r + z \\ 2x + a & 2y + b & 2z + c \end{vmatrix} = \begin{vmatrix} p - 4x & q - 4y & r - 4z \\ 2p + x & 2q + y & 2r + z \\ 2x + a & 2y + b & 2z + c \end{vmatrix}$$

Solution

$$\begin{aligned}\det B &= \begin{vmatrix} 2a + p & 2b + q & 2c + r \\ 2p + x & 2q + y & 2r + z \\ 2x + a & 2y + b & 2z + c \end{vmatrix} = \begin{vmatrix} p - 4x & q - 4y & r - 4z \\ 2p + x & 2q + y & 2r + z \\ 2x + a & 2y + b & 2z + c \end{vmatrix} \\ &= \begin{vmatrix} p - 4x & q - 4y & r - 4z \\ 9x & 9y & 9z \\ 2x + a & 2y + b & 2z + c \end{vmatrix} = 9 \begin{vmatrix} p - 4x & q - 4y & r - 4z \\ x & y & z \\ 2x + a & 2y + b & 2z + c \end{vmatrix}\end{aligned}$$

Solution

$$\begin{aligned}\det B &= \begin{vmatrix} 2a + p & 2b + q & 2c + r \\ 2p + x & 2q + y & 2r + z \\ 2x + a & 2y + b & 2z + c \end{vmatrix} = \begin{vmatrix} p - 4x & q - 4y & r - 4z \\ 2p + x & 2q + y & 2r + z \\ 2x + a & 2y + b & 2z + c \end{vmatrix} \\&= \begin{vmatrix} p - 4x & q - 4y & r - 4z \\ 9x & 9y & 9z \\ 2x + a & 2y + b & 2z + c \end{vmatrix} = 9 \begin{vmatrix} p - 4x & q - 4y & r - 4z \\ x & y & z \\ 2x + a & 2y + b & 2z + c \end{vmatrix} \\&= 9 \begin{vmatrix} p & q & r \\ x & y & z \\ 2x + a & 2y + b & 2z + c \end{vmatrix} = 9 \begin{vmatrix} p & q & r \\ x & y & z \\ a & b & c \end{vmatrix} = -9 \begin{vmatrix} a & b & c \\ x & y & z \\ p & q & r \end{vmatrix}\end{aligned}$$

Solution

$$\begin{aligned}\det B &= \begin{vmatrix} 2a + p & 2b + q & 2c + r \\ 2p + x & 2q + y & 2r + z \\ 2x + a & 2y + b & 2z + c \end{vmatrix} = \begin{vmatrix} p - 4x & q - 4y & r - 4z \\ 2p + x & 2q + y & 2r + z \\ 2x + a & 2y + b & 2z + c \end{vmatrix} \\&= \begin{vmatrix} p - 4x & q - 4y & r - 4z \\ 9x & 9y & 9z \\ 2x + a & 2y + b & 2z + c \end{vmatrix} = 9 \begin{vmatrix} p - 4x & q - 4y & r - 4z \\ x & y & z \\ 2x + a & 2y + b & 2z + c \end{vmatrix} \\&= 9 \begin{vmatrix} p & q & r \\ x & y & z \\ 2x + a & 2y + b & 2z + c \end{vmatrix} = 9 \begin{vmatrix} p & q & r \\ x & y & z \\ a & b & c \end{vmatrix} = -9 \begin{vmatrix} a & b & c \\ x & y & z \\ p & q & r \end{vmatrix} \\&= 9 \begin{vmatrix} a & b & c \\ p & q & r \\ x & y & z \end{vmatrix} = 9 \det A.\end{aligned}$$



Copyright

Determinant of Small Matrices

The Cofactor Expansion

Elementary Row Operations and Determinants

Determinant and Scalar Multiple

Determinant of Triangular Matrices

Determinant of Block Matrices

Some More Exercises

Determinant of Triangular Matrices

Theorem

If $A = [a_{ij}]$ is an $n \times n$ (square, upper or lower) triangular matrix, then

$$\det A = a_{11}a_{22}a_{33} \cdots a_{nn},$$

i.e., $\det A$ is the product of the entries of the main diagonal of A .



Determinants of Upper Triangular Matrices

$$U = \begin{bmatrix} u_{11} & u_{12} & \cdots & u_{1n} \\ 0 & u_{22} & \cdots & u_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & u_{nn} \end{bmatrix}$$

\Downarrow

$$\det(U) = u_{11} u_{22} \cdots u_{nn}$$

Determinants of **lower** Triangular Matrices

$$L = \begin{bmatrix} \ell_{11} & 0 & \cdots & 0 \\ \ell_{21} & \ell_{22} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ \ell_{n1} & \ell_{n2} & \cdots & \ell_{nn} \end{bmatrix}$$

\Downarrow

$$\det(L) = \ell_{11}\ell_{22}\cdots\ell_{nn}$$

Determinants of **diagonal** Matrices

$$D = \begin{bmatrix} d_{11} & 0 & \cdots & 0 \\ 0 & d_{22} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & d_{nn} \end{bmatrix}$$

\Downarrow

$$\det(D) = d_{11}d_{22} \cdots d_{nn}$$

Copyright

Determinant of Small Matrices

The Cofactor Expansion

Elementary Row Operations and Determinants

Determinant and Scalar Multiple

Determinant of Triangular Matrices

Determinant of Block Matrices

Some More Exercises

Determinant of Block Matrices

Determinant of Block Matrices

Theorem

Consider the matrices

$$\begin{bmatrix} A & X \\ 0 & B \end{bmatrix} \quad \text{and} \quad \begin{bmatrix} A & 0 \\ Y & B \end{bmatrix}$$

where A and B are square matrices. Then

$$\det \begin{bmatrix} A & X \\ 0 & B \end{bmatrix} = \det A \det B \quad \text{and} \quad \det \begin{bmatrix} A & 0 \\ Y & B \end{bmatrix} = \det A \det B.$$

Example

$$\det \begin{bmatrix} 1 & -1 & 2 & 0 & -2 \\ 0 & 1 & 0 & 4 & 1 \\ 1 & 1 & 5 & 0 & 0 \\ 0 & 0 & 0 & 3 & -1 \\ 0 & 0 & 0 & 1 & -1 \end{bmatrix} =$$

Example

$$\det \begin{bmatrix} 1 & -1 & 2 & 0 & -2 \\ 0 & 1 & 0 & 4 & 1 \\ 1 & 1 & 5 & 0 & 0 \\ 0 & 0 & 0 & 3 & -1 \\ 0 & 0 & 0 & 1 & -1 \end{bmatrix} = \det \left[\begin{array}{ccc|cc} 1 & -1 & 2 & 0 & -2 \\ 0 & 1 & 0 & 4 & 1 \\ 1 & 1 & 5 & 0 & 0 \\ \hline 0 & 0 & 0 & 3 & -1 \\ 0 & 0 & 0 & 1 & -1 \end{array} \right]$$

Example

$$\det \begin{bmatrix} 1 & -1 & 2 & 0 & -2 \\ 0 & 1 & 0 & 4 & 1 \\ 1 & 1 & 5 & 0 & 0 \\ 0 & 0 & 0 & 3 & -1 \\ 0 & 0 & 0 & 1 & -1 \end{bmatrix} = \det \left[\begin{array}{ccc|cc} 1 & -1 & 2 & 0 & -2 \\ 0 & 1 & 0 & 4 & 1 \\ 1 & 1 & 5 & 0 & 0 \\ \hline 0 & 0 & 0 & 3 & -1 \\ 0 & 0 & 0 & 1 & -1 \end{array} \right]$$
$$= \det \begin{bmatrix} 1 & -1 & 2 \\ 0 & 1 & 0 \\ 1 & 1 & 5 \end{bmatrix} \det \begin{bmatrix} 3 & -1 \\ 1 & -1 \end{bmatrix}$$

Example

$$\begin{aligned} \det \begin{bmatrix} 1 & -1 & 2 & 0 & -2 \\ 0 & 1 & 0 & 4 & 1 \\ 1 & 1 & 5 & 0 & 0 \\ 0 & 0 & 0 & 3 & -1 \\ 0 & 0 & 0 & 1 & -1 \end{bmatrix} &= \det \left[\begin{array}{ccc|cc} 1 & -1 & 2 & 0 & -2 \\ 0 & 1 & 0 & 4 & 1 \\ 1 & 1 & 5 & 0 & 0 \\ \hline 0 & 0 & 0 & 3 & -1 \\ 0 & 0 & 0 & 1 & -1 \end{array} \right] \\ &= \det \begin{bmatrix} 1 & -1 & 2 \\ 0 & 1 & 0 \\ 1 & 1 & 5 \end{bmatrix} \det \begin{bmatrix} 3 & -1 \\ 1 & -1 \end{bmatrix} \\ &= \det \begin{bmatrix} 1 & 2 \\ 1 & 5 \end{bmatrix} \det \begin{bmatrix} 3 & -1 \\ 1 & -1 \end{bmatrix} \end{aligned}$$

Example

$$\begin{aligned} \det \begin{bmatrix} 1 & -1 & 2 & 0 & -2 \\ 0 & 1 & 0 & 4 & 1 \\ 1 & 1 & 5 & 0 & 0 \\ 0 & 0 & 0 & 3 & -1 \\ 0 & 0 & 0 & 1 & -1 \end{bmatrix} &= \det \left[\begin{array}{ccc|cc} 1 & -1 & 2 & 0 & -2 \\ 0 & 1 & 0 & 4 & 1 \\ 1 & 1 & 5 & 0 & 0 \\ \hline 0 & 0 & 0 & 3 & -1 \\ 0 & 0 & 0 & 1 & -1 \end{array} \right] \\ &= \det \begin{bmatrix} 1 & -1 & 2 \\ 0 & 1 & 0 \\ 1 & 1 & 5 \end{bmatrix} \det \begin{bmatrix} 3 & -1 \\ 1 & -1 \end{bmatrix} \\ &= \det \begin{bmatrix} 1 & 2 \\ 1 & 5 \end{bmatrix} \det \begin{bmatrix} 3 & -1 \\ 1 & -1 \end{bmatrix} \\ &= 3 \times (-2) = -6. \end{aligned}$$

Copyright

Determinant of Small Matrices

The Cofactor Expansion

Elementary Row Operations and Determinants

Determinant and Scalar Multiple

Determinant of Triangular Matrices

Determinant of Block Matrices

Some More Exercises

Some More Exercises

Some More Exercises

Example (From Exercise)

Evaluate by inspection.

$$\det \begin{bmatrix} a & b & c \\ a+1 & b+1 & c+1 \\ a-1 & b-1 & c-1 \end{bmatrix} = ?$$

Some More Exercises

Example (From Exercise)

Evaluate by inspection.

$$\det \begin{bmatrix} a & b & c \\ a+1 & b+1 & c+1 \\ a-1 & b-1 & c-1 \end{bmatrix} = ?$$

$$\text{row2} + \text{row3} - 2(\text{row1}) = \begin{bmatrix} 0 & 0 & 0 \end{bmatrix}$$

Example (From Exercise)

(a) Find $\det A$ if A is 3×3 and $\det(2A) = 6$.

Example (From Exercise)

- (a) Find $\det A$ if A is 3×3 and $\det(2A) = 6$.
- (b) Let A be an $n \times n$ matrix. Under what conditions is $\det(-A) = \det A$?

Example (From Exercise)

In each case, prove the statement is true or give a counterexample showing that the statement is false.

(a) $\det(A + B) = \det A + \det B$.

Example (From Exercise)

In each case, prove the statement is true or give a counterexample showing that the statement is false.

(a) $\det(A + B) = \det A + \det B$.

(c) If A is 2×2 , then $\det(A^T) = \det A$.

Example (From Exercise)

In each case, prove the statement is true or give a counterexample showing that the statement is false.

- (a) $\det(A + B) = \det A + \det B$.
- (c) If A is 2×2 , then $\det(A^T) = \det A$.
- (e) If A is 2×2 , then $\det(7A) = 49 \det A$.

Example (From Exercise)

In each case, prove the statement is true or give a counterexample showing that the statement is false.

- (a) $\det(A + B) = \det A + \det B$.
- (c) If A is 2×2 , then $\det(A^T) = \det A$.
- (e) If A is 2×2 , then $\det(7A) = 49 \det A$.
- (g) $\det(-A) = -\det A$.