

# Math 221: LINEAR ALGEBRA

## Chapter 5. Vector Space $\mathbb{R}^n$

### §5-1. Subspaces and Spanning

Le Chen<sup>1</sup>

Emory University, 2021 Spring

(last updated on 01/12/2023)



Creative Commons License  
(CC BY-NC-SA)

<sup>1</sup>Slides are adapted from those by Karen Seyffarth from University of Calgary.



# Linear Algebra with Applications

## Lecture Notes

### Current Lecture Notes Revision: Version 2018 — Revision B

These lecture notes were originally developed by Karen Seyffarth of the University of Calgary. Edits, additions, and revisions have been made to these notes by the editorial team at Lyryx Learning to accompany their text [Linear Algebra with Applications](#) based on W. K. Nicholson's original text.

In addition we recognize the following contributors. All new content contributed is released under the same license as noted below.

- Ilijas Farah, York University

### BE A CHAMPION OF OER!

Contribute suggestions for improvements, new content, or errata:

A new topic

A new example or problem

A new or better proof to an existing theorem

Any other suggestions to improve the material

Contact Lyryx at [info@lyryx.com](mailto:info@lyryx.com) with your ideas.

### License



Attribution-NonCommercial-ShareAlike (CC BY-NC-SA)

This license lets others remix, tweak, and build upon your work non-commercially, as long as they credit you and license their new creations under the identical terms.

Copyright

Subspaces of  $\mathbb{R}^n$

The null space and the image space

The Eigenspace

Linear Combinations and Spanning Sets

Spanning sets of  $\text{null}(A)$  and  $\text{im}(A)$

Copyright

## Subspaces of $\mathbb{R}^n$

The null space and the image space

The Eigenspace

Linear Combinations and Spanning Sets

Spanning sets of  $\text{null}(A)$  and  $\text{im}(A)$

# Subspaces of $\mathbb{R}^n$

## Definitions

1.  $\mathbb{R}$  denotes the set of **real** numbers, and is an example of a set of **scalars**.
2.  $\mathbb{R}^n$  is the set of all n-tuples of real numbers, i.e.,

$$\mathbb{R}^n = \{(x_1, x_2, \dots, x_n) \mid x_i \in \mathbb{R}, 1 \leq i \leq n\}.$$

3. The **vector space**  $\mathbb{R}^n$  consists of the set  $\mathbb{R}^n$  written as **column matrices**, along with the (matrix) operations of addition and scalar multiplication. Unless stated otherwise,  $\mathbb{R}^n$  means the vector space  $\mathbb{R}^n$ .

## Remark

$\mathbb{R}^n$  is a concrete example of the abstract vector space will be studied in the next chapter.

A vectors is denoted by a lower case letter with an arrow written over it; for example,  $\vec{u}$ ,  $\vec{v}$ , and  $\vec{x}$  denote vectors.

Another example:  $\vec{u} = \begin{bmatrix} -2 \\ 3 \\ 0.7 \\ 5 \\ \pi \end{bmatrix}$  is a vector in  $\mathbb{R}^5$ , written  $\vec{u} \in \mathbb{R}^5$ .

To save space on the page, the same vector  $\vec{u}$  may be written instead as a row matrix by taking the transpose of the column:

$$\vec{u} = \begin{bmatrix} -2, & 3, & 0.7, & 5, & \pi \end{bmatrix}^T.$$

We are interested in nice subsets of  $\mathbb{R}^n$ , defined as follows.

### Definition (Subspaces)

A subset  $U$  of  $\mathbb{R}^n$  is a **subspace** of  $\mathbb{R}^n$  if


- S1. The zero vector of  $\mathbb{R}^n$ ,  $\vec{0}_n$ , is in  $U$ ;
- S2.  $U$  is closed under addition, i.e., for all  $\vec{u}, \vec{w} \in U$ ,  $\vec{u} + \vec{w} \in U$ ;
- S3.  $U$  is closed under scalar multiplication, i.e., for all  $\vec{u} \in U$  and  $k \in \mathbb{R}$ ,  $k\vec{u} \in U$ .

Both subset  $U = \{\vec{0}_n\}$  and  $\mathbb{R}^n$  itself are subspaces of  $\mathbb{R}^n$ . Any other subspace of  $\mathbb{R}^n$  is called a **proper** subspace of  $\mathbb{R}^n$ .

### Notation

If  $U$  is a subset of  $\mathbb{R}^n$ , we write  $U \subseteq \mathbb{R}^n$ .





## Example

In  $\mathbb{R}^3$ , the line  $L$  through the origin that is parallel to the vector  $\vec{d} = \begin{bmatrix} -5 \\ 1 \\ -4 \end{bmatrix}$  has (vector) equation  $\begin{bmatrix} x \\ y \\ z \end{bmatrix} = t \begin{bmatrix} -5 \\ 1 \\ -4 \end{bmatrix}, t \in \mathbb{R}$ , so

$$L = \left\{ t\vec{d} \mid t \in \mathbb{R} \right\}.$$

**Claim.**  $L$  is a subspace of  $\mathbb{R}^3$ .

- First:  $\vec{0}_3 \in L$  since  $0\vec{d} = \vec{0}_3$ .
- Suppose  $\vec{u}, \vec{v} \in L$ . Then by definition,  $\vec{u} = s\vec{d}$  and  $\vec{v} = t\vec{d}$ , for some  $s, t \in \mathbb{R}$ . Thus

$$\vec{u} + \vec{v} = s\vec{d} + t\vec{d} = (s + t)\vec{d}.$$

Since  $s + t \in \mathbb{R}$ ,  $\vec{u} + \vec{v} \in L$ ; i.e.,  $L$  is closed under addition.

### Example (continued)

- Suppose  $\vec{u} \in L$  and  $k \in \mathbb{R}$  ( $k$  is a scalar). Then  $\vec{u} = t\vec{d}$ , for some  $t \in \mathbb{R}$ , so

$$k\vec{u} = k(t\vec{d}) = (kt)\vec{d}.$$

Since  $kt \in \mathbb{R}$ ,  $k\vec{u} \in L$ ; i.e.,  $L$  is closed under scalar multiplication.

- Therefore,  $L$  is a subspace of  $\mathbb{R}^3$ .

### Remark

Note that there is nothing special about the vector  $\vec{d}$  used in this example; the same proof works for any **nonzero** vector  $\vec{d} \in \mathbb{R}^3$ , so any line through the origin is a subspace of  $\mathbb{R}^3$ .

## Example

In  $\mathbb{R}^3$ , let  $M$  denote the plane through the origin having equation

$3x - 2y + z = 0$ ; then  $M$  has normal vector  $\vec{n} = \begin{bmatrix} 3 \\ -2 \\ 1 \end{bmatrix}$ . If  $\vec{u} = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$ , then

$$M = \{ \vec{u} \in \mathbb{R}^3 \mid \vec{n} \cdot \vec{u} = 0 \},$$

where  $\vec{n} \cdot \vec{u}$  is the dot product of vectors  $\vec{n}$  and  $\vec{u}$ .

**Claim.**  $M$  is a subspace of  $\mathbb{R}^3$ .

- First:  $\vec{0}_3 \in M$  since  $\vec{n} \cdot \vec{0}_3 = 0$ .
- Suppose  $\vec{u}, \vec{v} \in M$ . Then by definition,  $\vec{n} \cdot \vec{u} = 0$  and  $\vec{n} \cdot \vec{v} = 0$ , so

$$\vec{n} \cdot (\vec{u} + \vec{v}) = \vec{n} \cdot \vec{u} + \vec{n} \cdot \vec{v} = 0 + 0 = 0,$$

and thus  $(\vec{u} + \vec{v}) \in M$ ; i.e.,  $M$  is closed under addition.

### Example (continued)

- Suppose  $\vec{u} \in M$  and  $k \in \mathbb{R}$ . Then  $\vec{n} \cdot \vec{u} = 0$ , so

$$\vec{n} \cdot (k\vec{u}) = k(\vec{n} \cdot \vec{u}) = k(0) = 0,$$

and thus  $k\vec{u} \in M$ ; i.e.,  $M$  is closed under scalar multiplication.

- Therefore,  $M$  is a subspace of  $\mathbb{R}^3$ .

### Remark

As in the previous example, there is nothing special about the plane chosen for this example; any plane through the origin is a subspace of  $\mathbb{R}^3$ .

### Problem

$$\text{Is } U = \left\{ \left[ \begin{array}{c} a \\ b \\ c \\ d \end{array} \right] \mid a, b, c, d \in \mathbb{R} \text{ and } 2a - b = c + 2d \right\} \text{ a subspace of } \mathbb{R}^4?$$

Justify your answer.

### Solution

The zero vector of  $\mathbb{R}^4$  is the vector  $\left[ \begin{array}{c} a \\ b \\ c \\ d \end{array} \right]$  with  $a = b = c = d = 0$ .

In this case,  $2a - b = 2(0) + 0 = 0$  and  $c + 2d = 0 + 2(0) = 0$ , so  $2a - b = c + 2d$ . Therefore,  $\vec{0}_4 \in U$ . ■

## Solution (continued)

Suppose

$$\vec{v}_1 = \begin{bmatrix} a_1 \\ b_1 \\ c_1 \\ d_1 \end{bmatrix} \quad \text{and} \quad \vec{v}_2 = \begin{bmatrix} a_2 \\ b_2 \\ c_2 \\ d_2 \end{bmatrix} \quad \text{are in } U.$$

Then  $2a_1 - b_1 = c_1 + 2d_1$  and  $2a_2 - b_2 = c_2 + 2d_2$ . Now

$$\vec{v}_1 + \vec{v}_2 = \begin{bmatrix} a_1 \\ b_1 \\ c_1 \\ d_1 \end{bmatrix} + \begin{bmatrix} a_2 \\ b_2 \\ c_2 \\ d_2 \end{bmatrix} = \begin{bmatrix} a_1 + a_2 \\ b_1 + b_2 \\ c_1 + c_2 \\ d_1 + d_2 \end{bmatrix},$$

and

$$\begin{aligned} 2(a_1 + a_2) - (b_1 + b_2) &= (2a_1 - b_1) + (2a_2 - b_2) \\ &= (c_1 + 2d_1) + (c_2 + 2d_2) \\ &= (c_1 + c_2) + 2(d_1 + d_2). \end{aligned}$$

Therefore,  $\vec{v}_1 + \vec{v}_2 \in U$ .

### Solution (continued)

Finally, suppose

$$\vec{v} = \begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix} \in U \quad \text{and} \quad k \in \mathbb{R}.$$

Then  $2a - b = c + 2d$ . Now

$$k\vec{v} = k \begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix} = \begin{bmatrix} ka \\ kb \\ kc \\ kd \end{bmatrix},$$

and

$$2ka - kb = k(2a - b) = k(c + 2d) = kc + 2kd.$$

Therefore,  $k\vec{v} \in U$ .

It follows from the **Subspace Test** that  $U$  is a subspace of  $\mathbb{R}^4$ .



### Problem

Is  $U = \left\{ \begin{bmatrix} 1 \\ s \\ t \end{bmatrix} \mid s, t \in \mathbb{R} \right\}$  a subspace of  $\mathbb{R}^3$ ? Justify your answer.

### Solution

Note that  $\vec{0}_3 \notin U$ , and thus  $U$  is not a subspace of  $\mathbb{R}^3$ .

(You could also show that  $U$  is not closed under addition, or not closed under scalar multiplication.)

### Problem

$$\text{Is } U = \left\{ \begin{bmatrix} r \\ 0 \\ s \end{bmatrix} \mid r, s \in \mathbb{R} \text{ and } r^2 + s^2 = 0 \right\} \text{ a subspace of } \mathbb{R}^3?$$

Justify your answer.

### Solution

Since  $r \in \mathbb{R}$ ,  $r^2 \geq 0$  with equality if and only if  $r = 0$ . Similarly,  $s \in \mathbb{R}$  implies  $s^2 \geq 0$ , and  $s^2 = 0$  if and only if  $s = 0$ . This means  $r^2 + s^2 = 0$  if and only if  $r^2 = s^2 = 0$ ; thus  $r^2 + s^2 = 0$  if and only if  $r = s = 0$ . Therefore  $U$  contains only  $\vec{0}_3$ , the zero vector, i.e.,  $U = \{\vec{0}_3\}$ . As we already observed,  $\{\vec{0}_n\}$  is a subspace of  $\mathbb{R}^n$ , and therefore  $U$  is a subspace of  $\mathbb{R}^3$ . ■

Copyright

Subspaces of  $\mathbb{R}^n$

The null space and the image space

The Eigenspace

Linear Combinations and Spanning Sets

Spanning sets of  $\text{null}(A)$  and  $\text{im}(A)$

# The null space and the image space

## Definitions (Null Space and Image Space)

Let  $A$  be an  $m \times n$  matrix. The **null space** of  $A$  is defined as

$$\text{null}(A) = \{\vec{x} \in \mathbb{R}^n \mid A\vec{x} = \vec{0}_m\},$$

and the **image space** of  $A$  is defined as

$$\text{im}(A) = \{A\vec{x} \mid \vec{x} \in \mathbb{R}^n\}.$$

## Remark

1. Since  $A$  is  $m \times n$  and  $\vec{x} \in \mathbb{R}^n$ ,  $A\vec{x} \in \mathbb{R}^m$ , so  $\text{im}(A) \subseteq \mathbb{R}^m$  while  $\text{null}(A) \subseteq \mathbb{R}^n$ .
2. Image space is also called **column space** of  $A$ , denoted as  $\text{col}(A)$ :

$$\text{col}(A) = \text{span}(\vec{a}_1, \dots, \vec{a}_n) = \text{im}(A).$$

## Problem

Prove that if  $A$  is an  $m \times n$  matrix, then  $\text{null}(A)$  is a subspace of  $\mathbb{R}^n$ .

## Proof.

S1. Since  $A\vec{0}_n = \vec{0}_m$ ,  $\vec{0}_n \in \text{null}(A)$ .

S2. Let  $\vec{x}, \vec{y} \in \text{null}(A)$ . Then  $A\vec{x} = \vec{0}_m$  and  $A\vec{y} = \vec{0}_m$ , so

$$A(\vec{x} + \vec{y}) = A\vec{x} + A\vec{y} = \vec{0}_m + \vec{0}_m = \vec{0}_m,$$

and thus  $\vec{x} + \vec{y} \in \text{null}(A)$ .

S3. Let  $\vec{x} \in \text{null}(A)$  and  $k \in \mathbb{R}$ . Then  $A\vec{x} = \vec{0}_m$ , so

$$A(k\vec{x}) = k(A\vec{x}) = k\vec{0}_m = \vec{0}_m,$$

and thus  $k\vec{x} \in \text{null}(A)$ .

Therefore,  $\text{null}(A)$  is a subspace of  $\mathbb{R}^n$ . ■

## Problem

Prove that if  $A$  is an  $m \times n$  matrix, then  $\text{im}(A)$  is a subspace of  $\mathbb{R}^m$ .

## Proof.

S1. Since  $\vec{0}_n \in \mathbb{R}^n$  and  $A\vec{0}_n = \vec{0}_m$ ,  $\vec{0}_m \in \text{im}(A)$ .

S2. Let  $\vec{x}, \vec{y} \in \text{im}(A)$ . Then  $\vec{x} = A\vec{u}$  and  $\vec{y} = A\vec{v}$  for some  $\vec{u}, \vec{v} \in \mathbb{R}^n$ , so

$$\vec{x} + \vec{y} = A\vec{u} + A\vec{v} = A(\vec{u} + \vec{v}).$$

Since  $\vec{u} + \vec{v} \in \mathbb{R}^n$ , it follows that  $\vec{x} + \vec{y} \in \text{im}(A)$ .

S3. Let  $\vec{x} \in \text{im}(A)$  and  $k \in \mathbb{R}$ . Then  $\vec{x} = A\vec{u}$  for some  $\vec{u} \in \mathbb{R}^n$ , and thus

$$k\vec{x} = k(A\vec{u}) = A(k\vec{u}).$$

Since  $k\vec{u} \in \mathbb{R}^n$ , it follows that  $k\vec{x} \in \text{im}(A)$ .

Therefore,  $\text{im}(A)$  is a subspace of  $\mathbb{R}^m$ . ■

Copyright

Subspaces of  $\mathbb{R}^n$

The null space and the image space

**The Eigenspace**

Linear Combinations and Spanning Sets

Spanning sets of  $\text{null}(A)$  and  $\text{im}(A)$

# The Eigenspace

## Definition (Eigenspace)

Let  $A$  be an  $n \times n$  matrix and  $\lambda \in \mathbb{R}$ . The eigenspace of  $A$  corresponding to  $\lambda$  is the set

$$E_{\lambda}(A) = \{\vec{x} \in \mathbb{R}^n \mid A\vec{x} = \lambda\vec{x}\}.$$



### Example


$A = \begin{pmatrix} 4 & -2 \\ -1 & 3 \end{pmatrix}$  has two eigenvalues:  $\lambda_1 = 2$  and  $\lambda_2 = 5$  with corresponding eigenvectors


$$\vec{v}_1 = \begin{pmatrix} 1 \\ 1 \end{pmatrix} \quad \text{and} \quad \vec{v}_2 = \begin{pmatrix} -1 \\ 1/2 \end{pmatrix}$$

Hence,

$$E_{\lambda_1}(A) = E_2(A) = \{t\vec{v}_1 | t \in \mathbb{R}\}$$

$$E_{\lambda_2}(A) = E_5(A) = \{t\vec{v}_2 | t \in \mathbb{R}\}$$





$$E_2(A) = \left\{ t \begin{pmatrix} 1 \\ 1 \end{pmatrix} \mid t \in \mathbb{R} \right\}$$

$$E_5(A) = \left\{ t \begin{pmatrix} -1 \\ 1/2 \end{pmatrix} \mid t \in \mathbb{R} \right\}$$

Note that

$$\begin{aligned} E_\lambda(A) &= \{\vec{x} \in \mathbb{R}^n \mid A\vec{x} = \lambda\vec{x}\} \\ &= \left\{ \vec{x} \in \mathbb{R}^n \mid \lambda\vec{x} - A\vec{x} = \vec{0}_n \right\} \\ &= \left\{ \vec{x} \in \mathbb{R}^n \mid (\lambda I - A)\vec{x} = \vec{0}_n \right\} \end{aligned}$$

showing that

$$E_\lambda(A) = \text{null}(\lambda I - A).$$

It follows that

- ▶ if  $\lambda$  is **not** an eigenvalue of  $A$ , then  $E_\lambda(A) = \{\vec{0}_n\}$ ;
- ▶ the nonzero vectors of  $E_\lambda(A)$  are the eigenvectors of  $A$  corresponding to  $\lambda$ ;
- ▶ the eigenspace of  $A$  corresponding to  $\lambda$  is a subspace of  $\mathbb{R}^n$ .

Copyright

Subspaces of  $\mathbb{R}^n$

The null space and the image space

The Eigenspace

**Linear Combinations and Spanning Sets**

Spanning sets of  $\text{null}(A)$  and  $\text{im}(A)$

# Linear Combinations and Spanning Sets

## Definition (Linear Combinations and Spanning)

Let  $\vec{x}_1, \vec{x}_2, \dots, \vec{x}_k \in \mathbb{R}^n$  and  $t_1, t_2, \dots, t_k \in \mathbb{R}$ . Then the vector

$$\vec{x} = t_1\vec{x}_1 + t_2\vec{x}_2 + \cdots + t_k\vec{x}_k$$

is called a **linear combination** of the vectors  $\vec{x}_1, \vec{x}_2, \dots, \vec{x}_k$ ; the (scalars)  $t_1, t_2, \dots, t_k \in \mathbb{R}$  are the coefficients. The set of all linear combinations of  $\vec{x}_1, \vec{x}_2, \dots, \vec{x}_k$  is called **the span of  $\vec{x}_1, \vec{x}_2, \dots, \vec{x}_k$** , and is written

$$\text{span}\{\vec{x}_1, \vec{x}_2, \dots, \vec{x}_k\} = \{t_1\vec{x}_1 + t_2\vec{x}_2 + \cdots + t_k\vec{x}_k \mid t_1, t_2, \dots, t_k \in \mathbb{R}\}.$$

Additional Terminology. If  $U = \text{span}\{\vec{x}_1, \vec{x}_2, \dots, \vec{x}_k\}$ , then

- ▶ **U is spanned by** the vectors  $\vec{x}_1, \vec{x}_2, \dots, \vec{x}_k$ .
- ▶ the vectors  $\vec{x}_1, \vec{x}_2, \dots, \vec{x}_k$  **span U**.
- ▶ the set of vectors  $\{\vec{x}_1, \vec{x}_2, \dots, \vec{x}_k\}$  is a **spanning set** for U.

### Example

Let  $\vec{x} \in \mathbb{R}^3$  be a nonzero vector. Then  $\text{span}\{\vec{x}\} = \{k\vec{x} \mid k \in \mathbb{R}\}$  is a line through the origin having direction vector  $\vec{x}$ .

### Example

Let  $\vec{x}, \vec{y} \in \mathbb{R}^3$  be nonzero vectors that are not parallel. Then

$$\text{span}\{\vec{x}, \vec{y}\} = \{k\vec{x} + t\vec{y} \mid k, t \in \mathbb{R}\}$$

is a plane through the origin containing  $\vec{x}$  and  $\vec{y}$ .

How would you find the equation of this plane?

### Problem

$$\text{Let } \vec{x} = \begin{bmatrix} 8 \\ 3 \\ -13 \\ 20 \end{bmatrix}, \vec{y} = \begin{bmatrix} 2 \\ 1 \\ -3 \\ 5 \end{bmatrix} \text{ and } \vec{z} = \begin{bmatrix} -1 \\ 0 \\ 2 \\ -3 \end{bmatrix}. \text{ Is } \vec{x} \in \text{span}\{\vec{y}, \vec{z}\}?$$

### Solution

An equivalent question is: can  $\vec{x}$  be expressed as a linear combination of  $\vec{y}$  and  $\vec{z}$ ?

Suppose there exist  $a, b \in \mathbb{R}$  so that  $\vec{x} = a\vec{y} + b\vec{z}$ . Then

$$\begin{bmatrix} 8 \\ 3 \\ -13 \\ 20 \end{bmatrix} = a \begin{bmatrix} 2 \\ 1 \\ -3 \\ 5 \end{bmatrix} + b \begin{bmatrix} -1 \\ 0 \\ 2 \\ -3 \end{bmatrix} = \begin{bmatrix} 2 & -1 \\ 1 & 0 \\ -3 & 2 \\ 5 & -3 \end{bmatrix} \begin{bmatrix} a \\ b \end{bmatrix}.$$

Solve this system of four linear equations in the two variables  $a$  and  $b$ .



Solution (continued)

$$\left[ \begin{array}{cc|c} 2 & -1 & 8 \\ 1 & 0 & 3 \\ -3 & 2 & -13 \\ 5 & -3 & 20 \end{array} \right] \rightarrow \left[ \begin{array}{cc|c} 1 & 0 & 3 \\ 0 & 1 & -2 \\ 0 & 0 & 0 \\ 0 & 0 & -1 \end{array} \right]$$

Since the system has no solutions,  $\vec{x} \notin \text{span}\{\vec{y}, \vec{z}\}$ .



## Problem

$$\text{Let } \vec{w} = \begin{bmatrix} 8 \\ 3 \\ -13 \\ \textcolor{red}{21} \end{bmatrix}, \vec{y} = \begin{bmatrix} 2 \\ 1 \\ -3 \\ 5 \end{bmatrix} \text{ and } \vec{z} = \begin{bmatrix} -1 \\ 0 \\ 2 \\ -3 \end{bmatrix}. \text{ Is } \vec{w} \in \text{span}\{\vec{y}, \vec{z}\}?$$

This is almost identical to a previous problem, except that  $\vec{w}$  (above) has one entry that is different from the vector  $\vec{x}$  of that problem.

## Solution

In this case, the system of linear equations is consistent, and gives us  $\vec{w} = 3\vec{y} - 2\vec{z}$ , so  $\vec{w} \in \text{span}\{\vec{y}, \vec{z}\}$ .

## Theorem

Let  $\vec{x}_1, \vec{x}_2, \dots, \vec{x}_k \in \mathbb{R}^n$  and let  $U = \text{span}\{\vec{x}_1, \vec{x}_2, \dots, \vec{x}_k\}$ . Then

1.  $U$  is a subspace of  $\mathbb{R}^n$  containing each  $\vec{x}_i$ ,  $1 \leq i \leq k$ ;
2. if  $W$  is a subspace of  $\mathbb{R}^n$  and  $\vec{x}_1, \vec{x}_2, \dots, \vec{x}_k \in W$ , then  $U \subseteq W$ .

## Remark

Property 2 is saying that  $U$  is the “smallest” subspace of  $\mathbb{R}^n$  that contains  $\vec{x}_1, \vec{x}_2, \dots, \vec{x}_k$ .

Proof. ( Part 1 of Theorem )

Since  $U = \text{span}\{\vec{x}_1, \vec{x}_2, \dots, \vec{x}_k\}$  and  $0\vec{x}_1 + 0\vec{x}_2 + \dots + 0\vec{x}_k = \vec{0}_n$ ,  $\vec{0}_n \in U$ .

Suppose  $\vec{x}, \vec{y} \in U$ . Then for some  $s_i, t_i \in \mathbb{R}$ ,  $1 \leq i \leq k$ ,

$$\vec{x} = s_1\vec{x}_1 + s_2\vec{x}_2 + \dots + s_k\vec{x}_k$$

$$\vec{y} = t_1\vec{x}_1 + t_2\vec{x}_2 + \dots + t_k\vec{x}_k$$

Thus

$$\begin{aligned}\vec{x} + \vec{y} &= (s_1\vec{x}_1 + s_2\vec{x}_2 + \dots + s_k\vec{x}_k) + (t_1\vec{x}_1 + t_2\vec{x}_2 + \dots + t_k\vec{x}_k) \\ &= (s_1 + t_1)\vec{x}_1 + (s_2 + t_2)\vec{x}_2 + \dots + (s_k + t_k)\vec{x}_k.\end{aligned}$$

Since  $s_i + t_i \in \mathbb{R}$  for all  $1 \leq i \leq k$ ,  $\vec{x} + \vec{y} \in U$ , i.e.,  **$U$  is closed under addition.**

Proof. ( Part 1 of Theorem – continued)

Suppose  $\vec{x} \in U$  and  $a \in \mathbb{R}$ . Then for some  $s_i \in \mathbb{R}$ ,  $1 \leq i \leq k$ ,

$$\vec{x} = s_1\vec{x}_1 + s_2\vec{x}_2 + \cdots + s_k\vec{x}_k$$

Thus

$$\begin{aligned} a\vec{x} &= a(s_1\vec{x}_1 + s_2\vec{x}_2 + \cdots + s_k\vec{x}_k) \\ &= (as_1)\vec{x}_1 + (as_2)\vec{x}_2 + \cdots + (as_k)\vec{x}_k. \end{aligned}$$

Since  $as_i \in \mathbb{R}$  for all  $1 \leq i \leq k$ ,  $a\vec{x} \in U$ . Hence, **U is closed under scalar multiplication.**

Therefore,  $U$  is a subspace of  $\mathbb{R}^n$ . Furthermore, since

$$\vec{x}_i = \sum_{j=1}^{i-1} 0\vec{x}_j + 1\vec{x}_i + \sum_{j=i+1}^k 0\vec{x}_j,$$

it follows that  $\vec{x}_i \in U$  for all  $i$ ,  $1 \leq i \leq k$ .

### Proof. (Part 2 of Theorem)

Let  $W \subset \mathbb{R}^n$  be a subspace that contains  $\vec{x}_1, \dots, \vec{x}_n$ . We need to prove that  $U \subseteq W$ .

Suppose  $\vec{x} \in U$ . Then  $\vec{x} = s_1\vec{x}_1 + s_2\vec{x}_2 + \dots + s_k\vec{x}_k$  for some  $s_i \in \mathbb{R}$ ,  $1 \leq i \leq k$ . Since  $W$  contain each  $\vec{x}_i$  and  $W$  is closed under scalar multiplication, it follows that  $s_i\vec{x}_i \in W$  for each  $i$ ,  $1 \leq i \leq k$ . Furthermore, since  $W$  is closed under addition,  $\vec{x} = s_1\vec{x}_1 + s_2\vec{x}_2 + \dots + s_k\vec{x}_k \in W$ . Therefore,  $U \subseteq W$ .

### Problem (revisited)

$$\text{Is } U = \left\{ \left[ \begin{array}{c} a \\ b \\ c \\ d \end{array} \right] \mid a, b, c, d \in \mathbb{R} \text{ and } 2a - b = c + 2d \right\} \text{ a subspace of } \mathbb{R}^4?$$

Justify your answer.

### Solution (Another)

$$\text{Let } \vec{v} = \left[ \begin{array}{c} a \\ b \\ c \\ d \end{array} \right] \in U. \text{ Since } 2a - b = c + 2d, c = 2a - b - 2d, \text{ and thus}$$

$$U = \left\{ \left[ \begin{array}{c} a \\ b \\ 2a - b - 2d \\ d \end{array} \right] \mid a, b, d \in \mathbb{R} \right\} = \text{span} \left\{ \left[ \begin{array}{c} 1 \\ 0 \\ 2 \\ 0 \end{array} \right], \left[ \begin{array}{c} 0 \\ 1 \\ -1 \\ 0 \end{array} \right], \left[ \begin{array}{c} 0 \\ 0 \\ -2 \\ 1 \end{array} \right] \right\}.$$

By a previous Theorem,  $U$  is a subspace of  $\mathbb{R}^4$ .

## Problem

Let  $\vec{x}, \vec{y} \in \mathbb{R}^n$ ,  $U_1 = \text{span}\{\vec{x}, \vec{y}\}$ , and  $U_2 = \text{span}\{2\vec{x} - \vec{y}, 2\vec{y} + \vec{x}\}$ . Prove that  $U_1 = U_2$ .

## Solution

To show that  $U_1 = U_2$ , prove that  $U_1 \subseteq U_2$ , and  $U_2 \subseteq U_1$ . We begin by noting that, by the first part of the previous Theorem,  $U_1$  and  $U_2$  are subspaces of  $\mathbb{R}^n$ .

Since  $2\vec{x} - \vec{y}, 2\vec{y} + \vec{x} \in U_1$ , it follows from the second part of the previous Theorem that  $\text{span}\{2\vec{x} - \vec{y}, 2\vec{y} + \vec{x}\} \subseteq U_1$ , i.e.,  $U_2 \subseteq U_1$ .

Also, since

$$\begin{aligned}\vec{x} &= \frac{2}{5}(2\vec{x} - \vec{y}) + \frac{1}{5}(2\vec{y} + \vec{x}), \\ \vec{y} &= -\frac{1}{5}(2\vec{x} - \vec{y}) + \frac{2}{5}(2\vec{y} + \vec{x}),\end{aligned}$$

$\vec{x}, \vec{y} \in U_2$ . Therefore, by the second part of the previous Theorem,  $\text{span}\{\vec{x}, \vec{y}\} \subseteq U_2$ , i.e.,  $U_1 \subseteq U_2$ . The result now follows.



## Problem

Show that  $\mathbb{R}^n = \text{span}\{\vec{e}_1, \vec{e}_2, \dots, \vec{e}_n\}$ , where  $\vec{e}_j$  denote the  $j^{\text{th}}$  column of  $I_n$ .

## Solution

Let  $\vec{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \in \mathbb{R}^n$ . Then  $\vec{x} = x_1\vec{e}_1 + x_2\vec{e}_2 + \dots + x_n\vec{e}_n$ , where

$x_1, x_2, \dots, x_n \in \mathbb{R}$ . Therefore,  $\vec{x} \in \text{span}\{\vec{e}_1, \vec{e}_2, \dots, \vec{e}_n\}$ , and thus  $\mathbb{R}^n \subseteq \text{span}\{\vec{e}_1, \vec{e}_2, \dots, \vec{e}_n\}$ .

Conversely, since  $\vec{e}_i \in \mathbb{R}^n$  for each  $i$ ,  $1 \leq i \leq n$  (and  $\mathbb{R}^n$  is a vector space), it follows that  $\text{span}\{\vec{e}_1, \vec{e}_2, \dots, \vec{e}_n\} \subseteq \mathbb{R}^n$ . The equality now follows.

## Problem

$$\text{Let } \vec{x}_1 = \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}, \vec{x}_2 = \begin{bmatrix} 0 \\ 1 \\ 1 \\ 1 \end{bmatrix}, \vec{x}_3 = \begin{bmatrix} 0 \\ 0 \\ 1 \\ 1 \end{bmatrix}, \vec{x}_4 = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 1 \end{bmatrix}.$$

Does  $\{\vec{x}_1, \vec{x}_2, \vec{x}_3, \vec{x}_4\}$  span  $\mathbb{R}^4$ ? (Equivalently, is  $\text{span}\{\vec{x}_1, \vec{x}_2, \vec{x}_3, \vec{x}_4\} = \mathbb{R}^4$ ?)

## Solution

To prove  $\text{span}\{\vec{x}_1, \vec{x}_2, \vec{x}_3, \vec{x}_4\} = \mathbb{R}^4$ , we need to prove two directions:

$$\text{span}\{\vec{x}_1, \vec{x}_2, \vec{x}_3, \vec{x}_4\} \subseteq \mathbb{R}^4 \quad \text{and} \quad \mathbb{R}^4 \subseteq \text{span}\{\vec{x}_1, \vec{x}_2, \vec{x}_3, \vec{x}_4\}.$$

For the first relation, since  $\vec{x}_1, \vec{x}_2, \vec{x}_3, \vec{x}_4 \in \mathbb{R}^4$  (and  $\mathbb{R}^4$  is a vector space),  $\text{span}\{\vec{x}_1, \vec{x}_2, \vec{x}_3, \vec{x}_4\} \subseteq \mathbb{R}^4$ .

### Solution (continued)

For the second relation, notice that

$$\vec{e}_1 = \vec{x}_1 - \vec{x}_2$$

$$\vec{e}_2 = \vec{x}_2 - \vec{x}_3$$

$$\vec{e}_3 = \vec{x}_3 - \vec{x}_4$$

$$\vec{e}_4 = \vec{x}_4,$$

showing that  $\vec{e}_1, \vec{e}_2, \vec{e}_3, \vec{e}_4 \in \text{span}\{\vec{x}_1, \vec{x}_2, \vec{x}_3, \vec{x}_4\}$ . Therefore, since  $\text{span}\{\vec{x}_1, \vec{x}_2, \vec{x}_3, \vec{x}_4\}$  is a vector space,

$$\mathbb{R}^4 = \text{span}\{\vec{e}_1, \vec{e}_2, \vec{e}_3, \vec{e}_4\} \subseteq \text{span}\{\vec{x}_1, \vec{x}_2, \vec{x}_3, \vec{x}_4\},$$

and the equality follows.

### Problem

$$\text{Let } \vec{u}_1 = \begin{bmatrix} 1 \\ -1 \\ 1 \\ -1 \end{bmatrix}, \vec{u}_2 = \begin{bmatrix} -1 \\ 1 \\ 1 \\ 1 \end{bmatrix}, \vec{u}_3 = \begin{bmatrix} 1 \\ -1 \\ -1 \\ 1 \end{bmatrix}, \vec{u}_4 = \begin{bmatrix} 1 \\ -1 \\ 1 \\ 1 \end{bmatrix}.$$

Show that  $\text{span}\{\vec{u}_1, \vec{u}_2, \vec{u}_3, \vec{u}_4\} \neq \mathbb{R}^4$ .

### Solution

If you check, you'll find that  $\vec{e}_2$  can not be written as a linear combination of  $\vec{u}_1, \vec{u}_2, \vec{u}_3$ , and  $\vec{u}_4$ .

Copyright

Subspaces of  $\mathbb{R}^n$

The null space and the image space

The Eigenspace

Linear Combinations and Spanning Sets

Spanning sets of  $\text{null}(A)$  and  $\text{im}(A)$

## Spanning sets of $\text{null}(A)$ and $\text{im}(A)$

### Lemma

Let  $A$  be an  $m \times n$  matrix, and let  $\{\vec{x}_1, \vec{x}_2, \dots, \vec{x}_k\}$  denote a set of basic solutions to  $A\vec{x} = \vec{0}_m$ . Then

$$\text{null}(A) = \text{span}\{\vec{x}_1, \dots, \vec{x}_k\}.$$

### Lemma

Let  $A$  be an  $m \times n$  matrix with columns  $\vec{c}_1, \vec{c}_2, \dots, \vec{c}_n$ . Then

$$\text{im}(A) = \text{span}\{\vec{c}_1, \vec{c}_2, \dots, \vec{c}_n\}.$$


Proof. (of  $\text{null}(A) = \text{span}\{\vec{x}_1, \dots, \vec{x}_k\}$ )

" $\supseteq$ :" Because  $\vec{x}_i \in \text{null}(A)$  for each  $i$ ,  $1 \leq i \leq k$ , it follows that

$$\text{span}\{\vec{x}_1, \vec{x}_2, \dots, \vec{x}_k\} \subseteq \text{null}(A).$$

" $\subseteq$ :" Every solution to  $A\vec{x} = \vec{0}_m$  can be expressed as a linear combination of basic solutions, implying that

$$\text{null}(A) \subseteq \text{span}\{\vec{x}_1, \vec{x}_2, \dots, \vec{x}_k\}.$$

Therefore,  $\text{null}(A) = \text{span}\{\vec{x}_1, \vec{x}_2, \dots, \vec{x}_k\}$ . 

Proof. (of  $\text{im}(A) = \text{span}\{\vec{c}_1, \vec{c}_2, \dots, \vec{c}_n\}$ )

" $\subseteq$ :" Suppose  $\vec{y} \in \text{im}(A)$ . Then (by definition) there is a vector  $\vec{x} \in \mathbb{R}^n$  so that  $\vec{y} = A\vec{x}$ . Write  $\vec{x} = \begin{bmatrix} x_1 & x_2 & \dots & x_n \end{bmatrix}^T$ . Then

$$\vec{y} = A\vec{x} = \begin{bmatrix} \vec{c}_1 & \vec{c}_2 & \dots & \vec{c}_n \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = x_1\vec{c}_1 + x_2\vec{c}_2 + \dots + x_n\vec{c}_n.$$

Therefore,  $\vec{y} \in \text{span}\{\vec{c}_1, \vec{c}_2, \dots, \vec{c}_n\}$ , implying that

$$\text{im}(A) \subseteq \text{span}\{\vec{c}_1, \vec{c}_2, \dots, \vec{c}_n\}.$$



Proof. (continued)

Notice that for each  $j$ ,  $1 \leq j \leq n$ ,

$$\begin{aligned} A\vec{e}_j &= \begin{bmatrix} \vec{c}_1 & \vec{c}_2 & \dots & \vec{c}_n \end{bmatrix} \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ 1 \\ 0 \\ \vdots \\ 0 \end{bmatrix} \leftarrow j\text{th row} \\ &= 0\vec{c}_1 + 0\vec{c}_2 + \dots + 0\vec{c}_{j-1} + 1\vec{c}_j + 0\vec{c}_{j+1} + \dots + 0\vec{c}_n \\ &= \vec{c}_j. \end{aligned}$$

Thus  $\vec{c}_j \in \text{im}(A)$  for each  $j$ ,  $1 \leq j \leq n$ . It follows that

$$\text{span}\{\vec{c}_1, \vec{c}_2, \dots, \vec{c}_n\} \subseteq \text{im}(A),$$

and therefore

$$\text{im}(A) = \text{span}\{\vec{c}_1, \vec{c}_2, \dots, \vec{c}_n\}.$$

