

Math 221: LINEAR ALGEBRA

Chapter 3. Determinants and Diagonalization

§3-4. Application to Linear Recurrences

Le Chen¹

Emory University, 2021 Spring

(last updated on 01/12/2023)



Creative Commons License
(CC BY-NC-SA)

¹Slides are adapted from those by Karen Seyffarth from University of Calgary.

Linear Algebra with Applications

Lecture Notes

Current Lecture Notes Revision: Version 2018 — Revision B

These lecture notes were originally developed by Karen Seyffarth of the University of Calgary. Edits, additions, and revisions have been made to these notes by the editorial team at Lyryx Learning to accompany their text [Linear Algebra with Applications](#) based on W. K. Nicholson's original text.

In addition we recognize the following contributors. All new content contributed is released under the same license as noted below.

- Ilijas Farah, York University

BE A CHAMPION OF OER!

Contribute suggestions for improvements, new content, or errata:

A new topic

A new example or problem

A new or better proof to an existing theorem

Any other suggestions to improve the material

Contact Lyryx at info@lyryx.com with your ideas.

License



Attribution-NonCommercial-ShareAlike (CC BY-NC-SA)

This license lets others remix, tweak, and build upon your work non-commercially, as long as they credit you and license their new creations under the identical terms.

Copyright

Linear Recurrences

Linear Recurrences

Example

The Fibonacci Numbers are the numbers in the sequence

$$1, 1, 2, 3, 5, 8, 13, 21, 34, 55, 89, \dots$$

and can be defined by the **linear recurrence relation**

$$f_{n+2} = f_{n+1} + f_n \text{ for all } n \geq 0,$$

with the initial conditions $f_0 = 1$ and $f_1 = 1$.

Linear Recurrences

Example

The Fibonacci Numbers are the numbers in the sequence

$$1, 1, 2, 3, 5, 8, 13, 21, 34, 55, 89, \dots$$

and can be defined by the **linear recurrence relation**

$$f_{n+2} = f_{n+1} + f_n \text{ for all } n \geq 0,$$

with the initial conditions $f_0 = 1$ and $f_1 = 1$.

Problem

Find f_{100} .

Linear Recurrences

Example

The Fibonacci Numbers are the numbers in the sequence

$$1, 1, 2, 3, 5, 8, 13, 21, 34, 55, 89, \dots$$

and can be defined by the **linear recurrence relation**

$$f_{n+2} = f_{n+1} + f_n \text{ for all } n \geq 0,$$

with the initial conditions $f_0 = 1$ and $f_1 = 1$.

Problem

Find f_{100} .

Instead of using the recurrence to compute f_{100} , we'd like to find a formula for f_n that holds for all $n \geq 0$.

Definitions

A sequence of numbers $x_0, x_1, x_2, x_3, \dots$ is defined **recursively** if each number in the sequence is determined by the numbers that occur before it in the sequence.

Definitions

A sequence of numbers $x_0, x_1, x_2, x_3, \dots$ is defined **recursively** if each number in the sequence is determined by the numbers that occur before it in the sequence.

A **linear recurrence** of **length k** has the form

$$x_{n+k} = a_1 x_{n+k-1} + a_2 x_{n+k-2} + \cdots + a_k x_n, n \geq 0,$$

for some real numbers a_1, a_2, \dots, a_k .

Example

The simplest linear recurrence has length one, so has the form

$$x_{n+1} = ax_n \text{ for } n \geq 0,$$

with $a \in \mathbb{R}$ and some initial value x_0 .

Example

The simplest linear recurrence has length one, so has the form

$$x_{n+1} = ax_n \text{ for } n \geq 0,$$

with $a \in \mathbb{R}$ and some initial value x_0 .

In this case,

$$x_1 = ax_0$$

$$x_2 = ax_1 = a^2 x_0$$

$$x_3 = ax_2 = a^3 x_0$$

$$\vdots \quad \vdots \quad \vdots$$

$$x_n = ax_{n-1} = a^n x_0$$

Therefore, $x_n = a^n x_0$.

Example

Find a formula for x_n if

$$x_{n+2} = 2x_{n+1} + 3x_n \text{ for } n \geq 0,$$

with $x_0 = 0$ and $x_1 = 1$.

Example

Find a formula for x_n if

$$x_{n+2} = 2x_{n+1} + 3x_n \text{ for } n \geq 0,$$

with $x_0 = 0$ and $x_1 = 1$.

Solution. Define $V_n = \begin{bmatrix} x_n \\ x_{n+1} \end{bmatrix}$ for each $n \geq 0$. Then

$$V_0 = \begin{bmatrix} x_0 \\ x_1 \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \end{bmatrix},$$

and for $n \geq 0$,

$$V_{n+1} = \begin{bmatrix} x_{n+1} \\ x_{n+2} \end{bmatrix} = \begin{bmatrix} x_{n+1} \\ 2x_{n+1} + 3x_n \end{bmatrix}$$

Example (continued)

Now express $V_{n+1} = \begin{bmatrix} x_{n+1} \\ 2x_{n+1} + 3x_n \end{bmatrix}$ as a matrix product:

Example (continued)

Now express $V_{n+1} = \begin{bmatrix} x_{n+1} \\ 2x_{n+1} + 3x_n \end{bmatrix}$ as a matrix product:

$$V_{n+1} = \begin{bmatrix} x_{n+1} \\ 2x_{n+1} + 3x_n \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ 3 & 2 \end{bmatrix} \begin{bmatrix} x_n \\ x_{n+1} \end{bmatrix} = AV_n$$

Example (continued)

Now express $V_{n+1} = \begin{bmatrix} x_{n+1} \\ 2x_{n+1} + 3x_n \end{bmatrix}$ as a matrix product:

$$V_{n+1} = \begin{bmatrix} x_{n+1} \\ 2x_{n+1} + 3x_n \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ 3 & 2 \end{bmatrix} \begin{bmatrix} x_n \\ x_{n+1} \end{bmatrix} = AV_n$$

This is a linear dynamical system, so we can apply the techniques from §3.3, provided that A is diagonalizable.

Example (continued)

Now express $V_{n+1} = \begin{bmatrix} x_{n+1} \\ 2x_{n+1} + 3x_n \end{bmatrix}$ as a matrix product:

$$V_{n+1} = \begin{bmatrix} x_{n+1} \\ 2x_{n+1} + 3x_n \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ 3 & 2 \end{bmatrix} \begin{bmatrix} x_n \\ x_{n+1} \end{bmatrix} = AV_n$$

This is a linear dynamical system, so we can apply the techniques from §3.3, provided that A is diagonalizable.

$$c_A(x) = \det(xI - A) = \begin{vmatrix} x & -1 \\ -3 & x - 2 \end{vmatrix} = x^2 - 2x - 3 = (x - 3)(x + 1)$$

Therefore A has eigenvalues $\lambda_1 = 3$ and $\lambda_2 = -1$, and **is diagonalizable**.

Example (continued)

$\vec{x}_1 = \begin{bmatrix} 1 \\ 3 \end{bmatrix}$ is a basic eigenvector corresponding to $\lambda_1 = 3$, and

$\vec{x}_2 = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$ is a basic eigenvector corresponding to $\lambda_2 = -1$.

Example (continued)

$\vec{x}_1 = \begin{bmatrix} 1 \\ 3 \end{bmatrix}$ is a basic eigenvector corresponding to $\lambda_1 = 3$, and

$\vec{x}_2 = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$ is a basic eigenvector corresponding to $\lambda_2 = -1$.

Furthermore $P = \begin{bmatrix} \vec{x}_1 & \vec{x}_2 \end{bmatrix} = \begin{bmatrix} 1 & -1 \\ 3 & 1 \end{bmatrix}$ is invertible and is the

diagonalizing matrix for A, and $P^{-1}AP = D = \begin{bmatrix} 3 & 0 \\ 0 & -1 \end{bmatrix}$

Example (continued)

$\vec{x}_1 = \begin{bmatrix} 1 \\ 3 \end{bmatrix}$ is a basic eigenvector corresponding to $\lambda_1 = 3$, and

$\vec{x}_2 = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$ is a basic eigenvector corresponding to $\lambda_2 = -1$.

Furthermore $P = \begin{bmatrix} \vec{x}_1 & \vec{x}_2 \end{bmatrix} = \begin{bmatrix} 1 & -1 \\ 3 & 1 \end{bmatrix}$ is invertible and is the

diagonalizing matrix for A, and $P^{-1}AP = D = \begin{bmatrix} 3 & 0 \\ 0 & -1 \end{bmatrix}$

Writing $P^{-1}V_0 = \begin{bmatrix} b_1 \\ b_2 \end{bmatrix}$, we get

$$\begin{bmatrix} b_1 \\ b_2 \end{bmatrix} = \frac{1}{4} \begin{bmatrix} 1 & 1 \\ -3 & 1 \end{bmatrix} \begin{bmatrix} 0 \\ 1 \end{bmatrix} = \begin{bmatrix} \frac{1}{4} \\ \frac{1}{4} \end{bmatrix}$$

Example (continued)

Therefore,

$$\begin{aligned} V_n = \begin{bmatrix} x_n \\ x_{n+1} \end{bmatrix} &= b_1 \lambda_1^n \vec{x}_1 + b_2 \lambda_2^n \vec{x}_2 \\ &= \frac{1}{4} 3^n \begin{bmatrix} 1 \\ 3 \end{bmatrix} + \frac{1}{4} (-1)^n \begin{bmatrix} -1 \\ 1 \end{bmatrix}, \end{aligned}$$

and so

$$x_n = \frac{1}{4} 3^n - \frac{1}{4} (-1)^n.$$

Example

Solve the recurrence relation

$$x_{k+2} = 5x_{k+1} - 6x_k, k \geq 0$$

with $x_0 = 0$ and $x_1 = 1$.

Example

Solve the recurrence relation

$$x_{k+2} = 5x_{k+1} - 6x_k, k \geq 0$$

with $x_0 = 0$ and $x_1 = 1$.

Solution. Write

$$V_{k+1} = \begin{bmatrix} x_{k+1} \\ x_{k+2} \end{bmatrix} = \begin{bmatrix} x_{k+1} \\ 5x_{k+1} - 6x_k \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ -6 & 5 \end{bmatrix} \begin{bmatrix} x_k \\ x_{k+1} \end{bmatrix}$$

Example

Solve the recurrence relation

$$x_{k+2} = 5x_{k+1} - 6x_k, k \geq 0$$

with $x_0 = 0$ and $x_1 = 1$.

Solution. Write

$$V_{k+1} = \begin{bmatrix} x_{k+1} \\ x_{k+2} \end{bmatrix} = \begin{bmatrix} x_{k+1} \\ 5x_{k+1} - 6x_k \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ -6 & 5 \end{bmatrix} \begin{bmatrix} x_k \\ x_{k+1} \end{bmatrix}$$

Find the eigenvalues and corresponding eigenvectors for

$$A = \begin{bmatrix} 0 & 1 \\ -6 & 5 \end{bmatrix}$$

Example (continued)

A has eigenvalues $\lambda_1 = 2$ with corresponding eigenvector $\vec{x}_1 = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$, and $\lambda_2 = 3$ with corresponding eigenvector $\vec{x}_2 = \begin{bmatrix} 1 \\ 3 \end{bmatrix}$.

Example (continued)

A has eigenvalues $\lambda_1 = 2$ with corresponding eigenvector $\vec{x}_1 = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$, and

$\lambda_2 = 3$ with corresponding eigenvector $\vec{x}_2 = \begin{bmatrix} 1 \\ 3 \end{bmatrix}$.

$$P = \begin{bmatrix} 1 & 1 \\ 2 & 3 \end{bmatrix}, P^{-1} = \begin{bmatrix} 3 & -1 \\ -2 & 1 \end{bmatrix},$$

and

$$\begin{bmatrix} b_1 \\ b_2 \end{bmatrix} = P^{-1}V_0 = \begin{bmatrix} 3 & -1 \\ -2 & 1 \end{bmatrix} \begin{bmatrix} 0 \\ 1 \end{bmatrix} = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

Example (continued)

A has eigenvalues $\lambda_1 = 2$ with corresponding eigenvector $\vec{x}_1 = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$, and $\lambda_2 = 3$ with corresponding eigenvector $\vec{x}_2 = \begin{bmatrix} 1 \\ 3 \end{bmatrix}$.

$$P = \begin{bmatrix} 1 & 1 \\ 2 & 3 \end{bmatrix}, P^{-1} = \begin{bmatrix} 3 & -1 \\ -2 & 1 \end{bmatrix},$$

and

$$\begin{bmatrix} b_1 \\ b_2 \end{bmatrix} = P^{-1}V_0 = \begin{bmatrix} 3 & -1 \\ -2 & 1 \end{bmatrix} \begin{bmatrix} 0 \\ 1 \end{bmatrix} = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

Finally,

$$V_k = \begin{bmatrix} x_k \\ x_{k+1} \end{bmatrix} = b_1 \lambda_1^k \vec{x}_1 + b_2 \lambda_2^k \vec{x}_2 = (-1)2^k \begin{bmatrix} 1 \\ 2 \end{bmatrix} + 3^k \begin{bmatrix} 1 \\ 3 \end{bmatrix}$$

Example

$$\begin{bmatrix} x_k \\ x_{k+1} \end{bmatrix} = (-1)2^k \begin{bmatrix} 1 \\ 2 \end{bmatrix} + 3^k \begin{bmatrix} 1 \\ 3 \end{bmatrix}$$

and therefore

$$x_k = 3^k - 2^k.$$