

Math 221: LINEAR ALGEBRA

Chapter 8. Orthogonality

§8-4. QR Factorization

Le Chen¹

Emory University, 2021 Spring

(last updated on 01/12/2023)



Creative Commons License
(CC BY-NC-SA)

¹Slides are adapted from those by Karen Seyffarth from University of Calgary.

Linear Algebra with Applications

Lecture Notes

Current Lecture Notes Revision: Version 2018 — Revision B

These lecture notes were originally developed by Karen Seyffarth of the University of Calgary. Edits, additions, and revisions have been made to these notes by the editorial team at Lyryx Learning to accompany their text [Linear Algebra with Applications](#) based on W. K. Nicholson's original text.

In addition we recognize the following contributors. All new content contributed is released under the same license as noted below.

- Ilijas Farah, York University

BE A CHAMPION OF OER!

Contribute suggestions for improvements, new content, or errata:

A new topic

A new example or problem

A new or better proof to an existing theorem

Any other suggestions to improve the material

Contact Lyryx at info@lyryx.com with your ideas.

License



Attribution-NonCommercial-ShareAlike (CC BY-NC-SA)

This license lets others remix, tweak, and build upon your work non-commercially, as long as they credit you and license their new creations under the identical terms.

Copyright

QR Factorization

Algorithm for the QR Factorization

Copyright

QR Factorization

Algorithm for the QR Factorization

The QR Factorization

Definition

Let A be a real $m \times n$ matrix. Then a **QR factorization** of A can be written as

$$A = QR$$

where Q is an orthogonal matrix and R is an upper (or right) triangular matrix.

The diagram illustrates the QR factorization of a matrix A into an orthogonal matrix Q and an upper triangular matrix R . Matrix A is shown as a 3x3 matrix with columns \mathbf{a}_1 , \mathbf{a}_2 , and \mathbf{a}_3 . Matrix Q is shown as a 3x3 matrix with columns \mathbf{e}_1 , \mathbf{e}_2 , and \mathbf{e}_3 , which are orthogonal unit vectors. Matrix R is shown as a 3x3 upper triangular matrix with elements $\mathbf{e}_1^T \cdot \mathbf{a}_1$, $\mathbf{e}_1^T \cdot \mathbf{a}_2$, $\mathbf{e}_1^T \cdot \mathbf{a}_3$ in the first row, $\mathbf{0}$, $\mathbf{e}_2^T \cdot \mathbf{a}_2$, $\mathbf{e}_2^T \cdot \mathbf{a}_3$ in the second row, and $\mathbf{0}$, $\mathbf{0}$, $\mathbf{e}_3^T \cdot \mathbf{a}_3$ in the third row. The equation is $A = QR$.

$$\begin{bmatrix} | & | & | \\ \mathbf{a}_1 & \mathbf{a}_2 & \mathbf{a}_3 \\ | & | & | \end{bmatrix} = \underbrace{\begin{bmatrix} | & | & | \\ \mathbf{e}_1 & \mathbf{e}_2 & \mathbf{e}_3 \\ | & | & | \end{bmatrix}}_{\text{orthogonal unit vector}} \underbrace{\begin{bmatrix} \mathbf{e}_1^T \cdot \mathbf{a}_1 & \mathbf{e}_1^T \cdot \mathbf{a}_2 & \mathbf{e}_1^T \cdot \mathbf{a}_3 \\ \mathbf{0} & \mathbf{e}_2^T \cdot \mathbf{a}_2 & \mathbf{e}_2^T \cdot \mathbf{a}_3 \\ \mathbf{0} & \mathbf{0} & \mathbf{e}_3^T \cdot \mathbf{a}_3 \end{bmatrix}}_{\text{upper diagonal matrix}}$$

Theorem


Let A be a real $m \times n$ matrix with linearly independent columns. Then A can be written

$$A = QR$$


with Q orthogonal and R upper triangular with positive entries on the main diagonal.

Proof.

Using columns of A to carry out the Gram-Schmidt algorithm to find an orthonormal basis for $\text{im}(A)$ or $\text{col}(A) \subseteq \mathbb{R}^m$ – columns of Q_1 . One may further extend this basis to an orthonormal basis for the whole space \mathbb{R}^m – columns of $Q = [Q_1, Q_2]$.



The Gram-Schmidt algorithm guarantees that the i th column of A is linear combinations of all j th columns of Q with $j = 1, \dots, i$, which gives the upper triangular structure of R . ■



Remark

$$A = QR = [Q_1, Q_2] \begin{bmatrix} R_1 \\ O \end{bmatrix} = Q_1 R_1 + Q_2 O = Q_1 R_1.$$

Both QR and $Q_1 R_1$ are called QR decompositions of A . The textbook refers $Q_1 R_1$.

Remark

Q is orthogonal matrix, namely, $QQ^T = Q^T Q = I_m$.

However, Q_1 is not orthogonal matrix (not a square matrix). But We have $Q_1^T Q_1 = I_n$ and $Q_1 Q_1^T \neq I_m$ (in general).

Copyright

QR Factorization

Algorithm for the QR Factorization

Algorithm for QR Factorization

Algorithm 1: QR Factorization Algorithm

Input : Independent columns of A: $\{\vec{c}_1, \vec{c}_2, \dots, \vec{c}_n\} \in \text{col}(A) \subseteq \mathbb{R}^m$

for $j \leftarrow 1$ to n do

$$\vec{f}_j \leftarrow \vec{c}_j - \frac{\vec{c}_j \cdot \vec{f}_1}{\|\vec{f}_1\|^2} \vec{f}_1 - \frac{\vec{c}_j \cdot \vec{f}_2}{\|\vec{f}_2\|^2} \vec{f}_2 - \dots - \frac{\vec{c}_j \cdot \vec{f}_{j-1}}{\|\vec{f}_{j-1}\|^2} \vec{f}_{j-1}.$$

$$\vec{q}_j \leftarrow \frac{\vec{f}_j}{\|\vec{f}_j\|}$$

for $i \leftarrow 1$ to j do

$$r_{ij} \leftarrow \vec{q}_i \cdot \vec{c}_j$$

end

end

Output: $Q = [\vec{q}_1, \dots, \vec{q}_n]$ and $R = [r_{ij}]$

Problem

Let

$$A = \begin{bmatrix} 4 & 1 \\ 2 & 3 \\ 0 & 1 \end{bmatrix}$$

Find the QR factorization of A.

Solution

Set $A = [\vec{c}_1, \vec{c}_2]$. When $j = 1$,

$$\vec{f}_1 = \vec{c}_1 = \begin{bmatrix} 4 \\ 2 \\ 0 \end{bmatrix} \quad \text{and} \quad \vec{q}_1 = \frac{\vec{f}_1}{\|\vec{f}_1\|} = \begin{bmatrix} \frac{4}{\sqrt{20}} \\ \frac{2}{\sqrt{20}} \\ 0 \end{bmatrix}.$$

For $i = 1$,

$$r_{11} = \vec{q}_1 \cdot \vec{c}_1 = \frac{\vec{f}_1}{\|\vec{f}_1\|} \cdot \vec{f}_1 = \|\vec{f}_1\| = \sqrt{20}.$$

Solution (continued)

When $j = 2$,

$$\vec{f}_2 = \vec{c}_2 - \frac{\vec{c}_2 \cdot \vec{f}_1}{\|\vec{f}_1\|^2} \vec{f}_1 = \begin{bmatrix} 1 \\ 3 \\ 1 \end{bmatrix} - \frac{10}{20} \begin{bmatrix} 4 \\ 2 \\ 0 \end{bmatrix} = \begin{bmatrix} -1 \\ 2 \\ 1 \end{bmatrix} \quad \text{and} \quad \vec{q}_2 = \frac{\vec{f}_2}{\|\vec{f}_2\|} = \begin{bmatrix} -\frac{1}{\sqrt{6}} \\ \frac{2}{\sqrt{6}} \\ \frac{1}{\sqrt{6}} \end{bmatrix}.$$

For $i = 1$,

$$r_{12} = \vec{q}_1 \cdot \vec{c}_2 = \begin{bmatrix} \frac{4}{\sqrt{20}} \\ \frac{2}{\sqrt{20}} \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 3 \\ 1 \end{bmatrix} = \sqrt{5}.$$

and for $i = 2$,

$$r_{22} = \vec{q}_2 \cdot \vec{c}_2 = \frac{\vec{f}_2}{\|\vec{f}_2\|} \cdot \left(\vec{f}_2 + \frac{\vec{c}_2 \cdot \vec{f}_1}{\|\vec{f}_1\|^2} \vec{f}_1 \right) = \frac{\vec{f}_2}{\|\vec{f}_2\|} \cdot \vec{f}_2 = \|\vec{f}_2\| = \sqrt{6}.$$

Solution (continued)

Therefore,

$$A = QR = [\vec{q}_1, \vec{q}_2] \begin{bmatrix} r_{11} & r_{12} \\ 0 & r_{22} \end{bmatrix}$$

$$\Updownarrow$$

$$\begin{bmatrix} 4 & 1 \\ 2 & 3 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} \frac{2}{\sqrt{5}} & -\frac{1}{\sqrt{6}} \\ \frac{1}{\sqrt{5}} & \frac{\sqrt{6}}{\sqrt{6}} \\ 0 & \frac{1}{\sqrt{6}} \end{bmatrix} \begin{bmatrix} \sqrt{20} & \sqrt{5} \\ 0 & \sqrt{6} \end{bmatrix}$$

