

Math 221: LINEAR ALGEBRA

Chapter 2. Matrix Algebra

§2-7. LU Factorization

Le Chen¹

Emory University, 2021 Spring

(last updated on 01/12/2023)



Creative Commons License
(CC BY-NC-SA)

¹Slides are adapted from those by Karen Seyffarth from University of Calgary.

Linear Algebra with Applications

Lecture Notes

Current Lecture Notes Revision: Version 2018 — Revision B

These lecture notes were originally developed by Karen Seyffarth of the University of Calgary. Edits, additions, and revisions have been made to these notes by the editorial team at Lyryx Learning to accompany their text [Linear Algebra with Applications](#) based on W. K. Nicholson's original text.

In addition we recognize the following contributors. All new content contributed is released under the same license as noted below.

- Ilijas Farah, York University

BE A CHAMPION OF OER!

Contribute suggestions for improvements, new content, or errata:

A new topic

A new example or problem

A new or better proof to an existing theorem

Any other suggestions to improve the material

Contact Lyryx at info@lyryx.com with your ideas.

License



Attribution-NonCommercial-ShareAlike (CC BY-NC-SA)

This license lets others remix, tweak, and build upon your work non-commercially, as long as they credit you and license their new creations under the identical terms.

Copyright

LU Factorization

Why do we need LU Factorization?

Finding the LU

Multiplier Method

LU-Algorithm

Copyright

LU Factorization

Why do we need LU Factorization?

Finding the LU

Multiplier Method

LU-Algorithm

LU Factorization

Definition

A matrix $A = [a_{ij}]$ is called **upper triangular** if $a_{ij} = 0$ whenever $i > j$. Thus the entries below the main diagonal equal 0.

LU Factorization

Definition

A matrix $A = [a_{ij}]$ is called **upper triangular** if $a_{ij} = 0$ whenever $i > j$. Thus the entries below the main diagonal equal 0.



LU Factorization

Definition

A matrix $A = [a_{ij}]$ is called **upper triangular** if $a_{ij} = 0$ whenever $i > j$. Thus the entries below the main diagonal equal 0.



A **lower triangular matrix** is defined similarly, as a matrix for which all entries above the main diagonal are equal to zero.


LU Factorization

Definition

A matrix $A = [a_{ij}]$ is called **upper triangular** if $a_{ij} = 0$ whenever $i > j$. Thus the entries below the main diagonal equal 0.



A **lower triangular matrix** is defined similarly, as a matrix for which all entries above the main diagonal are equal to zero.



An **LU factorization** of a matrix A is written

$$A = LU$$

where L is lower triangular matrix and U is upper triangular.

An **LU factorization** of a matrix A is written

$$A = LU$$

where L is lower triangular matrix and U is upper triangular.

We often require either L or U to have only 1's on the main diagonal.

An **LU factorization** of a matrix A is written

$$A = LU$$

where L is lower triangular matrix and U is upper triangular.

We often require either L or U to have only 1's on the main diagonal.

$$A = \begin{pmatrix} \textcolor{red}{1} & 0 & \cdots & 0 \\ * & \textcolor{red}{1} & \ddots & \vdots \\ \vdots & \ddots & \ddots & 0 \\ * & \cdots & * & \textcolor{red}{1} \end{pmatrix} \begin{pmatrix} * & * & \cdots & * \\ 0 & * & \ddots & \vdots \\ \vdots & \ddots & \ddots & * \\ 0 & \cdots & 0 & * \end{pmatrix}$$

Copyright

LU Factorization

Why do we need LU Factorization?

Finding the LU

Multiplier Method

LU-Algorithm

Why do we need LU Factorization?

Why do we need LU Factorization?

The LU factorization often helps to quickly solve equations of the form $A\vec{x} = \vec{b}$.

Why do we need LU Factorization?

The LU factorization often helps to quickly solve equations of the form $A\vec{x} = \vec{b}$.

Suppose we wish to find all solutions \vec{x} to the system $A\vec{x} = B$. The LU factorization of A can assist in this process.

Why do we need LU Factorization?

The LU factorization often helps to quickly solve equations of the form $A\vec{x} = \vec{b}$.

Suppose we wish to find all solutions \vec{x} to the system $A\vec{x} = B$. The LU factorization of A can assist in this process.

Consider the following reduction:

$$\begin{aligned} A\vec{x} &= B \\ (LU)\vec{x} &= B \\ L(U\vec{x}) &= B \\ L\vec{y} &= B \end{aligned}$$

Why do we need LU Factorization?

The LU factorization often helps to quickly solve equations of the form $A\vec{x} = \vec{b}$.

Suppose we wish to find all solutions \vec{x} to the system $A\vec{x} = B$. The LU factorization of A can assist in this process.

Consider the following reduction:

$$\begin{aligned}A\vec{x} &= B \\(LU)\vec{x} &= B \\L(U\vec{x}) &= B \\L\vec{y} &= B\end{aligned}$$

Therefore, if we can solve $L\vec{y} = B$ for \vec{y} , then all that remains is to solve $U\vec{x} = \vec{y}$ for \vec{x} .

Example

Find all solutions to

$$\begin{bmatrix} 1 & 3 & 2 & 0 \\ 3 & 10 & 5 & 1 \\ 0 & -1 & 2 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} 2 \\ 4 \\ 6 \end{bmatrix}$$

Example

Find all solutions to

$$\begin{bmatrix} 1 & 3 & 2 & 0 \\ 3 & 10 & 5 & 1 \\ 0 & -1 & 2 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} 2 \\ 4 \\ 6 \end{bmatrix}$$

Solution

Using a method of your choice, verify that the LU factorization of A gives

$$L = \begin{bmatrix} 1 & 0 & 0 \\ 3 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix}, U = \begin{bmatrix} 1 & 3 & 2 & 0 \\ 0 & 1 & -1 & 1 \\ 0 & 0 & 1 & 2 \end{bmatrix}$$

Solution (continued)

Let $\vec{y} = \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix}$ and solve $L\vec{y} = \vec{b}$.

Solution (continued)

Let $\vec{y} = \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix}$ and solve $L\vec{y} = \vec{b}$.

$$\begin{bmatrix} 1 & 0 & 0 \\ 3 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix} \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix} = \begin{bmatrix} 2 \\ 4 \\ 6 \end{bmatrix}$$

Solution (continued)

Let $\vec{y} = \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix}$ and solve $L\vec{y} = \vec{b}$.

$$\begin{bmatrix} 1 & 0 & 0 \\ 3 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix} \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix} = \begin{bmatrix} 2 \\ 4 \\ 6 \end{bmatrix}$$

The solution is $\vec{y} = \begin{bmatrix} 2 \\ -2 \\ 4 \end{bmatrix}$.

Solution (continued)

Let $\vec{y} = \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix}$ and solve $L\vec{y} = \vec{b}$.

$$\begin{bmatrix} 1 & 0 & 0 \\ 3 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix} \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix} = \begin{bmatrix} 2 \\ 4 \\ 6 \end{bmatrix}$$

The solution is $\vec{y} = \begin{bmatrix} 2 \\ -2 \\ 4 \end{bmatrix}$.

Now we solve $U\vec{x} = \vec{y}$.

$$\begin{bmatrix} 1 & 3 & 2 & 0 \\ 0 & 1 & -1 & 1 \\ 0 & 0 & 1 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} 2 \\ -2 \\ 4 \end{bmatrix}$$

Solution (continued)

Multiplying and solving (or finding the reduced row-echelon form), the general solution is given by

$$\vec{x} = \begin{bmatrix} -12 \\ 2 \\ 4 \\ 0 \end{bmatrix} + \begin{bmatrix} 13 \\ -3 \\ -2 \\ 1 \end{bmatrix} t, \quad \forall t \in \mathbb{R}.$$



Copyright

LU Factorization

Why do we need LU Factorization?

Finding the LU

Multiplier Method

LU-Algorithm

Finding the LU Factorization

Finding the LU Factorization

Condition for the existence of LU factorization: A matrix A has LU factorization provided that A can be **lower reduced**, namely, the row-echelon form of A can be calculated without interchanging rows.

Finding the LU Factorization

Condition for the existence of LU factorization: A matrix A has LU factorization provided that A can be **lower reduced**, namely, the row-echelon form of A can be calculated without interchanging rows.

Example

Determine if the LU factorization of A exists, and if so, find it.

$$A = \begin{bmatrix} 1 & 1 & 2 \\ 2 & 3 & 0 \\ 1 & 0 & 5 \end{bmatrix}$$

Finding the LU Factorization

Condition for the existence of LU factorization: A matrix A has LU factorization provided that A can be **lower reduced**, namely, the row-echelon form of A can be calculated without interchanging rows.

Example

Determine if the LU factorization of A exists, and if so, find it.

$$A = \begin{bmatrix} 1 & 1 & 2 \\ 2 & 3 & 0 \\ 1 & 0 & 5 \end{bmatrix}$$

Solution

Because the row-echelon form can be obtained without interchanging rows:

Finding the LU Factorization

Condition for the existence of LU factorization: A matrix A has LU factorization provided that A can be **lower reduced**, namely, the row-echelon form of A can be calculated without interchanging rows.

Example

Determine if the LU factorization of A exists, and if so, find it.

$$A = \begin{bmatrix} 1 & 1 & 2 \\ 2 & 3 & 0 \\ 1 & 0 & 5 \end{bmatrix}$$

Solution

Because the row-echelon form can be obtained without interchanging rows:

$$\begin{bmatrix} 1 & 1 & 2 \\ 2 & 3 & 0 \\ 1 & 0 & 5 \end{bmatrix} \xrightarrow{r_2 - 2r_1} \begin{bmatrix} 1 & 1 & 2 \\ 0 & 1 & -4 \\ 1 & 0 & 5 \end{bmatrix} \xrightarrow{r_3 - r_1} \begin{bmatrix} 1 & 1 & 2 \\ 0 & 1 & -4 \\ 0 & -1 & 3 \end{bmatrix} \xrightarrow{r_3 + r_2} \begin{bmatrix} 1 & 1 & 2 \\ 0 & 1 & -4 \\ 0 & 0 & -1 \end{bmatrix}$$

the LU factorization exists, or A can be lower reduced.

Solution (continued)

We proceed to finding L and U. Assign variables to the unknown entries and multiply.

$$\begin{aligned} A = \begin{bmatrix} 1 & 1 & 2 \\ 2 & 3 & 0 \\ 1 & 0 & 5 \end{bmatrix} &= \begin{bmatrix} 1 & 0 & 0 \\ x & 1 & 0 \\ y & z & 1 \end{bmatrix} \begin{bmatrix} a & d & e \\ 0 & b & f \\ 0 & 0 & c \end{bmatrix} \\ &= \begin{bmatrix} a & d & e \\ ax & dx + b & ex + f \\ ay & dy + bz & ey + fz + c \end{bmatrix} \end{aligned}$$

Solution (continued)

We proceed to finding L and U. Assign variables to the unknown entries and multiply.

$$\begin{aligned} A = \begin{bmatrix} 1 & 1 & 2 \\ 2 & 3 & 0 \\ 1 & 0 & 5 \end{bmatrix} &= \begin{bmatrix} 1 & 0 & 0 \\ x & 1 & 0 \\ y & z & 1 \end{bmatrix} \begin{bmatrix} a & d & e \\ 0 & b & f \\ 0 & 0 & c \end{bmatrix} \\ &= \begin{bmatrix} a & d & e \\ ax & dx + b & ex + f \\ ay & dy + bz & ey + fz + c \end{bmatrix} \end{aligned}$$

Solving each entry will give us values for the unknown entries.

Solution (continued)

$$\begin{bmatrix} 1 & 1 & 2 \\ 2 & 3 & 0 \\ 1 & 0 & 5 \end{bmatrix} = \begin{bmatrix} a & d & e \\ ax & dx + b & ex + f \\ ay & dy + bz & ey + fz + c \end{bmatrix}$$

Solution (continued)

$$\begin{bmatrix} 1 & 1 & 2 \\ 2 & 3 & 0 \\ 1 & 0 & 5 \end{bmatrix} = \begin{bmatrix} a & d & e \\ ax & dx + b & ex + f \\ ay & dy + bz & ey + fz + c \end{bmatrix}$$

We see easily that $a = 1$, $d = 1$, and $e = 2$. Continuing to solve the first column gives $x = 2$, $y = 1$. The other values are calculated as follows.

Solution (continued)

$$\begin{bmatrix} 1 & 1 & 2 \\ 2 & 3 & 0 \\ 1 & 0 & 5 \end{bmatrix} = \begin{bmatrix} a & d & e \\ ax & dx + b & ex + f \\ ay & dy + bz & ey + fz + c \end{bmatrix}$$

We see easily that $a = 1$, $d = 1$, and $e = 2$. Continuing to solve the first column gives $x = 2$, $y = 1$. The other values are calculated as follows.

$$\begin{array}{rcl} dx + b & = & 3 \\ (1)(2) + b & = & 3 \\ b & = & 1 \end{array} \qquad \begin{array}{rcl} ex + f & = & 0 \\ (2)(2) + f & = & 0 \\ f & = & -4 \end{array}$$

$$\begin{array}{rcl} dy + bz & = & 0 \\ (1)(1) + (1)z & = & 0 \\ z & = & -1 \end{array} \qquad \begin{array}{rcl} ey + fz + c & = & 5 \\ (2)(1) + (-4)(-1) + c & = & 5 \\ c & = & -1 \end{array}$$

Solution (continued)

Therefore,

$$\begin{aligned} & \mathbf{L} \\ & \parallel \\ & \begin{bmatrix} 1 & 0 & 0 \\ x & 1 & 0 \\ y & z & 1 \end{bmatrix} \\ & \parallel \\ & \begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 1 & -1 & 1 \end{bmatrix} \end{aligned}$$

Solution (continued)

Therefore,

$$\begin{array}{c} L \\ || \\ \left[\begin{array}{ccc} 1 & 0 & 0 \\ x & 1 & 0 \\ y & z & 1 \end{array} \right] \\ || \\ \left[\begin{array}{ccc} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 1 & -1 & 1 \end{array} \right] \end{array}$$

$$\begin{array}{c} U \\ || \\ \left[\begin{array}{ccc} a & d & e \\ 0 & b & f \\ 0 & 0 & c \end{array} \right] \\ || \\ \left[\begin{array}{ccc} 1 & 1 & 2 \\ 0 & 1 & -4 \\ 0 & 0 & -1 \end{array} \right] \end{array}$$



Remark

If you want the diagonal terms of U to be all 1's:

$$\begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 1 & -1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{bmatrix} \begin{bmatrix} 1 & 1 & 2 \\ 0 & 1 & -4 \\ 0 & 0 & -1 \end{bmatrix}$$

||

$$\underbrace{\begin{bmatrix} 1 & 0 & -0 \\ 2 & 1 & -0 \\ 1 & -1 & -1 \end{bmatrix}}_L \quad \underbrace{\begin{bmatrix} 1 & 1 & 2 \\ 0 & 1 & -4 \\ -0 & -0 & 1 \end{bmatrix}}_U$$



Copyright

LU Factorization

Why do we need LU Factorization?

Finding the LU

Multiplier Method

LU-Algorithm

Multiplier Method

The following process for finding L and U , called the **multiplier method**, can be more efficient.

Multiplier Method

The following process for finding L and U, called the **multiplier method**, can be more efficient.

Example

Find the LU factorization of $A = \begin{bmatrix} 1 & 1 & 2 \\ 2 & 3 & 0 \\ 1 & 0 & 5 \end{bmatrix}$

Multiplier Method

The following process for finding L and U, called the **multiplier method**, can be more efficient.

Example

Find the LU factorization of $A = \begin{bmatrix} 1 & 1 & 2 \\ 2 & 3 & 0 \\ 1 & 0 & 5 \end{bmatrix}$

Solution

First, write A as

$$\begin{bmatrix} 1 & 1 & 2 \\ 2 & 3 & 0 \\ 1 & 0 & 5 \end{bmatrix}$$

||

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 & 2 \\ 2 & 3 & 0 \\ 1 & 0 & 5 \end{bmatrix}$$

Multiplier Method

The following process for finding L and U, called the **multiplier method**, can be more efficient.

Example

Find the LU factorization of $A = \begin{bmatrix} 1 & 1 & 2 \\ 2 & 3 & 0 \\ 1 & 0 & 5 \end{bmatrix}$

Solution

First, write A as

$$\begin{bmatrix} 1 & 1 & 2 \\ 2 & 3 & 0 \\ 1 & 0 & 5 \end{bmatrix}$$

||

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 & 2 \\ 2 & 3 & 0 \\ 1 & 0 & 5 \end{bmatrix}$$

Solution (continued)

To do so, we use row operations to remove the entries of A below the main diagonal. For every operation we apply to A (the matrix on the right), we apply the inverse operation to the identity matrix (on the left). This ensures the product remains the same.

Solution (continued)

To do so, we use row operations to remove the entries of A below the main diagonal. For every operation we apply to A (the matrix on the right), we apply the inverse operation to the identity matrix (on the left). This ensures the product remains the same.

The first step is to add (-2) times the first row of A to the second row. To preserve the product, add (2) times the second column to the first column, for the matrix on the left.

||

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 & 2 \\ 2 & 3 & 0 \\ 1 & 0 & 5 \end{bmatrix}$$

||

$$c_1 + 2c_2 \rightarrow c_1 \quad \begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \quad \begin{bmatrix} 1 & 1 & 2 \\ 0 & 1 & -4 \\ 1 & 0 & 5 \end{bmatrix} \quad r_2 - 2r_1 \rightarrow r_2$$

Solution (continued)

We proceed in the same way.

$$c_1 + c_3 \rightarrow c_1 \quad \begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix} \quad \begin{bmatrix} 1 & 1 & 2 \\ 0 & 1 & -4 \\ 0 & -1 & 3 \end{bmatrix} \quad r_3 - r_1 \rightarrow r_3$$

Solution (continued)

We proceed in the same way.

$$c_1 + c_3 \rightarrow c_1 \quad \left[\begin{array}{ccc} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 1 & 0 & 1 \end{array} \right] \left[\begin{array}{ccc} 1 & 1 & 2 \\ 0 & 1 & -4 \\ 0 & -1 & 3 \end{array} \right] \quad r_3 - r_1 \rightarrow r_3$$

$$c_2 - c_3 \rightarrow c_2 \quad \left[\begin{array}{ccc} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 1 & -1 & 1 \end{array} \right] \left[\begin{array}{ccc} 1 & 1 & 2 \\ 0 & 1 & -4 \\ 0 & 0 & -1 \end{array} \right] \quad r_3 + r_2 \rightarrow r_3$$

Solution (continued)

We proceed in the same way.

$$c_1 + c_3 \rightarrow c_1 \quad \begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 & 2 \\ 0 & 1 & -4 \\ 0 & -1 & 3 \end{bmatrix} \quad r_3 - r_1 \rightarrow r_3$$

$$c_2 - c_3 \rightarrow c_2 \quad \begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 1 & -1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 & 2 \\ 0 & 1 & -4 \\ 0 & 0 & -1 \end{bmatrix} \quad r_3 + r_2 \rightarrow r_3$$

At this point we have a lower triangular matrix L on the left, and an upper triangular matrix U on the right so we are done. You can (and should!) check that this product equals A.

If you want the diagonal terms of U to be all 1's:

$$-1 \times c_3 \rightarrow c_3 \quad \begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 1 & -1 & -1 \end{bmatrix} \begin{bmatrix} 1 & 1 & 2 \\ 0 & 1 & -4 \\ 0 & 0 & 1 \end{bmatrix} \quad -1 \times r_3 \rightarrow r_3$$



Problem

Use the multiplier method to verify the LU factorization for

$$A = \begin{bmatrix} 1 & 4 & 2 \\ 3 & 13 & 5 \\ -2 & -7 & -4 \end{bmatrix}$$

Problem

Use the multiplier method to verify the LU factorization for

$$A = \begin{bmatrix} 1 & 4 & 2 \\ 3 & 13 & 5 \\ -2 & -7 & -4 \end{bmatrix}$$

Solution

$$A = \begin{bmatrix} 1 & 4 & 2 \\ 3 & 13 & 5 \\ -2 & -7 & -4 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 3 & 1 & 0 \\ -2 & 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 4 & 2 \\ 0 & 1 & -1 \\ 0 & 0 & 1 \end{bmatrix} = LU$$



Copyright

LU Factorization

Why do we need LU Factorization?

Finding the LU

Multiplier Method

LU-Algorithm

Theorem (LU-Algorithm)

Let A be an $m \times n$ matrix of rank r , and suppose that A can be lower reduced to a row-echelon matrix U . Then $A = LU$ where the lower triangular, invertible matrix L is constructed as follows:

1. If $A = 0$, take $L = I_m$ and $U = 0$.
2. If $A \neq 0$, write $A_1 = A$ and let \vec{c}_1 be the **leading column** of A_1 . Use \vec{c}_1 to create the first leading 1 and make its below all zeros. When this is completed, let A_2 denote the matrix consisting of rows 2 to m of the matrix just created.
3. If $A_2 \neq 0$, let \vec{c}_2 be the leading column of A_2 and repeat Step 2 on A_2 to create A_3 .
4. Continue in this way until U is reached, where all rows below the last leading 1 consist of zeros. This will happen after r steps.
5. Create L by placing $\vec{c}_1, \vec{c}_2, \dots, \vec{c}_r$ at the bottom of the first r columns of I_m .

Problem

Find an LU-factorization for $A = \begin{bmatrix} 2 & 4 & 2 \\ 1 & 1 & 2 \\ -1 & 0 & 2 \end{bmatrix}$.

Solution

$$\begin{bmatrix} 2 & 4 & 2 \\ 1 & 1 & 2 \\ -1 & 0 & 2 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 2 & 1 \\ 0 & -1 & 1 \\ 0 & 2 & 3 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 2 & 1 \\ 0 & 1 & -1 \\ 0 & 0 & 5 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 2 & 1 \\ 0 & 1 & -1 \\ 0 & 0 & 1 \end{bmatrix} = U$$

Problem

Find an LU-factorization for $A = \begin{bmatrix} 2 & 4 & 2 \\ 1 & 1 & 2 \\ -1 & 0 & 2 \end{bmatrix}$.

Solution

$$\begin{bmatrix} \boxed{2} & 4 & 2 \\ 1 & 1 & 2 \\ -1 & 0 & 2 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & \boxed{2} & 1 \\ 0 & \boxed{-1} & 1 \\ 0 & \boxed{2} & 3 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 2 & 1 \\ 0 & 1 & -1 \\ 0 & 0 & \boxed{5} \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 2 & 1 \\ 0 & 1 & -1 \\ 0 & 0 & 1 \end{bmatrix} = U$$

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \rightarrow \begin{bmatrix} \boxed{2} & 0 & 0 \\ \boxed{1} & 1 & 0 \\ -\boxed{1} & 0 & 1 \end{bmatrix} \rightarrow \begin{bmatrix} \boxed{2} & 0 & 0 \\ \boxed{1} & -\boxed{1} & 0 \\ -\boxed{1} & \boxed{2} & 1 \end{bmatrix} \rightarrow \begin{bmatrix} \boxed{2} & 0 & 0 \\ \boxed{1} & -\boxed{1} & 0 \\ -\boxed{1} & \boxed{2} & \boxed{5} \end{bmatrix} = L$$



Problem

Find an LU-factorization for $A = \begin{bmatrix} 5 & -5 & 10 & 0 & 5 \\ -3 & 3 & 2 & 2 & 1 \\ -2 & 2 & 0 & -1 & 0 \\ 1 & -1 & 10 & 2 & 5 \end{bmatrix}$.

Solution

$$\begin{bmatrix} 5 & -5 & 10 & 0 & 5 \\ -3 & 3 & 2 & 2 & 1 \\ -2 & 2 & 0 & -1 & 0 \\ 1 & -1 & 10 & 2 & 5 \end{bmatrix}$$

↓

$$\begin{bmatrix} 1 & -1 & 2 & 0 & 1 \\ 0 & 0 & 8 & 2 & 4 \\ 0 & 0 & 4 & -1 & 2 \\ 0 & 0 & 8 & 2 & 4 \end{bmatrix}$$

↓

$$\begin{bmatrix} 1 & -1 & 2 & 0 & 1 \\ 0 & 0 & 1 & 1/4 & 1/2 \\ 0 & 0 & 0 & -2 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

↓

$$\begin{bmatrix} 1 & -1 & 2 & 0 & 1 \\ 0 & 0 & 1 & 1/4 & 1/2 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

||

U

Solution

$$\begin{bmatrix} 5 & -5 & 10 & 0 & 5 \\ -3 & 3 & 2 & 2 & 1 \\ -2 & 2 & 0 & -1 & 0 \\ 1 & -1 & 10 & 2 & 5 \end{bmatrix}$$

↓

$$\begin{bmatrix} 1 & -1 & 2 & 0 & 1 \\ 0 & 0 & 8 & 2 & 4 \\ 0 & 0 & 4 & -1 & 2 \\ 0 & 0 & 8 & 2 & 4 \end{bmatrix}$$

↓

$$\begin{bmatrix} 1 & -1 & 2 & 0 & 1 \\ 0 & 0 & 1 & 1/4 & 1/2 \\ 0 & 0 & 0 & -2 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

↓

$$\begin{bmatrix} 1 & -1 & 2 & 0 & 1 \\ 0 & 0 & 1 & 1/4 & 1/2 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

||

U

⇓

L

||

$$\begin{bmatrix} 5 & 0 & 0 & 0 \\ -3 & 8 & 0 & 0 \\ -2 & 4 & -2 & 0 \\ 1 & 8 & 0 & 1 \end{bmatrix}$$

