

# Math 221: LINEAR ALGEBRA

## Chapter 1. Systems of Linear Equations

### §1-3. Homogeneous Equations

Le Chen<sup>1</sup>

Emory University, 2021 Spring

(last updated on 01/12/2023)



Creative Commons License  
(CC BY-NC-SA)

<sup>1</sup>Slides are adapted from those by Karen Seyffarth from University of Calgary.



# Linear Algebra with Applications

## Lecture Notes

### Current Lecture Notes Revision: Version 2018 — Revision B

These lecture notes were originally developed by Karen Seyffarth of the University of Calgary. Edits, additions, and revisions have been made to these notes by the editorial team at Lyryx Learning to accompany their text [Linear Algebra with Applications](#) based on W. K. Nicholson's original text.

In addition we recognize the following contributors. All new content contributed is released under the same license as noted below.

- Ilijas Farah, York University

### BE A CHAMPION OF OER!

Contribute suggestions for improvements, new content, or errata:

A new topic

A new example or problem

A new or better proof to an existing theorem

Any other suggestions to improve the material

Contact Lyryx at [info@lyryx.com](mailto:info@lyryx.com) with your ideas.

### License



Attribution-NonCommercial-ShareAlike (CC BY-NC-SA)

This license lets others remix, tweak, and build upon your work non-commercially, as long as they credit you and license their new creations under the identical terms.

Copyright

Homogeneous Equations

Linear Combination

Copyright

## Homogeneous Equations

Linear Combination

# Homogeneous Equations

# Homogeneous Equations

## Definition

A **homogeneous linear equation** is one whose constant term is equal to zero. A system of linear equations is called **homogeneous** if each equation in the system is homogeneous. A **homogeneous system** has the form

$$\left\{ \begin{array}{cccccc} a_{11}x_1 & + & a_{12}x_2 & + & \cdots & + & a_{1n}x_n & = & 0 \\ a_{21}x_1 & + & a_{22}x_2 & + & \cdots & + & a_{2n}x_n & = & 0 \\ & & & & \vdots & & & & \\ a_{m1}x_1 & + & a_{m2}x_2 & + & \cdots & + & a_{mn}x_n & = & 0 \end{array} \right.$$

where  $a_{ij}$  are scalars and  $x_i$  are variables,  $1 \leq i \leq m$ ,  $1 \leq j \leq n$ .

# Homogeneous Equations

## Definition

A **homogeneous linear equation** is one whose constant term is equal to zero. A system of linear equations is called **homogeneous** if each equation in the system is homogeneous. A **homogeneous system** has the form

$$\begin{cases} a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n = 0 \\ a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n = 0 \\ \vdots \\ a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n = 0 \end{cases}$$

where  $a_{ij}$  are scalars and  $x_i$  are variables,  $1 \leq i \leq m$ ,  $1 \leq j \leq n$ .

## Remark

1. Notice that  $x_1 = 0, x_2 = 0, \dots, x_n = 0$  is always a solution to a homogeneous system of equations. We call this the **trivial solution**.
2. We are interested in finding, if possible, **nontrivial solutions** (ones with at least one variable not equal to zero) to homogeneous systems.



### Example

$$\text{Solve the system } \begin{cases} x_1 + x_2 - x_3 + 3x_4 = 0 \\ -x_1 + 4x_2 + 5x_3 - 2x_4 = 0 \\ x_1 + 6x_2 + 3x_3 + 4x_4 = 0 \end{cases}$$

### Example

$$\text{Solve the system } \begin{cases} x_1 + x_2 - x_3 + 3x_4 = 0 \\ -x_1 + 4x_2 + 5x_3 - 2x_4 = 0 \\ x_1 + 6x_2 + 3x_3 + 4x_4 = 0 \end{cases}$$

### Solution

$$\left[ \begin{array}{cccc|c} 1 & 1 & -1 & 3 & 0 \\ -1 & 4 & 5 & -2 & 0 \\ 1 & 6 & 3 & 4 & 0 \end{array} \right]$$

### Example

$$\text{Solve the system } \begin{cases} x_1 + x_2 - x_3 + 3x_4 = 0 \\ -x_1 + 4x_2 + 5x_3 - 2x_4 = 0 \\ x_1 + 6x_2 + 3x_3 + 4x_4 = 0 \end{cases}$$

### Solution

$$\left[ \begin{array}{cccc|c} 1 & 1 & -1 & 3 & 0 \\ -1 & 4 & 5 & -2 & 0 \\ 1 & 6 & 3 & 4 & 0 \end{array} \right] \rightarrow \cdots \rightarrow \left[ \begin{array}{cccc|c} 1 & 0 & -9/5 & 14/5 & 0 \\ 0 & 1 & 4/5 & 1/5 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right]$$

### Example

$$\text{Solve the system } \begin{cases} x_1 + x_2 - x_3 + 3x_4 = 0 \\ -x_1 + 4x_2 + 5x_3 - 2x_4 = 0 \\ x_1 + 6x_2 + 3x_3 + 4x_4 = 0 \end{cases}$$

### Solution

$$\left[ \begin{array}{cccc|c} 1 & 1 & -1 & 3 & 0 \\ -1 & 4 & 5 & -2 & 0 \\ 1 & 6 & 3 & 4 & 0 \end{array} \right] \rightarrow \cdots \rightarrow \left[ \begin{array}{cccc|c} 1 & 0 & -9/5 & 14/5 & 0 \\ 0 & 1 & 4/5 & 1/5 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right]$$

The system has infinitely many solutions, and the general solution is

$$\begin{cases} x_1 = \frac{9}{5}s - \frac{14}{5}t \\ x_2 = -\frac{4}{5}s - \frac{1}{5}t \\ x_3 = s \\ x_4 = t \end{cases}$$

### Example

$$\text{Solve the system } \begin{cases} x_1 + x_2 - x_3 + 3x_4 = 0 \\ -x_1 + 4x_2 + 5x_3 - 2x_4 = 0 \\ x_1 + 6x_2 + 3x_3 + 4x_4 = 0 \end{cases}$$

### Solution

$$\left[ \begin{array}{cccc|c} 1 & 1 & -1 & 3 & 0 \\ -1 & 4 & 5 & -2 & 0 \\ 1 & 6 & 3 & 4 & 0 \end{array} \right] \rightarrow \cdots \rightarrow \left[ \begin{array}{cccc|c} 1 & 0 & -9/5 & 14/5 & 0 \\ 0 & 1 & 4/5 & 1/5 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right]$$

The system has infinitely many solutions, and the general solution is

$$\begin{cases} x_1 = \frac{9}{5}s - \frac{14}{5}t \\ x_2 = -\frac{4}{5}s - \frac{1}{5}t \\ x_3 = s \\ x_4 = t \end{cases} \quad \text{or} \quad \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} \frac{9}{5}s - \frac{14}{5}t \\ -\frac{4}{5}s - \frac{1}{5}t \\ s \\ t \end{bmatrix}, \forall s, t \in \mathbb{R}.$$

### Theorem

If a homogeneous system of linear equations has more variables than equations, then it has a nontrivial solution (in fact, infinitely many).

Copyright

Homogeneous Equations

Linear Combination





# Linear Combination

## Definition

If  $X_1, X_2, \dots, X_p$  are columns with the same number of entries, and if  $a_1, a_2, \dots, a_p \in \mathbb{R}$  (are scalars) then  $a_1X_1 + a_2X_2 + \dots + a_pX_p$  is a **linear combination** of columns  $X_1, X_2, \dots, X_p$ .

# Linear Combination

## Definition

If  $X_1, X_2, \dots, X_p$  are columns with the same number of entries, and if  $a_1, a_2, \dots, a_p \in \mathbb{R}$  (are scalars) then  $a_1X_1 + a_2X_2 + \dots + a_pX_p$  is a **linear combination** of columns  $X_1, X_2, \dots, X_p$ .

## Example (continued)

In the previous example,

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} \frac{9}{5}s - \frac{14}{5}t \\ -\frac{4}{5}s - \frac{1}{5}t \\ s \\ t \end{bmatrix}$$

# Linear Combination

## Definition

If  $X_1, X_2, \dots, X_p$  are columns with the same number of entries, and if  $a_1, a_2, \dots, a_p \in \mathbb{R}$  (are scalars) then  $a_1X_1 + a_2X_2 + \dots + a_pX_p$  is a **linear combination** of columns  $X_1, X_2, \dots, X_p$ .

## Example (continued)

In the previous example,

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} \frac{9}{5}s - \frac{14}{5}t \\ -\frac{4}{5}s - \frac{1}{5}t \\ s \\ t \end{bmatrix} = \begin{bmatrix} \frac{9}{5}s \\ -\frac{4}{5}s \\ s \\ 0 \end{bmatrix} + \begin{bmatrix} -\frac{14}{5}t \\ -\frac{1}{5}t \\ 0 \\ t \end{bmatrix}$$

# Linear Combination

## Definition

If  $X_1, X_2, \dots, X_p$  are columns with the same number of entries, and if  $a_1, a_2, \dots, a_p \in \mathbb{R}$  (are scalars) then  $a_1X_1 + a_2X_2 + \dots + a_pX_p$  is a **linear combination** of columns  $X_1, X_2, \dots, X_p$ .

## Example (continued)

In the previous example,

$$\begin{aligned} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} &= \begin{bmatrix} \frac{9}{5}s - \frac{14}{5}t \\ -\frac{4}{5}s - \frac{1}{5}t \\ s \\ t \end{bmatrix} = \begin{bmatrix} \frac{9}{5}s \\ -\frac{4}{5}s \\ s \\ 0 \end{bmatrix} + \begin{bmatrix} -\frac{14}{5}t \\ -\frac{1}{5}t \\ 0 \\ t \end{bmatrix} \\ &= s \begin{bmatrix} 9/5 \\ -4/5 \\ 1 \\ 0 \end{bmatrix} + t \begin{bmatrix} -14/5 \\ -1/5 \\ 0 \\ 1 \end{bmatrix} \end{aligned}$$

### Example (continued)

This gives us

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = s \begin{bmatrix} 9/5 \\ -4/5 \\ 1 \\ 0 \end{bmatrix} + t \begin{bmatrix} -14/5 \\ -1/5 \\ 0 \\ 1 \end{bmatrix} = sX_1 + tX_2,$$

$$\text{with } X_1 = \begin{bmatrix} 9/5 \\ -4/5 \\ 1 \\ 0 \end{bmatrix} \quad \text{and} \quad X_2 = \begin{bmatrix} -14/5 \\ -1/5 \\ 0 \\ 1 \end{bmatrix}.$$

### Example (continued)

This gives us

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = s \begin{bmatrix} 9/5 \\ -4/5 \\ 1 \\ 0 \end{bmatrix} + t \begin{bmatrix} -14/5 \\ -1/5 \\ 0 \\ 1 \end{bmatrix} = sX_1 + tX_2,$$

$$\text{with } X_1 = \begin{bmatrix} 9/5 \\ -4/5 \\ 1 \\ 0 \end{bmatrix} \quad \text{and} \quad X_2 = \begin{bmatrix} -14/5 \\ -1/5 \\ 0 \\ 1 \end{bmatrix}.$$

The columns  $X_1$  and  $X_2$  are called **basic solutions** to the original homogeneous system.

### Example (continued)

Notice that

$$\begin{aligned} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} &= s \begin{bmatrix} 9/5 \\ -4/5 \\ 1 \\ 0 \end{bmatrix} + t \begin{bmatrix} -14/5 \\ -1/5 \\ 0 \\ 1 \end{bmatrix} = \frac{s}{5} \begin{bmatrix} 9 \\ -4 \\ 5 \\ 0 \end{bmatrix} + \frac{t}{5} \begin{bmatrix} -14 \\ -1 \\ 0 \\ 5 \end{bmatrix} \\ &= r \begin{bmatrix} 9 \\ -4 \\ 5 \\ 0 \end{bmatrix} + q \begin{bmatrix} -14 \\ -1 \\ 0 \\ 5 \end{bmatrix} \\ &= r(5X_1) + q(5X_2) \end{aligned}$$

where  $r, q \in \mathbb{R}$ .

### Example (continued)

The columns  $5X_1 = \begin{bmatrix} 9 \\ -4 \\ 5 \\ 0 \end{bmatrix}$  and  $5X_2 = \begin{bmatrix} -14 \\ -1 \\ 0 \\ 5 \end{bmatrix}$  are also basic solutions to the original homogeneous system.



### Example (continued)

The columns  $5X_1 = \begin{bmatrix} 9 \\ -4 \\ 5 \\ 0 \end{bmatrix}$  and  $5X_2 = \begin{bmatrix} -14 \\ -1 \\ 0 \\ 5 \end{bmatrix}$  are also basic solutions to the original homogeneous system.

### Remark

In general, any nonzero multiple of a basic solution (to a homogeneous system of linear equations) is also a basic solution.

What does the rank tell us in the homogeneous case?

Suppose  $A$  is the augmented matrix of an homogeneous system of  $m$  linear equations in  $n$  variables, and  $\text{rank } A = r$ .

$$\begin{array}{c} m \\ \left\{ \begin{array}{c} \left[ \begin{array}{cccc|c} * & * & * & * & 0 \\ * & * & * & * & 0 \\ * & * & * & * & 0 \\ * & * & * & * & 0 \\ * & * & * & * & 0 \end{array} \right] \end{array} \right. \end{array} \rightarrow \begin{array}{c} \left[ \begin{array}{cccc|c} 1 & * & * & * & 0 \\ 0 & 0 & 1 & * & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right] \end{array}$$

$\underbrace{\hspace{10em}}_n$

$\underbrace{\hspace{10em}}_{r \text{ leading } 1's}$

What does the rank tell us in the homogeneous case?

Suppose  $A$  is the augmented matrix of an homogeneous system of  $m$  linear equations in  $n$  variables, and  $\text{rank } A = r$ .

$$\begin{array}{c} m \\ \left\{ \begin{array}{c} \left[ \begin{array}{cccc|c} * & * & * & * & 0 \\ * & * & * & * & 0 \\ * & * & * & * & 0 \\ * & * & * & * & 0 \\ * & * & * & * & 0 \end{array} \right] \end{array} \right\} \rightarrow \begin{array}{c} \left[ \begin{array}{cccc|c} 1 & * & * & * & 0 \\ 0 & 0 & 1 & * & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right] \end{array} \\ \underbrace{\hspace{10em}}_n \hspace{10em} \underbrace{\hspace{10em}}_{r \text{ leading } 1\text{'s}} \end{array}$$

There is always a solution, and the set of solutions to the system has  $n - r$  parameters, so

What does the rank tell us in the homogeneous case?

Suppose  $A$  is the augmented matrix of an homogeneous system of  $m$  linear equations in  $n$  variables, and  $\text{rank } A = r$ .

$$\begin{array}{c} m \\ \left\{ \begin{array}{c} \left[ \begin{array}{cccc|c} * & * & * & * & 0 \\ * & * & * & * & 0 \\ * & * & * & * & 0 \\ * & * & * & * & 0 \\ * & * & * & * & 0 \end{array} \right] \end{array} \right\} \rightarrow \begin{array}{c} \left[ \begin{array}{cccc|c} 1 & * & * & * & 0 \\ 0 & 0 & 1 & * & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right] \end{array} \\ \underbrace{\hspace{10em}}_n \hspace{10em} \underbrace{\hspace{10em}}_{r \text{ leading } 1\text{'s}} \end{array}$$

There is always a solution, and the set of solutions to the system has  $n - r$  parameters, so

- if  $r < n$ , there is at least one parameter, and the system has infinitely many solutions;

What does the rank tell us in the homogeneous case?

Suppose  $A$  is the augmented matrix of an homogeneous system of  $m$  linear equations in  $n$  variables, and  $\text{rank } A = r$ .

$$\begin{array}{c} m \\ \left\{ \begin{array}{c} \left[ \begin{array}{cccc|c} * & * & * & * & 0 \\ * & * & * & * & 0 \\ * & * & * & * & 0 \\ * & * & * & * & 0 \\ * & * & * & * & 0 \end{array} \right] \end{array} \right\} \rightarrow \begin{array}{c} \left[ \begin{array}{cccc|c} 1 & * & * & * & 0 \\ 0 & 0 & 1 & * & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right] \end{array} \\ \underbrace{\hspace{10em}}_n \hspace{10em} \underbrace{\hspace{10em}}_{r \text{ leading } 1\text{'s}} \end{array}$$

There is always a solution, and the set of solutions to the system has  $n - r$  parameters, so

- ▶ if  $r < n$ , there is at least one parameter, and the system has infinitely many solutions;
- ▶ if  $r = n$ , there are no parameters, and the system has a unique solution, the trivial solution.

## Theorem

Let  $A$  be an  $m \times n$  matrix of rank  $r$ , and consider the homogeneous system in  $n$  variables with  $A$  as coefficient matrix. Then:

1. The system has exactly  $n - r$  basic solutions, one for each parameter.
2. Every solution is a **linear combination** of these **basic solutions**.

## Problem

Find all values of  $a$  for which the system

$$\begin{cases} x + y = 0 \\ ay + z = 0 \\ x + y + az = 0 \end{cases}$$

has nontrivial solutions, and determine the solutions.

## Problem

Find all values of  $a$  for which the system

$$\begin{cases} x + y = 0 \\ ay + z = 0 \\ x + y + az = 0 \end{cases}$$

has nontrivial solutions, and determine the solutions.

## Solution

Non-trivial solutions occur only when  $a = 0$ , and the solutions when  $a = 0$  are given by (rank  $r = 2$ ,  $n - r = 3 - 2 = 1$  parameter)

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = s \begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}, \quad \forall s \in \mathbb{R}.$$