

# Math 221: LINEAR ALGEBRA

## Chapter 8. Orthogonality

### §8-3. Positive Definite Matrices

Le Chen<sup>1</sup>

Emory University, 2021 Spring

(last updated on 01/12/2023)



Creative Commons License  
(CC BY-NC-SA)

<sup>1</sup>Slides are adapted from those by Karen Seyffarth from University of Calgary.



# Linear Algebra with Applications

## Lecture Notes

### Current Lecture Notes Revision: Version 2018 — Revision B

These lecture notes were originally developed by Karen Seyffarth of the University of Calgary. Edits, additions, and revisions have been made to these notes by the editorial team at Lyryx Learning to accompany their text [Linear Algebra with Applications](#) based on W. K. Nicholson's original text.

In addition we recognize the following contributors. All new content contributed is released under the same license as noted below.

- Ilijas Farah, York University

### BE A CHAMPION OF OER!

Contribute suggestions for improvements, new content, or errata:

A new topic

A new example or problem

A new or better proof to an existing theorem

Any other suggestions to improve the material

Contact Lyryx at [info@lyryx.com](mailto:info@lyryx.com) with your ideas.

### License



Attribution-NonCommercial-ShareAlike (CC BY-NC-SA)

This license lets others remix, tweak, and build upon your work non-commercially, as long as they credit you and license their new creations under the identical terms.

Copyright

Positive Definite Matrices

Cholesky factorization – Square Root of a Matrix

Copyright

## Positive Definite Matrices

Cholesky factorization – Square Root of a Matrix



# Positive Definite Matrices

## Definition

An  $n \times n$  matrix  $A$  is **positive definite** if it is **symmetric** and has **positive** eigenvalues, i.e., if  $\lambda$  is a eigenvalue of  $A$ , then  $\lambda > 0$ .

# Positive Definite Matrices

## Definition

An  $n \times n$  matrix  $A$  is **positive definite** if it is **symmetric** and has **positive** eigenvalues, i.e., if  $\lambda$  is a eigenvalue of  $A$ , then  $\lambda > 0$ .

## Theorem

If  $A$  is a positive definite matrix, then  $\det(A) > 0$  and  $A$  is invertible.



# Positive Definite Matrices

## Definition

An  $n \times n$  matrix  $A$  is **positive definite** if it is **symmetric** and has **positive** eigenvalues, i.e., if  $\lambda$  is a eigenvalue of  $A$ , then  $\lambda > 0$ .

## Theorem

If  $A$  is a positive definite matrix, then  $\det(A) > 0$  and  $A$  is invertible.

## Proof.

Let  $\lambda_1, \lambda_2, \dots, \lambda_n$  denote the (not necessarily distinct) eigenvalues of  $A$ . Since  $A$  is symmetric,  $A$  is orthogonally diagonalizable. In particular,  $A \sim D$ , where  $D = \text{diag}(\lambda_1, \lambda_2, \dots, \lambda_n)$ . Similar matrices have the same determinant, so

$$\det(A) = \det(D) = \lambda_1 \lambda_2 \cdots \lambda_n.$$

Since  $A$  is positive definite,  $\lambda_i > 0$  for all  $i$ ,  $1 \leq i \leq n$ ; it follows that  $\det(A) > 0$ , and therefore  $A$  is invertible. ■

## Theorem

A symmetric matrix  $A$  is positive definite if and only if  $\vec{x}^T A \vec{x} > 0$  for all  $\vec{x} \in \mathbb{R}^n$ ,  $\vec{x} \neq \vec{0}$ .

## Theorem

A symmetric matrix  $A$  is positive definite if and only if  $\vec{x}^T A \vec{x} > 0$  for all  $\vec{x} \in \mathbb{R}^n$ ,  $\vec{x} \neq \vec{0}$ .

## Proof.

Since  $A$  is symmetric, there exists an orthogonal matrix  $P$  so that

$$P^T A P = \text{diag}(\lambda_1, \lambda_2, \dots, \lambda_n) = D,$$

where  $\lambda_1, \lambda_2, \dots, \lambda_n$  are the (not necessarily distinct) eigenvalues of  $A$ . Let  $\vec{x} \in \mathbb{R}^n$ ,  $\vec{x} \neq \vec{0}$ , and define  $\vec{y} = P^T \vec{x}$ . Then

$$\vec{x}^T A \vec{x} = \vec{x}^T (P D P^T) \vec{x} = (\vec{x}^T P) D (P^T \vec{x}) = (P^T \vec{x})^T D (P^T \vec{x}) = \vec{y}^T D \vec{y}.$$

Writing  $\vec{y}^T = \begin{bmatrix} y_1 & y_2 & \cdots & y_n \end{bmatrix}$ ,

$$\begin{aligned} \vec{x}^T A \vec{x} &= \begin{bmatrix} y_1 & y_2 & \cdots & y_n \end{bmatrix} \text{diag}(\lambda_1, \lambda_2, \dots, \lambda_n) \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{bmatrix} \\ &= \lambda_1 y_1^2 + \lambda_2 y_2^2 + \cdots + \lambda_n y_n^2. \end{aligned}$$

Proof. (continued)

( $\Rightarrow$ ) Suppose  $A$  is positive definite, and  $\vec{x} \in \mathbb{R}^n$ ,  $\vec{x} \neq \vec{0}$ . Since  $P^T$  is invertible,  $\vec{y} = P^T \vec{x} \neq \vec{0}$ , and thus  $y_j \neq 0$  for some  $j$ , implying  $y_j^2 > 0$  for some  $j$ . Furthermore, since all eigenvalues of  $A$  are positive,  $\lambda_i y_i^2 \geq 0$  for all  $i$ ; in particular  $\lambda_j y_j^2 > 0$ . Therefore,  $\vec{x}^T A \vec{x} > 0$ .

### Proof. (continued)

( $\Rightarrow$ ) Suppose  $A$  is positive definite, and  $\vec{x} \in \mathbb{R}^n$ ,  $\vec{x} \neq \vec{0}$ . Since  $P^T$  is invertible,  $\vec{y} = P^T \vec{x} \neq \vec{0}$ , and thus  $y_j \neq 0$  for some  $j$ , implying  $y_j^2 > 0$  for some  $j$ . Furthermore, since all eigenvalues of  $A$  are positive,  $\lambda_i y_i^2 \geq 0$  for all  $i$ ; in particular  $\lambda_j y_j^2 > 0$ . Therefore,  $\vec{x}^T A \vec{x} > 0$ .

( $\Leftarrow$ ) Conversely, if  $\vec{x}^T A \vec{x} > 0$  whenever  $\vec{x} \neq \vec{0}$ , choose  $\vec{x} = P \vec{e}_j$ , where  $\vec{e}_j$  is the  $j^{\text{th}}$  column of  $I_n$ . Since  $P$  is invertible,  $\vec{x} \neq \vec{0}$ , and thus

$$\vec{y} = P^T \vec{x} = P^T (P \vec{e}_j) = \vec{e}_j.$$

Thus  $y_j = 1$  and  $y_i = 0$  when  $i \neq j$ , so

$$\lambda_1 y_1^2 + \lambda_2 y_2^2 + \cdots + \lambda_n y_n^2 = \lambda_j,$$

i.e.,  $\lambda_j = \vec{x}^T A \vec{x} > 0$ . Therefore,  $A$  is positive definite. ■

## Theorem (Constructing Positive Definite Matrices)

Let  $U$  be an  $n \times n$  invertible matrix, and let  $A = U^T U$ . Then  $A$  is positive definite.


## Theorem (Constructing Positive Definite Matrices)

Let  $U$  be an  $n \times n$  invertible matrix, and let  $A = U^T U$ . Then  $A$  is positive definite.

Proof.

Let  $\vec{x} \in \mathbb{R}^n$ ,  $\vec{x} \neq \vec{0}$ . Then

$$\begin{aligned}\vec{x}^T A \vec{x} &= \vec{x}^T (U^T U) \vec{x} \\ &= (\vec{x}^T U^T)(U \vec{x}) \\ &= (U \vec{x})^T (U \vec{x}) \\ &= \|U \vec{x}\|^2.\end{aligned}$$

Since  $U$  is invertible and  $\vec{x} \neq \vec{0}$ ,  $U \vec{x} \neq \vec{0}$ , and hence  $\|U \vec{x}\|^2 > 0$ , i.e.,  $\vec{x}^T A \vec{x} = \|U \vec{x}\|^2 > 0$ . Therefore,  $A$  is positive definite. 

## Definition

Let  $A = [a_{ij}]$  be an  $n \times n$  matrix. For  $1 \leq r \leq n$ ,  ${}^{(r)}A$  denotes the  $r \times r$  submatrix in the upper left corner of  $A$ , i.e.,

$${}^{(r)}A = [a_{ij}], \quad 1 \leq i, j \leq r.$$

${}^{(1)}A, {}^{(2)}A, \dots, {}^{(n)}A$  are called the **principal submatrices** of  $A$ .



## Definition

Let  $A = [a_{ij}]$  be an  $n \times n$  matrix. For  $1 \leq r \leq n$ ,  $^{(r)}A$  denotes the  $r \times r$  submatrix in the upper left corner of  $A$ , i.e.,

$$^{(r)}A = [a_{ij}], \quad 1 \leq i, j \leq r.$$

$^{(1)}A, ^{(2)}A, \dots, ^{(n)}A$  are called the **principal submatrices** of  $A$ .

## Lemma

If  $A$  is an  $n \times n$  positive definite matrix, then each principal submatrix of  $A$  is positive definite.

**Proof.**

Suppose  $A$  is an  $n \times n$  positive definite matrix. For any integer  $r$ ,  $1 \leq r \leq n$ , write  $A$  in block form as

$$A = \begin{bmatrix} {}^{(r)}A & B \\ C & D \end{bmatrix},$$

where  $B$  is an  $r \times (n - r)$  matrix,  $C$  is an  $(n - r) \times r$  matrix, and  $D$  is an


$$(n - r) \times (n - r) \text{ matrix. Let } \vec{y} = \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_r \end{bmatrix} \neq \vec{0} \text{ and let } \vec{x} = \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_r \\ 0 \\ \vdots \\ 0 \end{bmatrix}. \text{ Then}$$

$\vec{x} \neq \vec{0}$ , and by the previous theorem,  $\vec{x}^T A \vec{x} > 0$ .

Proof. (continued)

But

$$\vec{x}^T A \vec{x} = \begin{bmatrix} y_1 & \cdots & y_r & 0 & \cdots & 0 \end{bmatrix} \begin{bmatrix} {}^{(r)}A & B \\ C & D \end{bmatrix} \begin{bmatrix} y_1 \\ \vdots \\ y_r \\ 0 \\ \vdots \\ 0 \end{bmatrix} = \vec{y}^T \left( {}^{(r)}A \right) \vec{y},$$

and therefore  $\vec{y}^T \left( {}^{(r)}A \right) \vec{y} > 0$ . Then  ${}^{(r)}A$  is positive definite again by the previous theorem. 

Copyright

Positive Definite Matrices

Cholesky factorization – Square Root of a Matrix

## Cholesky factorization – Square Root of a Matrix

## Cholesky factorization – Square Root of a Matrix

$$4 = 2 \times 2^T$$

$$\begin{bmatrix} 4 & 12 & -16 \\ 12 & 37 & -43 \\ -16 & -43 & 98 \end{bmatrix} = \begin{bmatrix} 2 & 0 & 0 \\ 6 & 1 & 0 \\ -8 & 5 & 3 \end{bmatrix} \begin{bmatrix} 2 & 6 & -8 \\ 0 & 1 & 5 \\ 0 & 0 & 3 \end{bmatrix}$$

## Cholesky factorization – Square Root of a Matrix

$$4 = 2 \times 2^T$$

$$\begin{bmatrix} 4 & 12 & -16 \\ 12 & 37 & -43 \\ -16 & -43 & 98 \end{bmatrix} = \begin{bmatrix} 2 & 0 & 0 \\ 6 & 1 & 0 \\ -8 & 5 & 3 \end{bmatrix} \begin{bmatrix} 2 & 6 & -8 \\ 0 & 1 & 5 \\ 0 & 0 & 3 \end{bmatrix}$$

### Theorem

Let  $A$  be an  $n \times n$  symmetric matrix. Then the following conditions are equivalent.

1.  $A$  is positive definite.
2.  $\det({}^{(r)}A) > 0$  for  $r = 1, 2, \dots, n$ .
3.  $A = U^T U$  where  $U$  is upper triangular and has positive entries on its main diagonal. Furthermore,  $U$  is unique. The expression  $A = U^T U$  is called the **Cholesky factorization** of  $A$ .

## Algorithm for Cholesky Factorization

Let  $A$  be a positive definite matrix. The Cholesky factorization  $A = U^T U$  can be obtained as follows.



## Algorithm for Cholesky Factorization

Let  $A$  be a positive definite matrix. The Cholesky factorization  $A = U^T U$  can be obtained as follows.

1. Using only type 3 elementary row operations, with multiples of rows added to lower rows, put  $A$  in upper triangular form. Call this matrix  $\hat{U}$ ; then  $\hat{U}$  has positive entries on its main diagonal (this can be proved by induction on  $n$ ).

## Algorithm for Cholesky Factorization

Let  $A$  be a positive definite matrix. The Cholesky factorization  $A = U^T U$  can be obtained as follows.

1. Using only type 3 elementary row operations, with multiples of rows added to lower rows, put  $A$  in upper triangular form. Call this matrix  $\hat{U}$ ; then  $\hat{U}$  has positive entries on its main diagonal (this can be proved by induction on  $n$ ).
2. Obtain  $U$  from  $\hat{U}$  by dividing each row of  $\hat{U}$  by the square root of the diagonal entry in that row.

## Problem

Show that  $A = \begin{bmatrix} 9 & -6 & 3 \\ -6 & 5 & -3 \\ 3 & -3 & 6 \end{bmatrix}$  is positive definite, and find the Cholesky factorization of  $A$ .

## Problem

Show that  $A = \begin{bmatrix} 9 & -6 & 3 \\ -6 & 5 & -3 \\ 3 & -3 & 6 \end{bmatrix}$  is positive definite, and find the Cholesky factorization of  $A$ .

## Solution

$$^{(1)}A = \begin{bmatrix} 9 \end{bmatrix} \quad \text{and} \quad ^{(2)}A = \begin{bmatrix} 9 & -6 \\ -6 & 5 \end{bmatrix},$$

so  $\det(^{(1)}A) = 9$  and  $\det(^{(2)}A) = 9$ . Since  $\det(A) = 36$ , it follows that  $A$  is positive definite.

## Problem

Show that  $A = \begin{bmatrix} 9 & -6 & 3 \\ -6 & 5 & -3 \\ 3 & -3 & 6 \end{bmatrix}$  is positive definite, and find the Cholesky factorization of  $A$ .

## Solution

$$^{(1)}A = \begin{bmatrix} 9 \end{bmatrix} \quad \text{and} \quad ^{(2)}A = \begin{bmatrix} 9 & -6 \\ -6 & 5 \end{bmatrix},$$

so  $\det(^{(1)}A) = 9$  and  $\det(^{(2)}A) = 9$ . Since  $\det(A) = 36$ , it follows that  $A$  is positive definite.

$$\begin{bmatrix} 9 & -6 & 3 \\ -6 & 5 & -3 \\ 3 & -3 & 6 \end{bmatrix} \rightarrow \begin{bmatrix} 9 & -6 & 3 \\ 0 & 1 & -1 \\ 0 & -1 & 5 \end{bmatrix} \rightarrow \begin{bmatrix} 9 & -6 & 3 \\ 0 & 1 & -1 \\ 0 & 0 & 4 \end{bmatrix}$$

## Problem

Show that  $A = \begin{bmatrix} 9 & -6 & 3 \\ -6 & 5 & -3 \\ 3 & -3 & 6 \end{bmatrix}$  is positive definite, and find the Cholesky factorization of  $A$ .

## Solution

$$^{(1)}A = \begin{bmatrix} 9 \end{bmatrix} \quad \text{and} \quad ^{(2)}A = \begin{bmatrix} 9 & -6 \\ -6 & 5 \end{bmatrix},$$

so  $\det(^{(1)}A) = 9$  and  $\det(^{(2)}A) = 9$ . Since  $\det(A) = 36$ , it follows that  $A$  is positive definite.

$$\begin{bmatrix} 9 & -6 & 3 \\ -6 & 5 & -3 \\ 3 & -3 & 6 \end{bmatrix} \rightarrow \begin{bmatrix} 9 & -6 & 3 \\ 0 & 1 & -1 \\ 0 & -1 & 5 \end{bmatrix} \rightarrow \begin{bmatrix} 9 & -6 & 3 \\ 0 & 1 & -1 \\ 0 & 0 & 4 \end{bmatrix}$$

Now divide the entries in each row by the square root of the diagonal entry in that row, to give

$$U = \begin{bmatrix} 3 & -2 & 1 \\ 0 & 1 & -1 \\ 0 & 0 & 2 \end{bmatrix} \quad \text{and} \quad U^T U = A.$$



## Problem

Verify that

$$A = \begin{bmatrix} 12 & 4 & 3 \\ 4 & 2 & -1 \\ 3 & -1 & 7 \end{bmatrix}$$

is positive definite, and find the Cholesky factorization of  $A$ .

### Problem

Verify that

$$A = \begin{bmatrix} 12 & 4 & 3 \\ 4 & 2 & -1 \\ 3 & -1 & 7 \end{bmatrix}$$

is positive definite, and find the Cholesky factorization of  $A$ .

### Solution ( Final Answer )

$\det \left( {}^{(1)}A \right) = 12$ ,  $\det \left( {}^{(2)}A \right) = 8$ ,  $\det (A) = 2$ ; by the previous theorem,  $A$  is positive definite.

$$U = \begin{bmatrix} 2\sqrt{3} & 2\sqrt{3}/3 & \sqrt{3}/2 \\ 0 & \sqrt{6}/3 & -\sqrt{6} \\ 0 & 0 & 1/2 \end{bmatrix}$$

and  $U^T U = A$ .