

Math 221: LINEAR ALGEBRA

Chapter 2. Matrix Algebra

§2-4. Matrix Inverses

Le Chen¹

Emory University, 2021 Spring

(last updated on 01/12/2023)



Creative Commons License
(CC BY-NC-SA)

¹Slides are adapted from those by Karen Seyffarth from University of Calgary.

Linear Algebra with Applications

Lecture Notes

Current Lecture Notes Revision: Version 2018 — Revision B

These lecture notes were originally developed by Karen Seyffarth of the University of Calgary. Edits, additions, and revisions have been made to these notes by the editorial team at Lyryx Learning to accompany their text [Linear Algebra with Applications](#) based on W. K. Nicholson's original text.

In addition we recognize the following contributors. All new content contributed is released under the same license as noted below.

- Ilijas Farah, York University

BE A CHAMPION OF OER!

Contribute suggestions for improvements, new content, or errata:

A new topic

A new example or problem

A new or better proof to an existing theorem

Any other suggestions to improve the material

Contact Lyryx at info@lyryx.com with your ideas.

License



Attribution-NonCommercial-ShareAlike (CC BY-NC-SA)

This license lets others remix, tweak, and build upon your work non-commercially, as long as they credit you and license their new creations under the identical terms.

Copyright

The Identity and Inverse Matrices

Finding the Inverse of a Matrix

Properties of the Inverse

Inverse of Transformations

Copyright

The Identity and Inverse Matrices

Finding the Inverse of a Matrix

Properties of the Inverse

Inverse of Transformations

The Identity and Inverse Matrices

Definition

For each $n \geq 2$, the $n \times n$ identity matrix, denoted \mathbf{I}_n , is the matrix having ones on its main diagonal and zeros elsewhere, and is defined for all $n \geq 2$.

Example

$$\mathbf{I}_2 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \quad \mathbf{I}_3 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Definition

Let $n \geq 2$. For each j , $1 \leq j \leq n$, we denote by \vec{e}_j the j^{th} column of I_n .

Example

When $n = 3$, $\vec{e}_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$, $\vec{e}_2 = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$, $\vec{e}_3 = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$.

Theorem

Let A be an $m \times n$ matrix. Then $AI_n = A$ and $I_m A = A$.

Proof.

The (i, j) -entry of AI_n is the product of the i^{th} row of $A = [a_{ij}]$, namely $\begin{bmatrix} a_{i1} & a_{i2} & \cdots & a_{ij} & \cdots & a_{in} \end{bmatrix}$ with the j^{th} column of I_n , namely \vec{e}_j . Since \vec{e}_j has a one in row j and zeros elsewhere,

$$\begin{bmatrix} a_{i1} & a_{i2} & \cdots & a_{ij} & \cdots & a_{in} \end{bmatrix} \vec{e}_j = a_{ij}$$

Since this is true for all $i \leq m$ and all $j \leq n$, $AI_n = A$.

The proof of $I_m A = A$ is analogous—work it out!



Instead of AI_n and $I_m A$ we often write AI and IA , respectively, since the size of the identity matrix is clear from the context: the sizes of A and I must be compatible for matrix multiplication.

Thus

$$AI = A \quad \text{and} \quad IA = A$$

which is why I is called an **identity** matrix – it is an identity for matrix multiplication.

Definition (Matrix Inverses)

Let A be an $n \times n$ matrix. Then B is **an inverse** of A if and only if $AB = I_n$ and $BA = I_n$.

Remark

Note that since A and I_n are both $n \times n$, B must also be an $n \times n$ matrix.

Example

Let $A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$ and $B = \begin{bmatrix} -2 & 1 \\ 3/2 & -1/2 \end{bmatrix}$. Then

$$AB = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \quad \text{and} \quad BA = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}.$$

Therefore, B is an inverse of A .

Problem

Does every square matrix have an inverse?

Solution

No! Take e.g. the zero matrix \mathbf{O}_n (all entries of \mathbf{O}_n are equal to 0)

$$\mathbf{A}\mathbf{O}_n = \mathbf{O}_n\mathbf{A} = \mathbf{O}_n$$

for all $n \times n$ matrices \mathbf{A} : The (i,j) -entry of $\mathbf{O}_n\mathbf{A}$ is equal to $\sum_{k=1}^n 0a_{kj} = 0$. ■

Problem

Does every nonzero square matrix have an inverse?

Problem

Does the following matrix A have an inverse?

$$A = \begin{bmatrix} 0 & 1 \\ 0 & 1 \end{bmatrix}$$

Solution

No! To see this, suppose

$$B = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

is an inverse of A. Then

$$AB = \begin{bmatrix} 0 & 1 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} c & d \\ c & d \end{bmatrix}$$

which is never equal to I_2 . (Why?)



Theorem (Uniqueness of an Inverse)

If A is a square matrix and B and C are inverses of A , then $B = C$.


Proof.

Since B and C are inverses of A , $AB = I = BA$ and $AC = I = CA$. Then

$$C = CI = C(AB) = CAB$$

and

$$B = IB = (CA)B = CAB$$

so $B = C$. 

Example (revisited)

For $A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$ and $B = \begin{bmatrix} -2 & 1 \\ 3/2 & -1/2 \end{bmatrix}$, we saw that

$$AB = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \quad \text{and} \quad BA = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

The preceding theorem tells us that B is **the inverse** of A, rather than just an inverse of A.

Remark (notation)

Let A be a square matrix, i.e., an $n \times n$ matrix.

- The inverse of A , if it exists, is denoted A^{-1} , and

$$AA^{-1} = I = A^{-1}A$$

- If A has an inverse, then we say that A is invertible.

Copyright

The Identity and Inverse Matrices

Finding the Inverse of a Matrix

Properties of the Inverse

Inverse of Transformations

Finding the inverse of a 2×2 matrix

Example

Suppose that $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. If $ad - bc \neq 0$, then there is a formula for A^{-1} :

$$A^{-1} = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}.$$

This can easily be verified by computing the products AA^{-1} and $A^{-1}A$.

$$\begin{aligned} AA^{-1} &= \begin{bmatrix} a & b \\ c & d \end{bmatrix} \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix} \\ &= \frac{1}{ad - bc} \left(\begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix} \right) \\ &= \frac{1}{ad - bc} \begin{bmatrix} ad - bc & 0 \\ 0 & -bc + ad \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \end{aligned}$$

Showing that $A^{-1}A = I_2$ is left as an exercise.

Remark

Here are some terminology related to this example:

1. **Determinant:**

$$\det \begin{pmatrix} a & b \\ c & d \end{pmatrix} := ad - cd$$

2. **Adjugate:**

$$\text{adj} \begin{pmatrix} a & b \\ c & d \end{pmatrix} := \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}$$



Problem

Suppose that A is any $n \times n$ matrix.

- ▶ How do we know whether or not A^{-1} exists?
- ▶ If A^{-1} exists, how do we find it?

Solution

The matrix inversion algorithm!

Although the formula for the inverse of a 2×2 matrix is quicker and easier to use than the matrix inversion algorithm, the general formula for the inverse of an $n \times n$ matrix, $n \geq 3$ (which we will see later), is more complicated and difficult to use than the matrix inversion algorithm. To find inverses of square matrices that are not 2×2 , the matrix inversion algorithm is the most efficient method to use.

The Matrix Inversion Algorithm

Let A be an $n \times n$ matrix. To find A^{-1} , if it exists,

Step 1 take the $n \times 2n$ matrix

$$\left[A \mid I_n \right]$$

obtained by augmenting A with the $n \times n$ identity matrix, I_n .

Step 2 Perform elementary row operations to transform $\left[A \mid I_n \right]$ into a reduced row-echelon matrix.

Theorem (Matrix Inverses)

Let A be an $n \times n$ matrix. Then the following conditions are equivalent.

1. A is invertible.
2. the reduced row-echelon form on A is I .
3. $\left[A \mid I_n \right]$ can be transformed into $\left[I_n \mid A^{-1} \right]$ using the Matrix Inversion Algorithm.

Problem

Find, if possible, the inverse of $\begin{bmatrix} 1 & 0 & -1 \\ -2 & 1 & 3 \\ -1 & 1 & 2 \end{bmatrix}$.

Solution

Using the matrix inversion algorithm

$$\left[\begin{array}{ccc|ccc} 1 & 0 & -1 & 1 & 0 & 0 \\ -2 & 1 & 3 & 0 & 1 & 0 \\ -1 & 1 & 2 & 0 & 0 & 1 \end{array} \right] \rightarrow \left[\begin{array}{ccc|ccc} 1 & 0 & -1 & 1 & 0 & 0 \\ 0 & 1 & 1 & 2 & 1 & 0 \\ 0 & 1 & 1 & 1 & 0 & 1 \end{array} \right] \rightarrow$$
$$\left[\begin{array}{ccc|ccc} 1 & 0 & -1 & 1 & 0 & 0 \\ 0 & 1 & 1 & 2 & 1 & 0 \\ 0 & 0 & 0 & -1 & -1 & 1 \end{array} \right]$$

From this, we see that **A has no inverse.**



Problem

Let $A = \begin{bmatrix} 3 & 1 & 2 \\ 1 & -1 & 3 \\ 1 & 2 & 4 \end{bmatrix}$. Find the inverse of A , if it exists.

Solution

Using the matrix inversion algorithm

$$\begin{aligned} [A | I] &= \left[\begin{array}{ccc|ccc} 3 & 1 & 2 & 1 & 0 & 0 \\ 1 & -1 & 3 & 0 & 1 & 0 \\ 1 & 2 & 4 & 0 & 0 & 1 \end{array} \right] \rightarrow \left[\begin{array}{ccc|ccc} 1 & -1 & 3 & 0 & 1 & 0 \\ 3 & 1 & 2 & 1 & 0 & 0 \\ 1 & 2 & 4 & 0 & 0 & 1 \end{array} \right] \\ &\rightarrow \left[\begin{array}{ccc|ccc} 1 & -1 & 3 & 0 & 1 & 0 \\ 0 & 4 & -7 & 1 & -3 & 0 \\ 0 & 3 & 1 & 0 & -1 & 1 \end{array} \right] \rightarrow \left[\begin{array}{ccc|ccc} 1 & -1 & 3 & 0 & 1 & 0 \\ 0 & 1 & -8 & 1 & -2 & -1 \\ 0 & 3 & 1 & 0 & -1 & 1 \end{array} \right] \\ &\rightarrow \left[\begin{array}{ccc|ccc} 1 & 0 & -5 & 1 & -1 & -1 \\ 0 & 1 & -8 & 1 & -2 & -1 \\ 0 & 0 & 25 & -3 & 5 & 4 \end{array} \right] \rightarrow \left[\begin{array}{ccc|ccc} 1 & 0 & -5 & 1 & -1 & -1 \\ 0 & 1 & -8 & 1 & -2 & -1 \\ 0 & 0 & 1 & -\frac{3}{25} & \frac{5}{25} & \frac{4}{25} \end{array} \right] \\ &\rightarrow \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & \frac{10}{25} & 0 & -\frac{5}{25} \\ 0 & 1 & 0 & \frac{1}{25} & -\frac{10}{25} & \frac{7}{25} \\ 0 & 0 & 1 & -\frac{3}{25} & \frac{5}{25} & \frac{4}{25} \end{array} \right] = [I | A^{-1}] \end{aligned}$$

Solution (continued)

Therefore, A^{-1} exists, and

$$A^{-1} = \begin{bmatrix} \frac{10}{25} & 0 & -\frac{5}{25} \\ \frac{1}{25} & -\frac{10}{25} & \frac{7}{25} \\ -\frac{3}{25} & \frac{5}{25} & \frac{4}{25} \end{bmatrix} = \frac{1}{25} \begin{bmatrix} 10 & 0 & -5 \\ 1 & -10 & 7 \\ -3 & 5 & 4 \end{bmatrix}.$$



Remark

It is always a good habit to check your answer by computing AA^{-1} and $A^{-1}A$.

One can use matrix inverse to solve $A\vec{x} = \vec{b}$ when there are n linear equations in n variables, i.e., A is a square matrix.

Example

The system of linear equations

$$\begin{aligned}2x - 7y &= 3 \\ 5x - 18y &= 8\end{aligned}$$

can be written in matrix form as $A\vec{x} = \vec{b}$:

$$\begin{bmatrix} 2 & -7 \\ 5 & -18 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 3 \\ 8 \end{bmatrix}$$

You can check that $A^{-1} = \begin{bmatrix} 18 & -7 \\ 5 & -2 \end{bmatrix}$.


Example (continued)

Since A^{-1} exists and has the property that $A^{-1}A = I$, we obtain the following.

$$\begin{aligned} A\vec{x} &= \vec{b} \\ A^{-1}(A\vec{x}) &= A^{-1}\vec{b} \\ (A^{-1}A)\vec{x} &= A^{-1}\vec{b} \\ I\vec{x} &= A^{-1}\vec{b} \\ \vec{x} &= A^{-1}\vec{b} \end{aligned}$$

i.e., $A\vec{x} = \vec{b}$ has the **unique solution** given by $\vec{x} = A^{-1}\vec{b}$. Therefore,

$$\vec{x} = A^{-1} \begin{bmatrix} 3 \\ 8 \end{bmatrix} = \begin{bmatrix} 18 & -7 \\ 5 & -2 \end{bmatrix} \begin{bmatrix} 3 \\ 8 \end{bmatrix} = \begin{bmatrix} -2 \\ -1 \end{bmatrix}$$

is the unique solution to the system. 

Remark

The last example illustrates another method for solving a system of linear equations when the coefficient matrix is square and invertible. Unless that coefficient matrix is 2×2 , this is generally NOT an efficient method for solving a system of linear equations.

Example

Let A, B and C be matrices, and suppose that A is invertible.

1. If $AB = AC$, then

$$A^{-1}(AB) = A^{-1}(AC)$$

$$(A^{-1}A)B = (A^{-1}A)C$$

$$IB = IC$$

$$B = C$$

2. If $BA = CA$, then

$$(BA)A^{-1} = (CA)A^{-1}$$

$$B(AA^{-1}) = C(AA^{-1})$$

$$BI = CI$$

$$B = C$$

Problem

Can you find square matrices A, B and C for which $AB = AC$ but $B \neq C$?

Copyright

The Identity and Inverse Matrices

Finding the Inverse of a Matrix

Properties of the Inverse

Inverse of Transformations

Properties of the Inverse

Example

Suppose A is an invertible matrix. What is the $(A^T)^{-1}$?

We need to find:

$$A^T \boxed{???} = \boxed{???} A^T = I.$$

Notice that

$$A^T (A^{-1})^T = (A^{-1} A)^T = I^T = I$$

and

$$(A^{-1})^T A^T = (A A^{-1})^T = I^T = I$$

Hence, $\boxed{???} = (A^{-1})^T$, i.e., $(A^T)^{-1} = (A^{-1})^T$.



Properties of the Inverse

Example

Suppose A and B are invertible $n \times n$ matrices. What is $(AB)^{-1}$?

We need to find:

$$(AB) \boxed{???} = \boxed{???} (AB) = I.$$

Notice that

$$(AB)(B^{-1}A^{-1}) = A(BB^{-1})A^{-1} = AIA^{-1} = AA^{-1} = I$$

and

$$(B^{-1}A^{-1})(AB) = B^{-1}(A^{-1}A)B = B^{-1}IB = B^{-1}B = I$$

Hence, $\boxed{B^{-1}A^{-1}} = B^{-1}A^{-1}$, i.e., $(AB)^{-1} = B^{-1}A^{-1}$.



The previous two examples prove the first two parts of the following theorem.

Theorem (Properties of Inverses)

1. If A is an invertible matrix, then $(A^T)^{-1} = (A^{-1})^T$.
2. If A and B are invertible matrices, then AB is invertible and

$$(AB)^{-1} = B^{-1}A^{-1}.$$

3. If A_1, A_2, \dots, A_k are invertible, then $A_1A_2 \cdots A_k$ is invertible and

$$(A_1A_2 \cdots A_k)^{-1} = A_k^{-1}A_{k-1}^{-1} \cdots A_2^{-1}A_1^{-1}.$$

Theorem (More Properties of Inverses)

1. I is invertible, and $I^{-1} = I$.
2. If A is invertible, so is A^{-1} , and $(A^{-1})^{-1} = A$.
3. If A is invertible, so is A^k , and $(A^k)^{-1} = (A^{-1})^k$.
(A^k means A multiplied by itself k times)
4. If A is invertible and $p \in \mathbb{R}$ is nonzero, then pA is invertible, and $(pA)^{-1} = \frac{1}{p}A^{-1}$.

Example

Given $(3\mathbf{I} - \mathbf{A}^T)^{-1} = 2 \begin{bmatrix} 1 & 1 \\ 2 & 3 \end{bmatrix}$, we wish to find the matrix \mathbf{A} . Taking inverses of both sides of the equation:

$$\begin{aligned} 3\mathbf{I} - \mathbf{A}^T &= \left(2 \begin{bmatrix} 1 & 1 \\ 2 & 3 \end{bmatrix} \right)^{-1} \\ &= \frac{1}{2} \begin{bmatrix} 1 & 1 \\ 2 & 3 \end{bmatrix}^{-1} \\ &= \frac{1}{2} \begin{bmatrix} 3 & -1 \\ -2 & 1 \end{bmatrix} \\ &= \begin{bmatrix} \frac{3}{2} & -\frac{1}{2} \\ -1 & \frac{1}{2} \end{bmatrix} \end{aligned}$$

Example (continued)

$$3\mathbf{I} - \mathbf{A}^T = \begin{bmatrix} \frac{3}{2} & -\frac{1}{2} \\ -1 & \frac{1}{2} \end{bmatrix}$$

$$-\mathbf{A}^T = \begin{bmatrix} \frac{3}{2} & -\frac{1}{2} \\ -1 & \frac{1}{2} \end{bmatrix} - 3\mathbf{I}$$

$$-\mathbf{A}^T = \begin{bmatrix} \frac{3}{2} & -\frac{1}{2} \\ -1 & \frac{1}{2} \end{bmatrix} - \begin{bmatrix} 3 & 0 \\ 0 & 3 \end{bmatrix}$$

$$-\mathbf{A}^T = \begin{bmatrix} -\frac{3}{2} & -\frac{1}{2} \\ -1 & -\frac{5}{2} \end{bmatrix}$$

$$\mathbf{A} = \begin{bmatrix} \frac{3}{2} & 1 \\ \frac{1}{2} & \frac{5}{2} \end{bmatrix}$$



Problem

True or false? Justify your answer.

If $A^3 = 4I$, then A is invertible.

Solution

To show A is invertible, We need to find:

$$A \boxed{???} = \boxed{???} A = I.$$

Because $A^3 = 4I$, we see that

$$\frac{1}{4}A^3 = I$$

so

$$\left(\frac{1}{4}A^2\right)A = I \quad \text{and} \quad A\left(\frac{1}{4}A^2\right) = I.$$

Therefore, A is invertible, and $\boxed{???} = \frac{1}{4}A^2$, i.e., $A^{-1} = \frac{1}{4}A^2$. ■

Theorem (Inverse Theorem)

Let A be an $n \times n$ matrix, and let \vec{x} , \vec{b} be $n \times 1$ vectors. The following conditions are equivalent.

1. A is invertible.
2. The rank of A is n .
3. The reduced row echelon form of A is I_n .
4. $A\vec{x} = \vec{0}$ has only the trivial solution, $\vec{x} = \vec{0}$.
5. A can be transformed to I_n by elementary row operations.
6. The system $A\vec{x} = \vec{b}$ has a unique solution \vec{x} for any choice of \vec{b} .
7. The system $A\vec{x} = \vec{b}$ has at least one solution \vec{x} for any choice of \vec{b} .
8. There exists an $n \times n$ matrix C with the property that $CA = I_n$.
9. There exists an $n \times n$ matrix C with the property that $AC = I_n$.

$$\begin{array}{ccccc}
(1) & \Leftrightarrow & (2) & \Leftrightarrow & (3) & \Leftrightarrow & (4) & \Leftrightarrow & (5) & \Leftrightarrow & (6) \\
\Downarrow & & & & & & \Uparrow & & & & \Downarrow \\
(9) & & & & & & (8) & \Leftarrow & & & (7)
\end{array}$$

Proof.

(1), (2), (4), (5) and (6) are all equivalent.

(6) \Rightarrow (7) is clear. As for (7) \Rightarrow (8), let \vec{c}_j be one of the solution of $A\vec{x} = \vec{e}_j$.
The

$$A[\vec{c}_1, \dots, \vec{c}_n] = [\vec{e}_1, \dots, \vec{e}_n] = I$$

Hence, (8) holds with $C = [\vec{c}_1, \dots, \vec{c}_n]$.

(1) \Rightarrow (8) and (9): Using $C = A^{-1}$.

(8) \Rightarrow (4): Whenever \vec{x} is a sol. i.e., $A\vec{x} = \vec{0}$, then $\vec{x} = I\vec{x} = CA\vec{x} = C\vec{0} = \vec{0}$.
Hence, $\vec{0}$ is the only solution. (4) holds true.

(9) \Rightarrow (1): By reversing the roles of A and C and apply (8) to see that C is invertible. Thus A is the inverse of C, and hence A is itself invertible. ■

Corollary

If A and B are $n \times n$ matrices such that $AB = I$, then $BA = I$. Furthermore, A and B are invertible, with $B = A^{-1}$ and $A = B^{-1}$.

$$\begin{array}{c} \begin{array}{cc} \text{red square} & \text{blue square} \end{array} = I_n \\ \Downarrow \\ \begin{array}{ccccc} \text{red square}^{-1} & = & \text{blue square} & \text{and} & \text{blue square}^{-1} = \text{red square} \end{array} \end{array}$$

Remark

Important Fact In Corollary, it is essential that the matrices be square.

Theorem

If A and B are matrices such that $AB = I$ and $BA = I$, then A and B are square matrices (of the same size).

Example

Let $A = \begin{bmatrix} 1 & 1 & 0 \\ -1 & 4 & 1 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & 0 \\ 0 & 0 \\ 1 & 1 \end{bmatrix}$. Then

$$AB = \begin{bmatrix} 1 & 1 & 0 \\ -1 & 4 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 0 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = I_2$$

and

$$BA = \begin{bmatrix} 1 & 0 \\ 0 & 0 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 & 0 \\ -1 & 4 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 5 & 1 \end{bmatrix} \neq I_3.$$

Remark

This example illustrates why “an inverse” of a non-square matrix doesn’t make sense. If A is $m \times n$ and B is $n \times m$, where $m \neq n$, then even if $AB = I$, it will never be the case that $BA = I$.

Copyright

The Identity and Inverse Matrices

Finding the Inverse of a Matrix

Properties of the Inverse

Inverse of Transformations

Inverse of Transformations

Definition

Suppose $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$ and $S : \mathbb{R}^n \rightarrow \mathbb{R}^n$ are transformations such that for each $\vec{x} \in \mathbb{R}^n$,

$$(S \circ T)(\vec{x}) = \vec{x} \quad \text{and} \quad (T \circ S)(\vec{x}) = \vec{x}.$$

Then T and S are **invertible** transformations; S is called **an inverse of T** , and T is called **an inverse of S** . (Geometrically, S reverses the action of T , and T reverses the action of S .)

Theorem

Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be a matrix transformation induced by matrix A . Then we have:

1. A is invertible if and only if T has an inverse.
2. In the case where T has an inverse, the inverse is unique and is denoted T^{-1} .
3. Furthermore, $T^{-1} : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is induced by the matrix A^{-1} .

Fundamental Identities relating T and T^{-1}

1. $T^{-1} \circ T = 1_{\mathbb{R}^n}$

2. $T \circ T^{-1} = 1_{\mathbb{R}^n}$

Example

Let $T : \mathbb{R}^2 \mapsto \mathbb{R}^2$ be a transformation given by

$$T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x + y \\ y \end{bmatrix}$$

Then T is a linear transformation induced by $A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$.

Notice that the matrix A is invertible. Therefore the transformation T has an inverse, T^{-1} , induced by

$$A^{-1} = \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix}$$

Example (continued)

Consider the action of T and T^{-1} :

$$T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x+y \\ y \end{bmatrix};$$

$$T^{-1} \begin{bmatrix} x+y \\ y \end{bmatrix} = \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x+y \\ y \end{bmatrix} = \begin{bmatrix} x \\ y \end{bmatrix}.$$

Therefore,

$$T^{-1} \left(T \begin{bmatrix} x \\ y \end{bmatrix} \right) = \begin{bmatrix} x \\ y \end{bmatrix}.$$

