

# Math 221: LINEAR ALGEBRA

## Chapter 2. Matrix Algebra §2-5. Elementary Matrices

Le Chen<sup>1</sup>

Emory University, 2021 Spring

(last updated on 01/12/2023)



Creative Commons License  
(CC BY-NC-SA)

<sup>1</sup>Slides are adapted from those by Karen Seyffarth from University of Calgary.



# Linear Algebra with Applications

## Lecture Notes

### Current Lecture Notes Revision: Version 2018 — Revision B

These lecture notes were originally developed by Karen Seyffarth of the University of Calgary. Edits, additions, and revisions have been made to these notes by the editorial team at Lyryx Learning to accompany their text [Linear Algebra with Applications](#) based on W. K. Nicholson's original text.

In addition we recognize the following contributors. All new content contributed is released under the same license as noted below.

- Ilijas Farah, York University

### BE A CHAMPION OF OER!

Contribute suggestions for improvements, new content, or errata:

A new topic

A new example or problem

A new or better proof to an existing theorem

Any other suggestions to improve the material

Contact Lyryx at [info@lyryx.com](mailto:info@lyryx.com) with your ideas.

### License



Attribution-NonCommercial-ShareAlike (CC BY-NC-SA)

This license lets others remix, tweak, and build upon your work non-commercially, as long as they credit you and license their new creations under the identical terms.

Copyright

Elementary Matrices

Inverses of elementary matrices

Smith Normal Form

Copyright

## Elementary Matrices

Inverses of elementary matrices

Smith Normal Form



# Elementary Matrices

## Definition

An **elementary matrix** is a matrix obtained from an identity matrix by performing a **single** elementary row operation.

# Elementary Matrices

## Definition

An **elementary matrix** is a matrix obtained from an identity matrix by performing a **single** elementary row operation.

## Remark ( Three Types of Elementary Row Operations )

( $\sim$  bases for genomic sequences)

- ▶ Type I: Interchange two rows.
- ▶ Type II: Multiply a row by a nonzero number.
- ▶ Type III: Add a (nonzero) multiple of one row to a different row.



### Example

Switch the 2nd row  
and the 4th row

Multiply  $-2$  to the  
3rd row

Add  $-3$  multiple of  
1st row to the 3rd row

$$E = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \end{bmatrix}, F = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -2 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}, G = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ -3 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix},$$

are examples of elementary matrices of types I, II and III, respectively.

Example (continued)

Let

$$A = \begin{bmatrix} 1 & 1 \\ 2 & 2 \\ 3 & 3 \\ 4 & 4 \end{bmatrix}$$

## Example (continued)

Let

$$A = \begin{bmatrix} 1 & 1 \\ 2 & 2 \\ 3 & 3 \\ 4 & 4 \end{bmatrix}$$

We are interested in the effect that (left) multiplication of A by E, F and G has on the matrix A.

## Example (continued)

Let

$$A = \begin{bmatrix} 1 & 1 \\ 2 & 2 \\ 3 & 3 \\ 4 & 4 \end{bmatrix}$$

We are interested in the effect that (left) multiplication of A by E, F and G has on the matrix A. Computing EA, FA, and GA ...

### Example (continued)

$$EA = \left[ \begin{array}{cccc} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \end{array} \right] \left[ \begin{array}{cc} 1 & 1 \\ 2 & 2 \\ 3 & 3 \\ 4 & 4 \end{array} \right] = \left[ \begin{array}{cc} 1 & 1 \\ 4 & 4 \\ 3 & 3 \\ 2 & 2 \end{array} \right]$$

Switch the 2nd row  
and the 4th row

### Example (continued)

$$EA = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 2 & 2 \\ 3 & 3 \\ 4 & 4 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 4 & 4 \\ 3 & 3 \\ 2 & 2 \end{bmatrix}$$

Switch the 2nd row  
and the 4th row

$$FA = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -2 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 2 & 2 \\ 3 & 3 \\ 4 & 4 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 2 & 2 \\ -6 & -6 \\ 4 & 4 \end{bmatrix}$$

Multiply  $-2$  to the  
3rd row

### Example (continued)

$$EA = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 2 & 2 \\ 3 & 3 \\ 4 & 4 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 4 & 4 \\ 3 & 3 \\ 2 & 2 \end{bmatrix}$$

Switch the 2nd row  
and the 4th row

$$FA = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -2 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 2 & 2 \\ 3 & 3 \\ 4 & 4 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 2 & 2 \\ -6 & -6 \\ 4 & 4 \end{bmatrix}$$

Multiply  $-2$  to the  
3rd row

$$GA = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ -3 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 2 & 2 \\ 3 & 3 \\ 4 & 4 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 2 & 2 \\ 0 & 0 \\ 4 & 4 \end{bmatrix}$$

Add  $-3$  multiple of  
1st row to the 3rd row



### Remark

The elementary matrices are the programmed receipts for your cooking!

## Theorem (Multiplication by an Elementary Matrix)

Let  $A$  be an  $m \times n$  matrix.

If  $B$  is obtained from  $A$  by performing **one single elementary** row operation,

then  $B = EA$

where  $E$  is the elementary matrix obtained from  $I_m$  by performing the same elementary operation on  $I_m$  as was performed on  $A$ .

$$A \longrightarrow B$$

$$\text{El. Op.} \quad \implies \quad A = EB$$

$$I \longrightarrow E$$



## Problem

Let

$$A = \begin{bmatrix} 4 & 1 \\ 1 & 3 \end{bmatrix} \quad \text{and} \quad C = \begin{bmatrix} 1 & 3 \\ 2 & -5 \end{bmatrix}$$

Find elementary matrices E and F so that  $C = FEA$ .

## Problem

Let

$$A = \begin{bmatrix} 4 & 1 \\ 1 & 3 \end{bmatrix} \quad \text{and} \quad C = \begin{bmatrix} 1 & 3 \\ 2 & -5 \end{bmatrix}$$

Find elementary matrices E and F so that  $C = FEA$ .

## Solution

Note. The statement of the problem implies that **C can be obtained from A by a sequence of two elementary row operations**, represented by elementary matrices E and F.

## Problem

Let

$$A = \begin{bmatrix} 4 & 1 \\ 1 & 3 \end{bmatrix} \quad \text{and} \quad C = \begin{bmatrix} 1 & 3 \\ 2 & -5 \end{bmatrix}$$

Find elementary matrices E and F so that  $C = FEA$ .

## Solution

Note. The statement of the problem implies that **C can be obtained from A by a sequence of two elementary row operations**, represented by elementary matrices E and F.

$$A = \begin{bmatrix} 4 & 1 \\ 1 & 3 \end{bmatrix}$$

## Problem

Let

$$A = \begin{bmatrix} 4 & 1 \\ 1 & 3 \end{bmatrix} \quad \text{and} \quad C = \begin{bmatrix} 1 & 3 \\ 2 & -5 \end{bmatrix}$$

Find elementary matrices E and F so that  $C = FEA$ .

## Solution

Note. The statement of the problem implies that **C can be obtained from A by a sequence of two elementary row operations**, represented by elementary matrices E and F.

$$A = \begin{bmatrix} 4 & 1 \\ 1 & 3 \end{bmatrix} \xrightarrow{E} \begin{bmatrix} 1 & 3 \\ 4 & 1 \end{bmatrix}$$

## Problem

Let

$$A = \begin{bmatrix} 4 & 1 \\ 1 & 3 \end{bmatrix} \quad \text{and} \quad C = \begin{bmatrix} 1 & 3 \\ 2 & -5 \end{bmatrix}$$

Find elementary matrices E and F so that  $C = FEA$ .

## Solution

Note. The statement of the problem implies that **C can be obtained from A by a sequence of two elementary row operations**, represented by elementary matrices E and F.

$$A = \begin{bmatrix} 4 & 1 \\ 1 & 3 \end{bmatrix} \xrightarrow{E} \begin{bmatrix} 1 & 3 \\ 4 & 1 \end{bmatrix} \xrightarrow{F} \begin{bmatrix} 1 & 3 \\ 2 & -5 \end{bmatrix} = C$$

## Problem

Let

$$A = \begin{bmatrix} 4 & 1 \\ 1 & 3 \end{bmatrix} \quad \text{and} \quad C = \begin{bmatrix} 1 & 3 \\ 2 & -5 \end{bmatrix}$$

Find elementary matrices E and F so that  $C = FEA$ .

## Solution

Note. The statement of the problem implies that **C can be obtained from A by a sequence of two elementary row operations**, represented by elementary matrices E and F.

$$A = \begin{bmatrix} 4 & 1 \\ 1 & 3 \end{bmatrix} \xrightarrow{E} \begin{bmatrix} 1 & 3 \\ 4 & 1 \end{bmatrix} \xrightarrow{F} \begin{bmatrix} 1 & 3 \\ 2 & -5 \end{bmatrix} = C$$

$$\text{where } E = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$$

## Problem

Let

$$A = \begin{bmatrix} 4 & 1 \\ 1 & 3 \end{bmatrix} \quad \text{and} \quad C = \begin{bmatrix} 1 & 3 \\ 2 & -5 \end{bmatrix}$$

Find elementary matrices E and F so that  $C = FEA$ .

## Solution

Note. The statement of the problem implies that **C can be obtained from A by a sequence of two elementary row operations**, represented by elementary matrices E and F.

$$A = \begin{bmatrix} 4 & 1 \\ 1 & 3 \end{bmatrix} \xrightarrow{E} \begin{bmatrix} 1 & 3 \\ 4 & 1 \end{bmatrix} \xrightarrow{F} \begin{bmatrix} 1 & 3 \\ 2 & -5 \end{bmatrix} = C$$

$$\text{where } E = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \text{ and } F = \begin{bmatrix} 1 & 0 \\ -2 & 1 \end{bmatrix}.$$

## Problem

Let

$$A = \begin{bmatrix} 4 & 1 \\ 1 & 3 \end{bmatrix} \quad \text{and} \quad C = \begin{bmatrix} 1 & 3 \\ 2 & -5 \end{bmatrix}$$

Find elementary matrices E and F so that  $C = FEA$ .

## Solution

Note. The statement of the problem implies that **C can be obtained from A by a sequence of two elementary row operations**, represented by elementary matrices E and F.

$$A = \begin{bmatrix} 4 & 1 \\ 1 & 3 \end{bmatrix} \xrightarrow{E} \begin{bmatrix} 1 & 3 \\ 4 & 1 \end{bmatrix} \xrightarrow{F} \begin{bmatrix} 1 & 3 \\ 2 & -5 \end{bmatrix} = C$$

where  $E = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$  and  $F = \begin{bmatrix} 1 & 0 \\ -2 & 1 \end{bmatrix}$ . Thus we have the sequence  
 $A \rightarrow EA \rightarrow F(EA) = C$ ,



## Problem

Let

$$A = \begin{bmatrix} 4 & 1 \\ 1 & 3 \end{bmatrix} \quad \text{and} \quad C = \begin{bmatrix} 1 & 3 \\ 2 & -5 \end{bmatrix}$$

Find elementary matrices E and F so that  $C = FEA$ .

## Solution

Note. The statement of the problem implies that **C can be obtained from A by a sequence of two elementary row operations**, represented by elementary matrices E and F.

$$A = \begin{bmatrix} 4 & 1 \\ 1 & 3 \end{bmatrix} \xrightarrow{E} \begin{bmatrix} 1 & 3 \\ 4 & 1 \end{bmatrix} \xrightarrow{F} \begin{bmatrix} 1 & 3 \\ 2 & -5 \end{bmatrix} = C$$

where  $E = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$  and  $F = \begin{bmatrix} 1 & 0 \\ -2 & 1 \end{bmatrix}$ . Thus we have the sequence  $A \rightarrow EA \rightarrow F(EA) = C$ , so  $C = FEA$ , i.e.,

$$\begin{bmatrix} 1 & 3 \\ 2 & -5 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ -2 & 1 \end{bmatrix} \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 4 & 1 \\ 1 & 3 \end{bmatrix}.$$



Copyright

Elementary Matrices

**Inverses of elementary matrices**

Smith Normal Form

## Inverses of Elementary Matrices

# Inverses of Elementary Matrices

## Lemma

Every elementary matrix  $E$  is invertible, and  $E^{-1}$  is also an elementary matrix (of the same type). Moreover,  $E^{-1}$  corresponds to the inverse of the row operation that produces  $E$ .

# Inverses of Elementary Matrices

## Lemma

Every elementary matrix  $E$  is invertible, and  $E^{-1}$  is also an elementary matrix (of the same type). Moreover,  $E^{-1}$  corresponds to the inverse of the row operation that produces  $E$ .

The following table gives the inverse of each type of elementary row operation:

Type	Operation	Inverse Operation
I	Interchange rows $p$ and $q$	Interchange rows $p$ and $q$
II	Multiply row $p$ by $k \neq 0$	Multiply row $p$ by $1/k$
III	Add $k$ times row $p$ to row $q \neq p$	Subtract $k$ times row $p$ from row $q$

Note that elementary matrices of type I are self-inverse.

# Inverses of Elementary Matrices

## Example

Without using the matrix inversion algorithm, find the inverse of the elementary matrix

$$G = \left[ \begin{array}{ccccc} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ -3 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{array} \right]$$

# Inverses of Elementary Matrices

## Example

Without using the matrix inversion algorithm, find the inverse of the elementary matrix

$$G = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ -3 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

Hint. What row operation can be applied to  $G$  to transform it to  $I_4$ ?

# Inverses of Elementary Matrices

## Example

Without using the matrix inversion algorithm, find the inverse of the elementary matrix

$$G = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ -3 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

Hint. What row operation can be applied to  $G$  to transform it to  $I_4$ ? The row operation  $G \rightarrow I_4$  is to **add** three times row one to row three,



# Inverses of Elementary Matrices

## Example

Without using the matrix inversion algorithm, find the inverse of the elementary matrix

$$G = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ -3 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

Hint. What row operation can be applied to  $G$  to transform it to  $I_4$ ? The row operation  $G \rightarrow I_4$  is to **add** three times row one to row three, and thus

$$G^{-1} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 3 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

# Inverses of Elementary Matrices

## Example

Without using the matrix inversion algorithm, find the inverse of the elementary matrix

$$G = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ -3 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

Hint. What row operation can be applied to  $G$  to transform it to  $I_4$ ? The row operation  $G \rightarrow I_4$  is to **add** three times row one to row three, and thus

$$G^{-1} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 3 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

**Check by computing  $G^{-1}G$ .**

Example (continued)

Similarly,

$$E^{-1} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \end{bmatrix}^{-1} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \end{bmatrix}$$

### Example (continued)

Similarly,

$$\mathbf{E}^{-1} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \end{bmatrix}^{-1} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \end{bmatrix}$$

and

$$\mathbf{F}^{-1} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -2 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}^{-1} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -\frac{1}{2} & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

Suppose  $A$  is an  $m \times n$  matrix and that  $B$  can be obtained from  $A$  by a sequence of  $k$  elementary row operations.

Suppose  $A$  is an  $m \times n$  matrix and that  $B$  can be obtained from  $A$  by a sequence of  $k$  elementary row operations. Then there exist elementary matrices  $E_1, E_2, \dots, E_k$  such that

$$B = E_k(E_{k-1}(\cdots(E_2(E_1A))\cdots))$$

Suppose  $A$  is an  $m \times n$  matrix and that  $B$  can be obtained from  $A$  by a sequence of  $k$  elementary row operations. Then there exist elementary matrices  $E_1, E_2, \dots, E_k$  such that

$$B = E_k(E_{k-1}(\cdots(E_2(E_1A))\cdots))$$

Since matrix multiplication is associative, we have

$$B = (E_k E_{k-1} \cdots E_2 E_1)A$$

Suppose  $A$  is an  $m \times n$  matrix and that  $B$  can be obtained from  $A$  by a sequence of  $k$  elementary row operations. Then there exist elementary matrices  $E_1, E_2, \dots, E_k$  such that

$$B = E_k(E_{k-1}(\cdots(E_2(E_1A))\cdots))$$

Since matrix multiplication is associative, we have

$$B = (E_k E_{k-1} \cdots E_2 E_1)A$$

or, more concisely,  $B = UA$  where  $U = E_k E_{k-1} \cdots E_2 E_1$ .



Suppose  $A$  is an  $m \times n$  matrix and that  $B$  can be obtained from  $A$  by a sequence of  $k$  elementary row operations. Then there exist elementary matrices  $E_1, E_2, \dots, E_k$  such that

$$B = E_k(E_{k-1}(\cdots(E_2(E_1A))\cdots))$$

Since matrix multiplication is associative, we have

$$B = (E_k E_{k-1} \cdots E_2 E_1)A$$

or, more concisely,  $B = UA$  where  $U = E_k E_{k-1} \cdots E_2 E_1$ .

To find  $U$  so that  $B = UA$ , we **could** find  $E_1, E_2, \dots, E_k$  and multiply these together (in the correct order), but there is an easier method for finding  $U$ .

## Definition

Let  $A$  be an  $m \times n$  matrix. We write

$$A \rightarrow B$$

if  $B$  can be obtained from  $A$  by a sequence of elementary row operations.

## Definition

Let  $A$  be an  $m \times n$  matrix. We write

$$A \rightarrow B$$

if  $B$  can be obtained from  $A$  by a sequence of elementary row operations. In this case, we call  $A$  and  $B$  are **row-equivalent**.

## Definition

Let  $A$  be an  $m \times n$  matrix. We write

$$A \rightarrow B$$

if  $B$  can be obtained from  $A$  by a sequence of elementary row operations. In this case, we call  $A$  and  $B$  are **row-equivalent**.

## Theorem

Suppose  $A$  is an  $m \times n$  matrix and that  $A \rightarrow B$ . Then

## Definition

Let  $A$  be an  $m \times n$  matrix. We write

$$A \rightarrow B$$

if  $B$  can be obtained from  $A$  by a sequence of elementary row operations. In this case, we call  $A$  and  $B$  are **row-equivalent**.

## Theorem

Suppose  $A$  is an  $m \times n$  matrix and that  $A \rightarrow B$ . Then

1. there exists an **invertible**  $m \times m$  matrix  $U$  such that  $B = UA$ ;

## Definition

Let  $A$  be an  $m \times n$  matrix. We write

$$A \rightarrow B$$

if  $B$  can be obtained from  $A$  by a sequence of elementary row operations. In this case, we call  $A$  and  $B$  are **row-equivalent**.

## Theorem

Suppose  $A$  is an  $m \times n$  matrix and that  $A \rightarrow B$ . Then

1. there exists an **invertible**  $m \times m$  matrix  $U$  such that  $B = UA$ ;
2.  $U$  can be computed by performing elementary row operations on  $\left[ \begin{array}{c|c} A & I_m \end{array} \right]$  to transform it into  $\left[ \begin{array}{c|c} B & U \end{array} \right]$ ;

## Definition

Let  $A$  be an  $m \times n$  matrix. We write

$$A \rightarrow B$$

if  $B$  can be obtained from  $A$  by a sequence of elementary row operations. In this case, we call  $A$  and  $B$  are **row-equivalent**.

## Theorem

Suppose  $A$  is an  $m \times n$  matrix and that  $A \rightarrow B$ . Then

1. there exists an **invertible**  $m \times m$  matrix  $U$  such that  $B = UA$ ;
2.  $U$  can be computed by performing elementary row operations on  $\left[ \begin{array}{c|c} A & I_m \end{array} \right]$  to transform it into  $\left[ \begin{array}{c|c} B & U \end{array} \right]$ ;
3.  $U = E_k E_{k-1} \cdots E_2 E_1$ , where  $E_1, E_2, \dots, E_k$  are elementary matrices corresponding, in order, to the elementary row operations used to obtain  $B$  from  $A$ .

### Problem

Let  $A = \begin{bmatrix} 3 & 0 & 1 \\ 2 & -1 & 0 \end{bmatrix}$ , and let  $R$  be the reduced row-echelon form of  $A$ .

Find a matrix  $U$  so that  $R = UA$ .



## Problem

Let  $A = \begin{bmatrix} 3 & 0 & 1 \\ 2 & -1 & 0 \end{bmatrix}$ , and let  $R$  be the reduced row-echelon form of  $A$ . Find a matrix  $U$  so that  $R = UA$ .

## Solution

$$\begin{aligned} \left[ \begin{array}{ccc|cc} 3 & 0 & 1 & 1 & 0 \\ 2 & -1 & 0 & 0 & 1 \end{array} \right] &\rightarrow \left[ \begin{array}{ccc|cc} 1 & 1 & 1 & 1 & -1 \\ 2 & -1 & 0 & 0 & 1 \end{array} \right] \rightarrow \left[ \begin{array}{ccc|cc} 1 & 1 & 1 & 1 & -1 \\ 0 & -3 & -2 & -2 & 3 \end{array} \right] \\ &\rightarrow \left[ \begin{array}{ccc|cc} 1 & 1 & 1 & 1 & -1 \\ 0 & 1 & 2/3 & 2/3 & -1 \end{array} \right] \rightarrow \left[ \begin{array}{ccc|cc} 1 & 0 & 1/3 & 1/3 & 0 \\ 0 & 1 & 2/3 & 2/3 & -1 \end{array} \right] \end{aligned}$$

## Problem

Let  $A = \begin{bmatrix} 3 & 0 & 1 \\ 2 & -1 & 0 \end{bmatrix}$ , and let  $R$  be the reduced row-echelon form of  $A$ . Find a matrix  $U$  so that  $R = UA$ .

## Solution

$$\begin{aligned} \left[ \begin{array}{ccc|cc} 3 & 0 & 1 & 1 & 0 \\ 2 & -1 & 0 & 0 & 1 \end{array} \right] &\rightarrow \left[ \begin{array}{ccc|cc} 1 & 1 & 1 & 1 & -1 \\ 2 & -1 & 0 & 0 & 1 \end{array} \right] \rightarrow \left[ \begin{array}{ccc|cc} 1 & 1 & 1 & 1 & -1 \\ 0 & -3 & -2 & -2 & 3 \end{array} \right] \\ &\rightarrow \left[ \begin{array}{ccc|cc} 1 & 1 & 1 & 1 & -1 \\ 0 & 1 & 2/3 & 2/3 & -1 \end{array} \right] \rightarrow \left[ \begin{array}{ccc|cc} 1 & 0 & 1/3 & 1/3 & 0 \\ 0 & 1 & 2/3 & 2/3 & -1 \end{array} \right] \end{aligned}$$

Starting with  $[A \mid I]$ , we've obtained  $[R \mid U]$ .

## Problem

Let  $A = \begin{bmatrix} 3 & 0 & 1 \\ 2 & -1 & 0 \end{bmatrix}$ , and let  $R$  be the reduced row-echelon form of  $A$ . Find a matrix  $U$  so that  $R = UA$ .

## Solution

$$\begin{aligned} \left[ \begin{array}{ccc|cc} 3 & 0 & 1 & 1 & 0 \\ 2 & -1 & 0 & 0 & 1 \end{array} \right] &\rightarrow \left[ \begin{array}{ccc|cc} 1 & 1 & 1 & 1 & -1 \\ 2 & -1 & 0 & 0 & 1 \end{array} \right] \rightarrow \left[ \begin{array}{ccc|cc} 1 & 1 & 1 & 1 & -1 \\ 0 & -3 & -2 & -2 & 3 \end{array} \right] \\ &\rightarrow \left[ \begin{array}{ccc|cc} 1 & 1 & 1 & 1 & -1 \\ 0 & 1 & 2/3 & 2/3 & -1 \end{array} \right] \rightarrow \left[ \begin{array}{ccc|cc} 1 & 0 & 1/3 & 1/3 & 0 \\ 0 & 1 & 2/3 & 2/3 & -1 \end{array} \right] \end{aligned}$$

Starting with  $[A \mid I]$ , we've obtained  $[R \mid U]$ .

Therefore  $R = UA$ , where

$$U = \begin{bmatrix} 1/3 & 0 \\ 2/3 & -1 \end{bmatrix}.$$



Example ( A Matrix as a product of elementary matrices )

Let

$$A = \begin{bmatrix} 1 & 2 & -4 \\ -3 & -6 & 13 \\ 0 & -1 & 2 \end{bmatrix}.$$

### Example ( A Matrix as a product of elementary matrices )

Let

$$A = \begin{bmatrix} 1 & 2 & -4 \\ -3 & -6 & 13 \\ 0 & -1 & 2 \end{bmatrix}.$$

Suppose we do row operations to put A in reduced row-echelon form, and write down the corresponding elementary matrices.

### Example ( A Matrix as a product of elementary matrices )

Let

$$A = \begin{bmatrix} 1 & 2 & -4 \\ -3 & -6 & 13 \\ 0 & -1 & 2 \end{bmatrix}.$$

Suppose we do row operations to put A in reduced row-echelon form, and write down the corresponding elementary matrices.

$$\begin{bmatrix} 1 & 2 & -4 \\ -3 & -6 & 13 \\ 0 & -1 & 2 \end{bmatrix}$$

### Example ( A Matrix as a product of elementary matrices )

Let

$$A = \begin{bmatrix} 1 & 2 & -4 \\ -3 & -6 & 13 \\ 0 & -1 & 2 \end{bmatrix}.$$

Suppose we do row operations to put A in reduced row-echelon form, and write down the corresponding elementary matrices.

$$\begin{bmatrix} 1 & 2 & -4 \\ -3 & -6 & 13 \\ 0 & -1 & 2 \end{bmatrix} \xrightarrow{E_1} \begin{bmatrix} 1 & 2 & -4 \\ 0 & 0 & 1 \\ 0 & -1 & 2 \end{bmatrix}$$

### Example ( A Matrix as a product of elementary matrices )

Let

$$A = \begin{bmatrix} 1 & 2 & -4 \\ -3 & -6 & 13 \\ 0 & -1 & 2 \end{bmatrix}.$$

Suppose we do row operations to put A in reduced row-echelon form, and write down the corresponding elementary matrices.

$$\begin{bmatrix} 1 & 2 & -4 \\ -3 & -6 & 13 \\ 0 & -1 & 2 \end{bmatrix} \xrightarrow{E_1} \begin{bmatrix} 1 & 2 & -4 \\ 0 & 0 & 1 \\ 0 & -1 & 2 \end{bmatrix} \xrightarrow{E_2} \begin{bmatrix} 1 & 2 & -4 \\ 0 & -1 & 2 \\ 0 & 0 & 1 \end{bmatrix}$$



### Example ( A Matrix as a product of elementary matrices )

Let

$$A = \begin{bmatrix} 1 & 2 & -4 \\ -3 & -6 & 13 \\ 0 & -1 & 2 \end{bmatrix}.$$

Suppose we do row operations to put A in reduced row-echelon form, and write down the corresponding elementary matrices.

$$\begin{bmatrix} 1 & 2 & -4 \\ -3 & -6 & 13 \\ 0 & -1 & 2 \end{bmatrix} \xrightarrow{E_1} \begin{bmatrix} 1 & 2 & -4 \\ 0 & 0 & 1 \\ 0 & -1 & 2 \end{bmatrix} \xrightarrow{E_2} \begin{bmatrix} 1 & 2 & -4 \\ 0 & -1 & 2 \\ 0 & 0 & 1 \end{bmatrix} \xrightarrow{E_3}$$
$$\begin{bmatrix} 1 & 2 & -4 \\ 0 & 1 & -2 \\ 0 & 0 & 1 \end{bmatrix}$$

### Example ( A Matrix as a product of elementary matrices )

Let

$$A = \begin{bmatrix} 1 & 2 & -4 \\ -3 & -6 & 13 \\ 0 & -1 & 2 \end{bmatrix}.$$

Suppose we do row operations to put A in reduced row-echelon form, and write down the corresponding elementary matrices.

$$\begin{bmatrix} 1 & 2 & -4 \\ -3 & -6 & 13 \\ 0 & -1 & 2 \end{bmatrix} \xrightarrow{E_1} \begin{bmatrix} 1 & 2 & -4 \\ 0 & 0 & 1 \\ 0 & -1 & 2 \end{bmatrix} \xrightarrow{E_2} \begin{bmatrix} 1 & 2 & -4 \\ 0 & -1 & 2 \\ 0 & 0 & 1 \end{bmatrix} \xrightarrow{E_3}$$

$$\begin{bmatrix} 1 & 2 & -4 \\ 0 & 1 & -2 \\ 0 & 0 & 1 \end{bmatrix} \xrightarrow{E_4} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & -2 \\ 0 & 0 & 1 \end{bmatrix}$$

### Example ( A Matrix as a product of elementary matrices )

Let

$$A = \begin{bmatrix} 1 & 2 & -4 \\ -3 & -6 & 13 \\ 0 & -1 & 2 \end{bmatrix}.$$

Suppose we do row operations to put A in reduced row-echelon form, and write down the corresponding elementary matrices.

$$\begin{bmatrix} 1 & 2 & -4 \\ -3 & -6 & 13 \\ 0 & -1 & 2 \end{bmatrix} \xrightarrow{E_1} \begin{bmatrix} 1 & 2 & -4 \\ 0 & 0 & 1 \\ 0 & -1 & 2 \end{bmatrix} \xrightarrow{E_2} \begin{bmatrix} 1 & 2 & -4 \\ 0 & -1 & 2 \\ 0 & 0 & 1 \end{bmatrix} \xrightarrow{E_3} \\ \begin{bmatrix} 1 & 2 & -4 \\ 0 & 1 & -2 \\ 0 & 0 & 1 \end{bmatrix} \xrightarrow{E_4} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & -2 \\ 0 & 0 & 1 \end{bmatrix} \xrightarrow{E_5} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

### Example ( A Matrix as a product of elementary matrices )

Let

$$A = \begin{bmatrix} 1 & 2 & -4 \\ -3 & -6 & 13 \\ 0 & -1 & 2 \end{bmatrix}.$$

Suppose we do row operations to put A in reduced row-echelon form, and write down the corresponding elementary matrices.

$$\begin{aligned} \begin{bmatrix} 1 & 2 & -4 \\ -3 & -6 & 13 \\ 0 & -1 & 2 \end{bmatrix} &\xrightarrow{E_1} \begin{bmatrix} 1 & 2 & -4 \\ 0 & 0 & 1 \\ 0 & -1 & 2 \end{bmatrix} \xrightarrow{E_2} \begin{bmatrix} 1 & 2 & -4 \\ 0 & -1 & 2 \\ 0 & 0 & 1 \end{bmatrix} \xrightarrow{E_3} \\ &\begin{bmatrix} 1 & 2 & -4 \\ 0 & 1 & -2 \\ 0 & 0 & 1 \end{bmatrix} \xrightarrow{E_4} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & -2 \\ 0 & 0 & 1 \end{bmatrix} \xrightarrow{E_5} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \end{aligned}$$

Notice that the reduced row-echelon form of A equals  $I_3$ . Now find the matrices  $E_1, E_2, E_3, E_4$  and  $E_5$ .

Example (continued)

$$E_1 = \begin{bmatrix} 1 & 0 & 0 \\ 3 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix},$$

Example (continued)

$$E_1 = \begin{bmatrix} 1 & 0 & 0 \\ 3 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, E_2 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix},$$

Example (continued)

$$E_1 = \begin{bmatrix} 1 & 0 & 0 \\ 3 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, E_2 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}, E_3 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Example (continued)

$$E_1 = \begin{bmatrix} 1 & 0 & 0 \\ 3 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, E_2 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}, E_3 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$E_4 = \begin{bmatrix} 1 & -2 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix},$$



Example (continued)

$$E_1 = \begin{bmatrix} 1 & 0 & 0 \\ 3 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, E_2 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}, E_3 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$E_4 = \begin{bmatrix} 1 & -2 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, E_5 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 2 \\ 0 & 0 & 1 \end{bmatrix}$$

Example (continued)

$$\begin{aligned} E_1 &= \begin{bmatrix} 1 & 0 & 0 \\ 3 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, E_2 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}, E_3 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \\ E_4 &= \begin{bmatrix} 1 & -2 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, E_5 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 2 \\ 0 & 0 & 1 \end{bmatrix} \end{aligned}$$

It follows that

$$\begin{aligned} (E_5(E_4(E_3(E_2(E_1A)))))) &= I \\ (E_5E_4E_3E_2E_1)A &= I \end{aligned}$$

and therefore

$$A^{-1} = E_5E_4E_3E_2E_1$$

### Example (continued)

Since  $A^{-1} = E_5 E_4 E_3 E_2 E_1$ ,

$$\begin{aligned} A^{-1} &= E_5 E_4 E_3 E_2 E_1 \\ (A^{-1})^{-1} &= (E_5 E_4 E_3 E_2 E_1)^{-1} \\ A &= E_1^{-1} E_2^{-1} E_3^{-1} E_4^{-1} E_5^{-1} \end{aligned}$$

### Example (continued)

Since  $A^{-1} = E_5 E_4 E_3 E_2 E_1$ ,

$$\begin{aligned} A^{-1} &= E_5 E_4 E_3 E_2 E_1 \\ (A^{-1})^{-1} &= (E_5 E_4 E_3 E_2 E_1)^{-1} \\ A &= E_1^{-1} E_2^{-1} E_3^{-1} E_4^{-1} E_5^{-1} \end{aligned}$$

This example illustrates the following result.

### Example (continued)

Since  $A^{-1} = E_5 E_4 E_3 E_2 E_1$ ,

$$\begin{aligned} A^{-1} &= E_5 E_4 E_3 E_2 E_1 \\ (A^{-1})^{-1} &= (E_5 E_4 E_3 E_2 E_1)^{-1} \\ A &= E_1^{-1} E_2^{-1} E_3^{-1} E_4^{-1} E_5^{-1} \end{aligned}$$

This example illustrates the following result.

### Theorem

Let  $A$  be an  $n \times n$  matrix. Then,  $A^{-1}$  exists if and only if  $A$  can be written as the product of elementary matrices.

Example ( revisited – Matrix inversion algorithm)

$$\left[ \begin{array}{ccc|c} 1 & 2 & -4 & I \end{array} \right] = \left[ \begin{array}{ccc|c} 1 & 2 & -4 & \\ -3 & -6 & 13 & \\ 0 & -1 & 2 & \end{array} \right]$$

$$E_1 \left[ \begin{array}{ccc|c} 1 & 2 & -4 & \\ 0 & 0 & 1 & E_1 \\ 0 & -1 & 2 & \end{array} \right] = \left[ \begin{array}{ccc|ccc} 1 & 2 & -4 & 1 & 0 & 0 \\ 0 & 0 & 1 & 3 & 1 & 0 \\ 0 & -1 & 2 & 0 & 0 & 1 \end{array} \right]$$

$$E_2 E_1 \left[ \begin{array}{ccc|c} 1 & 2 & -4 & \\ 0 & -1 & 2 & E_2 E_1 \\ 0 & 0 & 1 & \end{array} \right] = \left[ \begin{array}{ccc|ccc} 1 & 2 & -4 & 1 & 0 & 0 \\ 0 & -1 & 2 & 0 & 0 & 1 \\ 0 & 0 & 1 & 3 & 1 & 0 \end{array} \right]$$

Example ( continued )

$$E_3E_2E_1[ A \mid I ] = \left[ \begin{array}{ccc|c} 1 & 2 & -4 & E_3E_2E_1 \\ 0 & 1 & -2 & \\ 0 & 0 & 1 & \end{array} \right] = \left[ \begin{array}{ccc|ccc} 1 & 2 & -4 & 1 & 0 & 0 \\ 0 & 1 & -2 & 0 & 0 & -1 \\ 0 & 0 & 1 & 3 & 1 & 0 \end{array} \right]$$

$$E_4E_3E_2E_1[ A \mid I ] = \left[ \begin{array}{ccc|c} 1 & 0 & 0 & E_4E_3E_2E_1 \\ 0 & 1 & -2 & \\ 0 & 0 & 1 & \end{array} \right] = \left[ \begin{array}{ccc|ccc} 1 & 0 & 0 & 1 & 0 & 2 \\ 0 & 1 & -2 & 0 & 0 & -1 \\ 0 & 0 & 1 & 3 & 1 & 0 \end{array} \right]$$

$$E_5E_4E_3E_2E_1[ A \mid I ] = \left[ \begin{array}{ccc|c} 1 & 0 & 0 & E_5E_4E_3E_2E_1 \\ 0 & 1 & 0 & \\ 0 & 0 & 1 & \end{array} \right] = \left[ \begin{array}{ccc|ccc} 1 & 0 & 0 & 1 & 0 & 2 \\ 0 & 1 & 0 & 6 & 2 & -1 \\ 0 & 0 & 1 & 3 & 1 & 0 \end{array} \right]$$

$$A^{-1} = E_5E_4E_3E_2E_1 = \left[ \begin{array}{ccc} 1 & 0 & 2 \\ 6 & 2 & -1 \\ 3 & 1 & 0 \end{array} \right]$$

### Problem

Express  $A = \begin{bmatrix} 4 & 1 \\ -3 & 2 \end{bmatrix}$  as a product of elementary matrices.



### Problem

Express  $A = \begin{bmatrix} 4 & 1 \\ -3 & 2 \end{bmatrix}$  as a product of elementary matrices.

### Solution

$$\begin{bmatrix} 4 & 1 \\ -3 & 2 \end{bmatrix}$$

### Problem

Express  $A = \begin{bmatrix} 4 & 1 \\ -3 & 2 \end{bmatrix}$  as a product of elementary matrices.

### Solution

$$\begin{bmatrix} 4 & 1 \\ -3 & 2 \end{bmatrix} \xrightarrow{E_1} \begin{bmatrix} 1 & 3 \\ -3 & 2 \end{bmatrix}$$

### Problem

Express  $A = \begin{bmatrix} 4 & 1 \\ -3 & 2 \end{bmatrix}$  as a product of elementary matrices.

### Solution

$$\begin{bmatrix} 4 & 1 \\ -3 & 2 \end{bmatrix} \xrightarrow{E_1} \begin{bmatrix} 1 & 3 \\ -3 & 2 \end{bmatrix} \xrightarrow{E_2} \begin{bmatrix} 1 & 3 \\ 0 & 11 \end{bmatrix}$$

### Problem

Express  $A = \begin{bmatrix} 4 & 1 \\ -3 & 2 \end{bmatrix}$  as a product of elementary matrices.

### Solution

$$\begin{bmatrix} 4 & 1 \\ -3 & 2 \end{bmatrix} \xrightarrow{E_1} \begin{bmatrix} 1 & 3 \\ -3 & 2 \end{bmatrix} \xrightarrow{E_2} \begin{bmatrix} 1 & 3 \\ 0 & 11 \end{bmatrix} \xrightarrow{E_3} \begin{bmatrix} 1 & 3 \\ 0 & 1 \end{bmatrix}$$

### Problem

Express  $A = \begin{bmatrix} 4 & 1 \\ -3 & 2 \end{bmatrix}$  as a product of elementary matrices.

### Solution

$$\begin{bmatrix} 4 & 1 \\ -3 & 2 \end{bmatrix} \xrightarrow{E_1} \begin{bmatrix} 1 & 3 \\ -3 & 2 \end{bmatrix} \xrightarrow{E_2} \begin{bmatrix} 1 & 3 \\ 0 & 11 \end{bmatrix} \xrightarrow{E_3} \begin{bmatrix} 1 & 3 \\ 0 & 1 \end{bmatrix} \xrightarrow{E_4} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

## Problem

Express  $A = \begin{bmatrix} 4 & 1 \\ -3 & 2 \end{bmatrix}$  as a product of elementary matrices.

## Solution

$$\begin{bmatrix} 4 & 1 \\ -3 & 2 \end{bmatrix} \xrightarrow{E_1} \begin{bmatrix} 1 & 3 \\ -3 & 2 \end{bmatrix} \xrightarrow{E_2} \begin{bmatrix} 1 & 3 \\ 0 & 11 \end{bmatrix} \xrightarrow{E_3} \begin{bmatrix} 1 & 3 \\ 0 & 1 \end{bmatrix} \xrightarrow{E_4} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

with

$$E_1 = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix},$$

## Problem

Express  $A = \begin{bmatrix} 4 & 1 \\ -3 & 2 \end{bmatrix}$  as a product of elementary matrices.

## Solution

$$\begin{bmatrix} 4 & 1 \\ -3 & 2 \end{bmatrix} \xrightarrow{E_1} \begin{bmatrix} 1 & 3 \\ -3 & 2 \end{bmatrix} \xrightarrow{E_2} \begin{bmatrix} 1 & 3 \\ 0 & 11 \end{bmatrix} \xrightarrow{E_3} \begin{bmatrix} 1 & 3 \\ 0 & 1 \end{bmatrix} \xrightarrow{E_4} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

with

$$E_1 = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}, E_2 = \begin{bmatrix} 1 & 0 \\ 3 & 1 \end{bmatrix},$$

## Problem

Express  $A = \begin{bmatrix} 4 & 1 \\ -3 & 2 \end{bmatrix}$  as a product of elementary matrices.

## Solution

$$\begin{bmatrix} 4 & 1 \\ -3 & 2 \end{bmatrix} \xrightarrow{E_1} \begin{bmatrix} 1 & 3 \\ -3 & 2 \end{bmatrix} \xrightarrow{E_2} \begin{bmatrix} 1 & 3 \\ 0 & 11 \end{bmatrix} \xrightarrow{E_3} \begin{bmatrix} 1 & 3 \\ 0 & 1 \end{bmatrix} \xrightarrow{E_4} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

with

$$E_1 = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}, E_2 = \begin{bmatrix} 1 & 0 \\ 3 & 1 \end{bmatrix}, E_3 = \begin{bmatrix} 1 & 0 \\ 0 & \frac{1}{11} \end{bmatrix},$$



## Problem

Express  $A = \begin{bmatrix} 4 & 1 \\ -3 & 2 \end{bmatrix}$  as a product of elementary matrices.

## Solution

$$\begin{bmatrix} 4 & 1 \\ -3 & 2 \end{bmatrix} \xrightarrow{E_1} \begin{bmatrix} 1 & 3 \\ -3 & 2 \end{bmatrix} \xrightarrow{E_2} \begin{bmatrix} 1 & 3 \\ 0 & 11 \end{bmatrix} \xrightarrow{E_3} \begin{bmatrix} 1 & 3 \\ 0 & 1 \end{bmatrix} \xrightarrow{E_4} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

with

$$E_1 = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}, E_2 = \begin{bmatrix} 1 & 0 \\ 3 & 1 \end{bmatrix}, E_3 = \begin{bmatrix} 1 & 0 \\ 0 & \frac{1}{11} \end{bmatrix}, E_4 = \begin{bmatrix} 1 & -3 \\ 0 & 1 \end{bmatrix}$$

## Problem

Express  $A = \begin{bmatrix} 4 & 1 \\ -3 & 2 \end{bmatrix}$  as a product of elementary matrices.

## Solution

$$\begin{bmatrix} 4 & 1 \\ -3 & 2 \end{bmatrix} \xrightarrow{E_1} \begin{bmatrix} 1 & 3 \\ -3 & 2 \end{bmatrix} \xrightarrow{E_2} \begin{bmatrix} 1 & 3 \\ 0 & 11 \end{bmatrix} \xrightarrow{E_3} \begin{bmatrix} 1 & 3 \\ 0 & 1 \end{bmatrix} \xrightarrow{E_4} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

with

$$E_1 = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}, E_2 = \begin{bmatrix} 1 & 0 \\ 3 & 1 \end{bmatrix}, E_3 = \begin{bmatrix} 1 & 0 \\ 0 & \frac{1}{11} \end{bmatrix}, E_4 = \begin{bmatrix} 1 & -3 \\ 0 & 1 \end{bmatrix}$$

Since  $E_4 E_3 E_2 E_1 A = I$ ,  $A^{-1} = E_4 E_3 E_2 E_1$ , and hence

$$A = E_1^{-1} E_2^{-1} E_3^{-1} E_4^{-1}$$

Solution (continued)

Therefore,

$$A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}^{-1} \begin{bmatrix} 1 & 0 \\ 3 & 1 \end{bmatrix}^{-1} \begin{bmatrix} 1 & 0 \\ 0 & 1/11 \end{bmatrix}^{-1} \begin{bmatrix} 1 & -3 \\ 0 & 1 \end{bmatrix}^{-1}$$

## Solution (continued)

Therefore,

$$A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}^{-1} \begin{bmatrix} 1 & 0 \\ 3 & 1 \end{bmatrix}^{-1} \begin{bmatrix} 1 & 0 \\ 0 & 1/11 \end{bmatrix}^{-1} \begin{bmatrix} 1 & -3 \\ 0 & 1 \end{bmatrix}^{-1}$$

i.e.,

$$A = \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ -3 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 11 \end{bmatrix} \begin{bmatrix} 1 & 3 \\ 0 & 1 \end{bmatrix}$$



One result that we have assumed in all our work involving reduced row-echelon matrices is the following.

One result that we have assumed in all our work involving reduced row-echelon matrices is the following.

**Theorem ( Uniqueness of the Reduced Echelon Form )**

If  $A$  is an  $m \times n$  matrix and  $R$  and  $S$  are reduced row-echelon forms of  $A$ , then  $R = S$ .

One result that we have assumed in all our work involving reduced row-echelon matrices is the following.

### Theorem ( Uniqueness of the Reduced Echelon Form )

If  $A$  is an  $m \times n$  matrix and  $R$  and  $S$  are reduced row-echelon forms of  $A$ , then  $R = S$ .

### Remark

This theorem ensures that the reduced row-echelon form of a matrix is unique,

One result that we have assumed in all our work involving reduced row-echelon matrices is the following.

### Theorem ( Uniqueness of the Reduced Echelon Form )

If  $A$  is an  $m \times n$  matrix and  $R$  and  $S$  are reduced row-echelon forms of  $A$ , then  $R = S$ .

### Remark

This theorem ensures that the reduced row-echelon form of a matrix is **unique**, and its proof follows from the results about elementary matrices.



Copyright

Elementary Matrices

Inverses of elementary matrices

**Smith Normal Form**

## Smith Normal Form

# Smith Normal Form

## Definition

If  $A$  is an  $m \times n$  matrix of rank  $r$ , then the matrix  $\begin{pmatrix} I_r & 0 \\ 0 & 0 \end{pmatrix}_{m \times n}$  is called the **Smith normal form** of  $A$ .

## Theorem

If  $A$  is an  $m \times n$  matrix of rank  $r$ , then there exist invertible matrices  $U$  and  $V$  of size  $m \times m$  and  $n \times n$ , respectively, such that

$$UAV = \begin{pmatrix} I_r & 0 \\ 0 & 0 \end{pmatrix}_{m \times n}$$

Proof.

1. Apply the elementary row operations:

$$[A|I_m] \xrightarrow{\text{e.r.o.}} [\text{rref}(A)|U]$$

2. Apply the elementary column operations:

$$\begin{pmatrix} \text{rref}(A) \\ I_n \end{pmatrix} \xrightarrow{\text{e.c.o.}} \begin{pmatrix} \begin{pmatrix} I_r & 0 \\ 0 & 0 \end{pmatrix}_{m \times n} \\ V \end{pmatrix}_{2m \times n}$$



Remark

The elementary column operations above are equivalent to the elementary row operations on the transpose:

$$[\text{rref}(A)^T|I_n] \xrightarrow{\text{e.r.o.}} \left[ \begin{pmatrix} I_r & 0 \\ 0 & 0 \end{pmatrix}_{n \times m} \middle| V^T \right]_{n \times 2m}$$

## Problem

Find the decomposition of  $A = \begin{bmatrix} 3 & 0 & 1 \\ 2 & -1 & 0 \end{bmatrix}$  into the Smith normal form:

$A = \tilde{U}N\tilde{V}$ , where  $N$  is the Smith normal form of  $A$  and  $\tilde{U}, \tilde{V}$  are some invertible matrices.

## Problem

Find the decomposition of  $A = \begin{bmatrix} 3 & 0 & 1 \\ 2 & -1 & 0 \end{bmatrix}$  into the Smith normal form:

$A = \tilde{U}N\tilde{V}$ , where  $N$  is the Smith normal form of  $A$  and  $\tilde{U}, \tilde{V}$  are some invertible matrices.

## Solution

We have seen that

$$[A|I_2] = \left[ \begin{array}{ccc|cc} 3 & 0 & 1 & 1 & 0 \\ 2 & -1 & 0 & 0 & 1 \end{array} \right] \rightarrow \left[ \begin{array}{ccc|cc} 1 & 0 & 1/3 & 1/3 & 0 \\ 0 & 1 & 2/3 & 2/3 & -1 \end{array} \right] = [\text{rref}(A)|U]$$

Now,

$$\left( \text{rref}(A)^T \mid I_3 \right) = \left[ \begin{array}{cc|ccc} 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 \\ \frac{1}{3} & \frac{2}{3} & 0 & 0 & 1 \end{array} \right] \rightarrow \left[ \begin{array}{cc|ccc} 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 \\ 0 & 0 & -\frac{1}{3} & -\frac{2}{3} & 1 \end{array} \right] = [N^T|V^T]$$

## Solution (Continued)

Hence, we find  $N = UAV$ , namely,

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix} = \begin{pmatrix} 1/3 & 0 \\ 2/3 & -1 \end{pmatrix} \begin{bmatrix} 3 & 0 & 1 \\ 2 & -1 & 0 \end{bmatrix} \begin{pmatrix} 1 & 0 & -1/3 \\ 0 & 1 & -2/3 \\ 0 & 0 & 1 \end{pmatrix}$$

Finally, since  $U$  and  $V$  are invertible, we see that

$$A = U^{-1}NV^{-1},$$

namely,

$$\begin{aligned} A = \begin{bmatrix} 3 & 0 & 1 \\ 2 & -1 & 0 \end{bmatrix} &= \begin{pmatrix} 1/3 & 0 \\ 2/3 & -1 \end{pmatrix}^{-1} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 0 & -1/3 \\ 0 & 1 & -2/3 \\ 0 & 0 & 1 \end{pmatrix}^{-1} \\ &= \begin{pmatrix} 3 & 0 \\ 2 & -1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 0 & 1/3 \\ 0 & 1 & 2/3 \\ 0 & 0 & 1 \end{pmatrix} \\ &= \tilde{U}N\tilde{V}. \end{aligned}$$

