

Math 221: LINEAR ALGEBRA

Chapter 2. Matrix Algebra §2-6. Linear Transformations

Le Chen¹

Emory University, 2021 Spring

(last updated on 01/12/2023)



Creative Commons License
(CC BY-NC-SA)

¹Slides are adapted from those by Karen Seyffarth from University of Calgary.

Linear Algebra with Applications

Lecture Notes

Current Lecture Notes Revision: Version 2018 — Revision B

These lecture notes were originally developed by Karen Seyffarth of the University of Calgary. Edits, additions, and revisions have been made to these notes by the editorial team at Lyryx Learning to accompany their text [Linear Algebra with Applications](#) based on W. K. Nicholson's original text.

In addition we recognize the following contributors. All new content contributed is released under the same license as noted below.

- Ilijas Farah, York University

BE A CHAMPION OF OER!

Contribute suggestions for improvements, new content, or errata:

A new topic

A new example or problem

A new or better proof to an existing theorem

Any other suggestions to improve the material

Contact Lyryx at info@lyryx.com with your ideas.

License



Attribution-NonCommercial-ShareAlike (CC BY-NC-SA)

This license lets others remix, tweak, and build upon your work non-commercially, as long as they credit you and license their new creations under the identical terms.

Copyright

Linear Transformations

Finding the Matrix of a Linear Transformation

Composition of Linear Transformations

Rotations and Reflections in \mathbb{R}^2

Copyright

Linear Transformations

Finding the Matrix of a Linear Transformation

Composition of Linear Transformations

Rotations and Reflections in \mathbb{R}^2

Linear Transformations

Definition

A transformation $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a **linear transformation** if it satisfies the following two properties for all $\vec{x}, \vec{y} \in \mathbb{R}^n$ and all (scalars) $a \in \mathbb{R}$.

Linear Transformations

Definition

A transformation $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a **linear transformation** if it satisfies the following two properties for all $\vec{x}, \vec{y} \in \mathbb{R}^n$ and all (scalars) $a \in \mathbb{R}$.

1. $T(\vec{x} + \vec{y}) = T(\vec{x}) + T(\vec{y})$ (preservation of addition)

Linear Transformations

Definition

A transformation $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a **linear transformation** if it satisfies the following two properties for all $\vec{x}, \vec{y} \in \mathbb{R}^n$ and all (scalars) $a \in \mathbb{R}$.

1. $T(\vec{x} + \vec{y}) = T(\vec{x}) + T(\vec{y})$ (preservation of addition)
2. $T(a\vec{x}) = aT(\vec{x})$ (preservation of scalar multiplication)

Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation, and let $\vec{x} \in \mathbb{R}^n$.

Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation, and let $\vec{x} \in \mathbb{R}^n$. Since T preserves scalar multiplication,

Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation, and let $\vec{x} \in \mathbb{R}^n$. Since T preserves scalar multiplication,

1. $T(0\vec{x}) = 0T(\vec{x})$ implying $T(\vec{0}) = \vec{0}$, so **T preserves the zero vector.**

Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation, and let $\vec{x} \in \mathbb{R}^n$. Since T preserves scalar multiplication,

1. $T(0\vec{x}) = 0T(\vec{x})$ implying $T(\vec{0}) = \vec{0}$, so **T preserves the zero vector.**
2. $T((-1)\vec{x}) = (-1)T(\vec{x})$, implying $T(-\vec{x}) = -T(\vec{x})$, so **T preserves the negative of a vector.**

Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation, and let $\vec{x} \in \mathbb{R}^n$. Since T preserves scalar multiplication,

1. $T(0\vec{x}) = 0T(\vec{x})$ implying $T(\vec{0}) = \vec{0}$, so **T preserves the zero vector.**
2. $T((-1)\vec{x}) = (-1)T(\vec{x})$, implying $T(-\vec{x}) = -T(\vec{x})$, so **T preserves the negative of a vector.**

Suppose $\vec{x}_1, \vec{x}_2, \dots, \vec{x}_k$ are vectors in \mathbb{R}^n and for some $a_1, a_2, \dots, a_k \in \mathbb{R}$.

$$\vec{y} = a_1\vec{x}_1 + a_2\vec{x}_2 + \cdots + a_k\vec{x}_k.$$

Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation, and let $\vec{x} \in \mathbb{R}^n$. Since T preserves scalar multiplication,

1. $T(0\vec{x}) = 0T(\vec{x})$ implying $T(\vec{0}) = \vec{0}$, so **T preserves the zero vector.**
2. $T((-1)\vec{x}) = (-1)T(\vec{x})$, implying $T(-\vec{x}) = -T(\vec{x})$, so **T preserves the negative of a vector.**

Suppose $\vec{x}_1, \vec{x}_2, \dots, \vec{x}_k$ are vectors in \mathbb{R}^n and for some $a_1, a_2, \dots, a_k \in \mathbb{R}$.

$$\vec{y} = a_1\vec{x}_1 + a_2\vec{x}_2 + \cdots + a_k\vec{x}_k.$$

\Downarrow

3.

$$\begin{aligned} T(\vec{y}) &= T(a_1\vec{x}_1 + a_2\vec{x}_2 + \cdots + a_k\vec{x}_k) \\ &= a_1T(\vec{x}_1) + a_2T(\vec{x}_2) + \cdots + a_kT(\vec{x}_k), \end{aligned}$$

i.e., **T preserves linear combinations.**

Problem

Let $T : \mathbb{R}^3 \rightarrow \mathbb{R}^4$ be a linear transformation such that

$$T \begin{bmatrix} 1 \\ 3 \\ 1 \end{bmatrix} = \begin{bmatrix} 4 \\ 4 \\ 0 \\ -2 \end{bmatrix} \quad \text{and} \quad T \begin{bmatrix} 4 \\ 0 \\ 5 \end{bmatrix} = \begin{bmatrix} 4 \\ 5 \\ -1 \\ 5 \end{bmatrix}.$$

Problem

Let $T : \mathbb{R}^3 \rightarrow \mathbb{R}^4$ be a linear transformation such that

$$T \begin{bmatrix} 1 \\ 3 \\ 1 \end{bmatrix} = \begin{bmatrix} 4 \\ 4 \\ 0 \\ -2 \end{bmatrix} \quad \text{and} \quad T \begin{bmatrix} 4 \\ 0 \\ 5 \end{bmatrix} = \begin{bmatrix} 4 \\ 5 \\ -1 \\ 5 \end{bmatrix}. \text{ Find } T \begin{bmatrix} -7 \\ 3 \\ -9 \end{bmatrix}.$$

Problem

Let $T : \mathbb{R}^3 \rightarrow \mathbb{R}^4$ be a linear transformation such that

$$T \begin{bmatrix} 1 \\ 3 \\ 1 \end{bmatrix} = \begin{bmatrix} 4 \\ 4 \\ 0 \\ -2 \end{bmatrix} \quad \text{and} \quad T \begin{bmatrix} 4 \\ 0 \\ 5 \end{bmatrix} = \begin{bmatrix} 4 \\ 5 \\ -1 \\ 5 \end{bmatrix}. \text{ Find } T \begin{bmatrix} -7 \\ 3 \\ -9 \end{bmatrix}.$$

Solution

The only way it is possible to solve this problem is if

$$\begin{bmatrix} -7 \\ 3 \\ -9 \end{bmatrix} \text{ is a linear combination of } \begin{bmatrix} 1 \\ 3 \\ 1 \end{bmatrix} \quad \text{and} \quad \begin{bmatrix} 4 \\ 0 \\ 5 \end{bmatrix},$$

Problem

Let $T : \mathbb{R}^3 \rightarrow \mathbb{R}^4$ be a linear transformation such that

$$T \begin{bmatrix} 1 \\ 3 \\ 1 \end{bmatrix} = \begin{bmatrix} 4 \\ 4 \\ 0 \\ -2 \end{bmatrix} \quad \text{and} \quad T \begin{bmatrix} 4 \\ 0 \\ 5 \end{bmatrix} = \begin{bmatrix} 4 \\ 5 \\ -1 \\ 5 \end{bmatrix}. \text{ Find } T \begin{bmatrix} -7 \\ 3 \\ -9 \end{bmatrix}.$$

Solution

The only way it is possible to solve this problem is if

$$\begin{bmatrix} -7 \\ 3 \\ -9 \end{bmatrix} \text{ is a linear combination of } \begin{bmatrix} 1 \\ 3 \\ 1 \end{bmatrix} \quad \text{and} \quad \begin{bmatrix} 4 \\ 0 \\ 5 \end{bmatrix},$$

i.e., if there exist $a, b \in \mathbb{R}$ so that

$$\begin{bmatrix} -7 \\ 3 \\ -9 \end{bmatrix} = a \begin{bmatrix} 1 \\ 3 \\ 1 \end{bmatrix} + b \begin{bmatrix} 4 \\ 0 \\ 5 \end{bmatrix}.$$

Solution (continued)

To find a and b, solve the system of three equations in two variables:

$$\left[\begin{array}{cc|c} 1 & 4 & -7 \\ 3 & 0 & 3 \\ 1 & 5 & -9 \end{array} \right] \rightarrow \cdots \rightarrow \left[\begin{array}{cc|c} 1 & 0 & 1 \\ 0 & 1 & -2 \\ 0 & 0 & 0 \end{array} \right]$$

Thus $a = 1$, $b = -2$, and

$$\left[\begin{array}{c} -7 \\ 3 \\ -9 \end{array} \right] = \left[\begin{array}{c} 1 \\ 3 \\ 1 \end{array} \right] - 2 \left[\begin{array}{c} 4 \\ 0 \\ 5 \end{array} \right].$$

Solution (continued)

We now use that fact that linear transformations preserve linear combinations, implying that

$$T \begin{bmatrix} -7 \\ 3 \\ -9 \end{bmatrix} = T \left(\begin{bmatrix} 1 \\ 3 \\ 1 \end{bmatrix} - 2 \begin{bmatrix} 4 \\ 0 \\ 5 \end{bmatrix} \right)$$

Solution (continued)

We now use that fact that linear transformations preserve linear combinations, implying that

$$\begin{aligned} T \begin{bmatrix} -7 \\ 3 \\ -9 \end{bmatrix} &= T \left(\begin{bmatrix} 1 \\ 3 \\ 1 \end{bmatrix} - 2 \begin{bmatrix} 4 \\ 0 \\ 5 \end{bmatrix} \right) \\ &= T \begin{bmatrix} 1 \\ 3 \\ 1 \end{bmatrix} - 2T \begin{bmatrix} 4 \\ 0 \\ 5 \end{bmatrix} \end{aligned}$$

Solution (continued)

We now use that fact that linear transformations preserve linear combinations, implying that

$$\begin{aligned} T \begin{bmatrix} -7 \\ 3 \\ -9 \end{bmatrix} &= T \left(\begin{bmatrix} 1 \\ 3 \\ 1 \end{bmatrix} - 2 \begin{bmatrix} 4 \\ 0 \\ 5 \end{bmatrix} \right) \\ &= T \begin{bmatrix} 1 \\ 3 \\ 1 \end{bmatrix} - 2T \begin{bmatrix} 4 \\ 0 \\ 5 \end{bmatrix} \\ &= \begin{bmatrix} 4 \\ 4 \\ 0 \\ -2 \end{bmatrix} - 2 \begin{bmatrix} 4 \\ 5 \\ -1 \\ 5 \end{bmatrix} = \begin{bmatrix} -4 \\ -6 \\ 2 \\ -12 \end{bmatrix} \end{aligned}$$

Solution (continued)

We now use that fact that linear transformations preserve linear combinations, implying that

$$\begin{aligned} T \begin{bmatrix} -7 \\ 3 \\ -9 \end{bmatrix} &= T \left(\begin{bmatrix} 1 \\ 3 \\ 1 \end{bmatrix} - 2 \begin{bmatrix} 4 \\ 0 \\ 5 \end{bmatrix} \right) \\ &= T \begin{bmatrix} 1 \\ 3 \\ 1 \end{bmatrix} - 2T \begin{bmatrix} 4 \\ 0 \\ 5 \end{bmatrix} \\ &= \begin{bmatrix} 4 \\ 4 \\ 0 \\ -2 \end{bmatrix} - 2 \begin{bmatrix} 4 \\ 5 \\ -1 \\ 5 \end{bmatrix} = \begin{bmatrix} -4 \\ -6 \\ 2 \\ -12 \end{bmatrix} \end{aligned}$$

Therefore, $T \begin{bmatrix} -7 \\ 3 \\ -9 \end{bmatrix} = \begin{bmatrix} -4 \\ -6 \\ 2 \\ -12 \end{bmatrix}.$



Problem

Let $T : \mathbb{R}^4 \rightarrow \mathbb{R}^3$ be a linear transformation such that

$$T \begin{bmatrix} 1 \\ 1 \\ 0 \\ -2 \end{bmatrix} = \begin{bmatrix} 2 \\ 3 \\ -1 \end{bmatrix} \quad \text{and} \quad T \begin{bmatrix} 0 \\ -1 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 5 \\ 0 \\ 1 \end{bmatrix}. \text{ Find } T \begin{bmatrix} 1 \\ 3 \\ -2 \\ -4 \end{bmatrix}.$$

Problem

Let $T : \mathbb{R}^4 \rightarrow \mathbb{R}^3$ be a linear transformation such that

$$T \begin{bmatrix} 1 \\ 1 \\ 0 \\ -2 \end{bmatrix} = \begin{bmatrix} 2 \\ 3 \\ -1 \end{bmatrix} \quad \text{and} \quad T \begin{bmatrix} 0 \\ -1 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 5 \\ 0 \\ 1 \end{bmatrix}. \text{ Find } T \begin{bmatrix} 1 \\ 3 \\ -2 \\ -4 \end{bmatrix}.$$

Solution (Final Answer)

$$T \begin{bmatrix} 1 \\ 3 \\ -2 \\ -4 \end{bmatrix} = \begin{bmatrix} -8 \\ 3 \\ -3 \end{bmatrix}.$$



Theorem

Every matrix transformation is a linear transformation.

Theorem

Every matrix transformation is a linear transformation.

Theorem

Every matrix transformation is a linear transformation.

Proof.

Suppose $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a matrix transformation induced by the $m \times n$ matrix A ,

Theorem

Every matrix transformation is a linear transformation.

Proof.

Suppose $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a matrix transformation induced by the $m \times n$ matrix A , i.e., $T(\vec{x}) = A\vec{x}$ for each $\vec{x} \in \mathbb{R}^n$.

Theorem

Every matrix transformation is a linear transformation.

Proof.

Suppose $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a matrix transformation induced by the $m \times n$ matrix A , i.e., $T(\vec{x}) = A\vec{x}$ for each $\vec{x} \in \mathbb{R}^n$. Let $\vec{x}, \vec{y} \in \mathbb{R}^n$ and let $a \in \mathbb{R}$.

Theorem

Every matrix transformation is a linear transformation.

Proof.

Suppose $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a matrix transformation induced by the $m \times n$ matrix A , i.e., $T(\vec{x}) = A\vec{x}$ for each $\vec{x} \in \mathbb{R}^n$. Let $\vec{x}, \vec{y} \in \mathbb{R}^n$ and let $a \in \mathbb{R}$. Then

$$T(\vec{x} + \vec{y}) = A(\vec{x} + \vec{y}) = A\vec{x} + A\vec{y} = T(\vec{x}) + T(\vec{y}),$$

Theorem

Every matrix transformation is a linear transformation.

Proof.

Suppose $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a matrix transformation induced by the $m \times n$ matrix A , i.e., $T(\vec{x}) = A\vec{x}$ for each $\vec{x} \in \mathbb{R}^n$. Let $\vec{x}, \vec{y} \in \mathbb{R}^n$ and let $a \in \mathbb{R}$. Then

$$T(\vec{x} + \vec{y}) = A(\vec{x} + \vec{y}) = A\vec{x} + A\vec{y} = T(\vec{x}) + T(\vec{y}),$$

proving that T preserves addition.

Theorem

Every matrix transformation is a linear transformation.

Proof.

Suppose $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a matrix transformation induced by the $m \times n$ matrix A , i.e., $T(\vec{x}) = A\vec{x}$ for each $\vec{x} \in \mathbb{R}^n$. Let $\vec{x}, \vec{y} \in \mathbb{R}^n$ and let $a \in \mathbb{R}$. Then

$$T(\vec{x} + \vec{y}) = A(\vec{x} + \vec{y}) = A\vec{x} + A\vec{y} = T(\vec{x}) + T(\vec{y}),$$

proving that T preserves addition. Also,

Theorem

Every matrix transformation is a linear transformation.

Proof.

Suppose $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a matrix transformation induced by the $m \times n$ matrix A , i.e., $T(\vec{x}) = A\vec{x}$ for each $\vec{x} \in \mathbb{R}^n$. Let $\vec{x}, \vec{y} \in \mathbb{R}^n$ and let $a \in \mathbb{R}$. Then

$$T(\vec{x} + \vec{y}) = A(\vec{x} + \vec{y}) = A\vec{x} + A\vec{y} = T(\vec{x}) + T(\vec{y}),$$

proving that T preserves addition. Also,

$$T(a\vec{x}) = A(a\vec{x}) = a(A\vec{x}) = aT(\vec{x}),$$

Theorem

Every matrix transformation is a linear transformation.

Proof.

Suppose $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a matrix transformation induced by the $m \times n$ matrix A , i.e., $T(\vec{x}) = A\vec{x}$ for each $\vec{x} \in \mathbb{R}^n$. Let $\vec{x}, \vec{y} \in \mathbb{R}^n$ and let $a \in \mathbb{R}$. Then

$$T(\vec{x} + \vec{y}) = A(\vec{x} + \vec{y}) = A\vec{x} + A\vec{y} = T(\vec{x}) + T(\vec{y}),$$

proving that T preserves addition. Also,

$$T(a\vec{x}) = A(a\vec{x}) = a(A\vec{x}) = aT(\vec{x}),$$

proving that T preserves scalar multiplication.

Theorem

Every matrix transformation is a linear transformation.

Proof.

Suppose $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a matrix transformation induced by the $m \times n$ matrix A , i.e., $T(\vec{x}) = A\vec{x}$ for each $\vec{x} \in \mathbb{R}^n$. Let $\vec{x}, \vec{y} \in \mathbb{R}^n$ and let $a \in \mathbb{R}$. Then

$$T(\vec{x} + \vec{y}) = A(\vec{x} + \vec{y}) = A\vec{x} + A\vec{y} = T(\vec{x}) + T(\vec{y}),$$

proving that T preserves addition. Also,

$$T(a\vec{x}) = A(a\vec{x}) = a(A\vec{x}) = aT(\vec{x}),$$

proving that T preserves scalar multiplication.

Since T preserves addition and scalar multiplication T is a linear transformation. ■

Example (The Zero Transformation)

If A is the $m \times n$ matrix of zeros, then the transformation $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ induced by A is called the **zero transformation** because for every vector \vec{x} in \mathbb{R}^n

$$T(\vec{x}) = A\vec{x} = \vec{0}\vec{x} = \vec{0}.$$

Example (The Zero Transformation)

If A is the $m \times n$ matrix of zeros, then the transformation $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ induced by A is called the **zero transformation** because for every vector \vec{x} in \mathbb{R}^n

$$T(\vec{x}) = A\vec{x} = \mathbf{0}\vec{x} = \vec{0}.$$

The zero transformation is usually written as **$T = \mathbf{0}$** .

Example (The Zero Transformation)

If A is the $m \times n$ matrix of zeros, then the transformation $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ induced by A is called the **zero transformation** because for every vector \vec{x} in \mathbb{R}^n

$$T(\vec{x}) = A\vec{x} = \mathbf{0}\vec{x} = \vec{0}.$$

The zero transformation is usually written as **$T = \mathbf{0}$** .

Example (The Identity Transformation)

The transformation of \mathbb{R}^n induced by I_n , the $n \times n$ identity matrix, is called the **identity transformation** because for every vector \vec{x} in \mathbb{R}^n ,

$$T(\vec{x}) = I_n\vec{x} = \vec{x}.$$

Example (The Zero Transformation)

If A is the $m \times n$ matrix of zeros, then the transformation $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ induced by A is called the **zero transformation** because for every vector \vec{x} in \mathbb{R}^n

$$T(\vec{x}) = A\vec{x} = \mathbf{0}\vec{x} = \vec{0}.$$

The zero transformation is usually written as **$T = \mathbf{0}$** .

Example (The Identity Transformation)

The transformation of \mathbb{R}^n induced by I_n , the $n \times n$ identity matrix, is called the **identity transformation** because for every vector \vec{x} in \mathbb{R}^n ,

$$T(\vec{x}) = I_n\vec{x} = \vec{x}.$$

The identity transformation on \mathbb{R}^n is usually written as **$\mathbf{1}_{\mathbb{R}^n}$** .

Problem (Revisited)

Is the following $T : \mathbb{R}^3 \rightarrow \mathbb{R}^4$ a matrix transformation?

$$T \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} a + b \\ b + c \\ a - c \\ c - b \end{bmatrix}$$

Problem (Revisited)

Is the following $T : \mathbb{R}^3 \rightarrow \mathbb{R}^4$ a matrix transformation?

$$T \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} a + b \\ b + c \\ a - c \\ c - b \end{bmatrix}$$

Solution

$$A \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 1 & 0 & -1 \\ 0 & -1 & 1 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} a + b \\ b + c \\ a - c \\ c - b \end{bmatrix} = T \begin{bmatrix} a \\ b \\ c \end{bmatrix}$$


Problem (Revisited)

Is the following $T : \mathbb{R}^3 \rightarrow \mathbb{R}^4$ a matrix transformation?

$$T \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} a + b \\ b + c \\ a - c \\ c - b \end{bmatrix}$$

Solution

$$A \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 1 & 0 & -1 \\ 0 & -1 & 1 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} a + b \\ b + c \\ a - c \\ c - b \end{bmatrix} = T \begin{bmatrix} a \\ b \\ c \end{bmatrix}$$

Yes, T is a matrix transformation. 

Problem (Not all transformations are matrix transformations)

Consider $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by

$$T(\vec{x}) = \vec{x} + \begin{bmatrix} 1 \\ -1 \end{bmatrix} \text{ for all } \vec{x} \in \mathbb{R}^2.$$

Show that T NOT a matrix transformation.

Solution

We have $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by

$$T(\vec{x}) = \vec{x} + \begin{bmatrix} 1 \\ -1 \end{bmatrix} \text{ for all } \vec{x} \in \mathbb{R}^2.$$

Solution

We have $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by

$$T(\vec{x}) = \vec{x} + \begin{bmatrix} 1 \\ -1 \end{bmatrix} \text{ for all } \vec{x} \in \mathbb{R}^2.$$

Since every matrix transformation is a linear transformation,

Solution

We have $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by

$$T(\vec{x}) = \vec{x} + \begin{bmatrix} 1 \\ -1 \end{bmatrix} \text{ for all } \vec{x} \in \mathbb{R}^2.$$

Since every matrix transformation is a linear transformation, we consider $T(0)$, where 0 is the zero vector of \mathbb{R}^2 .

Solution

We have $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by

$$T(\vec{x}) = \vec{x} + \begin{bmatrix} 1 \\ -1 \end{bmatrix} \text{ for all } \vec{x} \in \mathbb{R}^2.$$

Since every matrix transformation is a linear transformation, we consider $T(0)$, where 0 is the zero vector of \mathbb{R}^2 .

$$T \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Solution

We have $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by

$$T(\vec{x}) = \vec{x} + \begin{bmatrix} 1 \\ -1 \end{bmatrix} \text{ for all } \vec{x} \in \mathbb{R}^2.$$

Since every matrix transformation is a linear transformation, we consider $T(0)$, where 0 is the zero vector of \mathbb{R}^2 .

$$T \begin{bmatrix} 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} + \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

Solution

We have $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by

$$T(\vec{x}) = \vec{x} + \begin{bmatrix} 1 \\ -1 \end{bmatrix} \text{ for all } \vec{x} \in \mathbb{R}^2.$$

Since every matrix transformation is a linear transformation, we consider $T(0)$, where 0 is the zero vector of \mathbb{R}^2 .

$$T \begin{bmatrix} 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} + \begin{bmatrix} 1 \\ -1 \end{bmatrix} = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

Solution

We have $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by

$$T(\vec{x}) = \vec{x} + \begin{bmatrix} 1 \\ -1 \end{bmatrix} \text{ for all } \vec{x} \in \mathbb{R}^2.$$

Since every matrix transformation is a linear transformation, we consider $T(0)$, where 0 is the zero vector of \mathbb{R}^2 .

$$T \begin{bmatrix} 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} + \begin{bmatrix} 1 \\ -1 \end{bmatrix} = \begin{bmatrix} 1 \\ -1 \end{bmatrix} \neq \begin{bmatrix} 0 \\ 0 \end{bmatrix},$$

violating one of the properties of a linear transformation.

Solution


We have $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by

$$T(\vec{x}) = \vec{x} + \begin{bmatrix} 1 \\ -1 \end{bmatrix} \text{ for all } \vec{x} \in \mathbb{R}^2.$$

Since every matrix transformation is a linear transformation, we consider $T(0)$, where 0 is the zero vector of \mathbb{R}^2 .

$$T \begin{bmatrix} 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} + \begin{bmatrix} 1 \\ -1 \end{bmatrix} = \begin{bmatrix} 1 \\ -1 \end{bmatrix} \neq \begin{bmatrix} 0 \\ 0 \end{bmatrix},$$

violating one of the properties of a linear transformation.

Therefore, T is not a linear transformation, and hence is not a matrix transformation. 

Remark

Recall that a transformation $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a **linear transformation** if it satisfies the following two properties for all $\vec{x}, \vec{y} \in \mathbb{R}^n$ and all (scalars) $a \in \mathbb{R}$.

Remark

Recall that a transformation $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a **linear transformation** if it satisfies the following two properties for all $\vec{x}, \vec{y} \in \mathbb{R}^n$ and all (scalars) $a \in \mathbb{R}$.

1. $T(\vec{x} + \vec{y}) = T(\vec{x}) + T(\vec{y})$ (preservation of addition)

Remark

Recall that a transformation $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a **linear transformation** if it satisfies the following two properties for all $\vec{x}, \vec{y} \in \mathbb{R}^n$ and all (scalars) $a \in \mathbb{R}$.

1. $T(\vec{x} + \vec{y}) = T(\vec{x}) + T(\vec{y})$ (preservation of addition)
2. $T(a\vec{x}) = aT(\vec{x})$ (preservation of scalar multiplication)

Remark

Recall that a transformation $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a **linear transformation** if it satisfies the following two properties for all $\vec{x}, \vec{y} \in \mathbb{R}^n$ and all (scalars) $a \in \mathbb{R}$.

1. $T(\vec{x} + \vec{y}) = T(\vec{x}) + T(\vec{y})$ (preservation of addition)
2. $T(a\vec{x}) = aT(\vec{x})$ (preservation of scalar multiplication)

Theorem (Every Linear Transformation is a Matrix Transformation)

Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation. Then we can find an $n \times m$ matrix A such that

$$T(\vec{x}) = A\vec{x}$$

Remark

Recall that a transformation $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a **linear transformation** if it satisfies the following two properties for all $\vec{x}, \vec{y} \in \mathbb{R}^n$ and all (scalars) $a \in \mathbb{R}$.

1. $T(\vec{x} + \vec{y}) = T(\vec{x}) + T(\vec{y})$ (preservation of addition)
2. $T(a\vec{x}) = aT(\vec{x})$ (preservation of scalar multiplication)

Theorem (Every Linear Transformation is a Matrix Transformation)

Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation. Then we can find an $n \times m$ matrix A such that

$$T(\vec{x}) = A\vec{x}$$

In this case, we say that T is induced, or determined, by A and we write

$$T_A(\vec{x}) = A\vec{x}$$

“linear” = “matrix”

Problem

The transformation $T : \mathbb{R}^3 \rightarrow \mathbb{R}^4$ defined by

$$T \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} a + b \\ b + c \\ a - c \\ c - b \end{bmatrix}$$

for each $\vec{x} \in \mathbb{R}^3$ is another matrix transformation, that is, $T(\vec{x}) = A\vec{x}$ for some matrix A . **Can you find a matrix A that works?**

Problem

The transformation $T : \mathbb{R}^3 \rightarrow \mathbb{R}^4$ defined by

$$T \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} a + b \\ b + c \\ a - c \\ c - b \end{bmatrix}$$

for each $\vec{x} \in \mathbb{R}^3$ is another matrix transformation, that is, $T(\vec{x}) = A\vec{x}$ for some matrix A . **Can you find a matrix A that works?**

Solution

First, since $T : \mathbb{R}^3 \rightarrow \mathbb{R}^4$, we know that A must have size

Solution

First, since $T : \mathbb{R}^3 \rightarrow \mathbb{R}^4$, we know that A must have size 4×3 .

Solution

First, since $T : \mathbb{R}^3 \rightarrow \mathbb{R}^4$, we know that A must have size 4×3 . Now consider the product

$$\begin{bmatrix} ? & ? & ? \\ ? & ? & ? \\ ? & ? & ? \\ ? & ? & ? \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} a + b \\ b + c \\ a - c \\ c - b \end{bmatrix},$$

and try to fill in the values of the matrix.

Solution

First, since $T : \mathbb{R}^3 \rightarrow \mathbb{R}^4$, we know that A must have size 4×3 . Now consider the product

$$\begin{bmatrix} ? & ? & ? \\ ? & ? & ? \\ ? & ? & ? \\ ? & ? & ? \end{bmatrix} \begin{bmatrix} a \\ b \\ c \end{bmatrix} = \begin{bmatrix} a + b \\ b + c \\ a - c \\ c - b \end{bmatrix},$$

and try to fill in the values of the matrix.

\vdots

We can deduce from the product that T is induced by the matrix

$$A = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 1 & 0 & -1 \\ 0 & -1 & 1 \end{bmatrix}.$$



Copyright

Linear Transformations

Finding the Matrix of a Linear Transformation

Composition of Linear Transformations

Rotations and Reflections in \mathbb{R}^2

Finding the Matrix of a Linear Transformation

Finding the Matrix of a Linear Transformation

Is there an easier way to find the matrix of T ?

Finding the Matrix of a Linear Transformation

Is there an easier way to find the matrix of T ? For some transformations guess and check will work, but this is not an efficient method. The next theorem gives a method for finding the matrix of T .

Finding the Matrix of a Linear Transformation

Is there an easier way to find the matrix of T ? For some transformations guess and check will work, but this is not an efficient method. The next theorem gives a method for finding the matrix of T .

Definition

The set of columns $\{\vec{e}_1, \vec{e}_2, \dots, \vec{e}_n\}$ of I_n is called the **standard basis of \mathbb{R}^n** .

Theorem (Matrix of a Linear Transformation)

Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation.

Theorem (Matrix of a Linear Transformation)

Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation. Then T is a matrix transformation.

Theorem (Matrix of a Linear Transformation)

Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation. Then T is a matrix transformation. Furthermore, T is induced by the **unique** matrix

$$A = \begin{bmatrix} T(\vec{e}_1) & T(\vec{e}_2) & \cdots & T(\vec{e}_n) \end{bmatrix},$$

where \vec{e}_j is the j -th column of I_n , and $T(\vec{e}_j)$ is the j -th column of A .

Theorem (Matrix of a Linear Transformation)

Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear transformation. Then T is a matrix transformation. Furthermore, T is induced by the **unique** matrix

$$A = \begin{bmatrix} T(\vec{e}_1) & T(\vec{e}_2) & \cdots & T(\vec{e}_n) \end{bmatrix},$$

where \vec{e}_j is the j -th column of I_n , and $T(\vec{e}_j)$ is the j -th column of A .

Corollary

A transformation $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a linear transformation if and only if it is a matrix transformation.

“linear” = “matrix”

Problem

Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a linear transformation defined by

$$T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x + 2y \\ x - y \end{bmatrix}$$

for each $\vec{x} \in \mathbb{R}^2$. Find the matrix, A , of T .

Problem

Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a linear transformation defined by

$$T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x + 2y \\ x - y \end{bmatrix}$$

for each $\vec{x} \in \mathbb{R}^2$. Find the matrix, A , of T .

Solution

$$T \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

Problem

Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a linear transformation defined by

$$T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x + 2y \\ x - y \end{bmatrix}$$

for each $\vec{x} \in \mathbb{R}^2$. Find the matrix, A , of T .

Solution

$$T \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 1 + 2(0) \\ 1 - 0 \end{bmatrix}$$

Problem

Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a linear transformation defined by

$$T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x + 2y \\ x - y \end{bmatrix}$$

for each $\vec{x} \in \mathbb{R}^2$. Find the matrix, A , of T .

Solution

$$T \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 1 + 2(0) \\ 1 - 0 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

Problem

Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a linear transformation defined by

$$T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x + 2y \\ x - y \end{bmatrix}$$

for each $\vec{x} \in \mathbb{R}^2$. Find the matrix, A , of T .

Solution

$$T \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 1 + 2(0) \\ 1 - 0 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \end{bmatrix} \quad \text{and} \quad T \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

Problem

Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a linear transformation defined by

$$T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x + 2y \\ x - y \end{bmatrix}$$

for each $\vec{x} \in \mathbb{R}^2$. Find the matrix, A , of T .

Solution

$$T \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 1 + 2(0) \\ 1 - 0 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \end{bmatrix} \quad \text{and} \quad T \begin{bmatrix} 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 + 2(1) \\ 0 - 1 \end{bmatrix}$$

Problem

Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a linear transformation defined by

$$T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x + 2y \\ x - y \end{bmatrix}$$

for each $\vec{x} \in \mathbb{R}^2$. Find the matrix, A , of T .

Solution

$$T \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 1 + 2(0) \\ 1 - 0 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \end{bmatrix} \quad \text{and} \quad T \begin{bmatrix} 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 + 2(1) \\ 0 - 1 \end{bmatrix} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$$

Problem

Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a linear transformation defined by

$$T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x + 2y \\ x - y \end{bmatrix}$$

for each $\vec{x} \in \mathbb{R}^2$. Find the matrix, A , of T .

Solution

$$T \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 1 + 2(0) \\ 1 - 0 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \end{bmatrix} \quad \text{and} \quad T \begin{bmatrix} 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 + 2(1) \\ 0 - 1 \end{bmatrix} = \begin{bmatrix} 2 \\ -1 \end{bmatrix}$$

\Downarrow

$$A = \begin{bmatrix} 1 & 2 \\ 1 & -1 \end{bmatrix}$$



Sometimes, T is defined through its actions several concrete vectors.

Problem

Find the matrix A of T where T is given as

$$T \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \end{bmatrix} \quad \text{and} \quad T \begin{bmatrix} 0 \\ -1 \end{bmatrix} = \begin{bmatrix} 3 \\ 2 \end{bmatrix}.$$

Solution (continued)

We need to write \vec{e}_1 and \vec{e}_2 as a linear combination of the vectors provided. First, find x and y such that

$$\begin{bmatrix} 1 \\ 0 \end{bmatrix} = x \begin{bmatrix} 1 \\ 1 \end{bmatrix} + y \begin{bmatrix} 0 \\ -1 \end{bmatrix}$$

Solution (continued)

We need to write \vec{e}_1 and \vec{e}_2 as a linear combination of the vectors provided. First, find x and y such that

$$\begin{bmatrix} 1 \\ 0 \end{bmatrix} = x \begin{bmatrix} 1 \\ 1 \end{bmatrix} + y \begin{bmatrix} 0 \\ -1 \end{bmatrix}$$

Once we find x and y we can compute

$$\begin{aligned} T \begin{bmatrix} 1 \\ 0 \end{bmatrix} &= xT \begin{bmatrix} 1 \\ 1 \end{bmatrix} + yT \begin{bmatrix} 0 \\ -1 \end{bmatrix} \\ &= x \begin{bmatrix} 1 \\ 2 \end{bmatrix} + y \begin{bmatrix} 3 \\ 2 \end{bmatrix} \end{aligned}$$

Solution (continued)

Finding x and y involves solving the following system of equations.

$$x = 1$$

$$x - y = 0$$

Solution (continued)

Finding x and y involves solving the following system of equations.

$$\begin{aligned}x &= 1 \\ x - y &= 0\end{aligned}$$

The solution is $x = 1, y = 1$.

Solution (continued)

Finding x and y involves solving the following system of equations.

$$\begin{aligned}x &= 1 \\ x - y &= 0\end{aligned}$$

The solution is $x = 1, y = 1$. Hence, we can find $T(\vec{e}_1)$ as follows.

$$T \begin{bmatrix} 1 \\ 0 \end{bmatrix} = 1 \begin{bmatrix} 1 \\ 2 \end{bmatrix} + 1 \begin{bmatrix} 3 \\ 2 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \end{bmatrix} + \begin{bmatrix} 3 \\ 2 \end{bmatrix} = \begin{bmatrix} 4 \\ 4 \end{bmatrix}.$$

As for $T(\vec{e}_2)$,

$$T \begin{bmatrix} 0 \\ 1 \end{bmatrix} = -T \begin{bmatrix} 0 \\ -1 \end{bmatrix} = \begin{bmatrix} -3 \\ -2 \end{bmatrix}.$$

\Downarrow

$$A = \begin{bmatrix} 4 & -3 \\ 4 & -2 \end{bmatrix}$$



Problem

Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^3$ be a transformation defined by $T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 2x \\ y \\ -x + 2y \end{bmatrix}$.

Is T a linear transformation?

Problem

Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^3$ be a transformation defined by $T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 2x \\ y \\ -x + 2y \end{bmatrix}$.

Is T a linear transformation?

Solution

If T were a linear transformation, then T would be induced by the matrix

Problem

Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^3$ be a transformation defined by $T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 2x \\ y \\ -x + 2y \end{bmatrix}$.

Is T a linear transformation?

Solution

If T were a linear transformation, then T would be induced by the matrix

$$A = \begin{bmatrix} T(\vec{e}_1) & T(\vec{e}_2) \end{bmatrix}$$

Problem

Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^3$ be a transformation defined by $T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 2x \\ y \\ -x + 2y \end{bmatrix}$.

Is T a linear transformation?

Solution

If T were a linear transformation, then T would be induced by the matrix

$$A = \begin{bmatrix} T(\vec{e}_1) & T(\vec{e}_2) \end{bmatrix} = \begin{bmatrix} T \begin{bmatrix} 1 \\ 0 \end{bmatrix} & T \begin{bmatrix} 0 \\ 1 \end{bmatrix} \end{bmatrix}$$

Problem

Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^3$ be a transformation defined by $T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 2x \\ y \\ -x + 2y \end{bmatrix}$.

Is T a linear transformation?

Solution

If T were a linear transformation, then T would be induced by the matrix

$$A = [T(\vec{e}_1) \quad T(\vec{e}_2)] = \left[T \begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad T \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right] = \begin{bmatrix} 2 & 0 \\ 0 & 1 \\ -1 & 2 \end{bmatrix}.$$

Problem

Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^3$ be a transformation defined by $T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 2x \\ y \\ -x + 2y \end{bmatrix}$.

Is T a linear transformation?

Solution

If T were a linear transformation, then T would be induced by the matrix

$$A = [T(\vec{e}_1) \quad T(\vec{e}_2)] = \left[T \begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad T \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right] = \begin{bmatrix} 2 & 0 \\ 0 & 1 \\ -1 & 2 \end{bmatrix}.$$

It remains to verify the matrix transform induced by A indeed coincides with T :

$$A \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 2 & 0 \\ 0 & 1 \\ -1 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 2x \\ y \\ -x + 2y \end{bmatrix} = T \begin{bmatrix} x \\ y \end{bmatrix}.$$

Therefore, T is a matrix transformation induced by A above. ■

Problem

Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a transformation defined by $T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} xy \\ x + y \end{bmatrix}$. Is T a linear transformation?

Problem

Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a transformation defined by $T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} xy \\ x + y \end{bmatrix}$. Is T a linear transformation?

Solution

If T were a linear transformation, then T would be induced by the matrix

Problem

Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a transformation defined by $T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} xy \\ x + y \end{bmatrix}$. Is T a linear transformation?

Solution

If T were a linear transformation, then T would be induced by the matrix

$$A = \begin{bmatrix} T(\vec{e}_1) & T(\vec{e}_2) \end{bmatrix}$$

Problem

Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a transformation defined by $T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} xy \\ x + y \end{bmatrix}$. Is T a linear transformation?

Solution

If T were a linear transformation, then T would be induced by the matrix

$$A = \begin{bmatrix} T(\vec{e}_1) & T(\vec{e}_2) \end{bmatrix} = \begin{bmatrix} T \begin{bmatrix} 1 \\ 0 \end{bmatrix} & T \begin{bmatrix} 0 \\ 1 \end{bmatrix} \end{bmatrix}$$

Problem

Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a transformation defined by $T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} xy \\ x + y \end{bmatrix}$. Is T a linear transformation?

Solution

If T were a linear transformation, then T would be induced by the matrix

$$A = [T(\vec{e}_1) \quad T(\vec{e}_2)] = \left[T \begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad T \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right] = \begin{bmatrix} 0 & 0 \\ 1 & 1 \end{bmatrix}.$$

Problem

Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a transformation defined by $T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} xy \\ x + y \end{bmatrix}$. Is T a linear transformation?

Solution

If T were a linear transformation, then T would be induced by the matrix

$$A = [T(\vec{e}_1) \quad T(\vec{e}_2)] = \left[T \begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad T \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right] = \begin{bmatrix} 0 & 0 \\ 1 & 1 \end{bmatrix}.$$

However, the matrix transform induced by A doesn't pass the verification:

$$A \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ x + y \end{bmatrix} \neq \begin{bmatrix} xy \\ x + y \end{bmatrix} = T \begin{bmatrix} x \\ y \end{bmatrix}$$

Problem

Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a transformation defined by $T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} xy \\ x + y \end{bmatrix}$. Is T a linear transformation?


Solution

If T were a linear transformation, then T would be induced by the matrix

$$A = [T(\vec{e}_1) \quad T(\vec{e}_2)] = \left[T \begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad T \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right] = \begin{bmatrix} 0 & 0 \\ 1 & 1 \end{bmatrix}.$$

However, the matrix transform induced by A doesn't pass the verification:

$$A \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 \\ x + y \end{bmatrix} \neq \begin{bmatrix} xy \\ x + y \end{bmatrix} = T \begin{bmatrix} x \\ y \end{bmatrix}$$

Therefore, T is NOT a linear transformation. 

Copyright

Linear Transformations

Finding the Matrix of a Linear Transformation

Composition of Linear Transformations

Rotations and Reflections in \mathbb{R}^2

Composition of Linear Transformations

Composition of Linear Transformations

Definition

Suppose $T : \mathbb{R}^k \rightarrow \mathbb{R}^n$ and $S : \mathbb{R}^n \rightarrow \mathbb{R}^m$ are linear transformations.

Composition of Linear Transformations

Definition

Suppose $T : \mathbb{R}^k \rightarrow \mathbb{R}^n$ and $S : \mathbb{R}^n \rightarrow \mathbb{R}^m$ are linear transformations. The **composite** (or composition) of S and T is

$$S \circ T : \mathbb{R}^k \rightarrow \mathbb{R}^m,$$

Composition of Linear Transformations

Definition

Suppose $T : \mathbb{R}^k \rightarrow \mathbb{R}^n$ and $S : \mathbb{R}^n \rightarrow \mathbb{R}^m$ are linear transformations. The **composite** (or composition) of S and T is

$$S \circ T : \mathbb{R}^k \rightarrow \mathbb{R}^m,$$

is defined by

$$(S \circ T)(\vec{x}) = S(T(\vec{x})) \text{ for all } \vec{x} \in \mathbb{R}^k.$$

Composition of Linear Transformations


Definition

Suppose $T : \mathbb{R}^k \rightarrow \mathbb{R}^n$ and $S : \mathbb{R}^n \rightarrow \mathbb{R}^m$ are linear transformations. The **composite** (or composition) of S and T is

$$S \circ T : \mathbb{R}^k \rightarrow \mathbb{R}^m,$$

is defined by

$$(S \circ T)(\vec{x}) = S(T(\vec{x})) \text{ for all } \vec{x} \in \mathbb{R}^k.$$



Composition of Linear Transformations


Definition

Suppose $T : \mathbb{R}^k \rightarrow \mathbb{R}^n$ and $S : \mathbb{R}^n \rightarrow \mathbb{R}^m$ are linear transformations. The **composite** (or composition) of S and T is

$$S \circ T : \mathbb{R}^k \rightarrow \mathbb{R}^m,$$

is defined by

$$(S \circ T)(\vec{x}) = S(T(\vec{x})) \text{ for all } \vec{x} \in \mathbb{R}^k.$$



Remark (Convention on the order)

$S \circ T$ means that the transformation T is applied first, followed by the transformation S .

Theorem

Let $\mathbb{R}^k \xrightarrow{T} \mathbb{R}^n \xrightarrow{S} \mathbb{R}^m$ be linear transformations, and suppose that S is induced by matrix A , and T is induced by matrix B . Then $S \circ T$ is a linear transformation, and $S \circ T$ is induced by the matrix AB .

Theorem

Let $\mathbb{R}^k \xrightarrow{T} \mathbb{R}^n \xrightarrow{S} \mathbb{R}^m$ be linear transformations, and suppose that S is induced by matrix A , and T is induced by matrix B . Then $S \circ T$ is a linear transformation, and $S \circ T$ is induced by the matrix AB .

Problem

Let $S : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ and $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be linear transformations defined by

$$S \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} x \\ -y \end{bmatrix} \quad \text{and} \quad T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} -y \\ x \end{bmatrix} \quad \text{for all} \quad \begin{bmatrix} x \\ y \end{bmatrix} \in \mathbb{R}^2.$$

Find $S \circ T$.

Solution

Then S and T are induced by matrices

$$A = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \quad \text{and} \quad B = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix},$$

respectively.

Solution

Then S and T are induced by matrices

$$A = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \quad \text{and} \quad B = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix},$$

respectively. The composite of S and T is the transformation
 $S \circ T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$

Solution

Then S and T are induced by matrices

$$A = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \quad \text{and} \quad B = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix},$$

respectively. The composite of S and T is the transformation $S \circ T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by

$$(S \circ T) \begin{bmatrix} x \\ y \end{bmatrix} = S \left(T \begin{bmatrix} x \\ y \end{bmatrix} \right),$$

Solution

Then S and T are induced by matrices

$$A = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \quad \text{and} \quad B = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix},$$

respectively. The composite of S and T is the transformation $S \circ T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by

$$(S \circ T) \begin{bmatrix} x \\ y \end{bmatrix} = S \left(T \begin{bmatrix} x \\ y \end{bmatrix} \right),$$

and has matrix (or is induced by the matrix)

$$AB = \begin{bmatrix} 0 & -1 \\ -1 & 0 \end{bmatrix}.$$

Example (continued)

Therefore the composite of S and T is the linear transformation

$$(S \circ T) \begin{bmatrix} x \\ y \end{bmatrix} = AB \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 & -1 \\ -1 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} -y \\ -x \end{bmatrix},$$

for all $\begin{bmatrix} x \\ y \end{bmatrix} \in \mathbb{R}^2$.



Example (continued)

Therefore the composite of S and T is the linear transformation

$$(S \circ T) \begin{bmatrix} x \\ y \end{bmatrix} = AB \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 0 & -1 \\ -1 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} -y \\ -x \end{bmatrix},$$

for all $\begin{bmatrix} x \\ y \end{bmatrix} \in \mathbb{R}^2$.



Remark

Compare this with the composite of T and S which is the linear transformation

$$(T \circ S) \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} y \\ x \end{bmatrix},$$

for all $\begin{bmatrix} x \\ y \end{bmatrix} \in \mathbb{R}^2$.

Copyright

Linear Transformations

Finding the Matrix of a Linear Transformation

Composition of Linear Transformations

Rotations and Reflections in \mathbb{R}^2

Rotations in \mathbb{R}^2

The rest part is an application of the linear transform to the study of the rotations in \mathbb{R}^2 . This is left your motivated students to study by themselves.

Rotations in \mathbb{R}^2

The rest part is an application of the linear transform to the study of the rotations in \mathbb{R}^2 . This is left your motivated students to study by themselves.

Definition

The transformation

$$R_\theta : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

denotes counterclockwise rotation about the origin through an angle of θ .

Rotations in \mathbb{R}^2

The rest part is an application of the linear transform to the study of the rotations in \mathbb{R}^2 . This is left your motivated students to study by themselves.

Definition

The transformation

$$R_\theta : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

denotes counterclockwise rotation about the origin through an angle of θ .

Rotation through an angle of θ preserves scalar multiplication.

Rotations in \mathbb{R}^2

The rest part is an application of the linear transform to the study of the rotations in \mathbb{R}^2 . This is left your motivated students to study by themselves.

Definition

The transformation

$$R_\theta : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

denotes counterclockwise rotation about the origin through an angle of θ .

Rotation through an angle of θ preserves scalar multiplication.

Rotation through an angle of θ preserves vector addition.

R_θ is a linear transformation

Since R_θ preserves addition and scalar multiplication, R_θ is a linear transformation, and hence a matrix transformation.

The matrix that induces R_θ can be found by computing $R_\theta(\vec{e}_1)$ and $R_\theta(\vec{e}_2)$, where

$$\vec{e}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad \text{and} \quad \vec{e}_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}.$$

R_θ is a linear transformation

Since R_θ preserves addition and scalar multiplication, R_θ is a linear transformation, and hence a matrix transformation.

The matrix that induces R_θ can be found by computing $R_\theta(\vec{e}_1)$ and $R_\theta(\vec{e}_2)$, where

$$\vec{e}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad \text{and} \quad \vec{e}_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}.$$

$$R_\theta(\vec{e}_1)$$

R_θ is a linear transformation

Since R_θ preserves addition and scalar multiplication, R_θ is a linear transformation, and hence a matrix transformation.

The matrix that induces R_θ can be found by computing $R_\theta(\vec{e}_1)$ and $R_\theta(\vec{e}_2)$, where

$$\vec{e}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad \text{and} \quad \vec{e}_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}.$$

$$R_\theta(\vec{e}_1) = R_\theta \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

R_θ is a linear transformation

Since R_θ preserves addition and scalar multiplication, R_θ is a linear transformation, and hence a matrix transformation.

The matrix that induces R_θ can be found by computing $R_\theta(\vec{e}_1)$ and $R_\theta(\vec{e}_2)$, where

$$\vec{e}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad \text{and} \quad \vec{e}_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}.$$

$$R_\theta(\vec{e}_1) = R_\theta \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \begin{bmatrix} \cos \theta \\ \sin \theta \end{bmatrix},$$

R_θ is a linear transformation

Since R_θ preserves addition and scalar multiplication, R_θ is a linear transformation, and hence a matrix transformation.

The matrix that induces R_θ can be found by computing $R_\theta(\vec{e}_1)$ and $R_\theta(\vec{e}_2)$, where

$$\vec{e}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad \text{and} \quad \vec{e}_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}.$$

$$R_\theta(\vec{e}_1) = R_\theta \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \begin{bmatrix} \cos \theta \\ \sin \theta \end{bmatrix},$$

and

$$R_\theta(\vec{e}_2)$$

R_θ is a linear transformation

Since R_θ preserves addition and scalar multiplication, R_θ is a linear transformation, and hence a matrix transformation.

The matrix that induces R_θ can be found by computing $R_\theta(\vec{e}_1)$ and $R_\theta(\vec{e}_2)$, where

$$\vec{e}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad \text{and} \quad \vec{e}_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}.$$

$$R_\theta(\vec{e}_1) = R_\theta \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \begin{bmatrix} \cos \theta \\ \sin \theta \end{bmatrix},$$

and

$$R_\theta(\vec{e}_2) = R_\theta \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

R_θ is a linear transformation

Since R_θ preserves addition and scalar multiplication, R_θ is a linear transformation, and hence a matrix transformation.

The matrix that induces R_θ can be found by computing $R_\theta(\vec{e}_1)$ and $R_\theta(\vec{e}_2)$, where

$$\vec{e}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad \text{and} \quad \vec{e}_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}.$$

$$R_\theta(\vec{e}_1) = R_\theta \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \begin{bmatrix} \cos \theta \\ \sin \theta \end{bmatrix},$$

and

$$R_\theta(\vec{e}_2) = R_\theta \begin{bmatrix} 0 \\ 1 \end{bmatrix} = \begin{bmatrix} -\sin \theta \\ \cos \theta \end{bmatrix}$$

The Matrix for R_θ

The rotation $R_\theta : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is a linear transformation, and is induced by the matrix

$$\begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}.$$

Example (Rotation through π)

We denote by

$$R_\pi : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$


counterclockwise rotation about the origin through an angle of π .

Example (Rotation through π)

We denote by

$$R_\pi : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

counterclockwise rotation about the origin through an angle of π .




We see that $R_\pi \begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} -a \\ -b \end{bmatrix} = \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} a \\ b \end{bmatrix}$, so R_π is a matrix transformation.

Example (Rotation through π)

We denote by

$$R_\pi : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

counterclockwise rotation about the origin through an angle of π .




We see that $R_\pi \begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} -a \\ -b \end{bmatrix} = \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} a \\ b \end{bmatrix}$, so R_π is a matrix transformation.

Example (Rotation through π)

We denote by

$$R_\pi : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

counterclockwise rotation about the origin through an angle of π .




Example (Rotation through π)

We denote by

$$R_\pi : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

counterclockwise rotation about the origin through an angle of π .




We see that $R_\pi \begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} -a \\ -b \end{bmatrix} =$

Example (Rotation through π)

We denote by

$$R_\pi : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

counterclockwise rotation about the origin through an angle of π .



We see that $R_\pi \begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} -a \\ -b \end{bmatrix} = \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} a \\ b \end{bmatrix}$, so R_π is a matrix transformation.

Problem

The transformation $R_{\frac{\pi}{2}} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ denotes a **counterclockwise** rotation about the origin through an angle of $\frac{\pi}{2}$ radians. Find the matrix of $R_{\frac{\pi}{2}}$.

Problem

The transformation $R_{\frac{\pi}{2}} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ denotes a **counterclockwise** rotation about the origin through an angle of $\frac{\pi}{2}$ radians. Find the matrix of $R_{\frac{\pi}{2}}$.

Solution

First,

$$R_{\frac{\pi}{2}} \begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} -b \\ a \end{bmatrix}$$

Problem

The transformation $R_{\frac{\pi}{2}} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ denotes a **counterclockwise** rotation about the origin through an angle of $\frac{\pi}{2}$ radians. Find the matrix of $R_{\frac{\pi}{2}}$.

Solution

First,

$$R_{\frac{\pi}{2}} \begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} -b \\ a \end{bmatrix}$$

Furthermore $R_{\frac{\pi}{2}}$ is a matrix transformation, and the matrix it is induced by is

$$\begin{bmatrix} -b \\ a \end{bmatrix} = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} a \\ b \end{bmatrix}.$$



Example (Rotation through $\pi/2$)

We denote by

$$R_{\pi/2} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$


counterclockwise rotation about the origin through an angle of $\pi/2$.

Example (Rotation through $\pi/2$)

We denote by

$$R_{\pi/2} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

counterclockwise rotation about the origin through an angle of $\pi/2$.




We see that $R_{\pi/2} \begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} -b \\ a \end{bmatrix} = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} a \\ b \end{bmatrix}$, so $R_{\pi/2}$ is a matrix transformation.

Example (Rotation through $\pi/2$)

We denote by

$$R_{\pi/2} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

counterclockwise rotation about the origin through an angle of $\pi/2$.




We see that $R_{\pi/2} \begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} -b \\ a \end{bmatrix} = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} a \\ b \end{bmatrix}$, so $R_{\pi/2}$ is a matrix transformation.

Example (Rotation through $\pi/2$)

We denote by

$$R_{\pi/2} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

counterclockwise rotation about the origin through an angle of $\pi/2$.




Example (Rotation through $\pi/2$)

We denote by

$$R_{\pi/2} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

counterclockwise rotation about the origin through an angle of $\pi/2$.




Example (Rotation through $\pi/2$)

We denote by

$$R_{\pi/2} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

counterclockwise rotation about the origin through an angle of $\pi/2$.




We see that $R_{\pi/2} \begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} -b \\ a \end{bmatrix} =$

Example (Rotation through $\pi/2$)

We denote by

$$R_{\pi/2} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$$

counterclockwise rotation about the origin through an angle of $\pi/2$.



We see that $R_{\pi/2} \begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} -b \\ a \end{bmatrix} = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} a \\ b \end{bmatrix}$, so $R_{\pi/2}$ is a matrix transformation.

Reflection in \mathbb{R}^2

Reflection in \mathbb{R}^2

Example

In \mathbb{R}^2 , reflection in the x-axis, which transforms $\begin{bmatrix} a \\ b \end{bmatrix}$ to $\begin{bmatrix} a \\ -b \end{bmatrix}$, is a matrix transformation because

$$\begin{bmatrix} a \\ -b \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} a \\ b \end{bmatrix}.$$

Reflection in \mathbb{R}^2

Example

In \mathbb{R}^2 , reflection in the x-axis, which transforms $\begin{bmatrix} a \\ b \end{bmatrix}$ to $\begin{bmatrix} a \\ -b \end{bmatrix}$, is a matrix transformation because

$$\begin{bmatrix} a \\ -b \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} a \\ b \end{bmatrix}.$$

Example

In \mathbb{R}^2 , reflection in the y-axis transforms $\begin{bmatrix} a \\ b \end{bmatrix}$ to $\begin{bmatrix} -a \\ b \end{bmatrix}$. This is a matrix transformation because


$$\begin{bmatrix} -a \\ b \end{bmatrix} = \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} a \\ b \end{bmatrix}.$$

Example

Reflection in the line $y = x$ transforms $\begin{bmatrix} a \\ b \end{bmatrix}$ to $\begin{bmatrix} b \\ a \end{bmatrix}$.


Example

Reflection in the line $y = x$ transforms $\begin{bmatrix} a \\ b \end{bmatrix}$ to $\begin{bmatrix} b \\ a \end{bmatrix}$.



Example

Reflection in the line $y = x$ transforms $\begin{bmatrix} a \\ b \end{bmatrix}$ to $\begin{bmatrix} b \\ a \end{bmatrix}$.



This is a matrix transformation because

$$\begin{bmatrix} b \\ a \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} a \\ b \end{bmatrix}.$$

Reflection in the line

Reflection in the line


Example (Reflection in $y = mx$ preserves scalar multiplication)

Let $Q_m : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ denote reflection in the line $y = mx$, and let $\vec{u} \in \mathbb{R}^2$.

Reflection in the line

Example (Reflection in $y = mx$ preserves scalar multiplication)



Let $Q_m : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ denote reflection in the line $y = mx$, and let $\vec{u} \in \mathbb{R}^2$.



Reflection in the line

Example (Reflection in $y = mx$ preserves scalar multiplication)



Let $Q_m : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ denote reflection in the line $y = mx$, and let $\vec{u} \in \mathbb{R}^2$.



Reflection in the line

Example (Reflection in $y = mx$ preserves scalar multiplication)



Let $Q_m : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ denote reflection in the line $y = mx$, and let $\vec{u} \in \mathbb{R}^2$.



Reflection in the line

Example (Reflection in $y = mx$ preserves scalar multiplication)

Let $Q_m : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ denote reflection in the line $y = mx$, and let $\vec{u} \in \mathbb{R}^2$.





The figure indicates that $Q_m(2\vec{u}) = 2Q_m(\vec{u})$.

Reflection in the line

Example (Reflection in $y = mx$ preserves scalar multiplication)

Let $Q_m : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ denote reflection in the line $y = mx$, and let $\vec{u} \in \mathbb{R}^2$.





The figure indicates that $Q_m(2\vec{u}) = 2Q_m(\vec{u})$. In general, for any scalar k ,

$$Q_m(kX) = kQ_m(X),$$

Reflection in the line

Example (Reflection in $y = mx$ preserves scalar multiplication)

Let $Q_m : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ denote reflection in the line $y = mx$, and let $\vec{u} \in \mathbb{R}^2$.



The figure indicates that $Q_m(2\vec{u}) = 2Q_m(\vec{u})$. In general, for any scalar k ,

$$Q_m(kX) = kQ_m(X),$$


i.e., Q_m preserves scalar multiplication.

Example (Reflection in $y = mx$ preserves vector addition)

Let $\vec{u}, \vec{v} \in \mathbb{R}^2$.


Example (Reflection in $y = mx$ preserves vector addition)

Let $\vec{u}, \vec{v} \in \mathbb{R}^2$.





Example (Reflection in $y = mx$ preserves vector addition)

Let $\vec{u}, \vec{v} \in \mathbb{R}^2$.





Example (Reflection in $y = mx$ preserves vector addition)

Let $\vec{u}, \vec{v} \in \mathbb{R}^2$.





Example (Reflection in $y = mx$ preserves vector addition)

Let $\vec{u}, \vec{v} \in \mathbb{R}^2$.




Example (Reflection in $y = mx$ preserves vector addition)

Let $\vec{u}, \vec{v} \in \mathbb{R}^2$.



Example (Reflection in $y = mx$ preserves vector addition)

Let $\vec{u}, \vec{v} \in \mathbb{R}^2$.




The figure indicates that

$$Q_m(\vec{u}) + Q_m(\vec{v}) = Q_m(\vec{u} + \vec{v}),$$

Example (Reflection in $y = mx$ preserves vector addition)

Let $\vec{u}, \vec{v} \in \mathbb{R}^2$.



The figure indicates that

$$Q_m(\vec{u}) + Q_m(\vec{v}) = Q_m(\vec{u} + \vec{v}),$$

i.e., Q_m preserves vector addition.


Since Q_m preserves addition and scalar multiplication, Q_m is a linear transformation, and hence a matrix transformation.

Since Q_m preserves addition and scalar multiplication, Q_m is a linear transformation, and hence a matrix transformation.

The matrix that induces Q_m can be found by computing $Q_m(\vec{e}_1)$ and $Q_m(\vec{e}_2)$, where


$$\vec{e}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad \text{and} \quad \vec{e}_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}.$$

$$Q_m(\vec{e}_1)$$




$$\cos \theta = \frac{1}{\sqrt{1+m^2}} \quad \text{and} \quad \sin \theta = \frac{m}{\sqrt{1+m^2}}$$

$$Q_m(\vec{e}_1)$$




$$\cos \theta = \frac{1}{\sqrt{1+m^2}} \quad \text{and} \quad \sin \theta = \frac{m}{\sqrt{1+m^2}}$$

$$Q_m(\vec{e}_1)$$



$$\cos \theta = \frac{1}{\sqrt{1 + m^2}} \quad \text{and} \quad \sin \theta = \frac{m}{\sqrt{1 + m^2}}$$


$$Q_m(\vec{e}_1)$$



$$\cos \theta = \frac{1}{\sqrt{1 + m^2}} \quad \text{and} \quad \sin \theta = \frac{m}{\sqrt{1 + m^2}}$$

$$Q_m(\vec{e}_1) = \begin{bmatrix} \cos(2\theta) \\ \sin(2\theta) \end{bmatrix}$$


$$Q_m(\vec{e}_1)$$



$$\cos \theta = \frac{1}{\sqrt{1+m^2}} \quad \text{and} \quad \sin \theta = \frac{m}{\sqrt{1+m^2}}$$

$$Q_m(\vec{e}_1) = \begin{bmatrix} \cos(2\theta) \\ \sin(2\theta) \end{bmatrix} = \begin{bmatrix} \cos^2 \theta - \sin^2 \theta \\ 2 \sin \theta \cos \theta \end{bmatrix}$$


$$Q_m(\vec{e}_1)$$



$$\cos \theta = \frac{1}{\sqrt{1+m^2}} \quad \text{and} \quad \sin \theta = \frac{m}{\sqrt{1+m^2}}$$


$$Q_m(\vec{e}_1) = \begin{bmatrix} \cos(2\theta) \\ \sin(2\theta) \end{bmatrix} = \begin{bmatrix} \cos^2 \theta - \sin^2 \theta \\ 2 \sin \theta \cos \theta \end{bmatrix} = \frac{1}{1+m^2} \begin{bmatrix} 1-m^2 \\ 2m \end{bmatrix}$$

$$Q_m(\vec{e}_2)$$




$$\cos \theta = \frac{m}{\sqrt{1+m^2}} \quad \text{and} \quad \sin \theta = \frac{1}{\sqrt{1+m^2}}$$

$$Q_m(\vec{e}_2)$$




$$\cos \theta = \frac{m}{\sqrt{1+m^2}} \quad \text{and} \quad \sin \theta = \frac{1}{\sqrt{1+m^2}}$$

$$Q_m(\vec{e}_2)$$



$$\cos \theta = \frac{m}{\sqrt{1+m^2}} \quad \text{and} \quad \sin \theta = \frac{1}{\sqrt{1+m^2}}$$


$$Q_m(\vec{e}_2)$$



$$\cos \theta = \frac{m}{\sqrt{1+m^2}} \quad \text{and} \quad \sin \theta = \frac{1}{\sqrt{1+m^2}}$$

$$Q_m(\vec{e}_2) = \begin{bmatrix} \cos(\frac{\pi}{2} - 2\theta) \\ \sin(\frac{\pi}{2} - 2\theta) \end{bmatrix}$$


$$Q_m(\vec{e}_2)$$



$$\cos \theta = \frac{m}{\sqrt{1+m^2}} \quad \text{and} \quad \sin \theta = \frac{1}{\sqrt{1+m^2}}$$

$$Q_m(\vec{e}_2) = \begin{bmatrix} \cos(\frac{\pi}{2} - 2\theta) \\ \sin(\frac{\pi}{2} - 2\theta) \end{bmatrix} = \begin{bmatrix} \cos \frac{\pi}{2} \cos(2\theta) + \sin \frac{\pi}{2} \sin(2\theta) \\ \sin \frac{\pi}{2} \cos(2\theta) - \cos \frac{\pi}{2} \sin(2\theta) \end{bmatrix}$$


$$Q_m(\vec{e}_2)$$



$$\cos \theta = \frac{m}{\sqrt{1+m^2}} \quad \text{and} \quad \sin \theta = \frac{1}{\sqrt{1+m^2}}$$

$$\begin{aligned} Q_m(\vec{e}_2) &= \begin{bmatrix} \cos(\frac{\pi}{2} - 2\theta) \\ \sin(\frac{\pi}{2} - 2\theta) \end{bmatrix} = \begin{bmatrix} \cos \frac{\pi}{2} \cos(2\theta) + \sin \frac{\pi}{2} \sin(2\theta) \\ \sin \frac{\pi}{2} \cos(2\theta) - \cos \frac{\pi}{2} \sin(2\theta) \end{bmatrix} \\ &= \begin{bmatrix} \sin(2\theta) \\ \cos(2\theta) \end{bmatrix} \end{aligned}$$


$$Q_m(\vec{e}_2)$$



$$\cos \theta = \frac{m}{\sqrt{1+m^2}} \quad \text{and} \quad \sin \theta = \frac{1}{\sqrt{1+m^2}}$$

$$\begin{aligned} Q_m(\vec{e}_2) &= \begin{bmatrix} \cos(\frac{\pi}{2} - 2\theta) \\ \sin(\frac{\pi}{2} - 2\theta) \end{bmatrix} = \begin{bmatrix} \cos \frac{\pi}{2} \cos(2\theta) + \sin \frac{\pi}{2} \sin(2\theta) \\ \sin \frac{\pi}{2} \cos(2\theta) - \cos \frac{\pi}{2} \sin(2\theta) \end{bmatrix} \\ &= \begin{bmatrix} \sin(2\theta) \\ \cos(2\theta) \end{bmatrix} = \begin{bmatrix} 2 \sin \theta \cos \theta \\ \cos^2 \theta - \sin^2 \theta \end{bmatrix} \end{aligned}$$

$$Q_m(\vec{e}_2)$$



$$\cos \theta = \frac{m}{\sqrt{1+m^2}} \quad \text{and} \quad \sin \theta = \frac{1}{\sqrt{1+m^2}}$$

$$\begin{aligned} Q_m(\vec{e}_2) &= \begin{bmatrix} \cos(\frac{\pi}{2} - 2\theta) \\ \sin(\frac{\pi}{2} - 2\theta) \end{bmatrix} = \begin{bmatrix} \cos \frac{\pi}{2} \cos(2\theta) + \sin \frac{\pi}{2} \sin(2\theta) \\ \sin \frac{\pi}{2} \cos(2\theta) - \cos \frac{\pi}{2} \sin(2\theta) \end{bmatrix} \\ &= \begin{bmatrix} \sin(2\theta) \\ \cos(2\theta) \end{bmatrix} = \begin{bmatrix} 2 \sin \theta \cos \theta \\ \cos^2 \theta - \sin^2 \theta \end{bmatrix} = \frac{1}{1+m^2} \begin{bmatrix} 2m \\ m^2 - 1 \end{bmatrix} \end{aligned}$$

Alternatively, we can use the following relation to find Q_m :

$$Q_m = R_\theta \circ Q_0 \circ R_{-\theta}$$

Alternatively, we can use the following relation to find Q_m :

$$Q_m = R_\theta \circ Q_0 \circ R_{-\theta}$$

$$R_\theta \sim \begin{bmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{bmatrix}, \quad Q_0 \sim \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \quad R_{-\theta} \sim \begin{bmatrix} \cos(\theta) & \sin(\theta) \\ -\sin(\theta) & \cos(\theta) \end{bmatrix},$$

Alternatively, we can use the following relation to find Q_m :

$$Q_m = R_\theta \circ Q_0 \circ R_{-\theta}$$

$$R_\theta \sim \begin{bmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{bmatrix}, \quad Q_0 \sim \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \quad R_{-\theta} \sim \begin{bmatrix} \cos(\theta) & \sin(\theta) \\ -\sin(\theta) & \cos(\theta) \end{bmatrix},$$

Then multiply these three matrices ...

The Matrix for Reflection in $y = mx$

The transformation $Q_m : \mathbb{R}^2 \rightarrow \mathbb{R}^2$, reflection in the line $y = mx$, is a linear transformation and is induced by the matrix

$$\frac{1}{1+m^2} \begin{bmatrix} 1-m^2 & 2m \\ 2m & m^2-1 \end{bmatrix}.$$

Problem (Multiple Actions)

Find the rotation or reflection that equals reflection in the x-axis followed by rotation through an angle of $\frac{\pi}{2}$.

Problem (Multiple Actions)

Find the rotation or reflection that equals reflection in the x-axis followed by rotation through an angle of $\frac{\pi}{2}$.

Solution

Let Q_0 denote the reflection in the x-axis, and $R_{\frac{\pi}{2}}$ denote the rotation through an angle of $\frac{\pi}{2}$. We want to find the matrix for the transformation $R_{\frac{\pi}{2}} \circ Q_0$.

Problem (Multiple Actions)

Find the rotation or reflection that equals reflection in the x-axis followed by rotation through an angle of $\frac{\pi}{2}$.

Solution

Let Q_0 denote the reflection in the x-axis, and $R_{\frac{\pi}{2}}$ denote the rotation through an angle of $\frac{\pi}{2}$. We want to find the matrix for the transformation $R_{\frac{\pi}{2}} \circ Q_0$.

Q_0 is induced by $A = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$, and $R_{\frac{\pi}{2}}$ is induced by

$$B = \begin{bmatrix} \cos \frac{\pi}{2} & -\sin \frac{\pi}{2} \\ \sin \frac{\pi}{2} & \cos \frac{\pi}{2} \end{bmatrix} = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$$

Solution

Hence $R_{\frac{\pi}{2}} \circ Q_0$ is induced by

$$BA = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}.$$

Solution

Hence $R_{\frac{\pi}{2}} \circ Q_0$ is induced by

$$BA = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}.$$

Notice that $BA = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ is a **reflection** matrix.

Solution

Hence $R_{\frac{\pi}{2}} \circ Q_0$ is induced by

$$BA = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}.$$

Notice that $BA = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ is a **reflection** matrix.

How do we know this?

Solution (continued)

Compare BA to

$$Q_m = \frac{1}{1+m^2} \begin{bmatrix} 1-m^2 & 2m \\ 2m & m^2-1 \end{bmatrix}$$

Solution (continued)

Compare BA to

$$Q_m = \frac{1}{1+m^2} \begin{bmatrix} 1-m^2 & 2m \\ 2m & m^2-1 \end{bmatrix}$$

Now, since $1-m^2=0$, we know that $m=1$ or $m=-1$. But $\frac{2m}{1+m^2}=1>0$, so $m>0$, implying $m=1$.

Solution (continued)

Compare BA to

$$Q_m = \frac{1}{1+m^2} \begin{bmatrix} 1-m^2 & 2m \\ 2m & m^2-1 \end{bmatrix}$$

Now, since $1-m^2=0$, we know that $m=1$ or $m=-1$. But $\frac{2m}{1+m^2}=1>0$, so $m>0$, implying $m=1$.

Therefore,

$$R_{\frac{\pi}{2}} \circ Q_0 = Q_1,$$

reflection in the line $y=x$. ■

Problem (Reflection followed by Reflection)

Find the rotation or reflection that equals reflection in the line $y = -x$ followed by reflection in the y -axis.

Problem (Reflection followed by Reflection)

Find the rotation or reflection that equals reflection in the line $y = -x$ followed by reflection in the y -axis.

Solution

We must find the matrix for the transformation $Q_Y \circ Q_{-1}$.

Problem (Reflection followed by Reflection)

Find the rotation or reflection that equals reflection in the line $y = -x$ followed by reflection in the y -axis.

Solution

We must find the matrix for the transformation $Q_Y \circ Q_{-1}$.

Q_{-1} is induced by

$$A = \frac{1}{2} \begin{bmatrix} 0 & -2 \\ -2 & 0 \end{bmatrix} = \begin{bmatrix} 0 & -1 \\ -1 & 0 \end{bmatrix},$$

and Q_Y is induced by

$$B = \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix}.$$

Problem (Reflection followed by Reflection)

Find the rotation or reflection that equals reflection in the line $y = -x$ followed by reflection in the y -axis.

Solution

We must find the matrix for the transformation $Q_Y \circ Q_{-1}$.

Q_{-1} is induced by

$$A = \frac{1}{2} \begin{bmatrix} 0 & -2 \\ -2 & 0 \end{bmatrix} = \begin{bmatrix} 0 & -1 \\ -1 & 0 \end{bmatrix},$$

and Q_Y is induced by

$$B = \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix}.$$

Therefore, $Q_Y \circ Q_{-1}$ is induced by BA .

Solution (continued)

$$\mathbf{BA} = \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 0 & -1 \\ -1 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}.$$

Solution (continued)

$$\mathbf{BA} = \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 0 & -1 \\ -1 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}.$$

What transformation does \mathbf{BA} induce?

Solution (continued)

$$\mathbf{BA} = \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 0 & -1 \\ -1 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}.$$

What transformation does \mathbf{BA} induce?

Rotation through an angle θ such that

$$\cos \theta = 0 \quad \text{and} \quad \sin \theta = -1.$$

Solution (continued)

$$\mathbf{BA} = \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 0 & -1 \\ -1 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}.$$

What transformation does \mathbf{BA} induce?

Rotation through an angle θ such that

$$\cos \theta = 0 \quad \text{and} \quad \sin \theta = -1.$$

Therefore, $\mathbf{Q}_Y \circ \mathbf{Q}_{-1} = \mathbf{R}_{-\frac{\pi}{2}} = \mathbf{R}_{\frac{3\pi}{2}}$.



Remark (Summary)

In general,

- ▶ The composite of two rotations is a

Remark (Summary)

In general,

- The composite of two rotations is a rotation

$$\mathbf{R}_\theta \circ \mathbf{R}_\eta = \mathbf{R}_{\theta+\eta}$$

Remark (Summary)

In general,

- ▶ The composite of two rotations is a rotation

$$R_\theta \circ R_\eta = R_{\theta+\eta}$$

- ▶ The composite of two reflections is a

Remark (Summary)

In general,

- The composite of two rotations is a rotation

$$R_\theta \circ R_\eta = R_{\theta+\eta}$$

- The composite of two reflections is a rotation.

$$Q_m \circ Q_n = R_\theta$$

where θ is $2\times$ the angle between lines $y = mx$ and $y = nx$.

Remark (Summary)

In general,

- The composite of two rotations is a rotation

$$R_\theta \circ R_\eta = R_{\theta+\eta}$$

- The composite of two reflections is a rotation.

$$Q_m \circ Q_n = R_\theta$$

where θ is $2\times$ the angle between lines $y = mx$ and $y = nx$.

- The composite of a reflection and a rotation is a

Remark (Summary)

In general,

- The composite of two rotations is a rotation

$$R_\theta \circ R_\eta = R_{\theta+\eta}$$

- The composite of two reflections is a rotation.

$$Q_m \circ Q_n = R_\theta$$

where θ is $2\times$ the angle between lines $y = mx$ and $y = nx$.

- The composite of a reflection and a rotation is a reflection.

$$R_\theta \circ Q_n = Q_m \circ Q_n \circ Q_n = Q_m$$