

Math 221: LINEAR ALGEBRA

Chapter 5. Vector Space \mathbb{R}^n

§5-4. Rank of a Matrix

Le Chen¹

Emory University, 2021 Spring

(last updated on 01/12/2023)



Creative Commons License
(CC BY-NC-SA)

¹Slides are adapted from those by Karen Seyffarth from University of Calgary.

Linear Algebra with Applications

Lecture Notes

Current Lecture Notes Revision: Version 2018 — Revision B

These lecture notes were originally developed by Karen Seyffarth of the University of Calgary. Edits, additions, and revisions have been made to these notes by the editorial team at Lyryx Learning to accompany their text [Linear Algebra with Applications](#) based on W. K. Nicholson's original text.

In addition we recognize the following contributors. All new content contributed is released under the same license as noted below.

- Ilijas Farah, York University

BE A CHAMPION OF OER!

Contribute suggestions for improvements, new content, or errata:

A new topic

A new example or problem

A new or better proof to an existing theorem

Any other suggestions to improve the material

Contact Lyryx at info@lyryx.com with your ideas.

License



Attribution-NonCommercial-ShareAlike (CC BY-NC-SA)

This license lets others remix, tweak, and build upon your work non-commercially, as long as they credit you and license their new creations under the identical terms.

Copyright

Row Space and Column Spaces

The Rank Theorem

Rank-Nullity Theorem

Full Rank Cases

Copyright

Row Space and Column Spaces

The Rank Theorem

Rank-Nullity Theorem

Full Rank Cases

Row Space and Column Spaces

Definitions

Let A be an $m \times n$ matrix.

- The **column space of A** , denoted $\text{col}(A)$ is the subspace of \mathbb{R}^m spanned by the columns of A .

$$\begin{bmatrix} 1 & 8 & 13 & 12 \\ 14 & 11 & 2 & 7 \\ 4 & 5 & 16 & 9 \\ 15 & 10 & 3 & 6 \end{bmatrix}$$

Row Space and Column Spaces

Definitions

Let A be an $m \times n$ matrix.

- The **column space of A** , denoted $\text{col}(A)$ is the subspace of \mathbb{R}^m spanned by the columns of A .

$$\begin{bmatrix} 1 & 8 & 13 & 12 \\ 14 & 11 & 2 & 7 \\ 4 & 5 & 16 & 9 \\ 15 & 10 & 3 & 6 \end{bmatrix}$$

- The **row space of A** , denoted $\text{row}(A)$ is the subspace of \mathbb{R}^n spanned by the rows of A (or the columns of A^T).

$$\begin{bmatrix} 1 & 8 & 13 & 12 \\ 14 & 11 & 2 & 7 \\ 4 & 5 & 16 & 9 \\ 15 & 10 & 3 & 6 \end{bmatrix}$$

We saw earlier that $\text{col}(A) = \text{im}(A)$.

Remark (Notation)

Let A and B be $m \times n$ matrices. We write $A \rightarrow B$ if B can be obtained from A by a sequence of elementary row (column) operations. Note that $A \rightarrow B$ if and only if $B \rightarrow A$.

Lemma

Let A and B be $m \times n$ matrices.

1. If $A \rightarrow B$ by elementary row operations, then $\text{row}(A) = \text{row}(B)$.
2. If $A \rightarrow B$ by elementary column operations, then $\text{col}(A) = \text{col}(B)$.

Lemma

Let A and B be $m \times n$ matrices.

1. If $A \rightarrow B$ by elementary row operations, then $\text{row}(A) = \text{row}(B)$.
2. If $A \rightarrow B$ by elementary column operations, then $\text{col}(A) = \text{col}(B)$.

Proof.

It suffices to prove only part one, and only for a single row operation.
(Why?)

Lemma

Let A and B be $m \times n$ matrices.

1. If $A \rightarrow B$ by elementary row operations, then $\text{row}(A) = \text{row}(B)$.
2. If $A \rightarrow B$ by elementary column operations, then $\text{col}(A) = \text{col}(B)$.

Proof.

It suffices to prove only part one, and only for a single row operation.

(Why?)

Thus let $\vec{r}_1, \vec{r}_2, \dots, \vec{r}_m$ denote the rows of A .

Lemma

Let A and B be $m \times n$ matrices.

1. If $A \rightarrow B$ by elementary row operations, then $\text{row}(A) = \text{row}(B)$.
2. If $A \rightarrow B$ by elementary column operations, then $\text{col}(A) = \text{col}(B)$.

Proof.

It suffices to prove only part one, and only for a single row operation.

(Why?)

Thus let $\vec{r}_1, \vec{r}_2, \dots, \vec{r}_m$ denote the rows of A .

- If B is obtained from A by interchanging two rows of A , then A and B have exactly the same rows, so $\text{row}(B) = \text{row}(A)$.

Proof. (continued)

- Suppose $p \neq 0$, and suppose that for some j , $1 \leq j \leq m$, B is obtained from A by multiplying row j by p . Then

$$\text{row}(B) = \text{span}\{\vec{r}_1, \dots, p\vec{r}_j, \dots, \vec{r}_m\}.$$

Since

$$\{\vec{r}_1, \dots, p\vec{r}_j, \dots, \vec{r}_m\} \subseteq \text{row}(A),$$

it follows that $\text{row}(B) \subseteq \text{row}(A)$.

Proof. (continued)

- Suppose $p \neq 0$, and suppose that for some j , $1 \leq j \leq m$, B is obtained from A by multiplying row j by p . Then

$$\text{row}(B) = \text{span}\{\vec{r}_1, \dots, p\vec{r}_j, \dots, \vec{r}_m\}.$$

Since

$$\{\vec{r}_1, \dots, p\vec{r}_j, \dots, \vec{r}_m\} \subseteq \text{row}(A),$$

it follows that $\text{row}(B) \subseteq \text{row}(A)$. Conversely, since

$$\{\vec{r}_1, \dots, \vec{r}_m\} \subseteq \text{row}(B),$$

it follows that $\text{row}(A) \subseteq \text{row}(B)$. Therefore, $\text{row}(B) = \text{row}(A)$.

Proof. (continued)

- Suppose $p \neq 0$, and suppose that for some i and j , $1 \leq i, j \leq m$, B is obtained from A by adding p times row j to row i . Without loss of generality, we may assume $i < j$.

Proof. (continued)

- Suppose $p \neq 0$, and suppose that for some i and j , $1 \leq i, j \leq m$, B is obtained from A by adding p times row j to row i . Without loss of generality, we may assume $i < j$. Then

$$\text{row}(B) = \text{span}\{\vec{r}_1, \dots, \vec{r}_{i-1}, \vec{r}_i + p\vec{r}_j, \dots, \vec{r}_j, \dots, \vec{r}_m\}.$$

Since

$$\{\vec{r}_1, \dots, \vec{r}_{i-1}, \vec{r}_i + p\vec{r}_j, \dots, \vec{r}_m\} \subseteq \text{row}(A),$$

it follows that $\text{row}(B) \subseteq \text{row}(A)$.

Proof. (continued)

- Suppose $p \neq 0$, and suppose that for some i and j , $1 \leq i, j \leq m$, B is obtained from A by adding p times row j to row i . Without loss of generality, we may assume $i < j$. Then

$$\text{row}(B) = \text{span}\{\vec{r}_1, \dots, \vec{r}_{i-1}, \vec{r}_i + p\vec{r}_j, \dots, \vec{r}_j, \dots, \vec{r}_m\}.$$

Since

$$\{\vec{r}_1, \dots, \vec{r}_{i-1}, \vec{r}_i + p\vec{r}_j, \dots, \vec{r}_m\} \subseteq \text{row}(A),$$

it follows that $\text{row}(B) \subseteq \text{row}(A)$. Conversely, since

$$\{\vec{r}_1, \dots, \vec{r}_m\} \subseteq \text{row}(B),$$

it follows that $\text{row}(A) \subseteq \text{row}(B)$. Therefore, $\text{row}(B) = \text{row}(A)$.



Corollary

Let A be an $m \times n$ matrix, U an invertible $m \times m$ matrix, and V an invertible $n \times n$ matrix. Then $\text{row}(UA) = \text{row}(A)$ and $\text{col}(AV) = \text{col}(A)$,

Corollary

Let A be an $m \times n$ matrix, U an invertible $m \times m$ matrix, and V an invertible $n \times n$ matrix. Then $\text{row}(UA) = \text{row}(A)$ and $\text{col}(AV) = \text{col}(A)$,

Proof.

Since U is invertible, U is a product of elementary matrices, implying that $A \rightarrow UA$ by a sequence of elementary row operations. By Lemma 2, $\text{row}(UA) = \text{row}(A)$.

Now consider AV : $\text{col}(AV) = \text{row}((AV)^T) = \text{row}(V^T A^T)$ and V^T is invertible (a matrix is invertible if and only if its transpose is invertible). It follows from the first part of this Corollary that

$$\text{row}(V^T A^T) = \text{row}(A^T).$$

But $\text{row}(A^T) = \text{col}(A)$, and therefore $\text{col}(AV) = \text{col}(A)$. ■

Lemma

If R is a row-echelon matrix then

1. the nonzero rows of R are a basis of $\text{row}(R)$;
2. the columns of R containing the leading ones are a basis of $\text{col}(R)$.

Example

Let

$$R = \begin{bmatrix} 1 & 2 & 2 & -2 & 0 & 0 \\ 0 & 1 & 3 & 1 & -1 & 2 \\ 0 & 0 & 0 & 1 & -2 & 5 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}.$$

1. Since the nonzero rows of R are linearly independent, they form a basis of $\text{row}(R)$.

Example

Let

$$R = \begin{bmatrix} 1 & 2 & 2 & -2 & 0 & 0 \\ 0 & 1 & 3 & 1 & -1 & 2 \\ 0 & 0 & 0 & 1 & -2 & 5 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}.$$

1. Since the nonzero rows of R are linearly independent, they form a basis of $\text{row}(R)$.
2. Let $B = \{\vec{e}_1, \vec{e}_2, \vec{e}_3, \vec{e}_4\} \subseteq \mathbb{R}^5$. Then B is linearly independent and spans $\text{col}(R)$, and thus is a basis of $\text{col}(R)$. This tells us that $\dim(\text{col}(R)) = 4$. Now let X denote the set of columns of R that contain the leading ones.

Example

Let

$$R = \begin{bmatrix} 1 & 2 & 2 & -2 & 0 & 0 \\ 0 & 1 & 3 & 1 & -1 & 2 \\ 0 & 0 & 0 & 1 & -2 & 5 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}.$$

1. Since the nonzero rows of R are linearly independent, they form a basis of $\text{row}(R)$.
2. Let $B = \{\vec{e}_1, \vec{e}_2, \vec{e}_3, \vec{e}_4\} \subseteq \mathbb{R}^5$. Then B is linearly independent and spans $\text{col}(R)$, and thus is a basis of $\text{col}(R)$. This tells us that $\dim(\text{col}(R)) = 4$. Now let X denote the set of columns of R that contain the leading ones. Then X is a linearly independent subset of $\text{col}(R)$ with $4 = \dim(\text{col}(R))$ vectors. It follows that X spans $\text{col}(R)$, and therefore is a basis of $\text{col}(R)$.

Problem

Find a basis of $U = \text{span} \left\{ \begin{bmatrix} 1 \\ -1 \\ 0 \\ 3 \end{bmatrix}, \begin{bmatrix} 2 \\ 1 \\ 5 \\ 1 \end{bmatrix}, \begin{bmatrix} 4 \\ -1 \\ 5 \\ 7 \end{bmatrix} \right\}$ and find $\dim(U)$.

Problem

Find a basis of $U = \text{span} \left\{ \begin{bmatrix} 1 \\ -1 \\ 0 \\ 3 \end{bmatrix}, \begin{bmatrix} 2 \\ 1 \\ 5 \\ 1 \end{bmatrix}, \begin{bmatrix} 4 \\ -1 \\ 5 \\ 7 \end{bmatrix} \right\}$ and find $\dim(U)$.

Solution

Let A be the 3×4 matrix whose **rows** are the three columns listed. Then $U = \text{row}(A)$, so it suffices to find a basis of $\text{row}(A)$.

$$A = \begin{bmatrix} 1 & -1 & 0 & 3 \\ 2 & 1 & 5 & 1 \\ 4 & -1 & 5 & 7 \end{bmatrix}.$$

Find R , a row-echelon form of A . Then the **nonzero rows of R** are a basis of $\text{row}(R)$. Since $\text{row}(A) = \text{row}(R)$, the nonzero rows of R are a basis of $\text{row}(A)$.

Solution (continued)

$$\begin{bmatrix} 1 & -1 & 0 & 3 \\ 2 & 1 & 5 & 1 \\ 4 & -1 & 5 & 7 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & -1 & 0 & 3 \\ 0 & 1 & 5/3 & -5/3 \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

Therefore, $B = \left\{ \begin{bmatrix} 1 \\ -1 \\ 0 \\ 3 \end{bmatrix}, \begin{bmatrix} 0 \\ 3 \\ 5 \\ -5 \end{bmatrix} \right\}$ is a basis of U and $\dim(U) = 2$.

Solution (continued)

$$\begin{bmatrix} 1 & -1 & 0 & 3 \\ 2 & 1 & 5 & 1 \\ 4 & -1 & 5 & 7 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & -1 & 0 & 3 \\ 0 & 1 & 5/3 & -5/3 \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

Therefore, $B = \left\{ \begin{bmatrix} 1 \\ -1 \\ 0 \\ 3 \end{bmatrix}, \begin{bmatrix} 0 \\ 3 \\ 5 \\ -5 \end{bmatrix} \right\}$ is a basis of U and $\dim(U) = 2$.

Solution (Another solution – usually more work.)

Take a linear combination of the three given vectors and set it equal to $\vec{0}_4$. If the vectors are independent, then they form a basis of U . Otherwise, delete vectors to cut the given set of vectors down to a basis.

Copyright

Row Space and Column Spaces

The Rank Theorem

Rank-Nullity Theorem

Full Rank Cases

The Rank Theorem

The Rank Theorem

$$\dim(\text{row}(A)) = \dim(\text{col}(A)) = \text{rank}(A)$$

Remark

Recall that $\text{rank}(A)$ is defined to be the nonzero rows in the row echelon form of A . From what we just learned, the **rank of A** can be equivalently defined as **$\text{rank}(A) = \dim(\text{row}(A))$** .

Remark

Recall that $\text{rank}(A)$ is defined to be the nonzero rows in the row echelon form of A . From what we just learned, the **rank of A** can be equivalently defined as **$\text{rank}(A) = \dim(\text{row}(A))$** .

Theorem (Rank Theorem)

Let $A = \begin{bmatrix} \vec{A}_1 & \vec{A}_2 & \cdots & \vec{A}_n \end{bmatrix}$ be an $m \times n$ matrix with columns $\{\vec{A}_1, \vec{A}_2, \dots, \vec{A}_n\}$, and suppose that $\text{rank}(A) = r$. Then

$$\dim(\text{row}(A)) = \dim(\text{col}(A)) = r.$$

Furthermore, if R is a row-echelon form of A then

1. the r nonzero rows of R are a basis of $\text{row}(A)$;
2. if $S = \{\vec{A}_{j_1}, \vec{A}_{j_2}, \dots, \vec{A}_{j_r}\}$ are the r columns of A corresponding to the columns of R containing leading ones, then S is basis of $\text{col}(A)$.

Problem

For the following matrix A , find $\text{rank}(A)$ and bases for $\text{row}(A)$ and $\text{col}(A)$.

$$A = \begin{bmatrix} 2 & -4 & 6 & 8 \\ 2 & -1 & 3 & 2 \\ 4 & -5 & 9 & 10 \\ 0 & -1 & 1 & 2 \end{bmatrix}.$$

Problem

For the following matrix A, find rank (A) and bases for row(A) and col(A).

$$A = \begin{bmatrix} 2 & -4 & 6 & 8 \\ 2 & -1 & 3 & 2 \\ 4 & -5 & 9 & 10 \\ 0 & -1 & 1 & 2 \end{bmatrix}.$$

Solution

$$\begin{bmatrix} 2 & -4 & 6 & 8 \\ 2 & -1 & 3 & 2 \\ 4 & -5 & 9 & 10 \\ 0 & -1 & 1 & 2 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & -2 & 3 & 4 \\ 0 & 1 & -1 & -2 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

Problem

For the following matrix A, find rank (A) and bases for row(A) and col(A).

$$A = \begin{bmatrix} 2 & -4 & 6 & 8 \\ 2 & -1 & 3 & 2 \\ 4 & -5 & 9 & 10 \\ 0 & -1 & 1 & 2 \end{bmatrix}.$$

Solution

$$\begin{bmatrix} 2 & -4 & 6 & 8 \\ 2 & -1 & 3 & 2 \\ 4 & -5 & 9 & 10 \\ 0 & -1 & 1 & 2 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & -2 & 3 & 4 \\ 0 & 1 & -1 & -2 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

► rank (A) = 2.

Problem

For the following matrix A, find rank (A) and bases for row(A) and col(A).

$$A = \begin{bmatrix} 2 & -4 & 6 & 8 \\ 2 & -1 & 3 & 2 \\ 4 & -5 & 9 & 10 \\ 0 & -1 & 1 & 2 \end{bmatrix}.$$

Solution

$$\begin{bmatrix} 2 & -4 & 6 & 8 \\ 2 & -1 & 3 & 2 \\ 4 & -5 & 9 & 10 \\ 0 & -1 & 1 & 2 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & -2 & 3 & 4 \\ 0 & 1 & -1 & -2 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

- rank (A) = 2.
- $\{[1 \ -2 \ 3 \ 4], [0 \ -1 \ -1 \ 2]\}$ is a basis of row(A).

Problem

For the following matrix A, find rank (A) and bases for row(A) and col(A).

$$A = \begin{bmatrix} 2 & -4 & 6 & 8 \\ 2 & -1 & 3 & 2 \\ 4 & -5 & 9 & 10 \\ 0 & -1 & 1 & 2 \end{bmatrix}.$$

Solution

$$\begin{bmatrix} 2 & -4 & 6 & 8 \\ 2 & -1 & 3 & 2 \\ 4 & -5 & 9 & 10 \\ 0 & -1 & 1 & 2 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & -2 & 3 & 4 \\ 0 & 1 & -1 & -2 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

► rank (A) = 2.

► $\left\{ \begin{bmatrix} 1 & -2 & 3 & 4 \end{bmatrix}, \begin{bmatrix} 0 & -1 & -1 & 2 \end{bmatrix} \right\}$ is a basis of row(A).

► $\left\{ \begin{bmatrix} 2 \\ 2 \\ 4 \\ 0 \end{bmatrix}, \begin{bmatrix} -4 \\ -1 \\ -5 \\ -1 \end{bmatrix} \right\}$ is a basis of col(A).



Problem (revisited)

Find a basis of $U = \text{span} \left\{ \begin{bmatrix} 1 \\ -1 \\ 0 \\ 3 \end{bmatrix}, \begin{bmatrix} 2 \\ 1 \\ 5 \\ 1 \end{bmatrix}, \begin{bmatrix} 4 \\ -1 \\ 5 \\ 7 \end{bmatrix} \right\}$ and find $\dim(U)$.

Problem (revisited)

Find a basis of $U = \text{span} \left\{ \begin{bmatrix} 1 \\ -1 \\ 0 \\ 3 \end{bmatrix}, \begin{bmatrix} 2 \\ 1 \\ 5 \\ 1 \end{bmatrix}, \begin{bmatrix} 4 \\ -1 \\ 5 \\ 7 \end{bmatrix} \right\}$ and find $\dim(U)$.

Solution

Let A denote the matrix whose columns are the three vectors listed, and let R denote a row-echelon form of A . Then

$$A = \begin{bmatrix} 1 & 2 & 4 \\ -1 & 1 & -1 \\ 0 & 5 & 5 \\ 3 & 1 & 7 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 2 & 4 \\ 0 & 1 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} = R.$$

By the Rank Theorem, $\left\{ \begin{bmatrix} 1 \\ -1 \\ 0 \\ 3 \end{bmatrix}, \begin{bmatrix} 2 \\ 1 \\ 5 \\ 1 \end{bmatrix} \right\}$ is a basis of $U = \text{col}(A)$, so

$$\dim(U) = 2.$$



Compare this to the basis found earlier.

Corollary

1. For any matrix A , $\text{rank}(A) = \text{rank}(A^T)$.
2. For any $m \times n$ matrix A , $\text{rank}(A) \leq m$ and $\text{rank}(A) \leq n$.
3. Let A be an $m \times n$ matrix. If U and V are invertible matrices (of sizes $m \times m$ and $n \times n$, respectively), then

$$\text{rank}(A) = \text{rank}(UA) = \text{rank}(AV).$$

Lemma

Let A be an $m \times n$ matrix, U a $p \times m$ matrix, and V an $n \times q$ matrix.

1. $\text{col}(AV) \subseteq \text{col}(A)$ with equality if $VV' = I_n$ for some V' .
2. $\text{row}(UA) \subseteq \text{row}(A)$ with equality if $U'U = I_m$ for some U' .

Lemma

Let A be an $m \times n$ matrix, U a $p \times m$ matrix, and V an $n \times q$ matrix.

1. $\text{col}(AV) \subseteq \text{col}(A)$ with equality if $VV' = I_n$ for some V' .
2. $\text{row}(UA) \subseteq \text{row}(A)$ with equality if $U'U = I_m$ for some U' .

Proof.

(1) Write $V = [\vec{v}_1 \ \vec{v}_2 \ \cdots \ \vec{v}_q]$, where \vec{v}_j denotes column j of V , $1 \leq j \leq q$. Then $AV = [A\vec{v}_1 \ A\vec{v}_2 \ \cdots \ A\vec{v}_q]$, where $A\vec{v}_j$ is column j of AV . By the definition of matrix-vector multiplication, $A\vec{v}_j$ is a linear combination of the columns of A , and thus $A\vec{v}_j \in \text{col}(A)$ for each j . Since $A\vec{v}_1, A\vec{v}_2, \dots, A\vec{v}_q \in \text{col}(A)$,

$$\text{span}\{A\vec{v}_1, A\vec{v}_2, \dots, A\vec{v}_q\} \subseteq \text{col}(A),$$

i.e., $\text{col}(AV) \subseteq \text{col}(A)$.

Lemma

Let A be an $m \times n$ matrix, U a $p \times m$ matrix, and V an $n \times q$ matrix.

1. $\text{col}(AV) \subseteq \text{col}(A)$ with equality if $VV' = I_n$ for some V' .
2. $\text{row}(UA) \subseteq \text{row}(A)$ with equality if $U'U = I_m$ for some U' .

Proof.

(1) Write $V = [\vec{v}_1 \ \vec{v}_2 \ \cdots \ \vec{v}_q]$, where \vec{v}_j denotes column j of V , $1 \leq j \leq q$. Then $AV = [A\vec{v}_1 \ A\vec{v}_2 \ \cdots \ A\vec{v}_q]$, where $A\vec{v}_j$ is column j of AV . By the definition of matrix-vector multiplication, $A\vec{v}_j$ is a linear combination of the columns of A , and thus $A\vec{v}_j \in \text{col}(A)$ for each j . Since $A\vec{v}_1, A\vec{v}_2, \dots, A\vec{v}_q \in \text{col}(A)$,

$$\text{span}\{A\vec{v}_1, A\vec{v}_2, \dots, A\vec{v}_q\} \subseteq \text{col}(A),$$

i.e., $\text{col}(AV) \subseteq \text{col}(A)$. If for some V' we have $VV' = I_n$, then

$$\text{col}(A) = \text{col}(AVV') \subseteq \text{col}(AV) \subseteq \text{col}(A).$$

Lemma

Let A be an $m \times n$ matrix, U a $p \times m$ matrix, and V an $n \times q$ matrix.

1. $\text{col}(AV) \subseteq \text{col}(A)$ with equality if $VV' = I_n$ for some V' .
2. $\text{row}(UA) \subseteq \text{row}(A)$ with equality if $U'U = I_m$ for some U' .

Proof.

(1) Write $V = [\vec{v}_1 \ \vec{v}_2 \ \cdots \ \vec{v}_q]$, where \vec{v}_j denotes column j of V , $1 \leq j \leq q$. Then $AV = [A\vec{v}_1 \ A\vec{v}_2 \ \cdots \ A\vec{v}_q]$, where $A\vec{v}_j$ is column j of AV . By the definition of matrix-vector multiplication, $A\vec{v}_j$ is a linear combination of the columns of A , and thus $A\vec{v}_j \in \text{col}(A)$ for each j . Since $A\vec{v}_1, A\vec{v}_2, \dots, A\vec{v}_q \in \text{col}(A)$,

$$\text{span}\{A\vec{v}_1, A\vec{v}_2, \dots, A\vec{v}_q\} \subseteq \text{col}(A),$$

i.e., $\text{col}(AV) \subseteq \text{col}(A)$. If for some V' we have $VV' = I_n$, then

$$\text{col}(A) = \text{col}(AVV') \subseteq \text{col}(AV) \subseteq \text{col}(A).$$

(2) This can be proved by part (1) and the fact that $\text{row}(A) = \text{col}(A^T)$. ■

Copyright


Row Space and Column Spaces

The Rank Theorem

Rank-Nullity Theorem

Full Rank Cases

Rank-Nullity Theorem



Theorem (Rank-Nullity Theorem)

Let A denote an $m \times n$ matrix of rank r . Then

1. The $n - r$ basic solutions to the system $A\vec{x} = \vec{0}_m$ provided by the Gaussian algorithm are a basis of $\text{null}(A)$, so

$$\dim(\text{null}(A)) = n - r.$$

2. The rank theorem provides a basis of $\text{im}(A) = \text{col}(A)$, and $\dim(\text{im}(A)) = r$.

Theorem (Rank-Nullity Theorem)

Let A denote an $m \times n$ matrix of rank r . Then

1. The $n - r$ basic solutions to the system $A\vec{x} = \vec{0}_m$ provided by the Gaussian algorithm are a basis of $\text{null}(A)$, so

$$\dim(\text{null}(A)) = n - r.$$

2. The rank theorem provides a basis of $\text{im}(A) = \text{col}(A)$, and $\dim(\text{im}(A)) = r$.

Remark (Common notation)

The nullspace A is also called kernel space of A , written as $\ker(A)$, i.e., $\ker(A) = \text{null}(A)$. Usually, the **nullity** of A is defined to be

$$\text{Nullity}(A) = \dim(\text{null}(A)) = \dim(\ker(A))$$

Let $T : V \mapsto W$ be the linear map from space V to W . Suppose $V = \mathbb{R}^n$ and $W = \mathbb{R}^m$ and let A be the induced matrix.

$$\begin{array}{ccccc}
 \text{Rank}(T) & + & \text{Nullity}(T) & = & \dim(V) \\
 \parallel & & \parallel & & \parallel \\
 \text{Rank}(A) & & \text{Nullity}(A) & & \dim(\mathbb{R}^n) \\
 \parallel & & \parallel & & \parallel \\
 \dim(\text{im}(A)) & & \dim(\text{null}(A)) & & n \\
 \parallel & & \parallel & & \\
 r & & \dim(\ker(A)) & &
 \end{array}$$

Proof. (Outline)

- ▶ We have already seen that $\text{null}(A)$ is spanned by any set of basic solutions to $A\vec{x} = \vec{0}_m$, so it is enough to prove that $\dim(\text{null}(A)) = n - r$, which will imply that the set of basic solutions is independent, hence this set forms a basis.
- ▶ Suppose $\{\vec{x}_1, \vec{x}_2, \dots, \vec{x}_k\}$ is a basis of $\text{null}(A)$
- ▶ Extend $\{\vec{x}_1, \vec{x}_2, \dots, \vec{x}_k\}$ to a basis $\{\vec{x}_1, \vec{x}_2, \dots, \vec{x}_k, \dots, \vec{x}_n\}$ of \mathbb{R}^n .
- ▶ Consider the set $\{A\vec{x}_1, A\vec{x}_2, \dots, A\vec{x}_k, \dots, A\vec{x}_n\} \subseteq \mathbb{R}^m$
- ▶ Then $A\vec{x}_j = \vec{0}_m$ for $1 \leq j \leq k$ since $\vec{x}_1, \dots, \vec{x}_k \in \text{null}(A)$.
- ▶ To complete the proof, show $S = \{A\vec{x}_{k+1}, \dots, A\vec{x}_n\}$ is a basis of $\text{im}(A)$, by showing that (exercise!)
 - (1) S is independent
 - (2) S spans $\text{im}(A)$
- ▶ Since $\text{im}(A) = \text{col}(A)$, $\dim(\text{im}(A)) = r$, implying $n - k = r$. Hence $k = n - r$. ■

Problem

For the following matrix A , find bases for $\text{null}(A)$ and $\text{im}(A)$, and find their dimensions.

$$A = \begin{bmatrix} 2 & -4 & 6 & 8 \\ 2 & -1 & 3 & 2 \\ 4 & -5 & 9 & 10 \\ 0 & -1 & 1 & 2 \end{bmatrix}.$$

Solution

Find the basic solutions to $A\vec{x} = \vec{0}_4$.

$$\left[\begin{array}{cccc|c} 2 & -4 & 6 & 8 & 0 \\ 2 & -1 & 3 & 2 & 0 \\ 4 & -5 & 9 & 10 & 0 \\ 0 & -1 & 1 & 2 & 0 \end{array} \right] \rightarrow \left[\begin{array}{cccc|c} 1 & -2 & 3 & 4 & 0 \\ 0 & 1 & -1 & -2 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right].$$

Hence,

$$\vec{x} = \begin{bmatrix} -s \\ s + 2t \\ s \\ t \end{bmatrix} \quad s, t \in \mathbb{R}.$$

Solution

Find the basic solutions to $A\vec{x} = \vec{0}_4$.

$$\left[\begin{array}{cccc|c} 2 & -4 & 6 & 8 & 0 \\ 2 & -1 & 3 & 2 & 0 \\ 4 & -5 & 9 & 10 & 0 \\ 0 & -1 & 1 & 2 & 0 \end{array} \right] \rightarrow \left[\begin{array}{cccc|c} 1 & -2 & 3 & 4 & 0 \\ 0 & 1 & -1 & -2 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right].$$

Hence,

$$\vec{x} = \begin{bmatrix} -s \\ s + 2t \\ s \\ t \end{bmatrix} \quad s, t \in \mathbb{R}.$$

Therefore,

$$\left\{ \begin{bmatrix} -1 \\ 1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 2 \\ 0 \\ 1 \end{bmatrix} \right\} \quad \text{and} \quad \left\{ \begin{bmatrix} 2 \\ 2 \\ 4 \\ 0 \end{bmatrix}, \begin{bmatrix} -4 \\ -1 \\ -5 \\ -1 \end{bmatrix} \right\}$$

are bases of $\text{null}(A)$ and $\text{im}(A)$, respectively, so

$$\dim(\text{null}(A)) = 2 \quad \text{and} \quad \dim(\text{im}(A)) = 2.$$



Problem

Can a 5×6 matrix have independent columns? Independent rows? Justify your answer.

Problem

Can a 5×6 matrix have independent columns? Independent rows? Justify your answer.

Solution

The rank of the matrix is at most five; since there are six columns, **the columns can not be independent**. However, the rows could be independent:

Problem

Can a 5×6 matrix have independent columns? Independent rows? Justify your answer.

Solution

The rank of the matrix is at most five; since there are six columns, **the columns can not be independent**. However, the rows could be independent: take a 5×6 matrix whose first five columns are the columns of the 5×5 identity matrix.

Problem

Let A be an $m \times n$ matrix with $\text{rank}(A) = m$. Prove that $m \leq n$.

Problem

Let A be an $m \times n$ matrix with $\text{rank}(A) = m$. Prove that $m \leq n$.

Proof.

As a consequence of the Rank Theorem, we have

$$\text{rank}(A) \leq m \quad \text{and} \quad \text{rank}(A) \leq n.$$

Since $\text{rank}(A) = m$, it follows that $m \leq n$. ■

Problem

Let A be an 5×9 matrix. Is it possible that $\dim(\text{null}(A)) = 3$? Justify your answer.

Problem

Let A be an 5×9 matrix. Is it possible that $\dim(\text{null}(A)) = 3$? Justify your answer.

Solution

As a consequence of the Rank Theorem, we have $\text{rank}(A) \leq 5$, so $\dim(\text{im}(A)) \leq 5$. Since $\dim(\text{null}(A)) = 9 - \dim(\text{im}(A))$, it follows that

$$\dim(\text{null}(A)) \geq 9 - 5 = 4.$$

Therefore, it is not possible that $\dim(\text{null}(A)) = 3$.



Copyright

Row Space and Column Spaces

The Rank Theorem

Rank-Nullity Theorem

Full Rank Cases

Full Rank Cases

Full Rank Cases

Theorem

Let A be an $m \times n$ matrix. The following are equivalent.

Full Rank Cases

Theorem

Let A be an $m \times n$ matrix. The following are equivalent.

1. $\text{rank}(A) = n$.

Full Rank Cases

Theorem

Let A be an $m \times n$ matrix. The following are equivalent.

1. $\text{rank}(A) = n$.
2. $\text{row}(A) = \mathbb{R}^n$, i.e., the rows of A span \mathbb{R}^n .

Full Rank Cases

Theorem

Let A be an $m \times n$ matrix. The following are equivalent.

1. $\text{rank}(A) = n$.
2. $\text{row}(A) = \mathbb{R}^n$, i.e., the rows of A span \mathbb{R}^n .
3. The columns of A are independent in \mathbb{R}^m .

Full Rank Cases

Theorem

Let A be an $m \times n$ matrix. The following are equivalent.

1. $\text{rank}(A) = n$.
2. $\text{row}(A) = \mathbb{R}^n$, i.e., the rows of A span \mathbb{R}^n .
3. The columns of A are independent in \mathbb{R}^m .
4. The $n \times n$ matrix $A^T A$ is invertible.

Full Rank Cases

Theorem

Let A be an $m \times n$ matrix. The following are equivalent.

1. $\text{rank}(A) = n$.
2. $\text{row}(A) = \mathbb{R}^n$, i.e., the rows of A span \mathbb{R}^n .
3. The columns of A are independent in \mathbb{R}^m .
4. The $n \times n$ matrix $A^T A$ is invertible.
5. There exists an $n \times m$ matrix C so that $CA = I_n$.

Full Rank Cases

Theorem

Let A be an $m \times n$ matrix. The following are equivalent.

1. $\text{rank}(A) = n$.
2. $\text{row}(A) = \mathbb{R}^n$, i.e., the rows of A span \mathbb{R}^n .
3. The columns of A are independent in \mathbb{R}^m .
4. The $n \times n$ matrix $A^T A$ is invertible.
5. There exists an $n \times m$ matrix C so that $CA = I_n$.
6. If $A\vec{x} = \vec{0}_m$ for some $\vec{x} \in \mathbb{R}^n$, then $\vec{x} = \vec{0}_n$.

Theorem

Let A be an $m \times n$ matrix. The following are equivalent.

Theorem

Let A be an $m \times n$ matrix. The following are equivalent.

1. $\text{rank}(A) = m$.

Theorem

Let A be an $m \times n$ matrix. The following are equivalent.

1. $\text{rank}(A) = m$.
2. $\text{col}(A) = \mathbb{R}^m$, i.e., the columns of A span \mathbb{R}^m .

Theorem

Let A be an $m \times n$ matrix. The following are equivalent.

1. $\text{rank}(A) = m$.
2. $\text{col}(A) = \mathbb{R}^m$, i.e., the columns of A span \mathbb{R}^m .
3. The rows of A are independent in \mathbb{R}^n .

Theorem

Let A be an $m \times n$ matrix. The following are equivalent.

1. $\text{rank}(A) = m$.
2. $\text{col}(A) = \mathbb{R}^m$, i.e., the columns of A span \mathbb{R}^m .
3. The rows of A are independent in \mathbb{R}^n .
4. The $m \times m$ matrix AA^T is invertible.

Theorem

Let A be an $m \times n$ matrix. The following are equivalent.

1. $\text{rank}(A) = m$.
2. $\text{col}(A) = \mathbb{R}^m$, i.e., the columns of A span \mathbb{R}^m .
3. The rows of A are independent in \mathbb{R}^n .
4. The $m \times m$ matrix AA^T is invertible.
5. There exists an $n \times m$ matrix C so that $AC = I_m$.

Theorem

Let A be an $m \times n$ matrix. The following are equivalent.

1. $\text{rank}(A) = m$.
2. $\text{col}(A) = \mathbb{R}^m$, i.e., the columns of A span \mathbb{R}^m .
3. The rows of A are independent in \mathbb{R}^n .
4. The $m \times m$ matrix AA^T is invertible.
5. There exists an $n \times m$ matrix C so that $AC = I_m$.
6. The system $A\vec{x} = \vec{b}$ is consistent for every $\vec{b} \in \mathbb{R}^m$.

Problem

Let $\vec{x} = (x_1, \dots, x_k)^T \in \mathbb{R}^k$. Show that the following matrix is invertible if and only if $\{x_i, i = 1, \dots, k\}$ are not all equal:

$$\begin{pmatrix} k & x_1 + \dots + x_k \\ x_1 + \dots + x_k & ||x||^2 \end{pmatrix}$$

Problem

Let $\vec{x} = (x_1, \dots, x_k)^T \in \mathbb{R}^k$. Show that the following matrix is invertible if and only if $\{x_i, i = 1, \dots, k\}$ are not all equal:

$$\begin{pmatrix} k & x_1 + \dots + x_k \\ x_1 + \dots + x_k & ||x||^2 \end{pmatrix}$$

Solution

Notice that

$$\begin{pmatrix} k & x_1 + \dots + x_k \\ x_1 + \dots + x_k & ||x||^2 \end{pmatrix} = A^T A$$

with

$$A = \begin{bmatrix} 1 & x_1 \\ 1 & x_2 \\ \vdots & \vdots \\ 1 & x_k \end{bmatrix}.$$

Now $A^T A$ is invertible iff the two columns of A are independent iff $\{x_i, i = 1, \dots, k\}$ are not all equal. ■