

# Math 221: LINEAR ALGEBRA

## §Appendix A. Complex Numbers

Le Chen<sup>1</sup>

Emory University, 2021 Spring

(last updated on 01/12/2023)



Creative Commons License  
(CC BY-NC-SA)

<sup>1</sup>Slides are adapted from those by Karen Seyffarth from University of Calgary.



# Linear Algebra with Applications

## Lecture Notes

### Current Lecture Notes Revision: Version 2018 — Revision B

These lecture notes were originally developed by Karen Seyffarth of the University of Calgary. Edits, additions, and revisions have been made to these notes by the editorial team at Lyryx Learning to accompany their text [Linear Algebra with Applications](#) based on W. K. Nicholson's original text.

In addition we recognize the following contributors. All new content contributed is released under the same license as noted below.

- Ilijas Farah, York University

### BE A CHAMPION OF OER!

Contribute suggestions for improvements, new content, or errata:

A new topic

A new example or problem

A new or better proof to an existing theorem

Any other suggestions to improve the material

Contact Lyryx at [info@lyryx.com](mailto:info@lyryx.com) with your ideas.

### License



Attribution-NonCommercial-ShareAlike (CC BY-NC-SA)

This license lets others remix, tweak, and build upon your work non-commercially, as long as they credit you and license their new creations under the identical terms.

Copyright

Complex Numbers

Modulus

Complex Numbers in Polar Form

Roots of Complex Numbers

Roots of Unity

The Quadratic Formula

Copyright

## Complex Numbers

Modulus

Complex Numbers in Polar Form

Roots of Complex Numbers

Roots of Unity

The Quadratic Formula



# Complex Numbers

## Why complex numbers?

- ▶ Counting numbers:  $1, 2, 3, 4, 5, \dots$
- ▶ Integers:  $0, 1, 2, 3, 4 \dots$  but also  $-1, -2, -3 \dots$

# Complex Numbers

## Why complex numbers?

- ▶ Counting numbers:  $1, 2, 3, 4, 5, \dots$
- ▶ Integers:  $0, 1, 2, 3, 4, \dots$  but also  $-1, -2, -3, \dots$
- ▶ To solve  $3x + 2 = 0$ , integers aren't enough, so we have **rational numbers** (fractions), i.e.,

$$\text{if } 3x + 2 = 0, \text{ then } x = -\frac{2}{3}.$$



# Complex Numbers

## Why complex numbers?

- ▶ Counting numbers:  $1, 2, 3, 4, 5, \dots$
- ▶ Integers:  $0, 1, 2, 3, 4, \dots$  but also  $-1, -2, -3, \dots$
- ▶ To solve  $3x + 2 = 0$ , integers aren't enough, so we have **rational numbers** (fractions), i.e.,

$$\text{if } 3x + 2 = 0, \text{ then } x = -\frac{2}{3}.$$

- ▶ We still can't solve  $x^2 - 2 = 0$  because there are no rational numbers  $x$  with the property that  $x^2 - 2 = 0$ , so we have **irrational numbers**, i.e.,

$$\text{if } x^2 - 2 = 0, \text{ then } x = \pm\sqrt{2}.$$

# Complex Numbers

## Why complex numbers?

- ▶ Counting numbers:  $1, 2, 3, 4, 5, \dots$
- ▶ Integers:  $0, 1, 2, 3, 4, \dots$  but also  $-1, -2, -3, \dots$
- ▶ To solve  $3x + 2 = 0$ , integers aren't enough, so we have **rational numbers** (fractions), i.e.,

$$\text{if } 3x + 2 = 0, \text{ then } x = -\frac{2}{3}.$$

- ▶ We still can't solve  $x^2 - 2 = 0$  because there are no rational numbers  $x$  with the property that  $x^2 - 2 = 0$ , so we have **irrational numbers**, i.e.,

$$\text{if } x^2 - 2 = 0, \text{ then } x = \pm\sqrt{2}.$$

- ▶ The set of **real numbers**,  $\mathbb{R}$ , consists of all rational and irrational numbers (note that integers are rational numbers). However, we still can't solve

$$x^2 + 1 = 0$$

because this requires  $x^2 = -1$ , but any **real** number  $x$  has the property that  $x^2 \geq 0$ .

## Definitions

- ▶ The **imaginary unit**, denoted  $i$ , is defined to be a number with the property that  $i^2 = -1$ .

## Definitions

- ▶ The **imaginary unit**, denoted  $i$ , is defined to be a number with the property that  $i^2 = -1$ .
- ▶ A **pure imaginary** number has the form  $bi$  where  $b \in \mathbb{R}$ ,  $b \neq 0$ , and  $i$  is the imaginary unit.

## Definitions

- ▶ The **imaginary unit**, denoted  $i$ , is defined to be a number with the property that  $i^2 = -1$ .
- ▶ A **pure imaginary** number has the form  $bi$  where  $b \in \mathbb{R}$ ,  $b \neq 0$ , and  $i$  is the imaginary unit.
- ▶ A **complex number** is any number  $z$  of the form

$$z = a + bi$$

where  $a, b \in \mathbb{R}$  and  $i$  is the imaginary unit.

## Definitions

- ▶ The **imaginary unit**, denoted  $i$ , is defined to be a number with the property that  $i^2 = -1$ .
- ▶ A **pure imaginary** number has the form  $bi$  where  $b \in \mathbb{R}$ ,  $b \neq 0$ , and  $i$  is the imaginary unit.
- ▶ A **complex number** is any number  $z$  of the form

$$z = a + bi$$

where  $a, b \in \mathbb{R}$  and  $i$  is the imaginary unit.

(1)  $a$  is called the **real part** of  $z$ .

## Definitions

- ▶ The **imaginary unit**, denoted  $i$ , is defined to be a number with the property that  $i^2 = -1$ .
- ▶ A **pure imaginary** number has the form  $bi$  where  $b \in \mathbb{R}$ ,  $b \neq 0$ , and  $i$  is the imaginary unit.
- ▶ A **complex number** is any number  $z$  of the form

$$z = a + bi$$

where  $a, b \in \mathbb{R}$  and  $i$  is the imaginary unit.

(1)  $a$  is called the **real part** of  $z$ .

(2)  $b$  is called the **imaginary part** of  $z$

## Definitions

- ▶ The **imaginary unit**, denoted  $i$ , is defined to be a number with the property that  $i^2 = -1$ .
- ▶ A **pure imaginary** number has the form  $bi$  where  $b \in \mathbb{R}$ ,  $b \neq 0$ , and  $i$  is the imaginary unit.
- ▶ A **complex number** is any number  $z$  of the form

$$z = a + bi$$

where  $a, b \in \mathbb{R}$  and  $i$  is the imaginary unit.

- (1)  $a$  is called the **real part** of  $z$ .
- (2)  $b$  is called the **imaginary part** of  $z$
- (3) If  $b = 0$ , then  $z$  is a real number.




# The Complex Plane

## The Complex Plane

A complex number  $z = a + bi$  can be represented geometrically by the point  $(a, b)$  in the  $xy$ -plane, where the  $x$ -axis is the **real axis** and the  $y$ -axis is the **imaginary axis**.


## The Complex Plane

A complex number  $z = a + bi$  can be represented geometrically by the point  $(a, b)$  in the  $xy$ -plane, where the  $x$ -axis is the **real axis** and the  $y$ -axis is the **imaginary axis**.



## The Complex Plane


A complex number  $z = a + bi$  can be represented geometrically by the point  $(a, b)$  in the  $xy$ -plane, where the  $x$ -axis is the **real axis** and the  $y$ -axis is the **imaginary axis**.



► Real numbers:  $a + 0i$  lie on the  $x$ -axis.

## The Complex Plane

A complex number  $z = a + bi$  can be represented geometrically by the point  $(a, b)$  in the  $xy$ -plane, where the  $x$ -axis is the **real axis** and the  $y$ -axis is the **imaginary axis**.



- Real numbers:  $a + 0i$  lie on the  $x$ -axis.
- Pure imaginary numbers:  $0 + bi$  ( $b \neq 0$ ) lie on the  $y$ -axis.

## Addition and Subtraction of Complex Numbers

## Addition and Subtraction of Complex Numbers

Let  $z = a + bi$  and  $w = c + di$  be complex numbers.

## Addition and Subtraction of Complex Numbers

Let  $z = a + bi$  and  $w = c + di$  be complex numbers.

- **Equality**  $z = w$  if and  $a = c$  and  $b = d$ .



## Addition and Subtraction of Complex Numbers

Let  $z = a + bi$  and  $w = c + di$  be complex numbers.

- ▶ **Equality**  $z = w$  if and  $a = c$  and  $b = d$ .
- ▶ **Addition**

## Addition and Subtraction of Complex Numbers

Let  $z = a + bi$  and  $w = c + di$  be complex numbers.

- ▶ **Equality**  $z = w$  if and  $a = c$  and  $b = d$ .
- ▶ **Addition**

$$z + w =$$

## Addition and Subtraction of Complex Numbers

Let  $z = a + bi$  and  $w = c + di$  be complex numbers.

► **Equality**  $z = w$  if and  $a = c$  and  $b = d$ .

► **Addition**

$$z + w = (a + bi) + (c + di) =$$

## Addition and Subtraction of Complex Numbers

Let  $z = a + bi$  and  $w = c + di$  be complex numbers.

► **Equality**  $z = w$  if and  $a = c$  and  $b = d$ .

► **Addition**

$$z + w = (a + bi) + (c + di) = (a + c) + (b + d)i.$$

## Addition and Subtraction of Complex Numbers

Let  $z = a + bi$  and  $w = c + di$  be complex numbers.

► **Equality**  $z = w$  if and  $a = c$  and  $b = d$ .

► **Addition**

$$z + w = (a + bi) + (c + di) = (a + c) + (b + d)i.$$

► **Subtraction**

## Addition and Subtraction of Complex Numbers

Let  $z = a + bi$  and  $w = c + di$  be complex numbers.

► **Equality**  $z = w$  if and  $a = c$  and  $b = d$ .

► **Addition**

$$z + w = (a + bi) + (c + di) = (a + c) + (b + d)i.$$

► **Subtraction**

$$z - w = (a + bi) - (c + di) = (a - c) + (b - d)i.$$

## Addition and Subtraction of Complex Numbers

Let  $z = a + bi$  and  $w = c + di$  be complex numbers.

► **Equality**  $z = w$  if and  $a = c$  and  $b = d$ .

► **Addition**

$$z + w = (a + bi) + (c + di) = (a + c) + (b + d)i.$$

► **Subtraction**

$$z - w = (a + bi) - (c + di) = (a - c) + (b - d)i.$$

## Examples

►  $(-3 + 6i) + (5 - i) =$

## Addition and Subtraction of Complex Numbers

Let  $z = a + bi$  and  $w = c + di$  be complex numbers.

► **Equality**  $z = w$  if and  $a = c$  and  $b = d$ .

► **Addition**

$$z + w = (a + bi) + (c + di) = (a + c) + (b + d)i.$$

► **Subtraction**

$$z - w = (a + bi) - (c + di) = (a - c) + (b - d)i.$$

## Examples

►  $(-3 + 6i) + (5 - i) = 2 + 5i.$



## Addition and Subtraction of Complex Numbers

Let  $z = a + bi$  and  $w = c + di$  be complex numbers.

► **Equality**  $z = w$  if and  $a = c$  and  $b = d$ .

► **Addition**

$$z + w = (a + bi) + (c + di) = (a + c) + (b + d)i.$$

► **Subtraction**

$$z - w = (a + bi) - (c + di) = (a - c) + (b - d)i.$$

## Examples

►  $(-3 + 6i) + (5 - i) = 2 + 5i.$

►  $(4 - 7i) + (6 - 2i) =$

## Addition and Subtraction of Complex Numbers

Let  $z = a + bi$  and  $w = c + di$  be complex numbers.

► **Equality**  $z = w$  if and  $a = c$  and  $b = d$ .

► **Addition**

$$z + w = (a + bi) + (c + di) = (a + c) + (b + d)i.$$

► **Subtraction**

$$z - w = (a + bi) - (c + di) = (a - c) + (b - d)i.$$

## Examples

►  $(-3 + 6i) + (5 - i) = 2 + 5i.$

►  $(4 - 7i) + (6 - 2i) = 10 - 9i.$

## Addition and Subtraction of Complex Numbers

Let  $z = a + bi$  and  $w = c + di$  be complex numbers.

► **Equality**  $z = w$  if and  $a = c$  and  $b = d$ .

► **Addition**

$$z + w = (a + bi) + (c + di) = (a + c) + (b + d)i.$$

► **Subtraction**

$$z - w = (a + bi) - (c + di) = (a - c) + (b - d)i.$$

## Examples

►  $(-3 + 6i) + (5 - i) = 2 + 5i.$

►  $(4 - 7i) + (6 - 2i) = 10 - 9i.$

►  $(-3 + 6i) - (5 - i) =$

## Addition and Subtraction of Complex Numbers

Let  $z = a + bi$  and  $w = c + di$  be complex numbers.

► **Equality**  $z = w$  if and  $a = c$  and  $b = d$ .

► **Addition**

$$z + w = (a + bi) + (c + di) = (a + c) + (b + d)i.$$

► **Subtraction**

$$z - w = (a + bi) - (c + di) = (a - c) + (b - d)i.$$

## Examples

►  $(-3 + 6i) + (5 - i) = 2 + 5i.$

►  $(4 - 7i) + (6 - 2i) = 10 - 9i.$

►  $(-3 + 6i) - (5 - i) = -8 + 7i.$

## Addition and Subtraction of Complex Numbers

Let  $z = a + bi$  and  $w = c + di$  be complex numbers.

► **Equality**  $z = w$  if and  $a = c$  and  $b = d$ .

► **Addition**

$$z + w = (a + bi) + (c + di) = (a + c) + (b + d)i.$$

► **Subtraction**

$$z - w = (a + bi) - (c + di) = (a - c) + (b - d)i.$$

## Examples

►  $(-3 + 6i) + (5 - i) = 2 + 5i.$

►  $(4 - 7i) + (6 - 2i) = 10 - 9i.$

►  $(-3 + 6i) - (5 - i) = -8 + 7i.$

►  $(4 - 7i) - (6 - 2i) =$

## Addition and Subtraction of Complex Numbers

Let  $z = a + bi$  and  $w = c + di$  be complex numbers.

► **Equality**  $z = w$  if and  $a = c$  and  $b = d$ .

► **Addition**

$$z + w = (a + bi) + (c + di) = (a + c) + (b + d)i.$$

► **Subtraction**

$$z - w = (a + bi) - (c + di) = (a - c) + (b - d)i.$$

## Examples

►  $(-3 + 6i) + (5 - i) = 2 + 5i.$

►  $(4 - 7i) + (6 - 2i) = 10 - 9i.$

►  $(-3 + 6i) - (5 - i) = -8 + 7i.$

►  $(4 - 7i) - (6 - 2i) = -2 - 5i.$

## Properties of Addition

## Properties of Addition

Let  $z$ ,  $w$ , and  $v$  be complex numbers.



## Properties of Addition

Let  $z$ ,  $w$ , and  $v$  be complex numbers.

1.  $z + w = w + z$ .

(addition is commutative)

## Properties of Addition

Let  $z$ ,  $w$ , and  $v$  be complex numbers.

1.  $z + w = w + z.$

(addition is commutative)

2.  $(z + w) + v = z + (w + v).$

(addition is associative)

## Properties of Addition

Let  $z$ ,  $w$ , and  $v$  be complex numbers.

1.  $z + w = w + z.$

(addition is commutative)

2.  $(z + w) + v = z + (w + v).$

(addition is associative)

3.  $z + 0 = z.$

(existence of an additive identity)

## Properties of Addition

Let  $z$ ,  $w$ , and  $v$  be complex numbers.

1.  $z + w = w + z.$  (addition is commutative)
2.  $(z + w) + v = z + (w + v).$  (addition is associative)
3.  $z + 0 = z.$  (existence of an additive identity)
4. For every  $z = a + bi$  there exists a complex number  $-z = -a - bi$  such that  $z + (-z) = 0.$  (existence of an additive inverse)


## Addition in the Complex Plane

## Addition in the Complex Plane

If  $z = a + bi$  and  $w = c + di$ , then  $z + w = (a + c) + (b + d)i$ .


## Addition in the Complex Plane

If  $z = a + bi$  and  $w = c + di$ , then  $z + w = (a + c) + (b + d)i$ . Geometrically, we have:



## Addition in the Complex Plane

If  $z = a + bi$  and  $w = c + di$ , then  $z + w = (a + c) + (b + d)i$ . Geometrically, we have:



$0$ ,  $z$ ,  $w$ , and  $z + w$  are the vertices of a parallelogram.



## Multiplication of Complex Numbers

Let  $z = a + bi$  and  $w = c + di$  be complex numbers. Then the **product** of  $z$  and  $w$  is

$$zw = (a + bi)(c + di) = (ac - bd) + (ad + bc)i.$$

## Multiplication of Complex Numbers

Let  $z = a + bi$  and  $w = c + di$  be complex numbers. Then the **product** of  $z$  and  $w$  is

$$zw = (a + bi)(c + di) = (ac - bd) + (ad + bc)i.$$

The multiplication is done essentially as the product of two linear polynomials, with  $i^2$  replaced by  $-1$ .

## Multiplication of Complex Numbers

Let  $z = a + bi$  and  $w = c + di$  be complex numbers. Then the **product** of  $z$  and  $w$  is

$$zw = (a + bi)(c + di) = (ac - bd) + (ad + bc)i.$$

The multiplication is done essentially as the product of two linear polynomials, with  $i^2$  replaced by  $-1$ .

### Example

$$\begin{aligned}(2 - 3i)(-3 + 4i) &= ((2)(-3) - (-3)(4)) + ((2)(4) + (-3)(-3))i \\ &= (-6 + 12) + (8 + 9)i \\ &= 6 + 17i\end{aligned}$$

## Properties of Multiplication

## Properties of Multiplication

Let  $z$ ,  $w$  and  $v$  be complex numbers.

## Properties of Multiplication

Let  $z$ ,  $w$  and  $v$  be complex numbers.

►  $zw = wz.$

(multiplication is commutative)

## Properties of Multiplication

Let  $z$ ,  $w$  and  $v$  be complex numbers.

►  $zw = wz.$

(multiplication is commutative)

►  $(zw)v = z(wv).$

(multiplication is associative)

## Properties of Multiplication

Let  $z$ ,  $w$  and  $v$  be complex numbers.

- ▶  $zw = wz.$  (multiplication is commutative)
- ▶  $(zw)v = z(wv).$  (multiplication is associative)
- ▶  $z(w + v) = zw + zv.$  (multiplication distributes over addition)



## Properties of Multiplication

Let  $z$ ,  $w$  and  $v$  be complex numbers.

- ▶  $zw = wz.$  (multiplication is commutative)
- ▶  $(zw)v = z(wv).$  (multiplication is associative)
- ▶  $z(w + v) = zw + zv.$  (multiplication distributes over addition)
- ▶  $1z = z.$  ('1' is the multiplicative identity)

## Properties of Multiplication

Let  $z$ ,  $w$  and  $v$  be complex numbers.

- ▶  $zw = wz.$  (multiplication is commutative)
- ▶  $(zw)v = z(wv).$  (multiplication is associative)
- ▶  $z(w + v) = zw + zv.$  (multiplication distributes over addition)
- ▶  $1z = z.$  ('1' is the multiplicative identity)
- ▶ For each  $z \neq 0$ , there exists  $z^{-1}$  such that  $zz^{-1} = 1.$   
(existence of a multiplicative inverse)

### Problem

Find all complex numbers  $z$  so that  $z^2 = -3 + 4i$ .

### Problem

Find all complex numbers  $z$  so that  $z^2 = -3 + 4i$ .

### Solution

Let  $z = a + bi$ . Then

$$z^2 = (a + bi)^2 = (a^2 - b^2) + 2abi = -3 + 4i,$$

## Problem

Find all complex numbers  $z$  so that  $z^2 = -3 + 4i$ .

## Solution

Let  $z = a + bi$ . Then

so 
$$z^2 = (a + bi)^2 = (a^2 - b^2) + 2abi = -3 + 4i,$$

$$a^2 - b^2 = -3 \quad \text{and} \quad 2ab = 4.$$

## Problem

Find all complex numbers  $z$  so that  $z^2 = -3 + 4i$ .

## Solution

Let  $z = a + bi$ . Then

so 
$$z^2 = (a + bi)^2 = (a^2 - b^2) + 2abi = -3 + 4i,$$

$$a^2 - b^2 = -3 \quad \text{and} \quad 2ab = 4.$$

Since  $2ab = 4$ ,  $a = \frac{2}{b}$ .

## Problem

Find all complex numbers  $z$  so that  $z^2 = -3 + 4i$ .

## Solution

Let  $z = a + bi$ . Then

so 
$$z^2 = (a + bi)^2 = (a^2 - b^2) + 2abi = -3 + 4i,$$

$$a^2 - b^2 = -3 \quad \text{and} \quad 2ab = 4.$$

Since  $2ab = 4$ ,  $a = \frac{2}{b}$ . Substituting this into the first equation gives us

## Problem

Find all complex numbers  $z$  so that  $z^2 = -3 + 4i$ .

## Solution

Let  $z = a + bi$ . Then

so 
$$z^2 = (a + bi)^2 = (a^2 - b^2) + 2abi = -3 + 4i,$$

Since  $2ab = 4$ ,  $a = \frac{2}{b}$ .  $a^2 - b^2 = -3$  and  $2ab = 4$ . Substituting this into the first equation gives us

$$\begin{aligned} a^2 - b^2 &= -3 \\ \left(\frac{2}{b}\right)^2 - b^2 &= -3 \end{aligned}$$



## Problem

Find all complex numbers  $z$  so that  $z^2 = -3 + 4i$ .

## Solution

Let  $z = a + bi$ . Then

so 
$$z^2 = (a + bi)^2 = (a^2 - b^2) + 2abi = -3 + 4i,$$

Since  $2ab = 4$ ,  $a = \frac{2}{b}$ .  $a^2 - b^2 = -3$  and  $2ab = 4$ . Substituting this into the first equation gives us

$$\begin{aligned} a^2 - b^2 &= -3 \\ \left(\frac{2}{b}\right)^2 - b^2 &= -3 \\ \frac{4}{b^2} - b^2 &= -3 \end{aligned}$$

## Problem

Find all complex numbers  $z$  so that  $z^2 = -3 + 4i$ .

## Solution

Let  $z = a + bi$ . Then

so 
$$z^2 = (a + bi)^2 = (a^2 - b^2) + 2abi = -3 + 4i,$$

Since  $2ab = 4$ ,  $a = \frac{2}{b}$ .  $a^2 - b^2 = -3$  and  $2ab = 4$ . Substituting this into the first equation gives us

$$\begin{aligned} a^2 - b^2 &= -3 \\ \left(\frac{2}{b}\right)^2 - b^2 &= -3 \\ \frac{4}{b^2} - b^2 &= -3 \\ 4 - b^4 &= -3b^2 \end{aligned}$$

## Problem

Find all complex numbers  $z$  so that  $z^2 = -3 + 4i$ .

## Solution

Let  $z = a + bi$ . Then

so 
$$z^2 = (a + bi)^2 = (a^2 - b^2) + 2abi = -3 + 4i,$$

Since  $2ab = 4$ ,  $a = \frac{2}{b}$ .  $a^2 - b^2 = -3$  and  $2ab = 4$ . Substituting this into the first equation gives us

$$\begin{aligned} a^2 - b^2 &= -3 \\ \left(\frac{2}{b}\right)^2 - b^2 &= -3 \\ \frac{4}{b^2} - b^2 &= -3 \\ 4 - b^4 &= -3b^2 \\ b^4 - 3b^2 - 4 &= 0. \end{aligned}$$

Solution (continued)

Now,  $b^4 - 3b^2 - 4 = 0$  can be factored

Solution (continued)

Now,  $b^4 - 3b^2 - 4 = 0$  can be factored into

$$(b^2 - 4)(b^2 + 1) = 0$$

### Solution (continued)

Now,  $b^4 - 3b^2 - 4 = 0$  can be factored into

$$\begin{aligned}(b^2 - 4)(b^2 + 1) &= 0 \\(b - 2)(b + 2)(b^2 + 1) &= 0.\end{aligned}$$

### Solution (continued)

Now,  $b^4 - 3b^2 - 4 = 0$  can be factored into

$$\begin{aligned}(b^2 - 4)(b^2 + 1) &= 0 \\ (b - 2)(b + 2)(b^2 + 1) &= 0.\end{aligned}$$

Since  $b \in \mathbb{R}$  and  $b^2 + 1$  has no real roots,  $b = 2$  or  $b = -2$ .

### Solution (continued)

Now,  $b^4 - 3b^2 - 4 = 0$  can be factored into

$$\begin{aligned}(b^2 - 4)(b^2 + 1) &= 0 \\ (b - 2)(b + 2)(b^2 + 1) &= 0.\end{aligned}$$

Since  $b \in \mathbb{R}$  and  $b^2 + 1$  has no real roots,  $b = 2$  or  $b = -2$ .

Since  $a = \frac{2}{b}$ , it follows that



### Solution (continued)

Now,  $b^4 - 3b^2 - 4 = 0$  can be factored into

$$\begin{aligned}(b^2 - 4)(b^2 + 1) &= 0 \\ (b - 2)(b + 2)(b^2 + 1) &= 0.\end{aligned}$$

Since  $b \in \mathbb{R}$  and  $b^2 + 1$  has no real roots,  $b = 2$  or  $b = -2$ .

Since  $a = \frac{2}{b}$ , it follows that

► when  $b = 2$ ,  $a = 1$ , and  $z = a + bi = 1 + 2i$ ;

### Solution (continued)

Now,  $b^4 - 3b^2 - 4 = 0$  can be factored into

$$\begin{aligned}(b^2 - 4)(b^2 + 1) &= 0 \\ (b - 2)(b + 2)(b^2 + 1) &= 0.\end{aligned}$$

Since  $b \in \mathbb{R}$  and  $b^2 + 1$  has no real roots,  $b = 2$  or  $b = -2$ .

Since  $a = \frac{2}{b}$ , it follows that

- ▶ when  $b = 2$ ,  $a = 1$ , and  $z = a + bi = 1 + 2i$ ;
- ▶ when  $b = -2$ ,  $a = -1$ , and  $z = a + bi = -1 - 2i$ .

### Solution (continued)

Now,  $b^4 - 3b^2 - 4 = 0$  can be factored into

$$\begin{aligned}(b^2 - 4)(b^2 + 1) &= 0 \\ (b - 2)(b + 2)(b^2 + 1) &= 0.\end{aligned}$$

Since  $b \in \mathbb{R}$  and  $b^2 + 1$  has no real roots,  $b = 2$  or  $b = -2$ .

Since  $a = \frac{2}{b}$ , it follows that

- ▶ when  $b = 2$ ,  $a = 1$ , and  $z = a + bi = 1 + 2i$ ;
- ▶ when  $b = -2$ ,  $a = -1$ , and  $z = a + bi = -1 - 2i$ .

Therefore, if  $z^2 = -3 + 4i$ , then  $z = 1 + 2i$  or  $z = -1 - 2i$ .

## The Conjugate of a Complex Number

## The Conjugate of a Complex Number

Let  $z = a + bi$  be a complex number.

## The Conjugate of a Complex Number


Let  $z = a + bi$  be a complex number. The **conjugate** of  $z$  is the complex number  $\bar{z} = a - bi$ .

## The Conjugate of a Complex Number

Let  $z = a + bi$  be a complex number. The **conjugate** of  $z$  is the complex number  $\bar{z} = a - bi$ . Geometrically,  $\bar{z}$  is the reflection of  $z$  in the  $x$ -axis.

## The Conjugate of a Complex Number


Let  $z = a + bi$  be a complex number. The **conjugate** of  $z$  is the complex number  $\bar{z} = a - bi$ . Geometrically,  $\bar{z}$  is the reflection of  $z$  in the  $x$ -axis.





## The Conjugate of a Complex Number


Let  $z = a + bi$  be a complex number. The **conjugate** of  $z$  is the complex number  $\bar{z} = a - bi$ . Geometrically,  $\bar{z}$  is the reflection of  $z$  in the  $x$ -axis.



## Examples

## The Conjugate of a Complex Number

Let  $z = a + bi$  be a complex number. The **conjugate** of  $z$  is the complex number  $\bar{z} = a - bi$ . Geometrically,  $\bar{z}$  is the reflection of  $z$  in the  $x$ -axis.




## Examples

- If  $z = 3 + 4i$ , then  $\bar{z} =$

## The Conjugate of a Complex Number

Let  $z = a + bi$  be a complex number. The **conjugate** of  $z$  is the complex number  $\bar{z} = a - bi$ . Geometrically,  $\bar{z}$  is the reflection of  $z$  in the  $x$ -axis.




## Examples

- If  $z = 3 + 4i$ , then  $\bar{z} = 3 - 4i$ ,

## The Conjugate of a Complex Number

Let  $z = a + bi$  be a complex number. The **conjugate** of  $z$  is the complex number  $\bar{z} = a - bi$ . Geometrically,  $\bar{z}$  is the reflection of  $z$  in the  $x$ -axis.




## Examples

- If  $z = 3 + 4i$ , then  $\bar{z} = 3 - 4i$ , i.e.,  $\overline{3 + 4i} = 3 - 4i$ .

## The Conjugate of a Complex Number

Let  $z = a + bi$  be a complex number. The **conjugate** of  $z$  is the complex number  $\bar{z} = a - bi$ . Geometrically,  $\bar{z}$  is the reflection of  $z$  in the  $x$ -axis.




## Examples

► If  $z = 3 + 4i$ , then  $\bar{z} = 3 - 4i$ , i.e.,  $\overline{3 + 4i} = 3 - 4i$ .

►  $\overline{-2 + 5i} =$

## The Conjugate of a Complex Number

Let  $z = a + bi$  be a complex number. The **conjugate** of  $z$  is the complex number  $\bar{z} = a - bi$ . Geometrically,  $\bar{z}$  is the reflection of  $z$  in the  $x$ -axis.




## Examples

- ▶ If  $z = 3 + 4i$ , then  $\bar{z} = 3 - 4i$ , i.e.,  $\overline{3 + 4i} = 3 - 4i$ .
- ▶  $\overline{-2 + 5i} = -2 - 5i$ .

## The Conjugate of a Complex Number

Let  $z = a + bi$  be a complex number. The **conjugate** of  $z$  is the complex number  $\bar{z} = a - bi$ . Geometrically,  $\bar{z}$  is the reflection of  $z$  in the  $x$ -axis.




### Examples

- ▶ If  $z = 3 + 4i$ , then  $\bar{z} = 3 - 4i$ , i.e.,  $\overline{3 + 4i} = 3 - 4i$ .
- ▶  $\overline{-2 + 5i} = -2 - 5i$ .
- ▶  $\bar{i} =$

## The Conjugate of a Complex Number

Let  $z = a + bi$  be a complex number. The **conjugate** of  $z$  is the complex number  $\bar{z} = a - bi$ . Geometrically,  $\bar{z}$  is the reflection of  $z$  in the  $x$ -axis.




### Examples

- ▶ If  $z = 3 + 4i$ , then  $\bar{z} = 3 - 4i$ , i.e.,  $\overline{3 + 4i} = 3 - 4i$ .
- ▶  $\overline{-2 + 5i} = -2 - 5i$ .
- ▶  $\bar{i} = -i$ .



## The Conjugate of a Complex Number

Let  $z = a + bi$  be a complex number. The **conjugate** of  $z$  is the complex number  $\bar{z} = a - bi$ . Geometrically,  $\bar{z}$  is the reflection of  $z$  in the  $x$ -axis.




### Examples

- ▶ If  $z = 3 + 4i$ , then  $\bar{z} = 3 - 4i$ , i.e.,  $\overline{3 + 4i} = 3 - 4i$ .
- ▶  $\overline{-2 + 5i} = -2 - 5i$ .
- ▶  $\bar{i} = -i$ .
- ▶  $\bar{\bar{z}} =$

## The Conjugate of a Complex Number

Let  $z = a + bi$  be a complex number. The **conjugate** of  $z$  is the complex number  $\bar{z} = a - bi$ . Geometrically,  $\bar{z}$  is the reflection of  $z$  in the  $x$ -axis.



### Examples

- ▶ If  $z = 3 + 4i$ , then  $\bar{z} = 3 - 4i$ , i.e.,  $\overline{3 + 4i} = 3 - 4i$ .
- ▶  $\overline{-2 + 5i} = -2 - 5i$ .
- ▶  $\bar{i} = -i$ .
- ▶  $\bar{7} = 7$ .

## Properties of the Conjugate

Let  $z$  and  $w$  be complex numbers.

►  $\overline{z \pm w} = \bar{z} \pm \bar{w}.$

## Properties of the Conjugate

Let  $z$  and  $w$  be complex numbers.

►  $\overline{z \pm w} = \bar{z} \pm \bar{w}.$

►  $\overline{(zw)} = \bar{z} \bar{w}.$

## Properties of the Conjugate

Let  $z$  and  $w$  be complex numbers.

►  $\overline{z \pm w} = \bar{z} \pm \bar{w}.$

►  $\overline{(zw)} = \bar{z} \bar{w}.$

►  $\overline{(\bar{z})} = z.$

## Properties of the Conjugate

Let  $z$  and  $w$  be complex numbers.

$$\blacktriangleright \overline{z \pm w} = \bar{z} \pm \bar{w}.$$

$$\blacktriangleright \overline{(zw)} = \bar{z} \bar{w}.$$

$$\blacktriangleright \overline{(\bar{z})} = z.$$

$$\blacktriangleright \overline{\left(\frac{z}{w}\right)} = \frac{\bar{z}}{\bar{w}}.$$

## Properties of the Conjugate

Let  $z$  and  $w$  be complex numbers.

$$\blacktriangleright \overline{z \pm w} = \bar{z} \pm \bar{w}.$$

$$\blacktriangleright \overline{(zw)} = \bar{z} \bar{w}.$$

$$\blacktriangleright \overline{(\bar{z})} = z.$$

$$\blacktriangleright \overline{\left(\frac{z}{w}\right)} = \frac{\bar{z}}{\bar{w}}.$$

$$\blacktriangleright z \text{ is real if and only if } \bar{z} = z.$$

## Properties of the Conjugate

Let  $z$  and  $w$  be complex numbers.

$$\blacktriangleright \overline{z \pm w} = \bar{z} \pm \bar{w}.$$

$$\blacktriangleright \overline{(zw)} = \bar{z} \bar{w}.$$

$$\blacktriangleright \overline{(\bar{z})} = z.$$

$$\blacktriangleright \overline{\left(\frac{z}{w}\right)} = \frac{\bar{z}}{\bar{w}}.$$

$$\blacktriangleright z \text{ is real if and only if } \bar{z} = z.$$

## Remark

If  $z = a + bi$ , then



## Properties of the Conjugate

Let  $z$  and  $w$  be complex numbers.

$$\blacktriangleright \overline{z \pm w} = \bar{z} \pm \bar{w}.$$

$$\blacktriangleright \overline{(zw)} = \bar{z} \bar{w}.$$

$$\blacktriangleright \overline{(\bar{z})} = z.$$

$$\blacktriangleright \overline{\left(\frac{z}{w}\right)} = \frac{\bar{z}}{\bar{w}}.$$

$$\blacktriangleright z \text{ is real if and only if } \bar{z} = z.$$

## Remark

If  $z = a + bi$ , then

$$z\bar{z} =$$

## Properties of the Conjugate

Let  $z$  and  $w$  be complex numbers.

$$\blacktriangleright \overline{z \pm w} = \bar{z} \pm \bar{w}.$$

$$\blacktriangleright \overline{(zw)} = \bar{z} \bar{w}.$$

$$\blacktriangleright \overline{(\bar{z})} = z.$$

$$\blacktriangleright \overline{\left(\frac{z}{w}\right)} = \frac{\bar{z}}{\bar{w}}.$$

$$\blacktriangleright z \text{ is real if and only if } \bar{z} = z.$$

## Remark

If  $z = a + bi$ , then

$$z\bar{z} = (a + bi)(a - bi) =$$

## Properties of the Conjugate

Let  $z$  and  $w$  be complex numbers.

$$\blacktriangleright \overline{z \pm w} = \bar{z} \pm \bar{w}.$$

$$\blacktriangleright \overline{(zw)} = \bar{z} \bar{w}.$$

$$\blacktriangleright \overline{(\bar{z})} = z.$$

$$\blacktriangleright \overline{\left(\frac{z}{w}\right)} = \frac{\bar{z}}{\bar{w}}.$$

$$\blacktriangleright z \text{ is real if and only if } \bar{z} = z.$$

## Remark

If  $z = a + bi$ , then

$$z\bar{z} = (a + bi)(a - bi) = a^2 + b^2.$$

## Division of Complex Numbers

## Division of Complex Numbers

Let  $z = a + bi$  and  $w = c + di$  be complex numbers.

## Division of Complex Numbers

Let  $z = a + bi$  and  $w = c + di$  be complex numbers. Suppose that  $c, d$  are not both zero.

## Division of Complex Numbers

Let  $z = a + bi$  and  $w = c + di$  be complex numbers. Suppose that  $c, d$  are not both zero. Then the **quotient  $z$  divided by  $w$**  is

$$\frac{z}{w} =$$

## Division of Complex Numbers

Let  $z = a + bi$  and  $w = c + di$  be complex numbers. Suppose that  $c, d$  are not both zero. Then the **quotient  $z$  divided by  $w$**  is

$$\frac{z}{w} = \frac{a + bi}{c + di} =$$



## Division of Complex Numbers

Let  $z = a + bi$  and  $w = c + di$  be complex numbers. Suppose that  $c, d$  are not both zero. Then the **quotient  $z$  divided by  $w$**  is

$$\begin{aligned}\frac{z}{w} &= \frac{a + bi}{c + di} = \frac{a + bi}{c + di} \times \frac{c - di}{c - di} \\ &= \end{aligned}$$

## Division of Complex Numbers

Let  $z = a + bi$  and  $w = c + di$  be complex numbers. Suppose that  $c, d$  are not both zero. Then the **quotient  $z$  divided by  $w$**  is

$$\begin{aligned}\frac{z}{w} &= \frac{a + bi}{c + di} = \frac{a + bi}{c + di} \times \frac{c - di}{c - di} \\ &= \frac{(ac + bd) + (bc - ad)i}{c^2 + d^2} \\ &= \end{aligned}$$

## Division of Complex Numbers

Let  $z = a + bi$  and  $w = c + di$  be complex numbers. Suppose that  $c, d$  are not both zero. Then the **quotient  $z$  divided by  $w$**  is

$$\begin{aligned}\frac{z}{w} = \frac{a + bi}{c + di} &= \frac{a + bi}{c + di} \times \frac{c - di}{c - di} \\ &= \frac{(ac + bd) + (bc - ad)i}{c^2 + d^2} \\ &= \frac{ac + bd}{c^2 + d^2} + \frac{bc - ad}{c^2 + d^2}i.\end{aligned}$$

## Division of Complex Numbers

Let  $z = a + bi$  and  $w = c + di$  be complex numbers. Suppose that  $c, d$  are not both zero. Then the **quotient  $z$  divided by  $w$**  is

$$\begin{aligned}\frac{z}{w} = \frac{a + bi}{c + di} &= \frac{a + bi}{c + di} \times \frac{c - di}{c - di} \\ &= \frac{(ac + bd) + (bc - ad)i}{c^2 + d^2} \\ &= \frac{ac + bd}{c^2 + d^2} + \frac{bc - ad}{c^2 + d^2}i.\end{aligned}$$

The quotient  $\frac{z}{w}$  is obtained by multiplying both top and bottom of  $\frac{z}{w}$  by  $\overline{w}$  and then simplifying the expression.

## Examples

## Examples



$$\frac{1}{i} =$$

## Examples



$$\frac{1}{i} = \frac{1}{i} \times \frac{-i}{-i} = \frac{-i}{-i^2} = -i.$$

## Examples



$$\frac{1}{i} = \frac{1}{i} \times \frac{-i}{-i} = \frac{-i}{-i^2} = -i.$$



$$\frac{2 - i}{3 + 4i} =$$



## Examples



$$\frac{1}{i} = \frac{1}{i} \times \frac{-i}{-i} = \frac{-i}{-i^2} = -i.$$



$$\frac{2-i}{3+4i} = \frac{2-i}{3+4i} \times \frac{3-4i}{3-4i} = \frac{(6-4) + (-3-8)i}{3^2+4^2} = \frac{2-11i}{25} = \frac{2}{25} - \frac{11}{25}i.$$

## Examples



$$\frac{1}{i} = \frac{1}{i} \times \frac{-i}{-i} = \frac{-i}{-i^2} = -i.$$



$$\frac{2-i}{3+4i} = \frac{2-i}{3+4i} \times \frac{3-4i}{3-4i} = \frac{(6-4) + (-3-8)i}{3^2+4^2} = \frac{2-11i}{25} = \frac{2}{25} - \frac{11}{25}i.$$



$$\frac{1-2i}{-2+5i} =$$

## Examples



$$\frac{1}{i} = \frac{1}{i} \times \frac{-i}{-i} = \frac{-i}{-i^2} = -i.$$



$$\frac{2-i}{3+4i} = \frac{2-i}{3+4i} \times \frac{3-4i}{3-4i} = \frac{(6-4) + (-3-8)i}{3^2+4^2} = \frac{2-11i}{25} = \frac{2}{25} - \frac{11}{25}i.$$



$$\frac{1-2i}{-2+5i} = \frac{1-2i}{-2+5i} \times \frac{-2-5i}{-2-5i} = \frac{(-2-10) + (4-5)i}{2^2+5^2} = -\frac{12}{29} - \frac{1}{29}i.$$

## The Multiplicative Inverse

Every nonzero complex number  $z = a + bi$  has a unique **multiplicative inverse**  $z^{-1} = \frac{1}{z}$  such that  $zz^{-1} = 1$ , and

## The Multiplicative Inverse

Every nonzero complex number  $z = a + bi$  has a unique **multiplicative inverse**  $z^{-1} = \frac{1}{z}$  such that  $zz^{-1} = 1$ , and

$$\frac{1}{z} =$$

## The Multiplicative Inverse

Every nonzero complex number  $z = a + bi$  has a unique **multiplicative inverse**  $z^{-1} = \frac{1}{z}$  such that  $zz^{-1} = 1$ , and

$$\frac{1}{z} = \frac{1}{z} \times \frac{\bar{z}}{\bar{z}} =$$

## The Multiplicative Inverse

Every nonzero complex number  $z = a + bi$  has a unique **multiplicative inverse**  $z^{-1} = \frac{1}{z}$  such that  $zz^{-1} = 1$ , and

$$\frac{1}{z} = \frac{1}{z} \times \frac{\bar{z}}{\bar{z}} = \frac{\bar{z}}{z\bar{z}} =$$

## The Multiplicative Inverse

Every nonzero complex number  $z = a + bi$  has a unique **multiplicative inverse**  $z^{-1} = \frac{1}{z}$  such that  $zz^{-1} = 1$ , and

$$\frac{1}{z} = \frac{1}{z} \times \frac{\bar{z}}{\bar{z}} = \frac{\bar{z}}{z\bar{z}} = \frac{a - bi}{a^2 + b^2} =$$



## The Multiplicative Inverse

Every nonzero complex number  $z = a + bi$  has a unique **multiplicative inverse**  $z^{-1} = \frac{1}{z}$  such that  $zz^{-1} = 1$ , and

$$\frac{1}{z} = \frac{1}{z} \times \frac{\bar{z}}{\bar{z}} = \frac{\bar{z}}{z\bar{z}} = \frac{a - bi}{a^2 + b^2} = \frac{a}{a^2 + b^2} - \frac{b}{a^2 + b^2}i.$$

## The Multiplicative Inverse

Every nonzero complex number  $z = a + bi$  has a unique **multiplicative inverse**  $z^{-1} = \frac{1}{z}$  such that  $zz^{-1} = 1$ , and

$$\frac{1}{z} = \frac{1}{z} \times \frac{\bar{z}}{\bar{z}} = \frac{\bar{z}}{z\bar{z}} = \frac{a - bi}{a^2 + b^2} = \frac{a}{a^2 + b^2} - \frac{b}{a^2 + b^2}i.$$

Since  $z$  is nonzero,  $a^2 + b^2 \neq 0$ , so the inverse is defined.

## The Multiplicative Inverse

Every nonzero complex number  $z = a + bi$  has a unique **multiplicative inverse**  $z^{-1} = \frac{1}{z}$  such that  $zz^{-1} = 1$ , and

$$\frac{1}{z} = \frac{1}{z} \times \frac{\bar{z}}{\bar{z}} = \frac{\bar{z}}{z\bar{z}} = \frac{a - bi}{a^2 + b^2} = \frac{a}{a^2 + b^2} - \frac{b}{a^2 + b^2}i.$$

Since  $z$  is nonzero,  $a^2 + b^2 \neq 0$ , so the inverse is defined.

## Example

When  $z = 2 + 6i$ ,  $z^{-1}$  is defined, and

## The Multiplicative Inverse

Every nonzero complex number  $z = a + bi$  has a unique **multiplicative inverse**  $z^{-1} = \frac{1}{z}$  such that  $zz^{-1} = 1$ , and

$$\frac{1}{z} = \frac{1}{z} \times \frac{\bar{z}}{\bar{z}} = \frac{\bar{z}}{z\bar{z}} = \frac{a - bi}{a^2 + b^2} = \frac{a}{a^2 + b^2} - \frac{b}{a^2 + b^2}i.$$

Since  $z$  is nonzero,  $a^2 + b^2 \neq 0$ , so the inverse is defined.

## Example

When  $z = 2 + 6i$ ,  $z^{-1}$  is defined, and

$$\frac{1}{z} = \frac{1}{2 + 6i} =$$

## The Multiplicative Inverse

Every nonzero complex number  $z = a + bi$  has a unique **multiplicative inverse**  $z^{-1} = \frac{1}{z}$  such that  $zz^{-1} = 1$ , and

$$\frac{1}{z} = \frac{1}{z} \times \frac{\bar{z}}{\bar{z}} = \frac{\bar{z}}{z\bar{z}} = \frac{a - bi}{a^2 + b^2} = \frac{a}{a^2 + b^2} - \frac{b}{a^2 + b^2}i.$$

Since  $z$  is nonzero,  $a^2 + b^2 \neq 0$ , so the inverse is defined.

## Example

When  $z = 2 + 6i$ ,  $z^{-1}$  is defined, and

$$\frac{1}{z} = \frac{1}{2 + 6i} = \frac{1}{2 + 6i} \times \frac{2 - 6i}{2 - 6i} =$$

## The Multiplicative Inverse

Every nonzero complex number  $z = a + bi$  has a unique **multiplicative inverse**  $z^{-1} = \frac{1}{z}$  such that  $zz^{-1} = 1$ , and

$$\frac{1}{z} = \frac{1}{z} \times \frac{\bar{z}}{\bar{z}} = \frac{\bar{z}}{z\bar{z}} = \frac{a - bi}{a^2 + b^2} = \frac{a}{a^2 + b^2} - \frac{b}{a^2 + b^2}i.$$

Since  $z$  is nonzero,  $a^2 + b^2 \neq 0$ , so the inverse is defined.

### Example

When  $z = 2 + 6i$ ,  $z^{-1}$  is defined, and

$$\frac{1}{z} = \frac{1}{2 + 6i} = \frac{1}{2 + 6i} \times \frac{2 - 6i}{2 - 6i} = \frac{2 - 6i}{2^2 + 6^2} =$$

## The Multiplicative Inverse

Every nonzero complex number  $z = a + bi$  has a unique **multiplicative inverse**  $z^{-1} = \frac{1}{z}$  such that  $zz^{-1} = 1$ , and

$$\frac{1}{z} = \frac{1}{z} \times \frac{\bar{z}}{\bar{z}} = \frac{\bar{z}}{z\bar{z}} = \frac{a - bi}{a^2 + b^2} = \frac{a}{a^2 + b^2} - \frac{b}{a^2 + b^2}i.$$

Since  $z$  is nonzero,  $a^2 + b^2 \neq 0$ , so the inverse is defined.

### Example

When  $z = 2 + 6i$ ,  $z^{-1}$  is defined, and

$$\frac{1}{z} = \frac{1}{2 + 6i} = \frac{1}{2 + 6i} \times \frac{2 - 6i}{2 - 6i} = \frac{2 - 6i}{2^2 + 6^2} = \frac{2 - 6i}{40} =$$

## The Multiplicative Inverse

Every nonzero complex number  $z = a + bi$  has a unique **multiplicative inverse**  $z^{-1} = \frac{1}{z}$  such that  $zz^{-1} = 1$ , and

$$\frac{1}{z} = \frac{1}{z} \times \frac{\bar{z}}{\bar{z}} = \frac{\bar{z}}{z\bar{z}} = \frac{a - bi}{a^2 + b^2} = \frac{a}{a^2 + b^2} - \frac{b}{a^2 + b^2}i.$$

Since  $z$  is nonzero,  $a^2 + b^2 \neq 0$ , so the inverse is defined.

### Example

When  $z = 2 + 6i$ ,  $z^{-1}$  is defined, and

$$\frac{1}{z} = \frac{1}{2 + 6i} = \frac{1}{2 + 6i} \times \frac{2 - 6i}{2 - 6i} = \frac{2 - 6i}{2^2 + 6^2} = \frac{2 - 6i}{40} = \frac{1}{20} - \frac{3}{20}i.$$



## The Multiplicative Inverse

Every nonzero complex number  $z = a + bi$  has a unique **multiplicative inverse**  $z^{-1} = \frac{1}{z}$  such that  $zz^{-1} = 1$ , and

$$\frac{1}{z} = \frac{1}{z} \times \frac{\bar{z}}{\bar{z}} = \frac{\bar{z}}{z\bar{z}} = \frac{a - bi}{a^2 + b^2} = \frac{a}{a^2 + b^2} - \frac{b}{a^2 + b^2}i.$$

Since  $z$  is nonzero,  $a^2 + b^2 \neq 0$ , so the inverse is defined.

## Example

When  $z = 2 + 6i$ ,  $z^{-1}$  is defined, and

$$\frac{1}{z} = \frac{1}{2 + 6i} = \frac{1}{2 + 6i} \times \frac{2 - 6i}{2 - 6i} = \frac{2 - 6i}{2^2 + 6^2} = \frac{2 - 6i}{40} = \frac{1}{20} - \frac{3}{20}i.$$

You can always check that  $zz^{-1} = 1$ .

Copyright

Complex Numbers

**Modulus**

Complex Numbers in Polar Form

Roots of Complex Numbers

Roots of Unity

The Quadratic Formula



# Modulus

## Definition

The **absolute value** or **modulus** of a complex number  $z = a + bi$  is

$$|z| = \sqrt{a^2 + b^2}$$

# Modulus

## Definition

The **absolute value** or **modulus** of a complex number  $z = a + bi$  is

$$|z| = \sqrt{a^2 + b^2}$$

Note that this is consistent with the definition of the absolute value of a real number.

# Modulus


## Definition

The **absolute value** or **modulus** of a complex number  $z = a + bi$  is

$$|z| = \sqrt{a^2 + b^2}$$

Note that this is consistent with the definition of the absolute value of a real number.

Geometrically,  $|z| = \sqrt{a^2 + b^2}$  is the distance from  $z$  to the origin.



## Examples

## Examples

1.  $|-3 + 4i| =$



## Examples

1.  $|-3 + 4i| = \sqrt{3^2 + 4^2} =$

## Examples

1.  $|-3 + 4i| = \sqrt{3^2 + 4^2} = \sqrt{25} =$

## Examples

1.  $|-3 + 4i| = \sqrt{3^2 + 4^2} = \sqrt{25} = 5.$

## Examples

1.  $|-3 + 4i| = \sqrt{3^2 + 4^2} = \sqrt{25} = 5.$

2.  $|3 - 2i| =$

## Examples

1.  $|-3 + 4i| = \sqrt{3^2 + 4^2} = \sqrt{25} = 5.$

2.  $|3 - 2i| = \sqrt{3^2 + 2^2} = \sqrt{13}.$

## Examples

1.  $|-3 + 4i| = \sqrt{3^2 + 4^2} = \sqrt{25} = 5.$

2.  $|3 - 2i| = \sqrt{3^2 + 2^2} = \sqrt{13}.$

3.  $|i| =$

## Examples

1.  $|-3 + 4i| = \sqrt{3^2 + 4^2} = \sqrt{25} = 5.$

2.  $|3 - 2i| = \sqrt{3^2 + 2^2} = \sqrt{13}.$

3.  $|i| = \sqrt{1^2} = 1.$

## Theorem ( Properties of the Modulus )



## Theorem ( Properties of the Modulus )

Let  $z$  and  $w$  be complex numbers.

## Theorem ( Properties of the Modulus )

Let  $z$  and  $w$  be complex numbers.

1.  $z \cdot \bar{z} = |z|^2$ .

## Theorem ( Properties of the Modulus )

Let  $z$  and  $w$  be complex numbers.

1.  $z \cdot \bar{z} = |z|^2.$

2.  $\frac{1}{z} = \frac{\bar{z}}{|z|^2}.$

## Theorem ( Properties of the Modulus )

Let  $z$  and  $w$  be complex numbers.

1.  $z \cdot \bar{z} = |z|^2$ .
2.  $\frac{1}{z} = \frac{\bar{z}}{|z|^2}$ .
3.  $|z| \geq 0$  for all  $z$ .

## Theorem ( Properties of the Modulus )

Let  $z$  and  $w$  be complex numbers.

1.  $z \cdot \bar{z} = |z|^2$ .
2.  $\frac{1}{z} = \frac{\bar{z}}{|z|^2}$ .
3.  $|z| \geq 0$  for all  $z$ .
4.  $|z| = 0$  if and only if  $z = 0$ .

## Theorem ( Properties of the Modulus )

Let  $z$  and  $w$  be complex numbers.

1.  $z \cdot \bar{z} = |z|^2$ .
2.  $\frac{1}{z} = \frac{\bar{z}}{|z|^2}$ .
3.  $|z| \geq 0$  for all  $z$ .
4.  $|z| = 0$  if and only if  $z = 0$ .
5.  $|zw| = |z| |w|$ .

## Theorem ( Properties of the Modulus )

Let  $z$  and  $w$  be complex numbers.

1.  $z \cdot \bar{z} = |z|^2$ .
2.  $\frac{1}{z} = \frac{\bar{z}}{|z|^2}$ .
3.  $|z| \geq 0$  for all  $z$ .
4.  $|z| = 0$  if and only if  $z = 0$ .
5.  $|zw| = |z| |w|$ .
6.  $\left| \frac{z}{w} \right| = \frac{|z|}{|w|}$ .

## Theorem ( Properties of the Modulus )

Let  $z$  and  $w$  be complex numbers.

1.  $z \cdot \bar{z} = |z|^2$ .
2.  $\frac{1}{z} = \frac{\bar{z}}{|z|^2}$ .
3.  $|z| \geq 0$  for all  $z$ .
4.  $|z| = 0$  if and only if  $z = 0$ .
5.  $|zw| = |z| |w|$ .
6.  $\left| \frac{z}{w} \right| = \frac{|z|}{|w|}$ .
7. **The Triangle Inequality**  
 $|z + w| \leq |z| + |w|$ .




Example ( The Triangle Inequality: Geometrically )

If  $z = a + bi$  and  $w = c + di$ , then  $|z - w| = \sqrt{(a - c)^2 + (b - d)^2}$ .

### Example ( The Triangle Inequality: Geometrically )

If  $z = a + bi$  and  $w = c + di$ , then  $|z - w| = \sqrt{(a - c)^2 + (b - d)^2}$ .




This shows that the **distance** between  $z$  and  $w$  in the complex plane is just the absolute value of their difference.

### Example (continued)

Now consider the points  $z$ ,  $z + w$ , and the origin  $0$  in the complex plane.


### Example (continued)

Now consider the points  $z$ ,  $z + w$ , and the origin  $0$  in the complex plane.



## Example (continued)


Now consider the points  $z$ ,  $z + w$ , and the origin  $0$  in the complex plane.



The triangle formed by these points has sides of length  $|z|$ ,  $|z + w|$  and  $|w|$  (the absolute value of the difference between  $z + w$  and  $z$ ).

### Example (continued)

Now consider the points  $z$ ,  $z + w$ , and the origin  $0$  in the complex plane.



The triangle formed by these points has sides of length  $|z|$ , and  $|z + w|$  and  $|w|$  (the absolute value of the difference between  $z + w$  and  $z$ ).

Since the length of **any** side of a triangle is at most the sum of the lengths of the other two sides, we get  $|z + w| \leq |z| + |w|$ .

Copyright

Complex Numbers

Modulus

**Complex Numbers in Polar Form**

Roots of Complex Numbers

Roots of Unity


The Quadratic Formula






## Complex Numbers in Polar Form

Suppose  $z = a + bi$ , and let  $r = |z| = \sqrt{a^2 + b^2}$ . Then  $r$  is the distance from  $z$  to the origin. Denote by  $\theta$  the angle that the line through 0 and  $z$  makes with the positive x-axis (measured clockwise).



## Complex Numbers in Polar Form


Suppose  $z = a + bi$ , and let  $r = |z| = \sqrt{a^2 + b^2}$ . Then  $r$  is the distance from  $z$  to the origin. Denote by  $\theta$  the angle that the line through 0 and  $z$  makes with the positive x-axis (measured clockwise).



Then  $\theta$  is an angle defined by  $\cos \theta = \frac{a}{r}$  and  $\sin \theta = \frac{b}{r}$ , so  
$$z = r \cos \theta + r \sin \theta i = r(\cos \theta + i \sin \theta).$$

## Complex Numbers in Polar Form

Suppose  $z = a + bi$ , and let  $r = |z| = \sqrt{a^2 + b^2}$ . Then  $r$  is the distance from  $z$  to the origin. Denote by  $\theta$  the angle that the line through 0 and  $z$  makes with the positive x-axis (measured clockwise).



Then  $\theta$  is an angle defined by  $\cos \theta = \frac{a}{r}$  and  $\sin \theta = \frac{b}{r}$ , so  
$$z = r \cos \theta + r \sin \theta i = r(\cos \theta + i \sin \theta).$$

$\theta$  is called an **argument of  $z$** , and is denoted  $\arg z$ .

### Definition (Polar Form of a Complex Number)

Let  $z$  be a complex number with  $|z| = r$  and  $\arg z = \theta$ . Then

$$z = re^{i\theta} = r(\cos \theta + i \sin \theta)$$

is called a polar form of  $z$ .

### Definition (Polar Form of a Complex Number)

Let  $z$  be a complex number with  $|z| = r$  and  $\arg z = \theta$ . Then

$$z = re^{i\theta} = r(\cos \theta + i \sin \theta)$$

is called a polar form of  $z$ .

### Remark

Since  $\arg z$  is not unique, we do not write the polar form of  $z$ .

### Definition (Polar Form of a Complex Number)

Let  $z$  be a complex number with  $|z| = r$  and  $\arg z = \theta$ . Then

$$z = re^{i\theta} = r(\cos \theta + i \sin \theta)$$

is called a **polar form** of  $z$ .

### Remark

Since  $\arg z$  is not unique, we do not write **the** polar form of  $z$ .

### Definition

Let  $z$  be a complex number with  $|z| = r$ . The **principal argument** of  $z$  is the unique angle  $\theta = \arg z$  (measured in radians) such that

$$-\pi < \theta \leq \pi.$$

### Example

Find the polar form for the number  $z = 1$ .

### Example

Find the polar form for the number  $z = 1$ .

### Solution

To convert  $z$  to polar form, we need to find  $r$  and  $\theta$  so that  $1 = re^{i\theta}$ .



### Example

Find the polar form for the number  $z = 1$ .

### Solution

To convert  $z$  to polar form, we need to find  $r$  and  $\theta$  so that  $1 = re^{i\theta}$ . Now  $r = |z| = \sqrt{1^2} = 1$ ,

### Example

Find the polar form for the number  $z = 1$ .

### Solution

To convert  $z$  to polar form, we need to find  $r$  and  $\theta$  so that  $1 = re^{i\theta}$ . Now  $r = |z| = \sqrt{1^2} = 1$ , and  $\theta = 0$  is an argument for  $z = 1$ .

### Example

Find the polar form for the number  $z = 1$ .

### Solution

To convert  $z$  to polar form, we need to find  $r$  and  $\theta$  so that  $1 = re^{i\theta}$ . Now  $r = |z| = \sqrt{1^2} = 1$ , and  $\theta = 0$  is an argument for  $z = 1$ . However, we may also write

$$1 = e^{2\pi i}, 1 = e^{-2\pi i}, e^{4\pi i}, e^{6\pi i}, \dots$$

### Example

Find the polar form for the number  $z = 1$ .

### Solution

To convert  $z$  to polar form, we need to find  $r$  and  $\theta$  so that  $1 = re^{i\theta}$ . Now  $r = |z| = \sqrt{1^2} = 1$ , and  $\theta = 0$  is an argument for  $z = 1$ . However, we may also write

$$1 = e^{2\pi i}, 1 = e^{-2\pi i}, e^{4\pi i}, e^{6\pi i}, \dots$$

Since sine and cosine have periodicity  $2\pi$ , we may add (or subtract) multiples of  $2\pi$  to any argument.

### Example ( Converting to Polar Form )

Convert the number  $z = -2 + 2\sqrt{3}i$  to polar form.

### Example ( Converting to Polar Form )

Convert the number  $z = -2 + 2\sqrt{3}i$  to polar form.

### Solution

To convert  $z$  to polar form, we need to find  $r$  and  $\theta$  so that  $-2 + 2\sqrt{3}i = re^{i\theta}$ .

### Example ( Converting to Polar Form )

Convert the number  $z = -2 + 2\sqrt{3}i$  to polar form.

#### Solution

To convert  $z$  to polar form, we need to find  $r$  and  $\theta$  so that  $-2 + 2\sqrt{3}i = re^{i\theta}$ . Since  $r = |z|$ ,

$$r = \sqrt{(-2)^2 + (2\sqrt{3})^2}$$

### Example ( Converting to Polar Form )

Convert the number  $z = -2 + 2\sqrt{3}i$  to polar form.

#### Solution

To convert  $z$  to polar form, we need to find  $r$  and  $\theta$  so that  $-2 + 2\sqrt{3}i = re^{i\theta}$ . Since  $r = |z|$ ,

$$r = \sqrt{(-2)^2 + (2\sqrt{3})^2} = \sqrt{4 + 4(3)}$$



### Example ( Converting to Polar Form )

Convert the number  $z = -2 + 2\sqrt{3}i$  to polar form.

#### Solution

To convert  $z$  to polar form, we need to find  $r$  and  $\theta$  so that  $-2 + 2\sqrt{3}i = re^{i\theta}$ . Since  $r = |z|$ ,

$$r = \sqrt{(-2)^2 + (2\sqrt{3})^2} = \sqrt{4 + 4(3)} = \sqrt{16}$$

### Example ( Converting to Polar Form )

Convert the number  $z = -2 + 2\sqrt{3}i$  to polar form.

#### Solution

To convert  $z$  to polar form, we need to find  $r$  and  $\theta$  so that  $-2 + 2\sqrt{3}i = re^{i\theta}$ . Since  $r = |z|$ ,

$$r = \sqrt{(-2)^2 + (2\sqrt{3})^2} = \sqrt{4 + 4(3)} = \sqrt{16} = 4.$$

### Example ( Converting to Polar Form )

Convert the number  $z = -2 + 2\sqrt{3}i$  to polar form.

#### Solution

To convert  $z$  to polar form, we need to find  $r$  and  $\theta$  so that  $-2 + 2\sqrt{3}i = re^{i\theta}$ . Since  $r = |z|$ ,

$$r = \sqrt{(-2)^2 + (2\sqrt{3})^2} = \sqrt{4 + 4(3)} = \sqrt{16} = 4.$$

There are two approaches to finding an argument,  $\theta$ .

## Example ( Converting to Polar Form )


Convert the number  $z = -2 + 2\sqrt{3}i$  to polar form.

### Solution

To convert  $z$  to polar form, we need to find  $r$  and  $\theta$  so that  $-2 + 2\sqrt{3}i = re^{i\theta}$ . Since  $r = |z|$ ,

$$r = \sqrt{(-2)^2 + (2\sqrt{3})^2} = \sqrt{4 + 4(3)} = \sqrt{16} = 4.$$

There are two approaches to finding an argument,  $\theta$ . One is to graph  $-2 + 2\sqrt{3}i$  in the complex plane.



### Solution (continued)

The triangle sitting on the negative half of the real axis has sides of length 2,  $2\sqrt{3}$ , and 4;

### Solution (continued)

The triangle sitting on the negative half of the real axis has sides of length 2,  $2\sqrt{3}$ , and 4; you should recognize this as a right triangle whose other two angles measure  $\frac{\pi}{3}$  and  $\frac{\pi}{6}$ .

### Solution (continued)

The triangle sitting on the negative half of the real axis has sides of length 2,  $2\sqrt{3}$ , and 4; you should recognize this as a right triangle whose other two angles measure  $\frac{\pi}{3}$  and  $\frac{\pi}{6}$ . From this, we see that  $\theta = \frac{2\pi}{3}$  is an argument of  $z$ .

### Solution (continued)

The triangle sitting on the negative half of the real axis has sides of length 2,  $2\sqrt{3}$ , and 4; you should recognize this as a right triangle whose other two angles measure  $\frac{\pi}{3}$  and  $\frac{\pi}{6}$ . From this, we see that  $\theta = \frac{2\pi}{3}$  is an argument of  $z$ . Therefore,  $z$  can be written in polar form as  $z = 4e^{i(2\pi/3)}$ .



### Solution (continued)

The triangle sitting on the negative half of the real axis has sides of length 2,  $2\sqrt{3}$ , and 4; you should recognize this as a right triangle whose other two angles measure  $\frac{\pi}{3}$  and  $\frac{\pi}{6}$ . From this, we see that  $\theta = \frac{2\pi}{3}$  is an argument of  $z$ .

Therefore,  $z$  can be written in polar form as  $z = 4e^{i(2\pi/3)}$ .

The other approach to finding an argument,  $\theta$ , for  $z = -2 + 2\sqrt{3}i$  is as follows. We've already calculated  $|z| = r = 4$ . By definition,  $\theta$  is an angle satisfying

### Solution (continued)

The triangle sitting on the negative half of the real axis has sides of length 2,  $2\sqrt{3}$ , and 4; you should recognize this as a right triangle whose other two angles measure  $\frac{\pi}{3}$  and  $\frac{\pi}{6}$ . From this, we see that  $\theta = \frac{2\pi}{3}$  is an argument of  $z$ .

Therefore,  $z$  can be written in polar form as  $z = 4e^{i(2\pi/3)}$ .

The other approach to finding an argument,  $\theta$ , for  $z = -2 + 2\sqrt{3}i$  is as follows. We've already calculated  $|z| = r = 4$ . By definition,  $\theta$  is an angle satisfying

$$\cos \theta = \frac{-2}{4}$$

### Solution (continued)

The triangle sitting on the negative half of the real axis has sides of length 2,  $2\sqrt{3}$ , and 4; you should recognize this as a right triangle whose other two angles measure  $\frac{\pi}{3}$  and  $\frac{\pi}{6}$ . From this, we see that  $\theta = \frac{2\pi}{3}$  is an argument of  $z$ .

Therefore,  $z$  can be written in polar form as  $z = 4e^{i(2\pi/3)}$ .

The other approach to finding an argument,  $\theta$ , for  $z = -2 + 2\sqrt{3}i$  is as follows. We've already calculated  $|z| = r = 4$ . By definition,  $\theta$  is an angle satisfying

$$\cos \theta = \frac{-2}{4} = -\frac{1}{2}$$

### Solution (continued)

The triangle sitting on the negative half of the real axis has sides of length 2,  $2\sqrt{3}$ , and 4; you should recognize this as a right triangle whose other two angles measure  $\frac{\pi}{3}$  and  $\frac{\pi}{6}$ . From this, we see that  $\theta = \frac{2\pi}{3}$  is an argument of  $z$ .

Therefore,  $z$  can be written in polar form as  $z = 4e^{i(2\pi/3)}$ .

The other approach to finding an argument,  $\theta$ , for  $z = -2 + 2\sqrt{3}i$  is as follows. We've already calculated  $|z| = r = 4$ . By definition,  $\theta$  is an angle satisfying

$$\cos \theta = \frac{-2}{4} = -\frac{1}{2} \quad \text{and} \quad \sin \theta = \frac{2\sqrt{3}}{4}$$

### Solution (continued)

The triangle sitting on the negative half of the real axis has sides of length 2,  $2\sqrt{3}$ , and 4; you should recognize this as a right triangle whose other two angles measure  $\frac{\pi}{3}$  and  $\frac{\pi}{6}$ . From this, we see that  $\theta = \frac{2\pi}{3}$  is an argument of  $z$ .

Therefore,  $z$  can be written in polar form as  $z = 4e^{i(2\pi/3)}$ .

The other approach to finding an argument,  $\theta$ , for  $z = -2 + 2\sqrt{3}i$  is as follows. We've already calculated  $|z| = r = 4$ . By definition,  $\theta$  is an angle satisfying

$$\cos \theta = \frac{-2}{4} = -\frac{1}{2} \quad \text{and} \quad \sin \theta = \frac{2\sqrt{3}}{4} = \frac{\sqrt{3}}{2}.$$


### Solution (continued)

The triangle sitting on the negative half of the real axis has sides of length 2,  $2\sqrt{3}$ , and 4; you should recognize this as a right triangle whose other two angles measure  $\frac{\pi}{3}$  and  $\frac{\pi}{6}$ . From this, we see that  $\theta = \frac{2\pi}{3}$  is an argument of  $z$ .

Therefore,  $z$  can be written in polar form as  $z = 4e^{i(2\pi/3)}$ .

The other approach to finding an argument,  $\theta$ , for  $z = -2 + 2\sqrt{3}i$  is as follows. We've already calculated  $|z| = r = 4$ . By definition,  $\theta$  is an angle satisfying

$$\cos \theta = \frac{-2}{4} = -\frac{1}{2} \quad \text{and} \quad \sin \theta = \frac{2\sqrt{3}}{4} = \frac{\sqrt{3}}{2}.$$

By graphing the point  $(-\frac{1}{2}, \frac{\sqrt{3}}{2})$ , we again determine that  $\theta = \frac{2\pi}{3}$ , and thus  $z$  can be written in polar form as  $z = 4e^{i(2\pi/3)}$ . 

### Problem

Convert each of the following complex numbers to polar form.

1.  $3i$

## Problem

Convert each of the following complex numbers to polar form.

1.  $3i = 3e^{(\pi/2)i}$ .



## Problem

Convert each of the following complex numbers to polar form.

1.  $3i = 3e^{(\pi/2)i}$ .

2.  $-1 - i$

## Problem

Convert each of the following complex numbers to polar form.

1.  $3i = 3e^{(\pi/2)i}$ .

2.  $-1 - i = \sqrt{2}e^{-(3\pi/4)i}$

## Problem

Convert each of the following complex numbers to polar form.

1.  $3i = 3e^{(\pi/2)i}$ .

2.  $-1 - i = \sqrt{2}e^{-(3\pi/4)i} = \sqrt{2}e^{(5\pi/4)i}$ .

## Problem

Convert each of the following complex numbers to polar form.

1.  $3i = 3e^{(\pi/2)i}$ .

2.  $-1 - i = \sqrt{2}e^{-(3\pi/4)i} = \sqrt{2}e^{(5\pi/4)i}$ .

3.  $\sqrt{3} - i$

## Problem

Convert each of the following complex numbers to polar form.

1.  $3i = 3e^{(\pi/2)i}$ .
2.  $-1 - i = \sqrt{2}e^{-(3\pi/4)i} = \sqrt{2}e^{(5\pi/4)i}$ .
3.  $\sqrt{3} - i = 2e^{-(\pi/6)i}$ .

## Problem

Convert each of the following complex numbers to polar form.

1.  $3i = 3e^{(\pi/2)i}$ .
2.  $-1 - i = \sqrt{2}e^{-(3\pi/4)i} = \sqrt{2}e^{(5\pi/4)i}$ .
3.  $\sqrt{3} - i = 2e^{-(\pi/6)i}$ .
4.  $\sqrt{3} + 3i$

## Problem

Convert each of the following complex numbers to polar form.

1.  $3i = 3e^{(\pi/2)i}$ .
2.  $-1 - i = \sqrt{2}e^{-(3\pi/4)i} = \sqrt{2}e^{(5\pi/4)i}$ .
3.  $\sqrt{3} - i = 2e^{-(\pi/6)i}$ .
4.  $\sqrt{3} + 3i = 2\sqrt{3}e^{(\pi/3)i}$ .

Problem ( Converting from Polar Form to Cartesian form )

Let  $z = 2e^{2\pi i/3}$ . Write  $z$  in the form  $z = a + bi$  (this is called **Cartesian form** or **Standard form**).



Problem ( Converting from Polar Form to Cartesian form )

Let  $z = 2e^{2\pi i/3}$ . Write  $z$  in the form  $z = a + bi$  (this is called **Cartesian form** or **Standard form**).

Solution

First, remember that  $e^{i\theta} = \cos \theta + i \sin \theta$ ,

Problem ( Converting from Polar Form to Cartesian form )

Let  $z = 2e^{2\pi i/3}$ . Write  $z$  in the form  $z = a + bi$  (this is called **Cartesian form** or **Standard form**).

Solution

First, remember that  $e^{i\theta} = \cos \theta + i \sin \theta$ , and thus

$$e^{2\pi i/3} = \cos(2\pi/3) + i \sin(2\pi/3)$$

Problem ( Converting from Polar Form to Cartesian form )

Let  $z = 2e^{2\pi i/3}$ . Write  $z$  in the form  $z = a + bi$  (this is called **Cartesian form** or **Standard form**).

Solution

First, remember that  $e^{i\theta} = \cos \theta + i \sin \theta$ , and thus

$$\begin{aligned} e^{2\pi i/3} &= \cos(2\pi/3) + i \sin(2\pi/3) \\ &= -\frac{1}{2} + i \frac{\sqrt{3}}{2}. \end{aligned}$$

### Problem ( Converting from Polar Form to Cartesian form )

Let  $z = 2e^{2\pi i/3}$ . Write  $z$  in the form  $z = a + bi$  (this is called **Cartesian form** or **Standard form**).

### Solution

First, remember that  $e^{i\theta} = \cos \theta + i \sin \theta$ , and thus

$$\begin{aligned} e^{2\pi i/3} &= \cos(2\pi/3) + i \sin(2\pi/3) \\ &= -\frac{1}{2} + i \frac{\sqrt{3}}{2}. \end{aligned}$$

Therefore

$$z = 2e^{2\pi i/3}$$

### Problem ( Converting from Polar Form to Cartesian form )

Let  $z = 2e^{2\pi i/3}$ . Write  $z$  in the form  $z = a + bi$  (this is called **Cartesian form** or **Standard form**).

### Solution

First, remember that  $e^{i\theta} = \cos \theta + i \sin \theta$ , and thus

$$\begin{aligned} e^{2\pi i/3} &= \cos(2\pi/3) + i \sin(2\pi/3) \\ &= -\frac{1}{2} + i \frac{\sqrt{3}}{2}. \end{aligned}$$

Therefore

$$z = 2e^{2\pi i/3} = 2 \left( -\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)$$

### Problem ( Converting from Polar Form to Cartesian form )

Let  $z = 2e^{2\pi i/3}$ . Write  $z$  in the form  $z = a + bi$  (this is called **Cartesian form** or **Standard form**).

### Solution

First, remember that  $e^{i\theta} = \cos \theta + i \sin \theta$ , and thus

$$\begin{aligned} e^{2\pi i/3} &= \cos(2\pi/3) + i \sin(2\pi/3) \\ &= -\frac{1}{2} + i \frac{\sqrt{3}}{2}. \end{aligned}$$

Therefore

$$\begin{aligned} z = 2e^{2\pi i/3} &= 2 \left( -\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) \\ &= -1 + \sqrt{3}i. \end{aligned}$$

### Problem

Express each of the following complex numbers in Cartesian form.

1.  $3e^{-\pi i}$

### Problem

Express each of the following complex numbers in Cartesian form.

1.  $3e^{-\pi i} = -3$



## Problem

Express each of the following complex numbers in Cartesian form.

1.  $3e^{-\pi i} = -3$

2.  $2e^{3\pi i/4}$

## Problem

Express each of the following complex numbers in Cartesian form.

1.  $3e^{-\pi i} = -3$

2.  $2e^{3\pi i/4} = -\sqrt{2} + i\sqrt{2}$

## Problem

Express each of the following complex numbers in Cartesian form.

1.  $3e^{-\pi i} = -3$

2.  $2e^{3\pi i/4} = -\sqrt{2} + i\sqrt{2}$

3.  $2\sqrt{3}e^{-2\pi i/6}$

## Problem

Express each of the following complex numbers in Cartesian form.

1.  $3e^{-\pi i} = -3$

2.  $2e^{3\pi i/4} = -\sqrt{2} + i\sqrt{2}$

3.  $2\sqrt{3}e^{-2\pi i/6} = \sqrt{3} - 3i$

Problems involving multiplication of complex numbers can often be solved more easily by using polar forms of the complex numbers.

Problems involving multiplication of complex numbers can often be solved more easily by using polar forms of the complex numbers.

### Theorem

If  $z_1 = r_1 e^{i\theta_1}$  and  $z_2 = r_2 e^{i\theta_2}$  are complex numbers, then

$$z_1 z_2 = r_1 r_2 e^{i(\theta_1 + \theta_2)}$$

Problems involving multiplication of complex numbers can often be solved more easily by using polar forms of the complex numbers.

### Theorem

If  $z_1 = r_1 e^{i\theta_1}$  and  $z_2 = r_2 e^{i\theta_2}$  are complex numbers, then

$$z_1 z_2 = r_1 r_2 e^{i(\theta_1 + \theta_2)}$$

### Theorem (De Moivre's Theorem)

If  $\theta$  is any angle and  $n$  is a positive integer,

$$\left( e^{i\theta} \right)^n = e^{in\theta}.$$

Problems involving multiplication of complex numbers can often be solved more easily by using polar forms of the complex numbers.

### Theorem

If  $z_1 = r_1 e^{i\theta_1}$  and  $z_2 = r_2 e^{i\theta_2}$  are complex numbers, then

$$z_1 z_2 = r_1 r_2 e^{i(\theta_1 + \theta_2)}$$

### Theorem (De Moivre's Theorem)

If  $\theta$  is any angle and  $n$  is a positive integer,

$$\left( e^{i\theta} \right)^n = e^{in\theta}.$$

As an immediate consequence of De Moivre's Theorem, we have that for any real number  $r > 0$  and any positive integer  $n$ ,

$$\begin{aligned} (r e^{i\theta})^n &= r^n e^{in\theta} \\ (r (\cos \theta + i \sin \theta))^n &= r^n (\cos n\theta + i \sin n\theta) \end{aligned}$$





### Problem

Express  $(1 - i)^6(\sqrt{3} + i)^3$  in the form  $a + bi$ .

### Solution

Let  $z = 1 - i = \sqrt{2}e^{-\pi i/4}$  and  $w = \sqrt{3} + i = 2e^{\pi i/6}$ .

## Problem

Express  $(1 - i)^6(\sqrt{3} + i)^3$  in the form  $a + bi$ .

## Solution

Let  $z = 1 - i = \sqrt{2}e^{-\pi i/4}$  and  $w = \sqrt{3} + i = 2e^{\pi i/6}$ . We want to compute  $z^6 w^3$ .

## Problem

Express  $(1 - i)^6(\sqrt{3} + i)^3$  in the form  $a + bi$ .

## Solution

Let  $z = 1 - i = \sqrt{2}e^{-\pi i/4}$  and  $w = \sqrt{3} + i = 2e^{\pi i/6}$ . We want to compute  $z^6 w^3$ .

$$z^6 w^3 = (\sqrt{2}e^{-\pi i/4})^6 (2e^{\pi i/6})^3$$

## Problem

Express  $(1 - i)^6(\sqrt{3} + i)^3$  in the form  $a + bi$ .

## Solution

Let  $z = 1 - i = \sqrt{2}e^{-\pi i/4}$  and  $w = \sqrt{3} + i = 2e^{\pi i/6}$ . We want to compute  $z^6 w^3$ .

$$\begin{aligned} z^6 w^3 &= (\sqrt{2}e^{-\pi i/4})^6 (2e^{\pi i/6})^3 \\ &= (2^3 e^{-6\pi i/4})(2^3 e^{3\pi i/6}) \end{aligned}$$

## Problem

Express  $(1 - i)^6(\sqrt{3} + i)^3$  in the form  $a + bi$ .

## Solution

Let  $z = 1 - i = \sqrt{2}e^{-\pi i/4}$  and  $w = \sqrt{3} + i = 2e^{\pi i/6}$ . We want to compute  $z^6 w^3$ .

$$\begin{aligned} z^6 w^3 &= (\sqrt{2}e^{-\pi i/4})^6 (2e^{\pi i/6})^3 \\ &= (2^3 e^{-6\pi i/4})(2^3 e^{3\pi i/6}) \\ &= (8e^{-3\pi i/2})(8e^{\pi i/2}) \end{aligned}$$

## Problem

Express  $(1 - i)^6(\sqrt{3} + i)^3$  in the form  $a + bi$ .

## Solution

Let  $z = 1 - i = \sqrt{2}e^{-\pi i/4}$  and  $w = \sqrt{3} + i = 2e^{\pi i/6}$ . We want to compute  $z^6 w^3$ .

$$\begin{aligned} z^6 w^3 &= (\sqrt{2}e^{-\pi i/4})^6 (2e^{\pi i/6})^3 \\ &= (2^3 e^{-6\pi i/4})(2^3 e^{3\pi i/6}) \\ &= (8e^{-3\pi i/2})(8e^{\pi i/2}) \\ &= 64e^{-\pi i} \end{aligned}$$

## Problem

Express  $(1 - i)^6(\sqrt{3} + i)^3$  in the form  $a + bi$ .

## Solution

Let  $z = 1 - i = \sqrt{2}e^{-\pi i/4}$  and  $w = \sqrt{3} + i = 2e^{\pi i/6}$ . We want to compute  $z^6 w^3$ .

$$\begin{aligned} z^6 w^3 &= (\sqrt{2}e^{-\pi i/4})^6 (2e^{\pi i/6})^3 \\ &= (2^3 e^{-6\pi i/4})(2^3 e^{3\pi i/6}) \\ &= (8e^{-3\pi i/2})(8e^{\pi i/2}) \\ &= 64e^{-\pi i} \\ &= 64e^{\pi i} \end{aligned}$$



## Problem

Express  $(1 - i)^6(\sqrt{3} + i)^3$  in the form  $a + bi$ .

## Solution

Let  $z = 1 - i = \sqrt{2}e^{-\pi i/4}$  and  $w = \sqrt{3} + i = 2e^{\pi i/6}$ . We want to compute  $z^6 w^3$ .

$$\begin{aligned} z^6 w^3 &= (\sqrt{2}e^{-\pi i/4})^6 (2e^{\pi i/6})^3 \\ &= (2^3 e^{-6\pi i/4})(2^3 e^{3\pi i/6}) \\ &= (8e^{-3\pi i/2})(8e^{\pi i/2}) \\ &= 64e^{-\pi i} \\ &= 64e^{\pi i} \\ &= 64(\cos \pi + i \sin \pi) \end{aligned}$$

## Problem

Express  $(1 - i)^6(\sqrt{3} + i)^3$  in the form  $a + bi$ .

## Solution

Let  $z = 1 - i = \sqrt{2}e^{-\pi i/4}$  and  $w = \sqrt{3} + i = 2e^{\pi i/6}$ . We want to compute  $z^6w^3$ .

$$\begin{aligned} z^6w^3 &= (\sqrt{2}e^{-\pi i/4})^6(2e^{\pi i/6})^3 \\ &= (2^3e^{-6\pi i/4})(2^3e^{3\pi i/6}) \\ &= (8e^{-3\pi i/2})(8e^{\pi i/2}) \\ &= 64e^{-\pi i} \\ &= 64e^{\pi i} \\ &= 64(\cos \pi + i \sin \pi) \\ &= -64. \end{aligned}$$

## Problem

Express  $\left(\frac{1}{2} - \frac{\sqrt{3}}{2}i\right)^{17}$  in the form  $a + bi$ .

### Problem

Express  $\left(\frac{1}{2} - \frac{\sqrt{3}}{2}i\right)^{17}$  in the form  $a + bi$ .

### Solution

Let  $z = \frac{1}{2} - \frac{\sqrt{3}}{2}i = e^{-\pi i/3}$ .

### Problem

Express  $\left(\frac{1}{2} - \frac{\sqrt{3}}{2}i\right)^{17}$  in the form  $a + bi$ .

### Solution

Let  $z = \frac{1}{2} - \frac{\sqrt{3}}{2}i = e^{-\pi i/3}$ . Then

$$z^{17} = \left(e^{-\pi i/3}\right)^{17}$$

### Problem

Express  $\left(\frac{1}{2} - \frac{\sqrt{3}}{2}i\right)^{17}$  in the form  $a + bi$ .

### Solution

Let  $z = \frac{1}{2} - \frac{\sqrt{3}}{2}i = e^{-\pi i/3}$ . Then

$$\begin{aligned} z^{17} &= \left(e^{-\pi i/3}\right)^{17} \\ &= e^{-17\pi i/3} \end{aligned}$$

### Problem

Express  $\left(\frac{1}{2} - \frac{\sqrt{3}}{2}i\right)^{17}$  in the form  $a + bi$ .

### Solution

Let  $z = \frac{1}{2} - \frac{\sqrt{3}}{2}i = e^{-\pi i/3}$ . Then

$$\begin{aligned} z^{17} &= \left(e^{-\pi i/3}\right)^{17} \\ &= e^{-17\pi i/3} \\ &= e^{\pi i/3} \end{aligned}$$

## Problem

Express  $\left(\frac{1}{2} - \frac{\sqrt{3}}{2}i\right)^{17}$  in the form  $a + bi$ .

## Solution

Let  $z = \frac{1}{2} - \frac{\sqrt{3}}{2}i = e^{-\pi i/3}$ . Then

$$\begin{aligned} z^{17} &= \left(e^{-\pi i/3}\right)^{17} \\ &= e^{-17\pi i/3} \\ &= e^{\pi i/3} \\ &= \cos \frac{\pi}{3} + i \sin \frac{\pi}{3} \end{aligned}$$



## Problem

Express  $\left(\frac{1}{2} - \frac{\sqrt{3}}{2}i\right)^{17}$  in the form  $a + bi$ .

## Solution

Let  $z = \frac{1}{2} - \frac{\sqrt{3}}{2}i = e^{-\pi i/3}$ . Then

$$\begin{aligned} z^{17} &= \left(e^{-\pi i/3}\right)^{17} \\ &= e^{-17\pi i/3} \\ &= e^{\pi i/3} \\ &= \cos \frac{\pi}{3} + i \sin \frac{\pi}{3} \\ &= \frac{1}{2} + \frac{\sqrt{3}}{2}i. \end{aligned}$$

Copyright

Complex Numbers

Modulus

Complex Numbers in Polar Form

**Roots of Complex Numbers**

Roots of Unity

The Quadratic Formula



# Roots of Complex Numbers

## Definition

Let  $z$  and  $q$  be complex numbers, and let  $n$  be a positive integer.

# Roots of Complex Numbers

## Definition

Let  $z$  and  $q$  be complex numbers, and let  $n$  be a positive integer. Then  $z$  is called **an  $n^{\text{th}}$  root of  $q$**  if  $z^n = q$ .

# Roots of Complex Numbers

## Definition

Let  $z$  and  $q$  be complex numbers, and let  $n$  be a positive integer. Then  $z$  is called an  $n^{\text{th}}$  root of  $q$  if  $z^n = q$ .

## De Moivre's Theorem and its implication

If  $\theta$  is any angle and  $n$  is a positive integer,  $(e^{i\theta})^n = e^{in\theta}$ . This implies that for any real number  $r > 0$  and any positive integer  $n$ ,

$$(re^{i\theta})^n = r^n e^{in\theta}.$$

This leads to the following result.

# Roots of Complex Numbers

## Definition

Let  $z$  and  $q$  be complex numbers, and let  $n$  be a positive integer. Then  $z$  is called an  $n^{\text{th}}$  root of  $q$  if  $z^n = q$ .

## De Moivre's Theorem and its implication

If  $\theta$  is any angle and  $n$  is a positive integer,  $(e^{i\theta})^n = e^{in\theta}$ . This implies that for any real number  $r > 0$  and any positive integer  $n$ ,

$$(re^{i\theta})^n = r^n e^{in\theta}.$$

This leads to the following result.

## Corollary

Let  $q$  be a nonzero complex number and  $n$  a positive integer. Then  $z^n = q$  has exactly  $n$  complex solutions, i.e.,  $q$  has exactly  $n$  complex  $n^{\text{th}}$  roots.

### Example

For any positive real number  $a$ ,  $z^2 = a$  has two complex (in this case, real) solutions,  $z = \sqrt{a}$  and  $z = -\sqrt{a}$ . This is equivalent to the statement that  $a$  has two complex (in this case, real) square roots.



## Example

For any positive real number  $a$ ,  $z^2 = a$  has two complex (in this case, real) solutions,  $z = \sqrt{a}$  and  $z = -\sqrt{a}$ . This is equivalent to the statement that  $a$  has two complex (in this case, real) square roots.

- One particular example: 25 has two square roots, 5 and  $-5$ , and these are the two solutions to  $z^2 = 25$ .

## Example

For any positive real number  $a$ ,  $z^2 = a$  has two complex (in this case, real) solution,  $z = \sqrt{a}$  and  $z = -\sqrt{a}$ . This is equivalent to the statement that  $a$  has two complex (in this case, real) square roots.

- ▶ One particular example: 25 has two square roots, 5 and  $-5$ , and these are the two solutions to  $z^2 = 25$ .
- ▶ But we all knew that. A more interesting example is that  $-1$  has no real square roots, but suddenly it has two (complex) square roots,  $i$  and  $-i$ . These are the two (complex) solutions to  $z^2 = 1$ .

### Example ( Cube Roots )

To find the (three) cube roots of  $i$ , we solve the equation  $z^3 = i$ .

### Example ( Cube Roots )

To find the (three) cube roots of  $i$ , we solve the equation  $z^3 = i$ . To do so, we express both  $z$  and  $i$  in polar form: convert  $i$  to polar form, and write  $z = re^{i\theta}$ , giving us  $(re^{i\theta})^3 =$

## Example ( Cube Roots )

To find the (three) cube roots of  $i$ , we solve the equation  $z^3 = i$ . To do so, we express both  $z$  and  $i$  in polar form: convert  $i$  to polar form, and write  $z = re^{i\theta}$ , giving us

$$(re^{i\theta})^3 = e^{\pi i/2}.$$

### Example ( Cube Roots )

To find the (three) cube roots of  $i$ , we solve the equation  $z^3 = i$ . To do so, we express both  $z$  and  $i$  in polar form: convert  $i$  to polar form, and write  $z = re^{i\theta}$ , giving us

$$(re^{i\theta})^3 = e^{\pi i/2}.$$

Thus  $r^3 e^{3i\theta} = 1e^{\pi i/2}$ ,

### Example ( Cube Roots )

To find the (three) cube roots of  $i$ , we solve the equation  $z^3 = i$ . To do so, we express both  $z$  and  $i$  in polar form: convert  $i$  to polar form, and write  $z = re^{i\theta}$ , giving us

$$(re^{i\theta})^3 = e^{\pi i/2}.$$

Thus  $r^3 e^{3i\theta} = 1e^{\pi i/2}$ , implying that  $r^3 = 1$

### Example ( Cube Roots )

To find the (three) cube roots of  $i$ , we solve the equation  $z^3 = i$ . To do so, we express both  $z$  and  $i$  in polar form: convert  $i$  to polar form, and write

$z = re^{i\theta}$ , giving us  $(re^{i\theta})^3 = e^{\pi i/2}$ .

Thus  $r^3 e^{3i\theta} = 1e^{\pi i/2}$ , implying that  $r^3 = 1$  and  $3\theta = \frac{\pi}{2}$ .



## Example ( Cube Roots )

To find the (three) cube roots of  $i$ , we solve the equation  $z^3 = i$ . To do so, we express both  $z$  and  $i$  in polar form: convert  $i$  to polar form, and write  $z = re^{i\theta}$ , giving us

$$(re^{i\theta})^3 = e^{\pi i/2}.$$

Thus  $r^3 e^{3i\theta} = 1e^{\pi i/2}$ , implying that  $r^3 = 1$  and  $3\theta = \frac{\pi}{2}$ .

► Since  $r$  is a real number,  $r^3 = 1$  implies that  $r = 1$ .

## Example ( Cube Roots )

To find the (three) cube roots of  $i$ , we solve the equation  $z^3 = i$ . To do so, we express both  $z$  and  $i$  in polar form: convert  $i$  to polar form, and write  $z = re^{i\theta}$ , giving us

$$(re^{i\theta})^3 = e^{\pi i/2}.$$

Thus  $r^3 e^{3i\theta} = 1e^{\pi i/2}$ , implying that  $r^3 = 1$  and  $3\theta = \frac{\pi}{2}$ .

- ▶ Since  $r$  is a real number,  $r^3 = 1$  implies that  $r = 1$ .
- ▶ The statement  $3\theta = \frac{\pi}{2}$  is not completely correct.

## Example ( Cube Roots )

To find the (three) cube roots of  $i$ , we solve the equation  $z^3 = i$ . To do so, we express both  $z$  and  $i$  in polar form: convert  $i$  to polar form, and write  $z = re^{i\theta}$ , giving us

$$(re^{i\theta})^3 = e^{\pi i/2}.$$

Thus  $r^3 e^{3i\theta} = 1 e^{\pi i/2}$ , implying that  $r^3 = 1$  and  $3\theta = \frac{\pi}{2}$ .

- Since  $r$  is a real number,  $r^3 = 1$  implies that  $r = 1$ .
- The statement  $3\theta = \frac{\pi}{2}$  is **not completely correct**. The problem that arises is that the argument for  $i$ ,  $\frac{\pi}{2}$  is not unique.

## Example ( Cube Roots )

To find the (three) cube roots of  $i$ , we solve the equation  $z^3 = i$ . To do so, we express both  $z$  and  $i$  in polar form: convert  $i$  to polar form, and write  $z = re^{i\theta}$ , giving us

$$(re^{i\theta})^3 = e^{\pi i/2}.$$

Thus  $r^3 e^{3i\theta} = 1e^{\pi i/2}$ , implying that  $r^3 = 1$  and  $3\theta = \frac{\pi}{2}$ .

- ▶ Since  $r$  is a real number,  $r^3 = 1$  implies that  $r = 1$ .
- ▶ The statement  $3\theta = \frac{\pi}{2}$  is **not completely correct**. The problem that arises is that the argument for  $i$ ,  $\frac{\pi}{2}$  is not unique. Instead, we could have written

$$i = e^{5\pi i/2} \text{ or } i = e^{9\pi i/2} \text{ or } i = e^{-3\pi i/2}.$$

## Example ( Cube Roots )

To find the (three) cube roots of  $i$ , we solve the equation  $z^3 = i$ . To do so, we express both  $z$  and  $i$  in polar form: convert  $i$  to polar form, and write  $z = re^{i\theta}$ , giving us

$$(re^{i\theta})^3 = e^{\pi i/2}.$$

Thus  $r^3 e^{3i\theta} = 1e^{\pi i/2}$ , implying that  $r^3 = 1$  and  $3\theta = \frac{\pi}{2}$ .

- ▶ Since  $r$  is a real number,  $r^3 = 1$  implies that  $r = 1$ .
- ▶ The statement  $3\theta = \frac{\pi}{2}$  is **not completely correct**. The problem that arises is that the argument for  $i$ ,  $\frac{\pi}{2}$  is not unique. Instead, we could have written

$$i = e^{5\pi i/2} \text{ or } i = e^{9\pi i/2} \text{ or } i = e^{-3\pi i/2}.$$

In fact, there are infinitely many choices for the argument of  $i$ .

## Example ( Cube Roots )

To find the (three) cube roots of  $i$ , we solve the equation  $z^3 = i$ . To do so, we express both  $z$  and  $i$  in polar form: convert  $i$  to polar form, and write  $z = re^{i\theta}$ , giving us

$$(re^{i\theta})^3 = e^{\pi i/2}.$$

Thus  $r^3 e^{3i\theta} = 1e^{\pi i/2}$ , implying that  $r^3 = 1$  and  $3\theta = \frac{\pi}{2}$ .

- ▶ Since  $r$  is a real number,  $r^3 = 1$  implies that  $r = 1$ .
- ▶ The statement  $3\theta = \frac{\pi}{2}$  is **not completely correct**. The problem that arises is that the argument for  $i$ ,  $\frac{\pi}{2}$  is not unique. Instead, we could have written

$$i = e^{5\pi i/2} \text{ or } i = e^{9\pi i/2} \text{ or } i = e^{-3\pi i/2}.$$

**In fact, there are infinitely many choices for the argument of  $i$ .** The important thing to notice is that any two different arguments differ by a multiple of  $2\pi$ ,

## Example ( Cube Roots )

To find the (three) cube roots of  $i$ , we solve the equation  $z^3 = i$ . To do so, we express both  $z$  and  $i$  in polar form: convert  $i$  to polar form, and write  $z = re^{i\theta}$ , giving us

$$(re^{i\theta})^3 = e^{\pi i/2}.$$

Thus  $r^3 e^{3i\theta} = 1e^{\pi i/2}$ , implying that  $r^3 = 1$  and  $3\theta = \frac{\pi}{2}$ .

- ▶ Since  $r$  is a real number,  $r^3 = 1$  implies that  $r = 1$ .
- ▶ The statement  $3\theta = \frac{\pi}{2}$  is **not completely correct**. The problem that arises is that the argument for  $i$ ,  $\frac{\pi}{2}$  is not unique. Instead, we could have written

$$i = e^{5\pi i/2} \text{ or } i = e^{9\pi i/2} \text{ or } i = e^{-3\pi i/2}.$$

**In fact, there are infinitely many choices for the argument of  $i$ .** The important thing to notice is that any two different arguments differ by a multiple of  $2\pi$ , and thus we may write

$$3\theta = \frac{\pi}{2} + 2\pi\ell, \ell \in \mathbb{Z}.$$

## Example ( Cube Roots )

To find the (three) cube roots of  $i$ , we solve the equation  $z^3 = i$ . To do so, we express both  $z$  and  $i$  in polar form: convert  $i$  to polar form, and write  $z = re^{i\theta}$ , giving us

$$(re^{i\theta})^3 = e^{\pi i/2}.$$

Thus  $r^3 e^{3i\theta} = 1e^{\pi i/2}$ , implying that  $r^3 = 1$  and  $3\theta = \frac{\pi}{2}$ .

- ▶ Since  $r$  is a real number,  $r^3 = 1$  implies that  $r = 1$ .
- ▶ The statement  $3\theta = \frac{\pi}{2}$  is **not completely correct**. The problem that arises is that the argument for  $i$ ,  $\frac{\pi}{2}$  is not unique. Instead, we could have written

$$i = e^{5\pi i/2} \text{ or } i = e^{9\pi i/2} \text{ or } i = e^{-3\pi i/2}.$$

**In fact, there are infinitely many choices for the argument of  $i$ .** The important thing to notice is that any two different arguments differ by a multiple of  $2\pi$ , and thus we may write

$$3\theta = \frac{\pi}{2} + 2\pi\ell, \ell \in \mathbb{Z}.$$

( $\mathbb{Z}$  denotes the set of integers:  $\{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\}$ ).



### Example (continued)

Dividing both sides of  $3\theta = \frac{\pi}{2} + 2\pi\ell$  by 3:

$$\theta = \frac{\pi}{6} + \frac{2}{3}\pi\ell,$$

where  $\ell$  is any integer.

### Example (continued)

Dividing both sides of  $3\theta = \frac{\pi}{2} + 2\pi\ell$  by 3:

$$\theta = \frac{\pi}{6} + \frac{2}{3}\pi\ell,$$

where  $\ell$  is any integer. The Corollary to De Moivre's Theorem tells us that there are only **three** different cube roots.

### Example (continued)

Dividing both sides of  $3\theta = \frac{\pi}{2} + 2\pi\ell$  by 3:

$$\theta = \frac{\pi}{6} + \frac{2}{3}\pi\ell,$$

where  $\ell$  is any integer. The Corollary to De Moivre's Theorem tells us that there are only **three** different cube roots. These are obtained by using  $\ell = 0$ ,  $\ell = 1$ , and  $\ell = 2$ , resulting in three values of  $\theta$ :

### Example (continued)

Dividing both sides of  $3\theta = \frac{\pi}{2} + 2\pi\ell$  by 3:

$$\theta = \frac{\pi}{6} + \frac{2}{3}\pi\ell,$$

where  $\ell$  is any integer. The Corollary to De Moivre's Theorem tells us that there are only **three** different cube roots. These are obtained by using  $\ell = 0$ ,  $\ell = 1$ , and  $\ell = 2$ , resulting in three values of  $\theta$ :

$$\frac{\pi}{6}, \frac{5\pi}{6}, \quad \text{and} \quad \frac{9\pi}{6} = \frac{3\pi}{2}.$$

### Example (continued)

Dividing both sides of  $3\theta = \frac{\pi}{2} + 2\pi\ell$  by 3:

$$\theta = \frac{\pi}{6} + \frac{2}{3}\pi\ell,$$

where  $\ell$  is any integer. The Corollary to De Moivre's Theorem tells us that there are only **three** different cube roots. These are obtained by using  $\ell = 0$ ,  $\ell = 1$ , and  $\ell = 2$ , resulting in three values of  $\theta$ :

$$\frac{\pi}{6}, \frac{5\pi}{6}, \quad \text{and} \quad \frac{9\pi}{6} = \frac{3\pi}{2}.$$

Thus the cube roots of  $i$  are

$$e^{\pi i/6}, e^{5\pi i/6}, \quad \text{and} \quad e^{3\pi i/2}.$$

### Example (continued)

Dividing both sides of  $3\theta = \frac{\pi}{2} + 2\pi\ell$  by 3:

$$\theta = \frac{\pi}{6} + \frac{2}{3}\pi\ell,$$

where  $\ell$  is any integer. The Corollary to De Moivre's Theorem tells us that there are only **three** different cube roots. These are obtained by using  $\ell = 0$ ,  $\ell = 1$ , and  $\ell = 2$ , resulting in three values of  $\theta$ :

$$\frac{\pi}{6}, \frac{5\pi}{6}, \quad \text{and} \quad \frac{9\pi}{6} = \frac{3\pi}{2}.$$

Thus the cube roots of  $i$  are

$$e^{\pi i/6}, e^{5\pi i/6}, \quad \text{and} \quad e^{3\pi i/2}.$$

We now convert these to Cartesian form.

Example (continued)

$$e^{\pi i/6} =$$

Example (continued)

$$e^{\pi i/6} = \frac{\sqrt{3}}{2} + \frac{1}{2}i,$$



Example (continued)

$$e^{\pi i/6} = \frac{\sqrt{3}}{2} + \frac{1}{2}i,$$

$$e^{\pi i/6} =$$

Example (continued)

$$e^{\pi i/6} = \frac{\sqrt{3}}{2} + \frac{1}{2}i,$$

$$e^{\pi i/6} = -\frac{\sqrt{3}}{2} + \frac{1}{2}i,$$

Example (continued)

$$e^{\pi i/6} = \frac{\sqrt{3}}{2} + \frac{1}{2}i,$$

$$e^{\pi i/6} = -\frac{\sqrt{3}}{2} + \frac{1}{2}i,$$

$$e^{3\pi i/2} =$$

Example (continued)

$$e^{\pi i/6} = \frac{\sqrt{3}}{2} + \frac{1}{2}i,$$

$$e^{\pi i/6} = -\frac{\sqrt{3}}{2} + \frac{1}{2}i,$$

$$e^{3\pi i/2} = -i.$$

Example (continued)

$$e^{\pi i/6} = \frac{\sqrt{3}}{2} + \frac{1}{2}i,$$

$$e^{\pi i/6} = -\frac{\sqrt{3}}{2} + \frac{1}{2}i,$$

$$e^{3\pi i/2} = -i.$$

You can check your work by computing the cube of each of these.

Example (continued)

$$e^{\pi i/6} = \frac{\sqrt{3}}{2} + \frac{1}{2}i,$$

$$e^{\pi i/6} = -\frac{\sqrt{3}}{2} + \frac{1}{2}i,$$

$$e^{3\pi i/2} = -i.$$

You can check your work by computing the cube of each of these.

This process is summarized in the following procedure.

## Finding Roots of a Complex Number

Let  $w$  be a complex number. We wish to find the  $n^{\text{th}}$  roots of  $w$ , that is all  $z$  such that  $z^n = w$ .

There are  $n$  distinct  $n^{\text{th}}$  roots and they can be found as follows:.

## Finding Roots of a Complex Number

Let  $w$  be a complex number. We wish to find the  $n^{\text{th}}$  roots of  $w$ , that is all  $z$  such that  $z^n = w$ .

There are  $n$  distinct  $n^{\text{th}}$  roots and they can be found as follows:.

1. Express both  $z$  and  $w$  in polar form  $z = re^{i\theta}$ ,  $w = se^{i\phi}$ . Then  $z^n = w$  becomes:

$$(re^{i\theta})^n = r^n e^{in\theta} = se^{i\phi}$$

We need to solve for  $r$  and  $\theta$ .



## Finding Roots of a Complex Number

Let  $w$  be a complex number. We wish to find the  $n^{\text{th}}$  roots of  $w$ , that is all  $z$  such that  $z^n = w$ .

There are  $n$  distinct  $n^{\text{th}}$  roots and they can be found as follows:.

1. Express both  $z$  and  $w$  in polar form  $z = re^{i\theta}$ ,  $w = se^{i\phi}$ . Then  $z^n = w$  becomes:

$$(re^{i\theta})^n = r^n e^{in\theta} = se^{i\phi}$$

We need to solve for  $r$  and  $\theta$ .

2. Solve the following two equations:

$$r^n = s$$

$$e^{in\theta} = e^{i\phi} \tag{1}$$

Continued

3. The solutions to  $r^n = s$  are given by  $r = \sqrt[n]{s}$ .

## Continued

3. The solutions to  $r^n = s$  are given by  $r = \sqrt[n]{s}$ .

4. The solutions to  $e^{in\theta} = e^{i\phi}$  are given by:

$$n\theta = \phi + 2\pi\ell, \text{ for } \ell = 0, 1, 2, \dots, n-1$$

or

$$\theta = \frac{\phi}{n} + \frac{2}{n}\pi\ell, \text{ for } \ell = 0, 1, 2, \dots, n-1$$

## Continued

3. The solutions to  $r^n = s$  are given by  $r = \sqrt[n]{s}$ .

4. The solutions to  $e^{in\theta} = e^{i\phi}$  are given by:

$$n\theta = \phi + 2\pi\ell, \text{ for } \ell = 0, 1, 2, \dots, n-1$$

or

$$\theta = \frac{\phi}{n} + \frac{2}{n}\pi\ell, \text{ for } \ell = 0, 1, 2, \dots, n-1$$

5. Using the solutions  $r, \theta$  to the equations given in (1) construct the  $n^{\text{th}}$  roots of the form  $z = re^{i\theta}$ .

## Problem

Find all complex numbers  $z$  such that  $z^4 = 2(\sqrt{3}i - 1)$ , and express each in the form  $a + bi$ .

## Problem

Find all complex numbers  $z$  such that  $z^4 = 2(\sqrt{3}i - 1)$ , and express each in the form  $a + bi$ .

## Solution

1. Convert  $2(\sqrt{3}i - 1) = -2 + 2\sqrt{3}i$  to polar form:

## Problem

Find all complex numbers  $z$  such that  $z^4 = 2(\sqrt{3}i - 1)$ , and express each in the form  $a + bi$ .

## Solution

1. Convert  $2(\sqrt{3}i - 1) = -2 + 2\sqrt{3}i$  to polar form:

$$|z^4| = \sqrt{(-2)^2 + (2\sqrt{3})^2} = \sqrt{16} = 4.$$

## Problem

Find all complex numbers  $z$  such that  $z^4 = 2(\sqrt{3}i - 1)$ , and express each in the form  $a + bi$ .

## Solution

1. Convert  $2(\sqrt{3}i - 1) = -2 + 2\sqrt{3}i$  to polar form:

$$|z^4| = \sqrt{(-2)^2 + (2\sqrt{3})^2} = \sqrt{16} = 4.$$

If  $\phi$  is an argument for  $-2 + 2\sqrt{3}i$ , then

$$\cos \phi = \frac{-2}{4} = -\frac{1}{2} \quad \text{and} \quad \sin \phi = \frac{2\sqrt{3}}{4} = \frac{\sqrt{3}}{2}, \text{ so } \phi = \frac{2\pi}{3}.$$



## Problem

Find all complex numbers  $z$  such that  $z^4 = 2(\sqrt{3}i - 1)$ , and express each in the form  $a + bi$ .

## Solution

1. Convert  $2(\sqrt{3}i - 1) = -2 + 2\sqrt{3}i$  to polar form:

$$|z^4| = \sqrt{(-2)^2 + (2\sqrt{3})^2} = \sqrt{16} = 4.$$

If  $\phi$  is an argument for  $-2 + 2\sqrt{3}i$ , then

$$\cos \phi = \frac{-2}{4} = -\frac{1}{2} \quad \text{and} \quad \sin \phi = \frac{2\sqrt{3}}{4} = \frac{\sqrt{3}}{2}, \quad \text{so } \phi = \frac{2\pi}{3}.$$

Thus  $z^4 = 4e^{2\pi i/3}$ .

## Problem

Find all complex numbers  $z$  such that  $z^4 = 2(\sqrt{3}i - 1)$ , and express each in the form  $a + bi$ .

## Solution

1. Convert  $2(\sqrt{3}i - 1) = -2 + 2\sqrt{3}i$  to polar form:

$$|z^4| = \sqrt{(-2)^2 + (2\sqrt{3})^2} = \sqrt{16} = 4.$$

If  $\phi$  is an argument for  $-2 + 2\sqrt{3}i$ , then

$$\cos \phi = \frac{-2}{4} = -\frac{1}{2} \quad \text{and} \quad \sin \phi = \frac{2\sqrt{3}}{4} = \frac{\sqrt{3}}{2}, \quad \text{so } \phi = \frac{2\pi}{3}.$$

Thus  $z^4 = 4e^{2\pi i/3}$ . Let  $z = re^{i\theta}$ .

## Problem

Find all complex numbers  $z$  such that  $z^4 = 2(\sqrt{3}i - 1)$ , and express each in the form  $a + bi$ .

## Solution

1. Convert  $2(\sqrt{3}i - 1) = -2 + 2\sqrt{3}i$  to polar form:

$$|z^4| = \sqrt{(-2)^2 + (2\sqrt{3})^2} = \sqrt{16} = 4.$$

If  $\phi$  is an argument for  $-2 + 2\sqrt{3}i$ , then

$$\cos \phi = \frac{-2}{4} = -\frac{1}{2} \quad \text{and} \quad \sin \phi = \frac{2\sqrt{3}}{4} = \frac{\sqrt{3}}{2}, \quad \text{so } \phi = \frac{2\pi}{3}.$$

Thus  $z^4 = 4e^{2\pi i/3}$ . Let  $z = re^{i\theta}$ .

2. The equation becomes  $r^4 e^{i4\theta} = 4e^{2\pi i/3}$ , so we need to solve

$$\begin{aligned} r^4 &= 4 \\ e^{i4\theta} &= e^{2\pi i/3} \end{aligned}$$

### Solution (continued)

3. Since  $r^4 = 4$ ,  $r^2 = \pm 2$ . But  $r$  is **real**, and so  $r^2 = 2$ , implying  $r = \pm\sqrt{2}$ .  
However  $r \geq 0$ , and therefore  $r = \sqrt{2}$ .

### Solution (continued)

3. Since  $r^4 = 4$ ,  $r^2 = \pm 2$ . But  $r$  is **real**, and so  $r^2 = 2$ , implying  $r = \pm\sqrt{2}$ . However  $r \geq 0$ , and therefore  $r = \sqrt{2}$ .
4. The solutions to  $e^{i4\theta} = e^{2\pi i/3}$  are given by

$$4\theta = \frac{2}{3}\pi + 2\pi\ell, \ell = 0, 1, 2, 3.$$

### Solution (continued)

3. Since  $r^4 = 4$ ,  $r^2 = \pm 2$ . But  $r$  is **real**, and so  $r^2 = 2$ , implying  $r = \pm\sqrt{2}$ . However  $r \geq 0$ , and therefore  $r = \sqrt{2}$ .
4. The solutions to  $e^{i4\theta} = e^{2\pi i/3}$  are given by

$$4\theta = \frac{2}{3}\pi + 2\pi\ell, \ell = 0, 1, 2, 3.$$

Therefore,

$$\theta = \frac{2\pi}{12} + \frac{2\pi\ell}{4} = \frac{\pi}{6} + \frac{\pi\ell}{2} = \frac{\pi(3\ell + 1)}{6}, \text{ for } \ell = 0, 1, 2, 3.$$

Solution (continued)

5. Thus  $r = \sqrt{2}$  and  $\theta = \left(\frac{3\ell+1}{6}\right)\pi$ ,  $\ell = 0, 1, 2, 3$ .

### Solution (continued)

5. Thus  $r = \sqrt{2}$  and  $\theta = \left(\frac{3\ell+1}{6}\right)\pi$ ,  $\ell = 0, 1, 2, 3$ . Converting to Cartesian form:



### Solution (continued)

5. Thus  $r = \sqrt{2}$  and  $\theta = \left(\frac{3\ell+1}{6}\right)\pi$ ,  $\ell = 0, 1, 2, 3$ . Converting to Cartesian form:

$$\ell = 0 : \quad z = \sqrt{2}e^{\pi i/6} \quad = \sqrt{2}\left(\frac{\sqrt{3}}{2} + \frac{1}{2}i\right) \quad = \frac{\sqrt{6}}{2} + \frac{\sqrt{2}}{2}i$$

### Solution (continued)

5. Thus  $r = \sqrt{2}$  and  $\theta = \left(\frac{3\ell+1}{6}\right)\pi$ ,  $\ell = 0, 1, 2, 3$ . Converting to Cartesian form:

$$\begin{aligned}\ell = 0 : \quad z &= \sqrt{2}e^{\pi i/6} &= \sqrt{2}\left(\frac{\sqrt{3}}{2} + \frac{1}{2}i\right) &= \frac{\sqrt{6}}{2} + \frac{\sqrt{2}}{2}i \\ \ell = 1 : \quad z &= \sqrt{2}e^{2\pi i/3} &= \sqrt{2}\left(-\frac{1}{2} + \frac{\sqrt{3}}{2}i\right) &= -\frac{\sqrt{2}}{2} + \frac{\sqrt{6}}{2}i\end{aligned}$$

### Solution (continued)

5. Thus  $r = \sqrt{2}$  and  $\theta = \left(\frac{3\ell+1}{6}\right)\pi$ ,  $\ell = 0, 1, 2, 3$ . Converting to Cartesian form:

$$\begin{aligned}\ell = 0 : \quad z &= \sqrt{2}e^{\pi i/6} &= \sqrt{2}\left(\frac{\sqrt{3}}{2} + \frac{1}{2}i\right) &= \frac{\sqrt{6}}{2} + \frac{\sqrt{2}}{2}i \\ \ell = 1 : \quad z &= \sqrt{2}e^{2\pi i/3} &= \sqrt{2}\left(-\frac{1}{2} + \frac{\sqrt{3}}{2}i\right) &= -\frac{\sqrt{2}}{2} + \frac{\sqrt{6}}{2}i \\ \ell = 2 : \quad z &= \sqrt{2}e^{7\pi i/6} &= \sqrt{2}\left(-\frac{\sqrt{3}}{2} - \frac{1}{2}i\right) &= -\frac{\sqrt{6}}{2} - \frac{\sqrt{2}}{2}i\end{aligned}$$

### Solution (continued)

5. Thus  $r = \sqrt{2}$  and  $\theta = \left(\frac{3\ell+1}{6}\right)\pi$ ,  $\ell = 0, 1, 2, 3$ . Converting to Cartesian form:

$$\begin{aligned}\ell = 0 : \quad z &= \sqrt{2}e^{\pi i/6} &= \sqrt{2}\left(\frac{\sqrt{3}}{2} + \frac{1}{2}i\right) &= \frac{\sqrt{6}}{2} + \frac{\sqrt{2}}{2}i \\ \ell = 1 : \quad z &= \sqrt{2}e^{2\pi i/3} &= \sqrt{2}\left(-\frac{1}{2} + \frac{\sqrt{3}}{2}i\right) &= -\frac{\sqrt{2}}{2} + \frac{\sqrt{6}}{2}i \\ \ell = 2 : \quad z &= \sqrt{2}e^{7\pi i/6} &= \sqrt{2}\left(-\frac{\sqrt{3}}{2} - \frac{1}{2}i\right) &= -\frac{\sqrt{6}}{2} - \frac{\sqrt{2}}{2}i \\ \ell = 3 : \quad z &= \sqrt{2}e^{5\pi i/3} &= \sqrt{2}\left(\frac{1}{2} - \frac{\sqrt{3}}{2}i\right) &= \frac{\sqrt{2}}{2} - \frac{\sqrt{6}}{2}i\end{aligned}$$

### Solution (continued)

5. Thus  $r = \sqrt{2}$  and  $\theta = \left(\frac{3\ell+1}{6}\right)\pi$ ,  $\ell = 0, 1, 2, 3$ . Converting to Cartesian form:

$$\begin{aligned}\ell = 0 : \quad z &= \sqrt{2}e^{\pi i/6} &= \sqrt{2}\left(\frac{\sqrt{3}}{2} + \frac{1}{2}i\right) &= \frac{\sqrt{6}}{2} + \frac{\sqrt{2}}{2}i \\ \ell = 1 : \quad z &= \sqrt{2}e^{2\pi i/3} &= \sqrt{2}\left(-\frac{1}{2} + \frac{\sqrt{3}}{2}i\right) &= -\frac{\sqrt{2}}{2} + \frac{\sqrt{6}}{2}i \\ \ell = 2 : \quad z &= \sqrt{2}e^{7\pi i/6} &= \sqrt{2}\left(-\frac{\sqrt{3}}{2} - \frac{1}{2}i\right) &= -\frac{\sqrt{6}}{2} - \frac{\sqrt{2}}{2}i \\ \ell = 3 : \quad z &= \sqrt{2}e^{5\pi i/3} &= \sqrt{2}\left(\frac{1}{2} - \frac{\sqrt{3}}{2}i\right) &= \frac{\sqrt{2}}{2} - \frac{\sqrt{6}}{2}i\end{aligned}$$

Therefore, the fourth roots of  $2(\sqrt{3}i - 1)$  are:

$$\frac{\sqrt{6}}{2} + \frac{\sqrt{2}}{2}i,$$

### Solution (continued)

5. Thus  $r = \sqrt{2}$  and  $\theta = \left(\frac{3\ell+1}{6}\right)\pi$ ,  $\ell = 0, 1, 2, 3$ . Converting to Cartesian form:

$$\begin{aligned}\ell = 0 : \quad z &= \sqrt{2}e^{\pi i/6} &= \sqrt{2}\left(\frac{\sqrt{3}}{2} + \frac{1}{2}i\right) &= \frac{\sqrt{6}}{2} + \frac{\sqrt{2}}{2}i \\ \ell = 1 : \quad z &= \sqrt{2}e^{2\pi i/3} &= \sqrt{2}\left(-\frac{1}{2} + \frac{\sqrt{3}}{2}i\right) &= -\frac{\sqrt{2}}{2} + \frac{\sqrt{6}}{2}i \\ \ell = 2 : \quad z &= \sqrt{2}e^{7\pi i/6} &= \sqrt{2}\left(-\frac{\sqrt{3}}{2} - \frac{1}{2}i\right) &= -\frac{\sqrt{6}}{2} - \frac{\sqrt{2}}{2}i \\ \ell = 3 : \quad z &= \sqrt{2}e^{5\pi i/3} &= \sqrt{2}\left(\frac{1}{2} - \frac{\sqrt{3}}{2}i\right) &= \frac{\sqrt{2}}{2} - \frac{\sqrt{6}}{2}i\end{aligned}$$

Therefore, the fourth roots of  $2(\sqrt{3}i - 1)$  are:

$$\frac{\sqrt{6}}{2} + \frac{\sqrt{2}}{2}i, -\frac{\sqrt{2}}{2} + \frac{\sqrt{6}}{2}i,$$

### Solution (continued)

5. Thus  $r = \sqrt{2}$  and  $\theta = \left(\frac{3\ell+1}{6}\right)\pi$ ,  $\ell = 0, 1, 2, 3$ . Converting to Cartesian form:

$$\begin{aligned}\ell = 0 : \quad z &= \sqrt{2}e^{\pi i/6} &= \sqrt{2}\left(\frac{\sqrt{3}}{2} + \frac{1}{2}i\right) &= \frac{\sqrt{6}}{2} + \frac{\sqrt{2}}{2}i \\ \ell = 1 : \quad z &= \sqrt{2}e^{2\pi i/3} &= \sqrt{2}\left(-\frac{1}{2} + \frac{\sqrt{3}}{2}i\right) &= -\frac{\sqrt{2}}{2} + \frac{\sqrt{6}}{2}i \\ \ell = 2 : \quad z &= \sqrt{2}e^{7\pi i/6} &= \sqrt{2}\left(-\frac{\sqrt{3}}{2} - \frac{1}{2}i\right) &= -\frac{\sqrt{6}}{2} - \frac{\sqrt{2}}{2}i \\ \ell = 3 : \quad z &= \sqrt{2}e^{5\pi i/3} &= \sqrt{2}\left(\frac{1}{2} - \frac{\sqrt{3}}{2}i\right) &= \frac{\sqrt{2}}{2} - \frac{\sqrt{6}}{2}i\end{aligned}$$

Therefore, the fourth roots of  $2(\sqrt{3}i - 1)$  are:

$$\frac{\sqrt{6}}{2} + \frac{\sqrt{2}}{2}i, -\frac{\sqrt{2}}{2} + \frac{\sqrt{6}}{2}i, -\frac{\sqrt{6}}{2} - \frac{\sqrt{2}}{2}i,$$

### Solution (continued)

5. Thus  $r = \sqrt{2}$  and  $\theta = \left(\frac{3\ell+1}{6}\right)\pi$ ,  $\ell = 0, 1, 2, 3$ . Converting to Cartesian form:

$$\begin{aligned}\ell = 0 : \quad z &= \sqrt{2}e^{\pi i/6} &= \sqrt{2}\left(\frac{\sqrt{3}}{2} + \frac{1}{2}i\right) &= \frac{\sqrt{6}}{2} + \frac{\sqrt{2}}{2}i \\ \ell = 1 : \quad z &= \sqrt{2}e^{2\pi i/3} &= \sqrt{2}\left(-\frac{1}{2} + \frac{\sqrt{3}}{2}i\right) &= -\frac{\sqrt{2}}{2} + \frac{\sqrt{6}}{2}i \\ \ell = 2 : \quad z &= \sqrt{2}e^{7\pi i/6} &= \sqrt{2}\left(-\frac{\sqrt{3}}{2} - \frac{1}{2}i\right) &= -\frac{\sqrt{6}}{2} - \frac{\sqrt{2}}{2}i \\ \ell = 3 : \quad z &= \sqrt{2}e^{5\pi i/3} &= \sqrt{2}\left(\frac{1}{2} - \frac{\sqrt{3}}{2}i\right) &= \frac{\sqrt{2}}{2} - \frac{\sqrt{6}}{2}i\end{aligned}$$

Therefore, the fourth roots of  $2(\sqrt{3}i - 1)$  are:

$$\frac{\sqrt{6}}{2} + \frac{\sqrt{2}}{2}i, -\frac{\sqrt{2}}{2} + \frac{\sqrt{6}}{2}i, -\frac{\sqrt{6}}{2} - \frac{\sqrt{2}}{2}i, \frac{\sqrt{2}}{2} - \frac{\sqrt{6}}{2}i.$$





Copyright

Complex Numbers

Modulus

Complex Numbers in Polar Form

Roots of Complex Numbers

**Roots of Unity**

The Quadratic Formula



# Roots of Unity

## Definition

A complex number  $z$  is a **root of unity** if there exists a positive integer  $n$  so that  $z^n = 1$ .

# Roots of Unity

## Definition

A complex number  $z$  is a **root of unity** if there exists a positive integer  $n$  so that  $z^n = 1$ .

## Problem

Find the sixth roots of unity, i.e., all solutions to  $z^6 = 1$ .

# Roots of Unity

## Definition

A complex number  $z$  is a **root of unity** if there exists a positive integer  $n$  so that  $z^n = 1$ .

## Problem

Find the sixth roots of unity, i.e., all solutions to  $z^6 = 1$ .

## Solution

Write  $z = re^{i\theta}$  and convert 1 to polar form to get

$$(re^{i\theta})^6 = e^{i0}, \text{ and so } r^6 e^{6\theta i} = e^{i0}.$$

# Roots of Unity

## Definition

A complex number  $z$  is a **root of unity** if there exists a positive integer  $n$  so that  $z^n = 1$ .

## Problem

Find the sixth roots of unity, i.e., all solutions to  $z^6 = 1$ .

## Solution

Write  $z = re^{i\theta}$  and convert 1 to polar form to get

$$(re^{i\theta})^6 = e^{i0}, \text{ and so } r^6 e^{6\theta i} = e^{i0}.$$

Equating the absolute values and arguments,

$$r^6 = 1 \quad \text{and} \quad 6\theta = 0 + 2\pi\ell, \quad \ell = 0, 1, 2, 3, 4, 5.$$

# Roots of Unity

## Definition

A complex number  $z$  is a **root of unity** if there exists a positive integer  $n$  so that  $z^n = 1$ .

## Problem

Find the sixth roots of unity, i.e., all solutions to  $z^6 = 1$ .

## Solution

Write  $z = re^{i\theta}$  and convert 1 to polar form to get

$$(re^{i\theta})^6 = e^{i0}, \text{ and so } r^6 e^{6\theta i} = e^{i0}.$$

Equating the absolute values and arguments,

$$r^6 = 1 \quad \text{and} \quad 6\theta = 0 + 2\pi\ell, \quad \ell = 0, 1, 2, 3, 4, 5.$$

Since  $r$  is real,  $r = 1$ .

# Roots of Unity

## Definition

A complex number  $z$  is a **root of unity** if there exists a positive integer  $n$  so that  $z^n = 1$ .

## Problem

Find the sixth roots of unity, i.e., all solutions to  $z^6 = 1$ .

## Solution

Write  $z = re^{i\theta}$  and convert 1 to polar form to get

$$(re^{i\theta})^6 = e^{i0}, \text{ and so } r^6 e^{6\theta i} = e^{i0}.$$

Equating the absolute values and arguments,

$$r^6 = 1 \quad \text{and} \quad 6\theta = 0 + 2\pi\ell, \quad \ell = 0, 1, 2, 3, 4, 5.$$

Since  $r$  is real,  $r = 1$ . The six arguments for the solutions are

$$\theta = \frac{2\pi\ell}{6} = \frac{\pi\ell}{3}, \quad \ell = 0, 1, 2, 3, 4, 5.$$



## Solution (continued)

The six arguments for the solutions are

$$\theta = \frac{2\pi\ell}{6} = \frac{\pi\ell}{3}, \ell = 0, 1, 2, 3, 4, 5.$$

### Solution (continued)

The six arguments for the solutions are

$$\theta = \frac{2\pi\ell}{6} = \frac{\pi\ell}{3}, \ell = 0, 1, 2, 3, 4, 5.$$

Converting these to Cartesian form:

### Solution (continued)

The six arguments for the solutions are

$$\theta = \frac{2\pi\ell}{6} = \frac{\pi\ell}{3}, \ell = 0, 1, 2, 3, 4, 5.$$

Converting these to Cartesian form:

$\ell$	$\theta$	$z$
0	0	$e^{0i} = 1$

## Solution (continued)

The six arguments for the solutions are

$$\theta = \frac{2\pi\ell}{6} = \frac{\pi\ell}{3}, \ell = 0, 1, 2, 3, 4, 5.$$

Converting these to Cartesian form:

$\ell$	$\theta$	$z$
0	0	$e^{0i} = 1$
1	$\frac{\pi}{3}$	$e^{\pi i/3} = \frac{1}{2} + \frac{\sqrt{3}}{2}i$

## Solution (continued)

The six arguments for the solutions are

$$\theta = \frac{2\pi\ell}{6} = \frac{\pi\ell}{3}, \ell = 0, 1, 2, 3, 4, 5.$$

Converting these to Cartesian form:

$\ell$	$\theta$	$z$
0	0	$e^{0i} = 1$
1	$\frac{\pi}{3}$	$e^{\pi i/3} = \frac{1}{2} + \frac{\sqrt{3}}{2}i$
2	$\frac{2\pi}{3}$	$e^{2\pi i/3} = -\frac{1}{2} + \frac{\sqrt{3}}{2}i$

## Solution (continued)

The six arguments for the solutions are

$$\theta = \frac{2\pi\ell}{6} = \frac{\pi\ell}{3}, \ell = 0, 1, 2, 3, 4, 5.$$

Converting these to Cartesian form:

$\ell$	$\theta$	$z$
0	0	$e^{0i} = 1$
1	$\frac{\pi}{3}$	$e^{\pi i/3} = \frac{1}{2} + \frac{\sqrt{3}}{2}i$
2	$\frac{2\pi}{3}$	$e^{2\pi i/3} = -\frac{1}{2} + \frac{\sqrt{3}}{2}i$
3	$\pi$	$e^{\pi i} = -1$

## Solution (continued)

The six arguments for the solutions are

$$\theta = \frac{2\pi\ell}{6} = \frac{\pi\ell}{3}, \ell = 0, 1, 2, 3, 4, 5.$$

Converting these to Cartesian form:

$\ell$	$\theta$	$z$
0	0	$e^{0i} = 1$
1	$\frac{\pi}{3}$	$e^{\pi i/3} = \frac{1}{2} + \frac{\sqrt{3}}{2}i$
2	$\frac{2\pi}{3}$	$e^{2\pi i/3} = -\frac{1}{2} + \frac{\sqrt{3}}{2}i$
3	$\pi$	$e^{\pi i} = -1$
4	$\frac{4\pi}{3}$	$e^{4\pi i/3} = -\frac{1}{2} - \frac{\sqrt{3}}{2}i$

## Solution (continued)

The six arguments for the solutions are

$$\theta = \frac{2\pi\ell}{6} = \frac{\pi\ell}{3}, \ell = 0, 1, 2, 3, 4, 5.$$

Converting these to Cartesian form:

$\ell$	$\theta$	$z$
0	0	$e^{0i} = 1$
1	$\frac{\pi}{3}$	$e^{\pi i/3} = \frac{1}{2} + \frac{\sqrt{3}}{2}i$
2	$\frac{2\pi}{3}$	$e^{2\pi i/3} = -\frac{1}{2} + \frac{\sqrt{3}}{2}i$
3	$\pi$	$e^{\pi i} = -1$
4	$\frac{4\pi}{3}$	$e^{4\pi i/3} = -\frac{1}{2} - \frac{\sqrt{3}}{2}i$
5	$\frac{5\pi}{3}$	$e^{5\pi i/3} = \frac{1}{2} - \frac{\sqrt{3}}{2}i$



## Solution (continued)

The six arguments for the solutions are

$$\theta = \frac{2\pi\ell}{6} = \frac{\pi\ell}{3}, \ell = 0, 1, 2, 3, 4, 5.$$

Converting these to Cartesian form:

$\ell$	$\theta$	$z$
0	0	$e^{0i} = 1$
1	$\frac{\pi}{3}$	$e^{\pi i/3} = \frac{1}{2} + \frac{\sqrt{3}}{2}i$
2	$\frac{2\pi}{3}$	$e^{2\pi i/3} = -\frac{1}{2} + \frac{\sqrt{3}}{2}i$
3	$\pi$	$e^{\pi i} = -1$
4	$\frac{4\pi}{3}$	$e^{4\pi i/3} = -\frac{1}{2} - \frac{\sqrt{3}}{2}i$
5	$\frac{5\pi}{3}$	$e^{5\pi i/3} = \frac{1}{2} - \frac{\sqrt{3}}{2}i$

If you graph these six point in the complex plane, you'll see that they result in six equally spaced points on the unit circle, one of them being  $(1, 0)$ . ■

## Definition

**Roots of Unity** For any integer  $n \geq 1$ , the (complex) solutions to  $z^n = 1$  are

$$z = e^{2\pi\ell i/n} \text{ for } \ell = 0, 1, 2, \dots, n-1.$$

## Definition

**Roots of Unity** For any integer  $n \geq 1$ , the (complex) solutions to  $z^n = 1$  are

$$z = e^{2\pi\ell i/n} \text{ for } \ell = 0, 1, 2, \dots, n-1.$$

Furthermore, the  $n^{\text{th}}$  roots of unity correspond to  $n$  equally spaced points on the unit circle, one of them being  $(1, 0)$ .

Copyright

Complex Numbers

Modulus

Complex Numbers in Polar Form

Roots of Complex Numbers

Roots of Unity

**The Quadratic Formula**



# The Quadratic Formula

## Definition

A **real** quadratic is an expression of the form  $ax^2 + bx + c$  where  $a, b, c \in \mathbb{R}$  and  $a \neq 0$ .

# The Quadratic Formula

## Definition

A **real** quadratic is an expression of the form  $ax^2 + bx + c$  where  $a, b, c \in \mathbb{R}$  and  $a \neq 0$ .

To find the roots of a real quadratic, we can either factor by inspection, or use the **quadratic formula**:

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

# The Quadratic Formula

## Definition

A **real** quadratic is an expression of the form  $ax^2 + bx + c$  where  $a, b, c \in \mathbb{R}$  and  $a \neq 0$ .

To find the roots of a real quadratic, we can either factor by inspection, or use the **quadratic formula**:

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

The expression  $b^2 - 4ac$  is called the **discriminant**,



# The Quadratic Formula

## Definition

A **real** quadratic is an expression of the form  $ax^2 + bx + c$  where  $a, b, c \in \mathbb{R}$  and  $a \neq 0$ .

To find the roots of a real quadratic, we can either factor by inspection, or use the **quadratic formula**:

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

The expression  $b^2 - 4ac$  is called the **discriminant**, and

- if  $b^2 - 4ac \geq 0$ , then the roots of the quadratic are **real**;

# The Quadratic Formula

## Definition

A **real** quadratic is an expression of the form  $ax^2 + bx + c$  where  $a, b, c \in \mathbb{R}$  and  $a \neq 0$ .

To find the roots of a real quadratic, we can either factor by inspection, or use the **quadratic formula**:

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

The expression  $b^2 - 4ac$  is called the **discriminant**, and

- ▶ if  $b^2 - 4ac \geq 0$ , then the roots of the quadratic are **real**;
- ▶ if  $b^2 - 4ac < 0$ , then the quadratic has **no real roots**.

## Definition

A real quadratic  $ax^2 + bx + c$  is called **irreducible** if the discriminant is less than zero, i.e.,  $b^2 - 4ac < 0$ .

## Definition

A real quadratic  $ax^2 + bx + c$  is called **irreducible** if the discriminant is less than zero, i.e.,  $b^2 - 4ac < 0$ .

Notice that if  $b^2 - 4ac < 0$ , then

$$\sqrt{b^2 - 4ac} = \sqrt{(-1)(4ac - b^2)} = (\pm)i\sqrt{4ac - b^2}.$$

## Definition

A real quadratic  $ax^2 + bx + c$  is called **irreducible** if the discriminant is less than zero, i.e.,  $b^2 - 4ac < 0$ .

Notice that if  $b^2 - 4ac < 0$ , then

$$\sqrt{b^2 - 4ac} = \sqrt{(-1)(4ac - b^2)} = (\pm)i\sqrt{4ac - b^2}.$$

It follows that the roots of an irreducible quadratic are

$$\frac{-b \pm i\sqrt{4ac - b^2}}{2a} = \left\{ \begin{array}{l} -\frac{b}{2a} + \frac{\sqrt{4ac - b^2}}{2a}i \\ -\frac{b}{2a} - \frac{\sqrt{4ac - b^2}}{2a}i \end{array} \right\},$$

## Definition

A real quadratic  $ax^2 + bx + c$  is called **irreducible** if the discriminant is less than zero, i.e.,  $b^2 - 4ac < 0$ .

Notice that if  $b^2 - 4ac < 0$ , then

$$\sqrt{b^2 - 4ac} = \sqrt{(-1)(4ac - b^2)} = (\pm)i\sqrt{4ac - b^2}.$$

It follows that the roots of an irreducible quadratic are

$$\frac{-b \pm i\sqrt{4ac - b^2}}{2a} = \left\{ \begin{array}{l} -\frac{b}{2a} + \frac{\sqrt{4ac - b^2}}{2a}i \\ -\frac{b}{2a} - \frac{\sqrt{4ac - b^2}}{2a}i \end{array} \right\},$$

and we see that the two roots are complex conjugates of each other.

## Definition

A real quadratic  $ax^2 + bx + c$  is called **irreducible** if the discriminant is less than zero, i.e.,  $b^2 - 4ac < 0$ .

Notice that if  $b^2 - 4ac < 0$ , then

$$\sqrt{b^2 - 4ac} = \sqrt{(-1)(4ac - b^2)} = (\pm)i\sqrt{4ac - b^2}.$$

It follows that the roots of an irreducible quadratic are

$$\frac{-b \pm i\sqrt{4ac - b^2}}{2a} = \left\{ \begin{array}{l} -\frac{b}{2a} + \frac{\sqrt{4ac - b^2}}{2a}i \\ -\frac{b}{2a} - \frac{\sqrt{4ac - b^2}}{2a}i \end{array} \right\},$$

and we see that the two roots are complex conjugates of each other. We denote the two roots by

$$u = -\frac{b}{2a} + \frac{\sqrt{4ac - b^2}}{2a}i \quad \text{and} \quad \bar{u} = -\frac{b}{2a} - \frac{\sqrt{4ac - b^2}}{2a}i.$$

### Example ( Real Quadratics with Complex Roots )

The quadratic  $x^2 - 14x + 58$  has roots

$$x = \frac{14 \pm \sqrt{196 - 4 \times 58}}{2}$$



### Example ( Real Quadratics with Complex Roots )

The quadratic  $x^2 - 14x + 58$  has roots

$$\begin{aligned}x &= \frac{14 \pm \sqrt{196 - 4 \times 58}}{2} \\&= \frac{14 \pm \sqrt{196 - 232}}{2}\end{aligned}$$

### Example ( Real Quadratics with Complex Roots )

The quadratic  $x^2 - 14x + 58$  has roots

$$\begin{aligned}x &= \frac{14 \pm \sqrt{196 - 4 \times 58}}{2} \\&= \frac{14 \pm \sqrt{196 - 232}}{2} \\&= \frac{14 \pm \sqrt{-36}}{2}\end{aligned}$$

### Example ( Real Quadratics with Complex Roots )

The quadratic  $x^2 - 14x + 58$  has roots

$$\begin{aligned}x &= \frac{14 \pm \sqrt{196 - 4 \times 58}}{2} \\&= \frac{14 \pm \sqrt{196 - 232}}{2} \\&= \frac{14 \pm \sqrt{-36}}{2} \\&= \frac{14 \pm 6i}{2}\end{aligned}$$

### Example ( Real Quadratics with Complex Roots )

The quadratic  $x^2 - 14x + 58$  has roots

$$\begin{aligned}x &= \frac{14 \pm \sqrt{196 - 4 \times 58}}{2} \\&= \frac{14 \pm \sqrt{196 - 232}}{2} \\&= \frac{14 \pm \sqrt{-36}}{2} \\&= \frac{14 \pm 6i}{2} \\&= 7 \pm 3i,\end{aligned}$$

### Example ( Real Quadratics with Complex Roots )

The quadratic  $x^2 - 14x + 58$  has roots

$$\begin{aligned}x &= \frac{14 \pm \sqrt{196 - 4 \times 58}}{2} \\&= \frac{14 \pm \sqrt{196 - 232}}{2} \\&= \frac{14 \pm \sqrt{-36}}{2} \\&= \frac{14 \pm 6i}{2} \\&= 7 \pm 3i,\end{aligned}$$

so the roots are  $7 + 3i$  and  $7 - 3i$ .

Conversely, given  $u = a + bi$  with  $b \neq 0$ , there is an irreducible quadratic having roots  $u$  and  $\bar{u}$ .

Conversely, given  $u = a + bi$  with  $b \neq 0$ , there is an irreducible quadratic having roots  $u$  and  $\bar{u}$ .

### Problem

Find an irreducible quadratic with  $u = 5 - 2i$  as a root. What is the other root?

Conversely, given  $u = a + bi$  with  $b \neq 0$ , there is an irreducible quadratic having roots  $u$  and  $\bar{u}$ .

### Problem

Find an irreducible quadratic with  $u = 5 - 2i$  as a root. What is the other root?

### Solution

$$(x - u)(x - \bar{u}) =$$



Conversely, given  $u = a + bi$  with  $b \neq 0$ , there is an irreducible quadratic having roots  $u$  and  $\bar{u}$ .

### Problem

Find an irreducible quadratic with  $u = 5 - 2i$  as a root. What is the other root?

### Solution

$$(x - u)(x - \bar{u}) = (x - (5 - 2i))(x - (5 + 2i))$$

Conversely, given  $u = a + bi$  with  $b \neq 0$ , there is an irreducible quadratic having roots  $u$  and  $\bar{u}$ .

### Problem

Find an irreducible quadratic with  $u = 5 - 2i$  as a root. What is the other root?

### Solution

$$\begin{aligned}(x - u)(x - \bar{u}) &= (x - (5 - 2i))(x - (5 + 2i)) \\ &= x^2 - (5 - 2i)x - (5 + 2i)x + (5 - 2i)(5 + 2i)\end{aligned}$$

Conversely, given  $u = a + bi$  with  $b \neq 0$ , there is an irreducible quadratic having roots  $u$  and  $\bar{u}$ .

### Problem

Find an irreducible quadratic with  $u = 5 - 2i$  as a root. What is the other root?

### Solution

$$\begin{aligned}(x - u)(x - \bar{u}) &= (x - (5 - 2i))(x - (5 + 2i)) \\&= x^2 - (5 - 2i)x - (5 + 2i)x + (5 - 2i)(5 + 2i) \\&= x^2 - 10x + 29.\end{aligned}$$

Conversely, given  $u = a + bi$  with  $b \neq 0$ , there is an irreducible quadratic having roots  $u$  and  $\bar{u}$ .

### Problem

Find an irreducible quadratic with  $u = 5 - 2i$  as a root. What is the other root?

### Solution

$$\begin{aligned}(x - u)(x - \bar{u}) &= (x - (5 - 2i))(x - (5 + 2i)) \\&= x^2 - (5 - 2i)x - (5 + 2i)x + (5 - 2i)(5 + 2i) \\&= x^2 - 10x + 29.\end{aligned}$$

Therefore,  $x^2 - 10x + 29$  is an irreducible quadratic with roots  $5 - 2i$  and  $5 + 2i$ .

Conversely, given  $u = a + bi$  with  $b \neq 0$ , there is an irreducible quadratic having roots  $u$  and  $\bar{u}$ .

### Problem

Find an irreducible quadratic with  $u = 5 - 2i$  as a root. What is the other root?

### Solution

$$\begin{aligned}(x - u)(x - \bar{u}) &= (x - (5 - 2i))(x - (5 + 2i)) \\&= x^2 - (5 - 2i)x - (5 + 2i)x + (5 - 2i)(5 + 2i) \\&= x^2 - 10x + 29.\end{aligned}$$

Therefore,  $x^2 - 10x + 29$  is an irreducible quadratic with roots  $5 - 2i$  and  $5 + 2i$ .

Notice that  $-10 = -(u + \bar{u})$  and  $29 = u\bar{u} = |u|^2$ .



### Problem

Find an irreducible quadratic with root  $u = -3 + 4i$ , and find the other root.

### Problem

Find an irreducible quadratic with root  $u = -3 + 4i$ , and find the other root.

### Solution ( answer )

$x^2 + 6x + 25$  has roots  $u = -3 + 4i$  and  $\bar{u} = -3 - 4i$ .

### Problem ( Quadratics with Complex Coefficients )

Find the roots of the quadratic  $x^2 - (3 - 2i)x + (5 - i) = 0$ .



### Problem ( Quadratics with Complex Coefficients )

Find the roots of the quadratic  $x^2 - (3 - 2i)x + (5 - i) = 0$ .

### Solution

Using the quadratic formula

$$x = \frac{3 - 2i \pm \sqrt{-(3 - 2i)^2 - 4(5 - i)}}{2}.$$

### Problem ( Quadratics with Complex Coefficients )

Find the roots of the quadratic  $x^2 - (3 - 2i)x + (5 - i) = 0$ .

### Solution

Using the quadratic formula

$$x = \frac{3 - 2i \pm \sqrt{-(3 - 2i)^2 - 4(5 - i)}}{2}.$$

Now,

$$-(3 - 2i)^2 - 4(5 - i) = 5 - 12i - 20 + 4i = -15 - 8i,$$

### Problem ( Quadratics with Complex Coefficients )

Find the roots of the quadratic  $x^2 - (3 - 2i)x + (5 - i) = 0$ .

### Solution

Using the quadratic formula

$$x = \frac{3 - 2i \pm \sqrt{-(3 - 2i)^2 - 4(5 - i)}}{2}.$$

Now,

$$-(3 - 2i)^2 - 4(5 - i) = 5 - 12i - 20 + 4i = -15 - 8i,$$

so

$$x = \frac{3 - 2i \pm \sqrt{-15 - 8i}}{2}.$$

### Problem ( Quadratics with Complex Coefficients )

Find the roots of the quadratic  $x^2 - (3 - 2i)x + (5 - i) = 0$ .

### Solution

Using the quadratic formula

$$x = \frac{3 - 2i \pm \sqrt{-(3 - 2i)^2 - 4(5 - i)}}{2}.$$

Now,

$$(-(3 - 2i))^2 - 4(5 - i) = 5 - 12i - 20 + 4i = -15 - 8i,$$

so

$$x = \frac{3 - 2i \pm \sqrt{-15 - 8i}}{2}.$$

To find  $\pm\sqrt{-15 - 8i}$ , solve  $z^2 = -15 - 8i$  for  $z$ .

Solution (continued)

Let  $z = a + bi$  and  $z^2 = -15 - 8i$ .

Solution (continued)

Let  $z = a + bi$  and  $z^2 = -15 - 8i$ . Then

$$(a^2 - b^2) + 2abi = -15 - 8i,$$

### Solution (continued)

Let  $z = a + bi$  and  $z^2 = -15 - 8i$ . Then

$$(a^2 - b^2) + 2abi = -15 - 8i,$$

so  $a^2 - b^2 = -15$  and  $2ab = -8$ .

### Solution (continued)

Let  $z = a + bi$  and  $z^2 = -15 - 8i$ . Then

$$(a^2 - b^2) + 2abi = -15 - 8i,$$

so  $a^2 - b^2 = -15$  and  $2ab = -8$ .

Solving for  $a$  and  $b$  gives us  $z = 1 - 4i, -1 + 4i$ , i.e.,  $z = \pm(1 - 4i)$ .



### Solution (continued)

Let  $z = a + bi$  and  $z^2 = -15 - 8i$ . Then

$$(a^2 - b^2) + 2abi = -15 - 8i,$$

so  $a^2 - b^2 = -15$  and  $2ab = -8$ .

Solving for  $a$  and  $b$  gives us  $z = 1 - 4i, -1 + 4i$ , i.e.,  $z = \pm(1 - 4i)$ .  
Therefore,

$$x = \frac{3 - 2i \pm (1 - 4i)}{2},$$

### Solution (continued)

Let  $z = a + bi$  and  $z^2 = -15 - 8i$ . Then

$$(a^2 - b^2) + 2abi = -15 - 8i,$$

so  $a^2 - b^2 = -15$  and  $2ab = -8$ .

Solving for  $a$  and  $b$  gives us  $z = 1 - 4i, -1 + 4i$ , i.e.,  $z = \pm(1 - 4i)$ .  
Therefore,

$$x = \frac{3 - 2i \pm (1 - 4i)}{2},$$

and

$$\frac{3-2i+(1-4i)}{2} = \frac{4-6i}{2} = 2 - 3i,$$

$$\frac{3-2i-(1-4i)}{2} = \frac{2+2i}{2} = 1 + i.$$

### Solution (continued)

Let  $z = a + bi$  and  $z^2 = -15 - 8i$ . Then

$$(a^2 - b^2) + 2abi = -15 - 8i,$$

so  $a^2 - b^2 = -15$  and  $2ab = -8$ .

Solving for  $a$  and  $b$  gives us  $z = 1 - 4i, -1 + 4i$ , i.e.,  $z = \pm(1 - 4i)$ .  
Therefore,

$$x = \frac{3 - 2i \pm (1 - 4i)}{2},$$

and

$$\frac{3-2i+(1-4i)}{2} = \frac{4-6i}{2} = 2 - 3i,$$

$$\frac{3-2i-(1-4i)}{2} = \frac{2+2i}{2} = 1 + i.$$

Thus the roots of  $x^2 - (3 - 2i)x + (5 - i)$  are  $2 - 3i$  and  $1 + i$ . ■



### Problem

Find the roots of  $x^2 - 3ix + (-3 + i)$ .

### Solution ( answer )

$1 + i$  and  $-1 + 2i$ .

### Problem

Verify that  $u_1 = (4 - i)$  is a root of

$$x^2 - (2 - 3i)x - (10 + 6i)$$

and find the other root,  $u_2$ .

## Problem

Verify that  $u_1 = (4 - i)$  is a root of

$$x^2 - (2 - 3i)x - (10 + 6i)$$

and find the other root,  $u_2$ .

## Solution

First,

$$\begin{aligned}u_1^2 - (2 - 3i)u_1 - (10 + 6i) &= (4 - i)^2 - (2 - 3i)(4 - i) - (10 + 6i) \\&= (15 - 8i) - (5 - 14i) - (10 + 6i) \\&= 0,\end{aligned}$$

so  $u_1 = (4 - i)$  is a root.

### Solution (continued)

Recall that if  $u_1$  and  $u_2$  are the roots of the quadratic, then

$$u_1 + u_2 = (2 - 3i) \quad \text{and} \quad u_1 u_2 = -(10 + 6i).$$

Solve for  $u_2$  using either one of these equations.



### Solution (continued)

Recall that if  $u_1$  and  $u_2$  are the roots of the quadratic, then

$$u_1 + u_2 = (2 - 3i) \quad \text{and} \quad u_1 u_2 = -(10 + 6i).$$

Solve for  $u_2$  using either one of these equations.

Since  $u_1 = 4 - i$  and  $u_1 + u_2 = 2 - 3i$ ,

$$u_2 = 2 - 3i - u_1 = 2 - 3i - (4 - i) = -2 - 2i.$$

### Solution (continued)

Recall that if  $u_1$  and  $u_2$  are the roots of the quadratic, then

$$u_1 + u_2 = (2 - 3i) \quad \text{and} \quad u_1 u_2 = -(10 + 6i).$$

Solve for  $u_2$  using either one of these equations.

Since  $u_1 = 4 - i$  and  $u_1 + u_2 = 2 - 3i$ ,

$$u_2 = 2 - 3i - u_1 = 2 - 3i - (4 - i) = -2 - 2i.$$

Therefore, the other root is  $u_2 = -2 - 2i$ .

### Solution (continued)

Recall that if  $u_1$  and  $u_2$  are the roots of the quadratic, then

$$u_1 + u_2 = (2 - 3i) \quad \text{and} \quad u_1 u_2 = -(10 + 6i).$$

Solve for  $u_2$  using either one of these equations.

Since  $u_1 = 4 - i$  and  $u_1 + u_2 = 2 - 3i$ ,

$$u_2 = 2 - 3i - u_1 = 2 - 3i - (4 - i) = -2 - 2i.$$

Therefore, the other root is  $u_2 = -2 - 2i$ .

You can easily verify your answer by computing  $u_1 u_2$ :

$$u_1 u_2 = (4 - i)(-2 - 2i) = -10 - 6i = -(10 + 6i).$$

