

RANDOM DETERMINANTS, MIXED VOLUMES OF ELLIPSOIDS, AND ZEROS OF GAUSSIAN RANDOM FIELDS

D. Zaporozhets* and Z. Kabluchko†

UDC 519.2+514

Consider a $d \times d$ matrix M whose rows are independent, centered, nondegenerate Gaussian vectors ξ_1, \dots, ξ_d with covariance matrices $\Sigma_1, \dots, \Sigma_d$. Denote by \mathcal{E}_i the dispersion ellipsoid of ξ_i : $\mathcal{E}_i = \{\mathbf{x} \in \mathbb{R}^d : \mathbf{x}^\top \Sigma_i^{-1} \mathbf{x} \leq 1\}$. We show that

$$\mathbb{E} |\det M| = \frac{d!}{(2\pi)^{d/2}} V_d(\mathcal{E}_1, \dots, \mathcal{E}_d),$$

where $V_d(\cdot, \dots, \cdot)$ denotes the mixed volume. We also generalize this result to the case of rectangular matrices. As a direct corollary, we get an analytic expression for the mixed volume of d arbitrary ellipsoids in \mathbb{R}^d .

As another application, we consider a smooth, centered, nondegenerate Gaussian random field $X = (X_1, \dots, X_k)^\top : \mathbb{R}^d \rightarrow \mathbb{R}^k$. Using the Kac–Rice formula, we obtain a geometric interpretation of the intensity of zeros of X in terms of the mixed volume of dispersion ellipsoids of the gradients of $X_i / \sqrt{\text{Var} X_i}$. This relates zero sets of equations to mixed volumes in a way which is reminiscent of the well-known Bernstein theorem about the number of solutions of a typical system of algebraic equations. Bibliography: 10 titles.

1. MAIN RESULTS

1.1. Random determinant and mixed volume of ellipsoids. Consider independent, centered, nondegenerate Gaussian random vectors $\xi_1, \dots, \xi_k \in \mathbb{R}^d$, $k \leq d$, with covariance matrices $\Sigma_1, \dots, \Sigma_k$. Denote by \mathcal{E}_i the dispersion ellipsoid of ξ_i :

$$\mathcal{E}_i = \left\{ \mathbf{x} = (x_1, \dots, x_d)^\top \in \mathbb{R}^d : \mathbf{x}^\top \Sigma_i^{-1} \mathbf{x} \leq 1 \right\}, \quad i = 1, \dots, k. \quad (1.1)$$

Denote by M the $k \times d$ matrix whose rows are ξ_1, \dots, ξ_k .

Theorem 1.1. *The following relation holds:*

$$\mathbb{E} \sqrt{\det(MM^\top)} = \frac{(d)_k}{(2\pi)^{k/2} \kappa_{d-k}} V_d(\mathcal{E}_1, \dots, \mathcal{E}_k, B, \dots, B), \quad (1.2)$$

where $V_d(\cdot, \dots, \cdot)$ denotes the mixed volume of d convex bodies in \mathbb{R}^d (see Sec. 2 for details), B is the unit ball in \mathbb{R}^d , $(d)_k = d(d-1) \cdots (d-k+1)$ is the Pochhammer symbol, and $\kappa_n = \pi^{n/2} / \Gamma(1+n/2)$ denotes the volume of the unit ball in \mathbb{R}^n .

The left-hand side of (1.2) can be interpreted as the average k -dimensional volume of a Gaussian random parallelotope.

Corollary 1.2. *In the case $k = d$, the following relation holds:*

$$\mathbb{E} |\det M| = \frac{d!}{(2\pi)^{d/2}} V_d(\mathcal{E}_1, \dots, \mathcal{E}_d).$$

As another direct corollary, we can calculate the mixed volume of d arbitrary ellipsoids in \mathbb{R}^d .

Corollary 1.3. *If $\mathcal{E}_1, \dots, \mathcal{E}_d$ are arbitrary ellipsoids defined by symmetric positive definite matrices $\Sigma_1, \dots, \Sigma_d$ as in (1.1), then*

$$V_d(\mathcal{E}_1, \dots, \mathcal{E}_d) = \frac{1}{d!} \prod_{i=1}^d (\det \Sigma_i)^{-1/2} \int_{\mathbb{R}^{d^2}} |\det(x_{ij})| \prod_{i=1}^d \exp\left(-\frac{1}{2} \mathbf{x}_i^\top \Sigma_i^{-1} \mathbf{x}_i\right) dx_{11} \dots dx_{dd},$$

where

$$\mathbf{x}_i = (x_{i1}, \dots, x_{id})^\top.$$

*St. Petersburg Department of the Steklov Mathematical Institute, St. Petersburg, Russia, e-mail: zap1979@gmail.com.

†Ulm University, Ulm, Germany, e-mail: zakhhar.kabluchko@uni-ulm.de.

The only estimate of the mixed volume of ellipsoids which we know is due to Barvinok [2]. He showed that

$$\frac{\kappa_d}{3^{(d-1)/2}} \sqrt{D_d(\Sigma_1, \dots, \Sigma_d)} \leq V_d(\mathcal{E}_1, \dots, \mathcal{E}_d) \leq \kappa_d \sqrt{D_d(\Sigma_1, \dots, \Sigma_d)},$$

where $D_d(\cdot, \dots, \cdot)$ denotes the mixed discriminant of d symmetric $d \times d$ matrices:

$$D_d(A_1, \dots, A_d) = \frac{1}{d!} \frac{\partial^d}{\partial \lambda_1 \dots \partial \lambda_d} \det(\lambda_1 A_1 + \dots + \lambda_d A_d) \Big|_{\lambda_1 = \dots = \lambda_d = 0}.$$

If ξ_1, \dots, ξ_k are independent, standard Gaussian vectors, then MM^\top is a Wishart matrix, and (1.2) turns into (see [5, 10])

$$\mathbf{E} \sqrt{\det(MM^\top)} = \frac{(d)_k \kappa_d}{(2\pi)^{k/2} \kappa_{d-k}}.$$

1.2. Intrinsic volumes. If $\xi_1, \xi_2, \dots, \xi_k \in \mathbb{R}^d, k \leq d$, are identically distributed with the common covariance matrix Σ and dispersion ellipsoid \mathcal{E} , then (1.2) turns into

$$\mathbf{E} \sqrt{\det(MM^\top)} = \frac{k!}{(2\pi)^{k/2}} V_k(\mathcal{E}), \quad (1.3)$$

where $V_k(\cdot)$ denotes the k th *intrinsic volume* of a convex body in \mathbb{R}^d :

$$V_k(K) = \frac{\binom{d}{k}}{\kappa_{d-k}} V_d(\underbrace{K, \dots, K}_{k \text{ times}}, B, \dots, B).$$

The normalization is chosen so that $V_k(K)$ depends only on K and not on the dimension of the surrounding space, i.e., if $\dim K < d$, then the computation of $V_k(K)$ in \mathbb{R}^d leads to the same result as the computation in the affine span of K . In particular, if $\dim K = k$, then $V_k(K) = \text{Vol}_k(K)$, the k -dimensional volume of K .

It is known that $V_1(K)$ is proportional to the mean width of K :

$$V_1(K) = \frac{d\kappa_d}{2\kappa_{d-1}} w(K).$$

Taking $k = 1$ in (1.3), we see that for any centered Gaussian vector ξ with dispersion ellipsoid \mathcal{E} ,

$$\mathbf{E} \|\xi\| = \frac{1}{\sqrt{2\pi}} V_1(\mathcal{E}). \quad (1.4)$$

It was pointed out by M. Lifshits that (1.4) is a special case of the following remarkable result of Sudakov.

1.3. Connection with Sudakov's result. For our purposes, the following finite-dimensional version of Sudakov's theorem suffices. The result in full generality can be found in [9, Proposition 14].

Proposition 1.4. *For an arbitrary subset $A \subset \mathbb{R}^d$,*

$$\mathbf{E} \sup_{\mathbf{x} \in A} \langle \mathbf{x}, \eta \rangle = \frac{1}{\sqrt{2\pi}} V_1(\text{conv}(A)), \quad (1.5)$$

where η is a standard Gaussian vector in \mathbb{R}^d and $\text{conv}(A)$ is the convex hull of A .

Let us deduce (1.4) from (1.5). Consider a matrix U such that $\Sigma = U^{-1}(U^{-1})^\top$ and $U\xi$ is a standard Gaussian vector. Using (1.5) with $A = \mathcal{E}$ and $\eta = U\xi$, we get

$$\mathbf{E} \|\xi\| = \mathbf{E} \sup_{\|\mathbf{x}\| \leq 1} \langle \mathbf{x}, \xi \rangle = \mathbf{E} \sup_{\|\mathbf{x}\| \leq 1} \langle (U^{-1})^\top \mathbf{x}, U\xi \rangle = \mathbf{E} \sup_{\|U^\top \mathbf{x}\| \leq 1} \langle \mathbf{x}, U\xi \rangle = \mathbf{E} \sup_{\mathbf{x} \in \mathcal{E}} \langle \mathbf{x}, U\xi \rangle = \frac{1}{\sqrt{2\pi}} V_1(\mathcal{E}).$$

1.4. Zeros of Gaussian random fields. Let $X(\mathbf{t}) = (X_1(\mathbf{t}), \dots, X_k(\mathbf{t}))^\top : \mathbb{R}^d \rightarrow \mathbb{R}^k$, $k \leq d$, be a random field. Following Azaïs and Wschebor [1], we always assume that the following conditions hold:

- (a) X is Gaussian;
- (b) almost surely, the function $X(\cdot)$ is of class \mathcal{C}^1 ;
- (c) for all $\mathbf{t} \in \mathbb{R}^d$, $X(\mathbf{t})$ has a nondegenerate distribution;
- (d) almost surely, if $X(\mathbf{t}) = 0$, then $X'(\mathbf{t})$, the Jacobian matrix of $X(\mathbf{t})$, has the full rank.

Then, almost surely, the level set $X^{-1}(0)$ is a \mathcal{C}^1 -manifold of dimension $d - k$, and for any Borel set F , the Lebesgue measure $\text{Vol}_{d-k}(X^{-1}(0) \cap F)$ is well-defined ($\text{Vol}_0(\cdot)$ denotes the counting measure).

It was shown in [1, p. 177] that

$$\mathbf{E} \text{Vol}_{d-k}(X^{-1}(0) \cap F) = \int_F \mathbf{E} \left(\sqrt{\det(X'(\mathbf{t})X'(\mathbf{t})^\top)} \mid X(\mathbf{t}) = 0 \right) p_{X(\mathbf{t})}(0) d\mathbf{t}, \quad (1.6)$$

where $p_{X(\mathbf{t})}(\cdot)$ is the density of $X(\mathbf{t})$. Thus, the integrand in (1.6) can be interpreted as the intensity of zeros of X .

In this paper, we consider the special case where X is centered and its coordinates X_1, \dots, X_k are independent. Denote by $\mathcal{E}_i(\mathbf{t})$ the dispersion ellipsoid of $\nabla[X_i(\mathbf{t})/\sqrt{\text{Var } X_i(\mathbf{t})}]$.

Theorem 1.5. *Let X be a centered random field with independent coordinates defined as above and satisfying conditions (a)–(d). Then*

$$\mathbf{E} \text{Vol}_{d-k}(X^{-1}(0) \cap F) = \frac{\binom{d}{k}}{(2\pi)^k \kappa_{d-k}} \int_F V_d(\mathcal{E}_1(\mathbf{t}), \dots, \mathcal{E}_k(\mathbf{t}), B, \dots, B) d\mathbf{t}. \quad (1.7)$$

Formula (1.7) relates zero sets of random equations to mixed volumes. In the case $k = d$, it is therefore reminiscent of the well-known fact from the algebraic geometry which we formulate in the next subsection.

1.5. Bernstein's theorem. Consider a complex polynomial in d variables,

$$f(z_1, \dots, z_d) = \sum c_{j_1, \dots, j_d} z_1^{j_1} \dots z_d^{j_d}.$$

The Newton polytope of f is the subset of \mathbb{R}^d defined as follows:

$$\text{Nw}(f) = \text{conv} \{ (j_1, \dots, j_d) \in \mathbb{Z}^d : c_{j_1, \dots, j_d} \neq 0 \}.$$

Let K_1, \dots, K_d be compact convex polytopes in \mathbb{R}^d with vertices in \mathbb{Z}^d . Consider a system of algebraic equations

$$\begin{cases} f_1(z_1, \dots, z_d) = 0, \\ \dots \\ f_d(z_1, \dots, z_d) = 0, \end{cases}$$

such that $\text{Nw}(f_i) = K_i$. Bernstein showed [3] that for almost all such systems (with respect to Lebesgue measure in the space of coefficients of the polynomials), the number of nonzero solutions is equal to

$$\text{Vol}_0(f_1^{-1}(0) \cap \dots \cap f_d^{-1}(0) \setminus \{\mathbf{0}\}) = d! V_d(K_1, \dots, K_d).$$

2. SOME ESSENTIAL TOOLS FROM GEOMETRY

For the basic facts from integral and convex geometry we refer the reader to [4] and [8].

2.1. Mixed volumes. Consider arbitrary convex bodies $K_1, \dots, K_d \subset \mathbb{R}^d$. Minkowski showed [7] that $\text{Vol}_d(\lambda_1 K_1 + \dots + \lambda_d K_d)$, where $\lambda_1, \dots, \lambda_d \geq 0$, is a homogeneous polynomial of degree d with nonnegative coefficients:

$$\text{Vol}_d(\lambda_1 K_1 + \dots + \lambda_d K_d) = \sum_{i_1=1}^d \dots \sum_{i_d=1}^d \lambda_{i_1} \dots \lambda_{i_d} V_d(K_{i_1}, \dots, K_{i_d}). \quad (2.1)$$

The coefficients $V_d(K_{i_1}, \dots, K_{i_d})$ are uniquely determined by the assumption that they are symmetric with respect to permutations of K_{i_1}, \dots, K_{i_d} . The coefficient $V_d(K_1, \dots, K_d)$ is called the mixed volume of K_1, \dots, K_d . Differentiating (2.1), we get an alternative definition of the mixed volume:

$$V_d(K_1, \dots, K_d) = \frac{1}{d!} \frac{\partial^d}{\partial \lambda_1 \dots \partial \lambda_d} \text{Vol}_d(\lambda_1 K_1 + \dots + \lambda_d K_d) \Big|_{\lambda_1 = \dots = \lambda_d = 0}.$$

For any affine transformation L ,

$$V_d(LK_1, \dots, LK_d) = |\det L| \cdot V_d(K_1, \dots, K_d). \quad (2.2)$$

The following relation can also be stated:

$$\int_{\mathbb{S}^{d-1}} V_{d-1}(P_{\mathbf{u}}K_1, \dots, P_{\mathbf{u}}K_{d-1}) d\mathbf{u} = \frac{\kappa_{d-1}}{\kappa_d} V_d(K_1, \dots, K_{d-1}, B), \quad (2.3)$$

where $d\mathbf{u}$ is the surface measure on \mathbb{S}^{d-1} normalized to have total mass 1 and $P_{\mathbf{u}}$ denotes the orthogonal projection to the linear hyperplane \mathbf{u}^\perp .

2.2. Volumes of parallelotopes. For any $A \subset \mathbb{R}^d$ and $\mathbf{x}_1, \dots, \mathbf{x}_k \in \mathbb{R}^d$ denote by $P_{\mathbf{x}_1, \dots, \mathbf{x}_k} A$ the orthogonal projection of A to $\text{span}^\perp\{\mathbf{x}_1, \dots, \mathbf{x}_k\}$ (the orthogonal complement of the linear span of $\mathbf{x}_1, \dots, \mathbf{x}_k$). Denote by $H_{\mathbf{x}_1, \dots, \mathbf{x}_k}$ the parallelotope generated by the vectors $\mathbf{x}_1, \dots, \mathbf{x}_k$. It is known that

$$\text{Vol}_k(H_{\mathbf{x}_1, \dots, \mathbf{x}_k}) = \sqrt{\det(AA^\top)}, \quad (2.4)$$

where A is the matrix whose rows are $\mathbf{x}_1, \dots, \mathbf{x}_k$.

For any $\mathbf{x}_1, \dots, \mathbf{x}_d \in \mathbb{R}^d$ and $k = 1, \dots, d-1$,

$$\text{Vol}_d(H_{\mathbf{x}_1, \dots, \mathbf{x}_d}) = \text{Vol}_k(H_{\mathbf{x}_1, \dots, \mathbf{x}_k}) \text{Vol}_{d-k}(P_{\mathbf{x}_1, \dots, \mathbf{x}_k} H_{\mathbf{x}_{k+1}, \dots, \mathbf{x}_d}). \quad (2.5)$$

2.3. Ellipsoids. There is a bijection $A \mapsto \mathcal{E}$ between $d \times d$ symmetric positive definite matrices and d -dimensional nondegenerate ellipsoids centered at the origin (see [6] for details):

$$\mathcal{E} = \{\mathbf{x} \in \mathbb{R}^d : \mathbf{x}^\top A^{-1} \mathbf{x} \leq 1\}.$$

Any nondegenerate linear coordinate transformation of the form $\mathbf{x} \mapsto L\mathbf{x}$ is reflected by a change of the corresponding representing matrix A to the matrix A_L given by

$$A_L = LAL^\top. \quad (2.6)$$

Let \mathcal{E}' be the orthogonal projection of \mathcal{E} onto an k -dimensional subspace with some orthonormal basis $\mathbf{x}_1, \dots, \mathbf{x}_k \in \mathbb{R}^d$. Denote by A' the $k \times k$ matrix representing the ellipsoid \mathcal{E}' in this basis. If C is the $k \times d$ matrix whose rows are $\mathbf{x}_1, \dots, \mathbf{x}_k$, then

$$A' = CAC^\top. \quad (2.7)$$

3. PROOFS

3.1. Proof of Theorem 1.1. Case $k = d$. We proceed by induction on d . First let us assume that ξ_d is a standard Gaussian vector. Denote by χ_d a random variable having the chi distribution with d degrees of freedom and independent from ξ_1, \dots, ξ_{d-1} . Using (2.4) and (2.5) with $k = 1$, we get the relations

$$\begin{aligned} \mathbf{E} |\det M| &= \mathbf{E} \text{Vol}_d(H_{\xi_1, \dots, \xi_d}) = \int_{\mathbb{S}^{d-1}} \mathbf{E} \text{Vol}_d(H_{\xi_1, \dots, \xi_{d-1}, \chi_d \mathbf{u}}) d\mathbf{u} \\ &= \mathbf{E} \chi_d \int_{\mathbb{S}^{d-1}} \mathbf{E} \text{Vol}_{d-1}(P_{\mathbf{u}} H_{\xi_1, \dots, \xi_{d-1}}) d\mathbf{u} \\ &= \frac{d\kappa_d}{\sqrt{2\pi\kappa_{d-1}}} \int_{\mathbb{S}^{d-1}} \mathbf{E} \text{Vol}_{d-1}(H_{P_{\mathbf{u}}\xi_1, \dots, P_{\mathbf{u}}\xi_{d-1}}) d\mathbf{u}. \end{aligned}$$

It follows from (2.7) that $P_{\mathbf{u}}\xi_i$ has dispersion ellipsoid $P_{\mathbf{u}}\mathcal{E}_i$. By the induction assumption,

$$\mathbf{E} \text{Vol}_{d-1}(H_{P_{\mathbf{u}}\xi_1, \dots, P_{\mathbf{u}}\xi_{d-1}}) = \frac{(d-1)!}{(2\pi)^{(d-1)/2}} V_{d-1}(P_{\mathbf{u}}\mathcal{E}_1, \dots, P_{\mathbf{u}}\mathcal{E}_{d-1}).$$

Combining the latter two relations with (2.3), we obtain the equality

$$\mathbf{E} |\det M| = \frac{d!}{(2\pi)^{d/2}} V_d(\mathcal{E}_1, \dots, \mathcal{E}_{d-1}, B). \quad (3.1)$$

If ξ_d is an arbitrary nondegenerate Gaussian vector, then there exists a linear transformation L such that $L\xi_d$ is a standard Gaussian vector. It follows from (2.6) that $L\xi_i$ is the dispersion ellipsoid of $L\xi_i$, and, in particular, $L\xi_d = B$. Applying (3.1) to the matrix LM^\top and using (2.2), we get the equalities

$$\begin{aligned} \mathbf{E} |\det M| &= |\det L|^{-1} \mathbf{E} |\det LM^\top| = \frac{d!}{(2\pi)^{d/2}} |\det L|^{-1} V_d(L\xi_1, \dots, L\xi_{d-1}, B) \\ &= \frac{d!}{(2\pi)^{d/2}} V_d(\mathcal{E}_1, \dots, \mathcal{E}_{d-1}, \mathcal{E}_d). \end{aligned} \quad \square$$

3.2. Proof of Theorem 1.1. Case $k < d$. Consider a $d \times d$ matrix M' whose first k rows form the matrix M and the last $d - k$ rows are independent standard Gaussian vectors ξ_{k+1}, \dots, ξ_d (independent from M). By the previous case,

$$\mathbf{E} |\det M'| = \frac{d!}{(2\pi)^{d/2}} V_d(\mathcal{E}_1, \dots, \mathcal{E}_k, B, \dots, B).$$

On the other hand, by (2.5),

$$\begin{aligned} \mathbf{E} |\det M'| &= \mathbf{E} \text{Vol}_d(H_{\xi_1, \dots, \xi_d}) = \mathbf{E} \text{Vol}_k(H_{\xi_1, \dots, \xi_k}) \text{Vol}_{d-k}(P_{\xi_1, \dots, \xi_k} H_{\xi_{k+1}, \dots, \xi_d}) \\ &= \mathbf{E} \sqrt{\det(MM^\top)} \mathbf{E} \text{Vol}_{d-k}(H_{\eta_1, \dots, \eta_{d-k}}), \end{aligned}$$

where $\eta_1, \dots, \eta_{d-k}$ are independent, standard Gaussian vectors in \mathbb{R}^{d-k} . By the previous case,

$$\mathbf{E} \text{Vol}_{d-k}(H_{\eta_1, \dots, \eta_{d-k}}) = \frac{(d-k)!}{(2\pi)^{(d-k)/2}} \kappa_{d-k}.$$

Combining the latter three relations completes the proof. \square

3.3. Proof of Theorem 1.5. First we assume that X_j has a unit variance: $\text{Var } X_j(\mathbf{t}) \equiv 1$ for all $j = 1, \dots, k$. Differentiating the relation $\mathbf{E} X_j(\mathbf{t}) X_j(\mathbf{t}) = 1$ with respect to t_i , we obtain the equality

$$\mathbf{E} \frac{\partial X_j}{\partial t_i}(\mathbf{t}) X_j(\mathbf{t}) = 0,$$

which, together with the independence of the coordinates of X , implies that $X'(\mathbf{t})$ and $X(\mathbf{t})$ are independent. This means that the conditioning on $X(\mathbf{t}) = 0$ in (1.6) may be dropped. To complete the proof of the theorem in the case $\text{Var } X_j(\mathbf{t}) \equiv 1$, it remains to combine (1.6) with (1.2).

To cover the general case, it suffices to note that $X_j/\sqrt{\text{Var } X_j}$ has the same zero set as X_j . \square

Acknowledgments. We are grateful to M. Lifshits for bringing our attention to Sudakov's result. We are also grateful to A. I. Barvinok for the useful discussion.

The second author is partially supported by the RFBR (project 10-01-00242), the Program "Leading Scientific Schools" (project-1216.2012.1), and DFG (grant 436 RUS 113/962/0-1 R).

Translated by D. Zaporozhets and Z. Kabluchko.

REFERENCES

1. J. M. Azaïs and M. Wschebor, *Level Sets and Extrema of Random Processes and Fields*, Wiley (2009).
2. A. Barvinok, "Computing mixed discriminants, mixed volumes, and permanents," *Discrete Comput. Geom.*, **18**, 205–237 (1997).
3. D. N. Bernshtein, "The number of roots of a system of equations," *Funct. Anal. Appl.*, **9**, 183–185 (1975).
4. Yu. D. Burago and V. A. Zalgaller, *Geometric Inequalities*, Grundlehren der Mathematischen Wissenschaften, Vol. 285, Springer–Verlag, Berlin (1988),
5. N. R. Goodman, "The distribution of the determinant of a complex Wishart distributed matrix," *Ann. Math. Statist.*, **34**, 178–180 (1963).
6. W. C. Karl, G. C. Verghese, and A. S. Willsky, "Reconstructing ellipsoids from projections," *CVGIP: Graphical Model and Image Processing*, **56**, 124–139 (1994).
7. H. Minkowski, "Theorie der konvexen Körper, insbesondere Begründung ihres Oberflächenbegriffs," in: *Gesammelte Abhandlungen*, Vol. 2 (1911), pp. 131–229.
8. R. Schneider and W. Weil, *Stochastic and Integral Geometry*, Springer–Verlag (2008).

9. V. N. Sudakov, "Geometric problems in the theory of infinite-dimensional probability distributions," *Trudy Mat. Inst. Akad. Nauk SSSR*, **141** (1976).
10. S. S. Wilks, "Moment-generating operators for determinants of product moments in samples from a normal system," *Ann. Math.*, **35**, 312–340 (1934).