

A Reflected Stochastic Heat Equation as Symmetric Dynamics with Respect to the 3-d Bessel Bridge

Lorenzo Zambotti

Scuola Normale Superiore, Piazza dei Cavalieri 7, 56126 Pisa, Italy
 E-mail: zambotti@cibs.sns.it

Communicated by Paul Malliavin

Received July 20, 2000; accepted September 6, 2000

We prove that a stochastic heat equation with reflection at 0, on the spatial interval $[0, 1]$ with Dirichlet boundary conditions and additive white-noise, admits an explicit symmetrizing invariant measure on $C([0, 1])$: the 3-d Bessel Bridge, i.e., the law of the modulus of a 3-dimensional Brownian motion conditioned to be 0 at time 1, a classical measure in probability theory, also connected with the theory of excursions of Brownian motion. This is a non-trivial example of a Gibbs-type measure being singular with respect to the reference Gaussian measure and concentrated on the convex set of positive, continuous functions on $[0, 1]$. © 2001

Academic Press

1. INTRODUCTION

In [6], Nualart and Pardoux studied existence and uniqueness of a solution to a reflected stochastic heat equation, namely of a pair (u, η) , where u is a continuous function of $(t, \xi) \in \mathcal{O} := [0, +\infty) \times [0, 1]$ and η is a positive measure on \mathcal{O} , satisfying

$$\begin{cases} \frac{\partial u}{\partial t} = \frac{1}{2} \frac{\partial^2 u}{\partial \xi^2} - f(\xi, u(t, \xi)) + \frac{\partial^2 W}{\partial t \partial \xi} + \frac{\eta(dt, d\xi)}{dt d\xi} \\ u(0, \xi) = x(\xi), \quad u(t, 0) = u(t, 1) = 0 \\ u \geq 0, \quad \int_{\mathcal{O}} u d\eta = 0. \end{cases} \quad (1)$$

where $x : [0, 1] \mapsto [0, \infty)$ is continuous with $x(0) = x(1) = 0$, $\{W(t, \xi) : (t, \xi) \in \mathcal{O}\}$ is a Brownian sheet, and $f : [0, 1] \times \mathbb{R} \mapsto \mathbb{R}$.

The aim of this paper is to prove that Eq. (1) admits, on the space $C_0(0, 1)$ of continuous functions of $\xi \in [0, 1]$, satisfying Dirichlet boundary conditions, an explicit invariant and symmetrizing measure, given by

$$\exp \left\{ - \int_0^1 d\xi \int_0^{x(\xi)} 2f(\xi, s) ds \right\} \nu(dx), \quad x \in C_0(0, 1), \quad (2)$$

where ν is a well-known probability measure on $C_0(0, 1)$: the 3-d Bessel bridge, namely the law of the modulus of a 3-dimensional Brownian motion $(B_\tau)_{\tau \in [0, 1]}$, conditioned to be 0 at $\tau = 1$. This measure plays an important role in the study of Brownian motion (see [7]).

On the other hand, Da Prato proved in [2] the existence of a symmetric semigroup $(P_t)_{t \geq 0}$ on $L^2(H, (\int_H e^{-2U} d\mu)^{-1} e^{-2U} d\mu)$, associated with the stochastic differential inclusion

$$dX \in (AX - \partial U(X)) dt + dW, \quad X(0) = x \in H, \quad (3)$$

where H is a separable Hilbert space, $A := D(A) \subset H \mapsto H$ is a strictly negative self-adjoint operator such that $Q := (-2A)^{-1}$ is trace-class, μ is the Gaussian measure $\mathcal{N}(0, Q)$ on H with 0 mean and covariance operator Q , and W is a cylindrical white-noise on H . Moreover, $U: H \mapsto \mathbb{R} \cup \{+\infty\}$ is a convex, lower semicontinuous function, satisfying suitable integrability conditions with respect to μ and

$$\mu(x : U(x) < +\infty \text{ and } \partial U(x) \neq \emptyset) = 1,$$

where $\partial U(x)$, the subdifferential of U at x , is defined as the subset of H :

$$\partial U(x) := \{y \in H : U(x+h) \geq U(x) + \langle h, y \rangle, \forall h \in H\}.$$

Equation (1) can be interpreted as an example of (3), setting $H := L^2(0, 1)$, $A := (1/2) d^2/d\xi^2$ on H with Dirichlet boundary conditions,

$$U(x) := \begin{cases} \int_0^1 d\xi \int_0^{x(\xi)} f(\xi, s) ds & \text{if } x \geq 0 \\ +\infty & \text{otherwise} \end{cases} \quad (4)$$

and defining, for all non-negative $x \in H$, $\partial U(x)$ as the subset of the dual space M of $C_0(0, 1)$, $M := \{\text{signed measures on } (0, 1)\}$,

$$\begin{aligned} \partial U(x) &:= \{m \in M : U(x+z) \geq U(x) + \langle z, m \rangle, \forall z \in C_0(0, 1)\} \\ &= f(\xi, x(\xi)) d\xi - \left\{ m \in M : m \geq 0, \int_{(0, 1)} x(\xi) m(d\xi) = 0 \right\}. \end{aligned}$$

Then (1) can be written formally as a differential inclusion of measures:

$$\left(\frac{\partial u}{\partial t} - Au - \frac{\partial^2 W}{\partial t \partial \xi} \right) d\xi \in \partial U(u(t, \cdot)).$$

However, in this case $\mu(U = +\infty) = 1$ (see Lemma 5 below) and the measure $(\int_H e^{-2U} d\mu)^{-1} e^{-2U} d\mu$ is not well defined. Our result shows that a natural Gibbs-type measure for (1) is provided by (2). Notice that ν is not Gaussian and is even singular with respect to the reference Gaussian measure μ . Moreover, the support of ν is a closed convex set having empty interior both in the topologies of $L^2(0, 1)$ and $C([0, 1])$.

The proof we give relies on the construction of solutions to (1) given by Nualart and Pardoux in [6] and on a result of Biane on a connection between the law of the Brownian Bridge and the law of the 3-d Bessel Bridge (see [1] and Theorem 4 below).

2. DEFINITIONS

Our aim is to find a symmetrizing invariant measure for the process $x \mapsto u(t, \cdot)$, $t \geq 0$, where (u, η) satisfies:

$$\begin{cases} \frac{\partial u}{\partial t} = \frac{1}{2} \frac{\partial^2 u}{\partial \xi^2} - f(\xi, u(t, \xi)) + \frac{\partial^2 W}{\partial t \partial \xi} + \frac{\eta(dt, d\xi)}{dt d\xi} \\ u(0, \xi) = x(\xi), \quad u(t, 0) = u(t, 1) = 0 \\ u + \alpha \geq 0, \quad \int_{\mathcal{O}} (u + \alpha) d\eta = 0, \end{cases} \quad (5)$$

where $\alpha \geq 0$, $x: [0, 1] \mapsto [-\alpha, +\infty)$ is continuous and $x(0) = x(1) = 0$.

We introduce the following notations: $(t, \xi) \in \mathcal{O} := [0, +\infty) \times [0, 1]$, $H := L^2(0, 1)$ with scalar product $\langle \cdot, \cdot \rangle$ and norm $\|\cdot\|$,

$$\langle h, k \rangle := \int_0^1 h(\xi) k(\xi) d\xi, \quad \|h\|^2 := \langle h, h \rangle,$$

$$C_0(0, 1) := \{c: [0, 1] \mapsto \mathbb{R} \text{ continuous, } c(0) = c(1) = 0\},$$

$$A: D(A) \subset H \mapsto H, \quad D(A) := H^2 \cap H_0^1(0, 1), \quad A := \frac{1}{2} \frac{d^2}{d\xi^2}.$$

We set $K_\alpha := \{h \in H : h \geq -\alpha\}$ with $\alpha \geq 0$, and we denote by $\Pi_{K_\alpha} : H \mapsto K_\alpha$ the projection from H onto the closed convex set $K_\alpha \subset H$. Recall that Π_{K_α} is 1-Lipschitz continuous. If $D \subseteq H$, we denote by $C_b(D)$ the space of all $\varphi : D \mapsto \mathbb{R}$ being bounded and uniformly continuous with respect to $\|\cdot\|$. If $D \subseteq H$ and $\varphi \in C_b(D)$, we denote by ω_φ the modulus of continuity of φ :

$$\omega_\varphi : [0, \infty) \mapsto [0, 1], \quad \omega_\varphi(r) := \sup\{|\varphi(x) - \varphi(x')| \wedge 1 : \|x - x'\| \leq r\}.$$

We identify $C_b(K_\alpha)$ with a subspace of $C_b(H)$ by means of the injection: $C_b(K_\alpha) \ni \varphi \mapsto \varphi \circ \Pi_{K_\alpha} \in C_b(H)$. If $0 \leq \alpha \leq \beta$, then $C_b(K_\alpha) \subseteq C_b(K_\beta)$.

If $\{m_n\}_n \cup \{m\}$ is a sequence of probability measures on $(H, \mathcal{B}(H))$, where $\mathcal{B}(H)$ is the Borel σ -field of H , we say that m_n converges weakly to m , if:

$$\lim_{n \rightarrow \infty} \int_H \varphi \, dm_n = \int_H \varphi \, dm, \quad \forall \varphi \in C_b(H).$$

Given a Markov process $\{Y(t, x) : t \geq 0, x \in D\}$ on $D \subseteq H$, we say that a probability measure m on D is symmetrizing for Y , if, setting for all $\varphi \in C_b(D) : R_t^Y \varphi(x) := \mathbb{E}[\varphi(Y(t, x))]$, $x \in D$, we have:

$$\int_D \varphi \, R_t^Y \psi \, dm = \int_D \psi \, R_t^Y \varphi \, dm, \quad \forall \varphi, \psi \in C_b(D).$$

A symmetrizing measure is in particular invariant; i.e.,

$$\int_D R_t^Y \varphi \, dm = \int_D \varphi \, dm, \quad \forall \varphi \in C_b(D).$$

We denote by $1_D(\cdot)$ the characteristic function of a set D . We sometimes write $m(\varphi)$ for $\int_H \varphi \, dm$, $\varphi \in C_b(H)$.

By $W = \{W(t, \xi) : (t, \xi) \in \mathcal{O}\}$ we denote a two-parameter Wiener process defined on a complete probability space $(\Omega, \mathcal{F}, \mathbb{P})$, i.e., W is a Gaussian process with zero mean and covariance function

$$\mathbb{E}[W(t, \xi) W(t', \xi')] = (t \wedge t')(\xi \wedge \xi'), \quad (t, \xi), (t', \xi') \in \mathcal{O}.$$

We denote by \mathcal{F}_t the σ -field generated by the random variables $\{W(s, \xi) : (s, \xi) \in [0, t] \times [0, 1]\}$ and by $C_c^\infty(0, 1)$ the subset of $C_0(0, 1)$ of all C^∞ functions with support being compact in $(0, 1)$.

We always assume in the following that:

(H1) $f = f_1 + f_2$, $f_1, f_2 : [0, 1] \times \mathbb{R} \mapsto \mathbb{R}$ are jointly measurable.

(H2) There exists $c > 0$ such that $|f_1(\xi, y) - f_1(\xi, y')| \leq c |y - y'|$ for all $\xi \in [0, 1]$, $y, y' \in \mathbb{R}$.

(H3) $f_2(\xi, \cdot)$ is non-decreasing and continuous, for all $\xi \in [0, 1]$.

(H4) There exist a constant $\lambda < \pi^2$, $a \in H$ and a polynomial p of degree $N \in \mathbb{N}$, such that for all $\xi \in [0, 1]$, $y \in \mathbb{R}$:

$$-a(\xi) - \lambda |y| \leq 2f(\xi, y) \leq a(\xi) + p(|y|).$$

Following [6], we set the following:

DEFINITION 1. A pair (u, η) is said to be a solution of Eq. (5) with reflection in $-\alpha \leq 0$ and initial value $x \in K_\alpha \cap C_0(0, 1)$, if:

(i) $\{u(t, \xi) : (t, \xi) \in \mathcal{O}\}$ is a continuous and adapted process, i.e., $u(t, \xi)$ is \mathcal{F}_t -measurable for all $(t, \xi) \in \mathcal{O}$, and a.s. $u(\cdot, \cdot)$ is continuous on \mathcal{O} , $u(t, \cdot) \in K_\alpha \cap C_0(0, 1)$ for all $t \geq 0$, and $u(0, \cdot) = x$.

(ii) $\eta(dt, d\xi)$ is a random positive measure on \mathcal{O} such that $\eta([0, T] \times [\delta, 1 - \delta]) < +\infty$ for all $T, \delta > 0$, and η is adapted; i.e., $\eta(B)$ is \mathcal{F}_t -measurable for every Borel set $B \subset [0, t] \times [0, 1]$.

(iii) For all $t \geq 0$ and $\varphi \in C_c^\infty(0, 1)$, setting $u_s := u(s, \cdot)$,

$$\begin{aligned} \langle u_t, \varphi \rangle - \int_0^t \langle u_s, A\varphi \rangle ds + \int_0^t \langle f(\cdot, u_s), \varphi \rangle ds \\ = \langle x, \varphi \rangle + \int_0^t \int_0^1 \varphi(\xi) dW_{s, \xi} + \int_0^t \int_0^1 \varphi(\xi) \eta(ds, d\xi). \end{aligned}$$

(iv) $\int_{\mathcal{O}} (u + \alpha) d\eta = 0$.

Finally, we will use the following:

LEMMA 1. Let T be a Polish metric space, and let $\{m_n\}_n \cup \{m\}$, respectively $\{\varphi_n\}_n$, be a sequence of probability measures, resp. of real-valued continuous functions, on T , satisfying:

- m_n converges weakly to m .
- The family $\{\varphi_n\}_n$ is uniformly bounded and equicontinuous on T .
- $\varphi_n(x)$ has a limit $\varphi(x)$ as $n \rightarrow \infty$, for all $x \in S$, with $S \subseteq T$ Borel and $m(S) = 1$.

Then:

$$\lim_{n \rightarrow \infty} \int_T \varphi_n dm_n = \int_S \varphi dm.$$

Proof. We can suppose that $0 \leq \varphi_n \leq 1$ for all n . By Prokhorov's theorem, there exists for every $\delta > 0$ a compact set $Q_\delta \subset T$ such that eventually $m_n(Q_\delta) \geq 1 - \delta$. Let $\{\varphi_{n_k}\}_k$ be any subsequence of $\{\varphi_n\}_n$. On Q_δ we can apply the Ascoli-Arzelà theorem and obtain uniform convergence of a sub-subsequence $\{\varphi_{n_{k(l)}}\}_l$ to a continuous function $f: Q_\delta \mapsto \mathbb{R}$. Then:

$$\begin{aligned} \int_T \varphi_{n_{k(l)}} dm_{n_{k(l)}} - \int_T \varphi_{n_{k(l)}} dm &\leq m_{n_{k(l)}}(T - Q_\delta) + \int_{Q_\delta} \varphi_{n_{k(l)}} [dm_{n_{k(l)}} - dm] \\ &\leq \delta + 2\delta + \int_{Q_\delta} f [dm_{n_{k(l)}} - dm], \end{aligned}$$

where for $l \geq l_0$, $\sup_{Q_\delta} |\varphi_{n_{k(l)}} - f| \leq \delta$. Since $m(T - S) = 0$ and Q_δ is closed,

$$\lim_{n \rightarrow \infty} \int_T \varphi_n dm = \int_S \varphi dm,$$

$$\limsup_{n \rightarrow \infty} \int_{Q_\delta} f dm_n \leq \int_{Q_\delta} f dm, \quad \text{and therefore:}$$

$$\limsup_{l \rightarrow \infty} \int_T \varphi_{n_{k(l)}} dm_{n_{k(l)}} \leq \int_S \varphi dm.$$

Changing φ_n with $1 - \varphi_n$, we obtain the thesis. ■

3. THE PROCESS X_α , $\alpha \geq 0$

In [6], the following theorem is proved:

THEOREM 1. Assume that f satisfies (H1), (H2), (H3), (H4) and let $x \in K_\alpha \cap C_0(0, 1)$. Then there exists a unique solution (u, η) to Eq. (5) with reflection in $-\alpha$ and initial value x .

We recall the strategy of the proof, given in [6], of the existence of solutions. First, the following approximating problem is introduced:

$$\begin{cases} \frac{\partial u_\alpha^\varepsilon}{\partial t} = \frac{1}{2} \frac{\partial^2 u_\alpha^\varepsilon}{\partial \xi^2} - f(\cdot, u_\alpha^\varepsilon(t, \cdot)) + \frac{\partial^2 W}{\partial t \partial \xi} + \frac{(\alpha + u_\alpha^\varepsilon)^-}{\varepsilon} \\ u_\alpha^\varepsilon(0, \cdot) = x \in H, \quad u_\alpha^\varepsilon(t, 0) = u_\alpha^\varepsilon(t, 1) = 0, \quad \forall t \geq 0. \end{cases} \quad (6)$$

with $\varepsilon > 0$, $(r)^- := \sup\{-r, 0\}$, and $\alpha \geq 0$. This is now a SPDE in $L^2(0, 1)$ with additive noise and monotone or Lipschitz-continuous drift terms, for which existence and uniqueness of a solution are well known (see [3]). Then, if $x \in K_\alpha \cap C_0(0, 1)$, the following is proved:

(a) $u_\alpha^\varepsilon(t, \cdot) \in C_0(0, 1)$ for all $t \geq 0$, and u_α^ε is continuous on \mathcal{O} .

(b) The map $0 < \varepsilon \mapsto u_\alpha^\varepsilon(t, \xi)$ is non-decreasing for all $(t, \xi) \in \mathcal{O}$. The limit $\lim_{\varepsilon \downarrow 0} u_\alpha^\varepsilon(t, \xi) = \sup_{\varepsilon > 0} u_\alpha^\varepsilon(t, \xi) =: u_\alpha(t, \xi)$ is finite for all $(t, \xi) \in \mathcal{O}$, $u_\alpha(t, \cdot) \in K_\alpha \cap C_0(0, 1)$ for all $t \geq 0$, and u_α is continuous on \mathcal{O} .

(c) The measure on \mathcal{O} , $\eta_\alpha^\varepsilon(dt, d\xi) := (1/\varepsilon)(\alpha + u_\alpha^\varepsilon)^- dt d\xi$, converges distributionally to a measure $\eta_\alpha(dt, d\xi)$ on \mathcal{O} .

(d) The pair (u_α, η_α) is the solution to (5) with reflection in $-\alpha$ and initial value $x \in K_\alpha \cap C_0(0, 1)$.

We set for all $t \geq 0$, $\alpha \geq 0$, $\varepsilon > 0$:

- $X_\alpha(t, x) \in C_0(0, 1)$, $X_\alpha(t, x)(\xi) := u_\alpha(t, \xi)$, $x \in K_\alpha \cap C_0(0, 1)$,
- $X_\alpha^\varepsilon(t, x) \in H$, $X_\alpha^\varepsilon(t, x)(\xi) := u_\alpha^\varepsilon(t, \xi)$, $x \in H$.

LEMMA 2. For all $\alpha \geq 0$, $\varepsilon > 0$, $t \geq 0$, we have: $\forall x, x' \in C_0(0, 1)$,

$$\|X_\alpha^\varepsilon(t, x) - X_\alpha^\varepsilon(t, x')\| \leq e^{ct} \|x - x'\|, \quad (7)$$

where $\|\cdot\|$ denotes the norm in H and $c > 0$ is the constant of (H2).

Proof. By the monotonicity properties of $(\cdot)^-$ and $f_2(\xi, \cdot)$ we have

$$\frac{1}{2} \frac{d}{dt} \|X_\alpha^\varepsilon(t, x) - X_\alpha^\varepsilon(t, x')\|^2 \leq c \|X_\alpha^\varepsilon(t, x) - X_\alpha^\varepsilon(t, x')\|^2$$

and the thesis follows from Gronwall's lemma. ■

Therefore, the same estimate holds for X_α , $\alpha \geq 0$: $\forall x, x' \in K_\alpha \cap C_0(0, 1)$,

$$\|X_\alpha(t, x) - X_\alpha(t, x')\| \leq e^{ct} \|x - x'\| \quad (8)$$

and we can uniquely extend $X_\alpha^\varepsilon(t, \cdot)$, respectively $X_\alpha(t, \cdot)$, to maps from H to H , resp. from K_α to K_α , that we denote by the same symbols, satisfying (7) for all $x, x' \in H$, resp. (8) for all $x, x' \in K_\alpha$. We set for all $\alpha \geq 0$, $\varepsilon > 0$, $\varphi \in C_b(H)$, $t \geq 0$:

$$P_\alpha^\varepsilon(t) \varphi: H \mapsto \mathbb{R}, \quad P_\alpha^\varepsilon(t) \varphi(x) := \mathbb{E}[\varphi(X_\alpha^\varepsilon(t, x))], \quad x \in H, \quad (9)$$

$$P_\alpha(t) \varphi: K_\alpha \mapsto \mathbb{R}, \quad P_\alpha(t) \varphi(x) := \mathbb{E}[\varphi(X_\alpha(t, x))], \quad x \in K_\alpha. \quad (10)$$

LEMMA 3. For all $\alpha \geq 0$, $\varepsilon > 0$, $\varphi \in C_b(H)$, $t \geq 0$, we have:

$$P_\alpha^\varepsilon(t) \varphi \in C_b(H), \quad \omega_{P_\alpha^\varepsilon(t) \varphi}(r) \leq \omega_\varphi(e^{ct}r) \quad \forall r \geq 0, \quad (11)$$

$$P_\alpha(t) \varphi \in C_b(K_\alpha), \quad \omega_{P_\alpha(t) \varphi}(r) \leq \omega_\varphi(e^{ct}r) \quad \forall r \geq 0, \quad (12)$$

$$\lim_{\varepsilon \downarrow 0} P_\alpha^\varepsilon(t) \varphi(x) = P_\alpha(t) \varphi(x) \quad \forall x \in K_\alpha, \quad (13)$$

$$P_\alpha(s) P_\alpha(t) \varphi(x) = P_\alpha(t+s) \varphi(x), \quad \forall x \in K_\alpha. \quad (14)$$

In particular, $(P_\alpha(t))_{t \geq 0}$ is a Markov semigroup acting on $C_b(K_\alpha)$.

Proof. For (12), notice that, by (8), for all $x, x' \in K_\alpha$:

$$\begin{aligned} |P_\alpha(t) \varphi(x) - P_\alpha(t) \varphi(x')| &\leq \mathbb{E}[|\varphi(X_\alpha(t, x)) - \varphi(X_\alpha(t, x'))|] \\ &\leq \mathbb{E}[\omega_\varphi(\|X_\alpha(t, x) - X_\alpha(t, x')\|)] \\ &\leq \omega_\varphi(e^{ct} \|x - x'\|), \end{aligned}$$

and (11) follows analogously. Equation (13) is a consequence of (b) in the proof of Theorem 1 and (11). It is well known that $(P_\alpha^\varepsilon(t))_{t \geq 0}$ is a semigroup acting on $C_b(H)$: since the family of probability measures $\{m^\varepsilon\}_{\varepsilon > 0}$, where m^ε is the law of $X_\alpha^\varepsilon(s, x)$, and the family of functions $\{P_\alpha^\varepsilon(t) \varphi\}_{\varepsilon > 0}$ satisfy the Hypothesis of Lemma 1; Eq. (14) follows. ■

LEMMA 4. For all $\varphi \in C_b(H)$, $\lim_{\alpha \downarrow 0} P_\alpha(t) \varphi(x) = P_0(t) \varphi(x)$, $t \geq 0$, $x \in K_0$.

Proof. If $x \in K_0 \cap C_0(0, 1)$, then the map $0 < \alpha \mapsto X_\alpha^\varepsilon(t, x)(\xi)$ is non-decreasing for all $(t, \xi) \in \mathcal{O}$, $\varepsilon > 0$. Therefore,

$$\begin{aligned} \lim_{\alpha \downarrow 0} X_\alpha(t, x)(\xi) &= \sup_{\alpha > 0} X_\alpha(t, x)(\xi) = \sup_{\alpha > 0} \sup_{\varepsilon > 0} X_\alpha^\varepsilon(t, x)(\xi) \\ &= \sup_{\varepsilon > 0} \sup_{\alpha > 0} X_\alpha^\varepsilon(t, x)(\xi) = \sup_{\varepsilon > 0} X_0^\varepsilon(t, x)(\xi) \\ &= X_0(t, x)(\xi), \end{aligned}$$

since $\sup_{\alpha > 0} X_\alpha^\varepsilon(t, x)(\cdot) = X_0^\varepsilon(t, x)(\cdot)$ by the uniqueness of solutions to (6). The general case follows by (12) and a density argument. ■

4. THE BROWNIAN BRIDGE RESTRICTED TO K_α AS SYMMETRIZING MEASURE FOR X_α , $\alpha > 0$

Recall that the Ornstein–Uhlenbeck process

$$Z(t, x) := e^{tA}x + \int_0^t e^{(t-s)A} dW_s \quad t \geq 0, \quad x \in H,$$

is, under our assumptions, a continuous Markov process with values in H , admitting as symmetrizing measure $\mu := \mathcal{N}(0, (-2A)^{-1})$, i.e., the Gaussian measure on H with 0 mean and covariance operator $(-2A)^{-1}$.

Our next lemma identifies μ with a well-known probability measure on $C_0(0, 1)$: the law of the Brownian bridge. Recall that the Brownian bridge is defined as a linear Brownian motion $(w_\tau)_{\tau \in [0, 1]}$, conditioned to be 0 at $\tau = 1$, and can be realized as $[0, 1] \ni \tau \mapsto w_\tau - \tau w_1$. The law of the Brownian bridge is concentrated on $C_0(0, 1)$ and is the unique Gaussian measure on $\mathbb{R}^{[0, 1]}$ with 0 mean and covariance function: $\Gamma(\tau, \sigma) = \tau \wedge \sigma - \tau\sigma$, $\sigma, \tau \in [0, 1]$, (see [7, Chap. I]).

LEMMA 5. *The measure μ coincides with the law of the Brownian bridge.*

Proof. Recall that the measure μ is concentrated on $C([0, 1]) \subset H$. By definition of Gaussian measures, the following holds for all $h, k \in H$:

$$\int_H \langle x, h \rangle \langle x, k \rangle \mathcal{N}(0, (-2A)^{-1})(dx) = \langle (-2A)^{-1} h, k \rangle. \quad (15)$$

Since the operator $(-2A)^{-1}$ can be expressed as an integral operator with kernel: $\xi \wedge \sigma - \xi\sigma$, $\xi, \sigma \in [0, 1]$, then setting in (19) $h = \chi_{[0, t]}$, $k = \chi_{[0, s]}$, $s, t \in [0, 1]$, and differentiating with respect to t and s , we obtain:

$$\int_{C([0, 1])} x(t) x(s) d\mu(x) = t \wedge s - ts. \quad \blacksquare$$

Lemma 5 allows us to calculate explicitly $\mu(K_\alpha)$:

$$\mu(K_\alpha) = 1 - \exp\{-2\alpha^2\}, \quad \alpha \geq 0, \quad (16)$$

(see [7, Chap. III, Exercise (3.14)]). We introduce the functions

$$F: L^{N+1}(0, 1) \subset H \mapsto \mathbb{R}, \quad F(x) := \int_0^1 d\xi \int_0^{x(\xi)} f(\xi, s) ds,$$

$$V_\alpha: H \mapsto [0, +\infty),$$

$$V_\alpha(x) := \frac{1}{2} \int_0^1 [(\alpha + x(\xi))^-]^2 d\xi = \frac{1}{2} [d(x, K_\alpha)]^2,$$

where N is the degree of p in (H4) and $d(x, K_\alpha)$ denotes the distance in H of x from the closed convex set K_α , $\alpha \geq 0$. Notice that $\mu(L^{N+1}(0, 1)) = 1$,

and by (H2), (H3), and (H4), $\exp(-F)$ is well defined and satisfies for all $x \in H$,

$$\exp(-2F(x)) \leq \exp\left(\langle a, x \rangle + \frac{\lambda}{2} \|x\|^2\right) \leq \exp(C \|a\|^2) \exp\left(\frac{\rho}{2} \|x\|^2\right) \quad (17)$$

for some $\lambda < \rho < \pi^2$, $C > 0$. Since $-2A \geq \pi^2 > \rho$, we have:

$$\begin{aligned} \int_H \exp(-2F(x)) \mu(dx) &\leq e^{C \|a\|^2} \int_H \exp\left(\frac{\rho}{2} \|x\|^2\right) \mathcal{N}(0, (-2A)^{-1}) \\ &= e^{C \|a\|^2} \frac{1}{\sqrt{\det(I + \rho(2A)^{-1})}} < \infty. \end{aligned} \quad (18)$$

V_α is Fréchet differentiable on H with Lipschitz-continuous gradient $\nabla V_\alpha(x) = -(\alpha + x(\cdot))^-$, $x \in H$. F belongs to the Sobolev space $W^{1,p}(H, \mu)$ for all $p \in [1, \infty[$, with gradient $\nabla F(x) = f(\cdot, x(\cdot))$, $x \in L^{2N}(0, 1)$, and $\mu(L^{2N}(0, 1)) = 1$. Equation (6) can now be written in the following form:

$$\begin{cases} dX_\varepsilon^\alpha = \left(AX_\varepsilon^\alpha - \nabla F(X_\varepsilon^\alpha) - \frac{1}{\varepsilon} \nabla V_\alpha(X_\varepsilon^\alpha) \right) dt + dW \\ X_\varepsilon^\alpha(0, x) = x \in H \end{cases} \quad (19)$$

If $\varepsilon > 0$, (19) is a gradient system, namely (see [4, Sect. 8.6], and [5, Chap. 2]):

PROPOSITION 1. *If $\varepsilon > 0$, then $\forall \alpha \geq 0$, setting*

$$Z_{\alpha, \varepsilon} := \int_H e^{-2F - (2V_\alpha/\varepsilon)} d\mu > 0,$$

the probability measure on H ,

$$\nu_{\alpha, \varepsilon}^F(dx) := \frac{1}{Z_{\alpha, \varepsilon}} \exp\left\{-2F(x) - \frac{2}{\varepsilon} V_\alpha(x)\right\} \mu(dx),$$

is symmetrizing for the process $\{X_\alpha^\varepsilon(t, x) : t \geq 0, x \in H\}$.

By (16), if $\alpha > 0$ then $\mu(K_\alpha) > 0$, and we can introduce the probability measures ν_α and ν_α^F on K_α , with $Z_\alpha := \int_{K_\alpha} e^{-2F} d\mu > 0$:

$$d\nu_\alpha := \frac{1}{\mu(K_\alpha)} 1_{K_\alpha} d\mu, \quad d\nu_\alpha^F := \frac{1}{Z_\alpha} 1_{K_\alpha} e^{-2F} d\mu. \quad (20)$$

THEOREM 2. *If $\alpha > 0$, ν_α^F is a symmetrizing measure for the processes $\{X_\alpha(t, x) : t \geq 0, x \in K_\alpha\}$ and $\{X_\alpha(t, x) : t \geq 0, x \in K_\alpha \cap C_0(0, 1)\}$.*

Proof. First, we have:

$$\frac{1}{Z_\alpha} 1_{K_\alpha}(x) e^{-2F(x)} = \lim_{\varepsilon \downarrow 0} \frac{1}{Z_{\alpha, \varepsilon}} \exp \left\{ -2F(x) - \frac{2}{\varepsilon} V_\alpha(x) \right\} \quad \forall x \in H,$$

$$\frac{1}{Z_{\alpha, \varepsilon}} \exp \left\{ -2F - \frac{2}{\varepsilon} V_\alpha \right\} \leq \frac{1}{Z_{\alpha, 1}} \exp \left\{ \langle a, x \rangle + \frac{\lambda}{2} \|x\|^2 \right\} \quad \forall \varepsilon \in]0, 1].$$

Then, by (13), (17), and (18) and by the dominated convergence theorem, we obtain:

$$v_\alpha^F(\psi P_\alpha(t) \varphi) = \lim_{\varepsilon \downarrow 0} v_{\alpha, \varepsilon}^F(\psi P_\alpha^\varepsilon(t) \varphi) = \lim_{\varepsilon \downarrow 0} v_{\alpha, \varepsilon}^F(\varphi P_\alpha^\varepsilon(t) \psi) = v_\alpha^F(\varphi P_\alpha(t) \psi)$$

for all $\varphi, \psi \in C_b(H)$. Therefore, v_α^F is symmetrizing measure for $\{X_\alpha(t, x) : t \geq 0, x \in K_\alpha\}$. Finally, $v_\alpha^F(K_\alpha \cap C_0(0, 1)) = 1$ and, by Theorem 1, the set $K_\alpha \cap C_0(0, 1)$ is invariant for $\{X_\alpha(t, \cdot) : t \geq 0\}$, i.e., $x \in K_\alpha \cap C_0(0, 1)$ implies $X_\alpha(t, x) \in K_\alpha \cap C_0(0, 1)$ for all $t \geq 0$, a.s. ■

5. CONVERGENCE OF v_α TO v

Let $(B_\tau)_{\tau \in [0, 1]}$ be a 3-dimensional Brownian motion. We denote by v the law of the 3-d Bessel Bridge, namely of the modulus of B , conditioned to be equal to 0 at $\tau = 1$. The probability measure v is concentrated on $K_0 \cap C_0(0, 1)$. In this section we prove the following:

THEOREM 3. *The measures $dv_\alpha = (1/\mu(K_\alpha)) 1_{K_\alpha} d\mu$, $\alpha > 0$, converge weakly as $\alpha \downarrow 0$ to the law v of the 3-d Bessel bridge.*

We recall the following result from [1]:

THEOREM 4. *Let $(e_\tau)_{\tau \in [0, 1]}$ be a 3-d Bessel bridge, and let ζ be a random variable with uniform distribution on $[0, 1]$ and independent of e . Then the process:*

$$(\beta_\tau)_{\tau \in [0, 1]}, \quad \beta_\tau := e_{\tau \oplus \zeta} - e_\zeta,$$

where \oplus denotes the sum mod 1, is a Brownian bridge.

Theorem 3 was proved in [8]. We give here a proof, based on Theorem 4, which seems to be promising for further developments.

We set $e_\tau: C_0(0, 1) \mapsto \mathbb{R}$, $e_\tau(x) := x(\tau)$, $\tau \in [0, 1]$. Then $(e_\tau)_{\tau \in [0, 1]}$ is a 3-d Bessel bridge under v . By Theorem 4 and (16), we have for $\alpha > 0$, $\varphi \in C_b(H)$,

$$v_\alpha(\varphi) = \frac{1}{1 - \exp\{-2\alpha^2\}} \int_0^1 v(\varphi(e_{(\cdot \oplus r)} - e_r) 1_{(e_r \leq \alpha)}) dr, \quad (21)$$

since $\{e_{(\cdot \oplus r)} - e_r \geq -\alpha\} = \{e_r \leq \alpha\}$.

LEMMA 6. For all $r \in [0, 1]$, there exists a regular conditional distribution $\{v(\cdot | e_r = y) : y \geq 0\}$ of v given e_r , such that, setting

$$\delta_\varphi(r, y) := v(\varphi(e_{(\cdot \oplus r)} - e_r) | e_r = y), \quad \varphi \in C_b(H), \quad r \in]0, 1[, \quad y \geq 0,$$

we have for all $y \geq 0$:

$$\lim_{r \downarrow 0} \delta_\varphi(r, \sqrt{r(1-r)} y) = \lim_{r \uparrow 1} \delta_\varphi(r, \sqrt{r(1-r)} y) = v(\varphi).$$

Proof. Let $(B_\tau)_{\tau \in [0, \infty)}$ and $(\hat{B}_\tau)_{\tau \in [0, \infty)}$ be two independent 3-d Brownian motions and $r \in]0, 1[$. Denoting by $|\cdot|$ the euclidean norm in \mathbb{R}^3 , we set $b := |B|$, $\hat{b} := |\hat{B}|$,

$$\begin{aligned} \beta(z)(\tau) &:= B_\tau - \tau B_1 + \tau z, & \hat{\beta}(z)(\tau) &:= \hat{B}_\tau - \tau \hat{B}_1 + \tau z, & \tau \in [0, 1], & z \in \mathbb{R}^3, \\ \pi_r, \hat{\pi}_r &: L^2(0, \infty) \times L^2(0, \infty) \mapsto L^2(0, 1), \\ \pi_r(c, d)(\tau) &:= 1_{[0, r]}(\tau) c(\tau) + 1_{]r, 1]}(\tau) d(1 - \tau), \\ \hat{\pi}_r(c, d)(\tau) &:= 1_{[0, 1-r]}(\tau) d(1 - r - \tau) + 1_{]1-r, 1]}(\tau) c(\tau + r - 1). \end{aligned} \tag{22}$$

For all $\varphi \in C_b(H)$, we set

$$v(\varphi(e) | e_r = y) := \mathbb{E}[\varphi(\pi_r(b, \hat{b})) | b_r = y = \hat{b}_{1-r}], \quad y \geq 0, \tag{23}$$

$$\mathbb{E}[\varphi(b) | b(1) = y] := \int_{S^2} \sigma(dn) \mathbb{E}[\varphi(|\beta(y n)|)], \quad y \geq 0, \tag{24}$$

where S^2 is the unitary sphere in \mathbb{R}^3 and $\sigma(dn)$ is the normalized uniform distribution on S^2 . Then (23), respectively (24), is a regular conditional distribution of v given e_r , resp. of $\mathbb{P}(b \in \cdot)$ given $b(1)$. In particular, the law of $|\beta(0)|$ is equal to v . By (22) and (23) we have:

$$v(\varphi(e_{(\cdot \oplus r)} - e_r) | e_r = y) = \mathbb{E}[\varphi(\hat{\pi}_r(b, \hat{b}) - y) | b_r = y = \hat{b}_{1-r}]. \tag{25}$$

Identifying $h \in L^2(0, 1)$ with $h 1_{[0, 1]} \in L^2(0, \infty)$, we set $\varphi_r: H \times H \mapsto \mathbb{R}$,

$$\varphi_r(h, k) := \varphi(\hat{\pi}_r(\sqrt{r} h(\cdot / r), \sqrt{1-r} k(\cdot / (1-r))) - \sqrt{r(1-r)} y).$$

Since for $\gamma > 0$, $\sqrt{\gamma} B_{(\cdot/\gamma)}$ is still a 3-d Brownian motion, we obtain by (25):

$$\begin{aligned} \delta_\varphi(r, \sqrt{r(1-r)} y) &= v(\varphi(e_{(\cdot \oplus r)} - e_r) | e_r = \sqrt{r(1-r)} y) \\ &= \mathbb{E}[\varphi(\hat{\pi}_r(b, \hat{b}) - \sqrt{r(1-r)} y) | b_r = \sqrt{r(1-r)} y = \hat{b}_{1-r}] \\ &= \mathbb{E}[\varphi_r(b, \hat{b}) | b_1 = \sqrt{1-r} y, \hat{b}_1 = \sqrt{r} y]. \end{aligned} \tag{26}$$

Since for all $n \in S^2$ and $y \geq 0$,

$$\lim_{r \downarrow 0} \varphi_r(|\beta(\sqrt{1-r} \, yn)|, |\hat{\beta}(\sqrt{r} \, yn)|) = \varphi(|\hat{\beta}(0)(1 - \cdot)|),$$

$$\lim_{r \uparrow 1} \varphi_r(|\beta(\sqrt{1-r} \, yn)|, |\hat{\beta}(\sqrt{r} \, yn)|) = \varphi(|\beta(0)|),$$

and since ν is invariant by the time-change $\tau \mapsto 1 - \tau$, the thesis follows by (24)–(26) and by the dominated convergence theorem. ■

Proof of Theorem 3. We split the integral on $[0, 1]$ in (21) into two integrals on $[0, 1/2]$ and $[1/2, 1]$, respectively. Conditioning with respect to e_r and setting $c_\alpha := (1 - \exp\{-2\alpha^2\})$, we obtain

$$\begin{aligned} & \frac{1}{c_\alpha} \int_0^{1/2} dr \, \nu(\varphi(e_{(\cdot \oplus r)} - e_r) 1_{(e_r \leq \alpha)}) \\ &= \frac{1}{c_\alpha} \int_0^{1/2} dr \int_0^\alpha dy \sqrt{\frac{2}{\pi[r(1-r)]^3}} y^2 \exp\left\{-\frac{y^2}{2r(1-r)}\right\} \delta_\varphi(r, y) \\ &= \frac{1}{c_\alpha} \int_0^{1/2} dr \int_0^{\alpha/\sqrt{r(1-r)}} dy \sqrt{\frac{2}{\pi}} y^2 \exp\left\{-\frac{y^2}{2}\right\} \delta_\varphi(r, \sqrt{r(1-r)} y) \\ &= \frac{1}{c_\alpha} \sqrt{\frac{2}{\pi}} \int_0^{2\alpha} dy \exp\left\{-\frac{y^2}{2}\right\} y^2 \int_0^{1/2} dr \delta_\varphi(r, \sqrt{r(1-r)} y) \\ & \quad + \frac{\alpha^2}{c_\alpha} \sqrt{\frac{2}{\pi}} \int_{2\alpha}^{+\infty} dy \exp\left\{-\frac{y^2}{2}\right\} \left(\frac{y}{\alpha}\right)^2 \int_0^{\rho(\alpha, y)} dr \delta_\varphi(r, \sqrt{r(1-r)} y) \\ &=: I_1(\alpha) + I_2(\alpha), \quad \rho(\alpha, y) := \frac{1}{2} \left(1 - \sqrt{1 - \left(\frac{2\alpha}{y}\right)^2}\right) \sim \left(\frac{\alpha}{y}\right)^2 \end{aligned}$$

as $\alpha \downarrow 0$, $y > 0$. It is easy to see that $\lim_{\alpha \downarrow 0} I_1(\alpha) = 0$, while $I_2(\alpha)$ tends to $(1/2) \nu(\varphi)$ by Lemma 6 and the dominated convergence theorem. Since analogous computations hold for the integral on $[1/2, 1]$, we obtain that $\nu_\alpha(\varphi)$ converges to $\nu(\varphi)$ and Theorem 3 is proved. ■

6. 3-D BESSEL BRIDGE AS SYMMETRIZING MEASURE FOR X_0

In this section we prove that the probability measure on K_0 :

$$d\nu^F := \frac{1}{\int_{K_0} \exp\{-2F\} d\nu} \exp\{-2F\} d\nu, \quad (27)$$

is well defined, that v_α^F , defined in (20), converges weakly to v^F as $\alpha \downarrow 0$, and that X_0 is symmetric with respect to v^F . The difficulty is that $\exp(-2F)$ is not bounded, so that we cannot apply directly Theorem 3.

LEMMA 7. $v(e^{-2F}) \in]0, \infty[$ and v_α^F converges weakly to v^F as $\alpha \downarrow 0$.

Proof. We retain the notations of the proof of Lemma 6. Moreover, we set for all $\varphi \in C_b(H)$, $\varphi^F := \varphi e^{-2F}$. By Lemma 5 the law of $\beta(0)$ on $H \times H \times H$ is $\mu \otimes \mu \otimes \mu = \mathcal{N}(0, Q)$, $Q := (-2A)^{-1} \oplus (-2A)^{-1} \oplus (-2A)^{-1}$. Since $-2A \geq \pi^2 > \lambda$, we have by (17), (H4) and by $\mu(L^{N+1}(0, 1)) = 1$:

$$\begin{aligned} \int \exp\{-2F\} dv &= \mathbb{E}[\exp\{-2F(|\beta(0)|)\}] \\ &= \int_{H^3} \exp\{-2F(|z|)\} \mathcal{N}(0, Q)(dz) \in]0, \infty[. \end{aligned}$$

Notice that

$$\exp\left\{\frac{\lambda}{2} \|\beta(yn)\|^2\right\} \leq e^{Cy^2} \exp\left\{\frac{\rho}{2} \|\beta(0)\|^2\right\} \quad (28)$$

for some $\lambda < \rho < \pi^2$, $C > 0$, and:

$$\mathbb{E}\left[\exp\left\{\frac{\rho}{2} \|\beta(0)\|^2\right\}\right] = \frac{1}{\sqrt{\det(I - \rho Q)}} < \infty. \quad (29)$$

By (23), (24), (28), and (29), $v(e^{-2F}(e) | e(r) = y) < \infty$, and therefore $\delta_{(\varphi^F)}(r, y)$ is well defined for all $r \in]0, 1[$ and $y \geq 0$. Arguing as in the proof of Lemma 6, by the dominated convergence theorem we have for all $\varphi \in C_b(H)$ and $y \geq 0$,

$$\lim_{r \downarrow 0} \delta_{(\varphi^F)}(r, \sqrt{r(1-r)} y) = \lim_{r \uparrow 1} \delta_{(\varphi^F)}(r, \sqrt{r(1-r)} y) = v(\varphi^F), \quad (30)$$

and the thesis follows proceeding as in the proof of Theorem 3. ■

THEOREM 5. v^F is a symmetrizing measure for the processes $\{X_0(t, x) : t \geq 0, x \in K_0\}$ and $\{X_0(t, x) : t \geq 0, x \in K_0 \cap C_0(0, 1)\}$.

Proof. Arguing as in the proof of Theorem 2, the thesis follows from Lemmas 1, 3, 4, and 7, Theorems 1 and 2, and from $v^F(K_0 \cap C_0(0, 1)) = v(K_0 \cap C_0(0, 1)) = 1$. ■

ACKNOWLEDGMENTS

The author thanks the University of Bielefeld for a very pleasant stay, during which part of this work was done.

Note added in proof. While correcting proofs, we were informed that the result of Theorem 5 in this paper was stated independently in a recent preprint [9] of T. Funaki and S. Olla.

REFERENCES

1. P. Biane, Relations entre pont et excursion du mouvement Brownien réel, *Ann. Inst. Henri Poincaré* **22** (1986), 1–7.
2. G. Da Prato, Monotone gradient systems in L^2 spaces, to appear in “Proceedings of the Ascona Conference on Stochastic Analysis, Random Fields and Applications,” 1999.
3. G. Da Prato and J. Zabczyk, “Stochastic Equations in Infinite Dimensions,” Encyclopedia of Mathematics and its Applications, Cambridge Univ. Press, Cambridge, UK, 1992.
4. G. Da Prato and J. Zabczyk, “Ergodicity for Infinite Dimensional Systems,” London Mathematical Society Lecture Notes, Vol. 229, Cambridge Univ. Press, Cambridge, UK, 1996.
5. Z. M. Ma and M. Röckner, “Introduction to the Theory of (Non Symmetric) Dirichlet Forms,” Springer-Verlag, Berlin/New York, 1992.
6. D. Nualart and E. Pardoux, White noise driven quasilinear SPDEs with reflection, *Probab. Theory Related Fields* **93** (1992), 77–89.
7. D. Revuz and M. Yor, “Continuous Martingales and Brownian Motion,” Springer-Verlag, Berlin/New York, 1991.
8. R. T. Durrett, D. L. Iglehart, and D. R. Miller, Weak convergence to Brownian meander and Brownian excursion, *Ann. Probab.* **5**, No. 1 (1977), 117–129.
9. T. Funaki and S. Olla, Fluctuations for $\nabla\phi$ interface model on a wall, preprint, 2000.