

Optimization under Uncertainty using Exponential Cones

CHEN Li

Institute of Operations Research and Analytics
National University of Singapore

June 16, 2022

Outline

Introduction

The Key Methodology of The ECP Approach to EV Charging

Robust CARA Optimization

► Deterministic optimization

$$\max_{\mathbf{x} \in \mathcal{X}} f(\mathbf{x}, \mathbf{z})$$

where \mathbf{z} is the known parameter.

Optimization under uncertainty

- ▶ Deterministic optimization

$$\max_{\mathbf{x} \in \mathcal{X}} f(\mathbf{x}, \mathbf{z})$$

where \mathbf{z} is the known parameter.

- ▶ Uncertainty in problem parameters
 - ▶ Risk: known distribution $\tilde{\mathbf{z}} \sim \mathbb{P}$

Optimization under uncertainty

- ▶ Deterministic optimization

$$\max_{\mathbf{x} \in \mathcal{X}} f(\mathbf{x}, \mathbf{z})$$

where \mathbf{z} is the known parameter.

- ▶ Uncertainty in problem parameters
 - ▶ Risk: known distribution $\tilde{\mathbf{z}} \sim \mathbb{P}$
 - ▶ Example: flip a fair coin

Optimization under uncertainty

- ▶ Deterministic optimization

$$\max_{\mathbf{x} \in \mathcal{X}} f(\mathbf{x}, \mathbf{z})$$

where \mathbf{z} is the known parameter.

- ▶ Uncertainty in problem parameters

- ▶ Risk: known distribution $\tilde{\mathbf{z}} \sim \mathbb{P}$

- ▶ Example: flip a fair coin

- ▶ Ambiguity: unknown distribution $\tilde{\mathbf{z}} \in \mathbb{P} \in \mathcal{F}$

Optimization under uncertainty

- ▶ Deterministic optimization

$$\max_{\mathbf{x} \in \mathcal{X}} f(\mathbf{x}, \mathbf{z})$$

where \mathbf{z} is the known parameter.

- ▶ Uncertainty in problem parameters

- ▶ Risk: known distribution $\tilde{\mathbf{z}} \sim \mathbb{P}$

- ▶ Example: flip a fair coin

- ▶ Ambiguity: unknown distribution $\tilde{\mathbf{z}} \in \mathbb{P} \in \mathcal{F}$

- ▶ Example: we do not know whether the coin is unbiased

Optimization under uncertainty

- ▶ Deterministic optimization

$$\max_{\mathbf{x} \in \mathcal{X}} f(\mathbf{x}, \mathbf{z})$$

where \mathbf{z} is the known parameter.

- ▶ Uncertainty in problem parameters
 - ▶ Risk: known distribution $\tilde{\mathbf{z}} \sim \mathbb{P}$
 - ▶ Example: flip a fair coin
 - ▶ Ambiguity: unknown distribution $\tilde{\mathbf{z}} \in \mathbb{P} \in \mathcal{F}$
 - ▶ Example: we do not know whether the coin is unbiased
- ▶ How to rank decisions under uncertainty?
 - ▶ Decision criteria

Decision-making under uncertainty

- ▶ Expected utility theory [VNM47]

$$\max_{x \in \mathcal{X}} \mathbb{E}_{\mathbb{P}} [u(f(x, \tilde{z}))]$$

where the utility function $u : \mathbb{R} \rightarrow \mathbb{R}$ is increasing.

Decision-making under uncertainty

- ▶ Expected utility theory [VNM47]

$$\max_{x \in \mathcal{X}} \mathbb{E}_{\mathbb{P}} [u(f(x, \tilde{z}))]$$

where the utility function $u : \mathbb{R} \rightarrow \mathbb{R}$ is increasing.

- ▶ Risk averse: $u(\cdot)$ is concave

Decision-making under uncertainty

- ▶ Expected utility theory [VNM47]

$$\max_{x \in \mathcal{X}} \mathbb{E}_{\mathbb{P}} [u(f(x, \tilde{z}))]$$

where the utility function $u : \mathbb{R} \rightarrow \mathbb{R}$ is increasing.

- ▶ Risk averse: $u(\cdot)$ is concave
- ▶ Maxmin expected utility [GS89]

$$\max_{x \in \mathcal{X}} \inf_{\mathbb{P} \in \mathcal{F}} \mathbb{E}_{\mathbb{P}} [u(f(x, \tilde{z}))]$$

- ▶ Many others...

How to solve optimization problems under uncertainty?

► Stochastic Optimization

$$\max_{\mathbf{x} \in \mathcal{X}} \mathbb{E}_{\mathbb{P}} [f(\mathbf{x}, \tilde{\mathbf{z}})]$$

Fact: evaluating $\mathbb{E}_{\mathbb{P}} [\max\{\mathbf{a}^{\top} \tilde{\mathbf{z}} - b, 0\}]$ is #P-hard for given $\mathbf{a} \in \mathbb{R}_+^n$, $b \in \mathbb{R}_+$, \tilde{z}_i 's are independent uniform on $[0, 1]$.

How to solve optimization problems under uncertainty?

- ▶ Stochastic Optimization

$$\max_{\mathbf{x} \in \mathcal{X}} \mathbb{E}_{\mathbb{P}} [f(\mathbf{x}, \tilde{\mathbf{z}})]$$

- ▶ Random approximation: Sample Average Approximation (SAA)

$$\max_{\mathbf{x} \in \mathcal{X}} \frac{1}{S} \sum_{s \in [S]} f(\mathbf{x}, \hat{\mathbf{z}}^s)$$

- ▶ General: sampling oracle
- ▶ Effective: statistics and optimization perspectives
- ▶ Disadvantage: curse of dimensionality

Computation challenges

How to solve optimization problems under uncertainty?

- ▶ Stochastic Optimization

$$\max_{\mathbf{x} \in \mathcal{X}} \mathbb{E}_{\mathbb{P}} [f(\mathbf{x}, \tilde{\mathbf{z}})]$$

- ▶ Random approximation: Sample Average Approximation (SAA)

$$\max_{\mathbf{x} \in \mathcal{X}} \frac{1}{S} \sum_{s \in [S]} f(\mathbf{x}, \hat{\mathbf{z}}^s)$$

- ▶ General: sampling oracle
 - ▶ Effective: statistics and optimization perspectives
 - ▶ Disadvantage: curse of dimensionality
- ▶ Deterministic approximation: bound $\mathbb{E}_{\mathbb{P}} [f(\mathbf{x}, \tilde{\mathbf{z}})]$
 - ▶ Ad hoc: utilize special structures of f and \mathbb{P} , e.g., moments
 - ▶ Advantages: scalability, handle decision-dependent uncertainty
 - ▶ Disadvantages: suboptimality

How to solve optimization problems under uncertainty?

- ▶ Stochastic Optimization

$$\max_{\mathbf{x} \in \mathcal{X}} \mathbb{E}_{\mathbb{P}} [f(\mathbf{x}, \tilde{\mathbf{z}})]$$

- ▶ Distributionally Robust Optimization (DRO)

$$\max_{\mathbf{x} \in \mathcal{X}} \inf_{\mathbb{P} \in \mathcal{F}} \mathbb{E}_{\mathbb{P}} [f(\mathbf{x}, \tilde{\mathbf{z}})]$$

- ▶ Tractability depends on underlying problem structures and choices of ambiguity sets
- ▶ Mostly duality-based reformulation techniques: DRO \rightarrow RO \rightarrow convex (conic) optimization

Computation challenges

How to solve optimization problems under uncertainty?

- ▶ Stochastic Optimization

$$\max_{\mathbf{x} \in \mathcal{X}} \mathbb{E}_{\mathbb{P}} [f(\mathbf{x}, \tilde{\mathbf{z}})]$$

- ▶ Distributionally Robust Optimization (DRO)

$$\max_{\mathbf{x} \in \mathcal{X}} \inf_{\mathbb{P} \in \mathcal{F}} \mathbb{E}_{\mathbb{P}} [f(\mathbf{x}, \tilde{\mathbf{z}})]$$

Why do we care tractability?

... can be treated as actual sources of “immunized against uncertainty” decisions only if these problems are computationally tractable; when that is not the case, these settings become more wishful thinking than actual decision-making tool.

— Robust Optimization by Ben-Tal, El Ghaoui and Nemirovski

Exponential Conic Optimization

- ▶ Exponential cone: 3-dimensional closed convex cone

$$\mathcal{K}_{\text{exp}} := \text{cl} \{ (x_1, x_2, x_3) \mid x_1 \geq x_2 \exp(x_3/x_2), x_2 > 0 \}$$

i.e. closure of epigraph of perspective of exponential function.

Exponential Conic Optimization

- ▶ Exponential cone: 3-dimensional closed convex cone

$$\mathcal{K}_{\text{exp}} := \text{cl} \{ (x_1, x_2, x_3) \mid x_1 \geq x_2 \exp(x_3/x_2), x_2 > 0 \}$$

i.e. closure of epigraph of perspective of exponential function.

- ▶ Exponential cone programming (ECP):

$$\begin{array}{ll} \min_{\mathbf{x}} & \mathbf{c}^\top \mathbf{x} \\ \text{s.t.} & \mathbf{Ax} - \mathbf{b} \in \mathcal{K}_{\text{exp}} \times \cdots \times \mathcal{K}_{\text{exp}} \\ & \mathbf{Fx} = \mathbf{g} \end{array}$$

- ▶ Generalization of linear programming (LP)
- ▶ Great modeling power:
 - ▶ exponential cone \succeq power cone \succeq second-order cone
 - ▶ Many convex constraints involving exponential or logarithm functions are \mathcal{K}_{exp} -representable, e.g.

$$t \leq -x \log x \iff (1, x, t) \in \mathcal{K}_{\text{exp}}$$

- ▶ Computational advances: efficient interior point algorithm based solvers such as MOSEK

What is this thesis about?

- ▶ Goal: Leverage modeling advantages and computational advances of ECP to solve optimization problems under uncertainty.

What is this thesis about?

- ▶ Goal: Leverage modeling advantages and computational advances of ECP to solve optimization problems under uncertainty.
- ▶ How does ECP related to optimization under uncertainty?
 - ▶ At a high level, we inject information of uncertain parameters into optimization models through moment generating functions (MGFs) and develop the deterministic ECP approximations.

What is this thesis about?

- ▶ Goal: Leverage modeling advantages and computational advances of ECP to solve optimization problems under uncertainty.
- ▶ How does ECP related to optimization under uncertainty?
 - ▶ At a high level, we inject information of uncertain parameters into optimization models through moment generating functions (MGFs) and develop the deterministic ECP approximations.
- ▶ Motivation of using MGFs:
 - ▶ Decomposability under independence of random variables
 - ▶ Close relation to exponential utility function and relative entropy based ambiguity set
 - ▶ The log-MGFs of many random variables are \mathcal{K}_{exp} -representable.

What is this thesis about?

- ▶ Goal: Leverage modeling advantages and computational advances of ECP to solve optimization problems under uncertainty.
- ▶ How does ECP related to optimization under uncertainty?
 - ▶ At a high level, we inject information of uncertain parameters into optimization models through moment generating functions (MGFs) and develop the deterministic ECP approximations.
- ▶ Motivation of using MGFs:
 - ▶ Decomposability under independence of random variables
 - ▶ Close relation to exponential utility function and relative entropy based ambiguity set
 - ▶ The log-MGFs of many random variables are \mathcal{K}_{exp} -representable.
- ▶ Structure:
 - ▶ An ECP approach to electric vehicle (EV) charging management
 - ▶ **Robust CARA Optimization**
 - ▶ Extensions and future work

Outline

Introduction

The Key Methodology of The ECP Approach to EV Charging

Robust CARA Optimization

A large-scale stochastic program

The EV charging scheduling problem is

$$\min_{\mathbf{x} \in \mathcal{X}} \mathbb{E}_{\mathbb{P}} [c(\mathbf{x}, \tilde{\mathbf{z}})] \quad (\text{EV-SP})$$

where

$$c(\mathbf{x}, \tilde{\mathbf{z}}) \triangleq \sum_{s \in [T]} e_s \sum_{v \in \mathcal{V}_s} x_{v,s} \tilde{z}_v + d \max_{t \in [T]} \left\{ \sum_{v \in \mathcal{V}_t} x_{v,t} \tilde{z}_v \right\}$$

A large-scale stochastic program

The EV charging scheduling problem is

$$\min_{\mathbf{x} \in \mathcal{X}} \mathbb{E}_{\mathbb{P}} [c(\mathbf{x}, \tilde{\mathbf{z}})] \quad (\text{EV-SP})$$

where

$$c(\mathbf{x}, \tilde{\mathbf{z}}) \triangleq \sum_{s \in [T]} e_s \sum_{v \in \mathcal{V}_s} x_{v,s} \tilde{z}_v + d \max_{t \in [T]} \left\{ \sum_{v \in \mathcal{V}_t} x_{v,t} \tilde{z}_v \right\}$$

Computationally challenging for SAA due to large scale of the problem:
more than 80,000 random variables and 700,000 decision variables!

ECP approximation in uncapacitated case

In uncapacitated case, $\tilde{z} \sim \mathbb{P}$ are independent Poisson with rate λ .

► Difficulty:

$$\sum_{s \in [T]} e_s f_s(\mathbf{x}, \lambda) + d \mathbb{E}_{\mathbb{P}} \left[\max_{t \in [T]} \{f_t(\mathbf{x}, \tilde{z})\} \right]$$

ECP approximation in uncapacitated case

In uncapacitated case, $\tilde{z} \sim \mathbb{P}$ are independent Poisson with rate λ .

- Difficulty:

$$\sum_{s \in [T]} e_s f_s(\mathbf{x}, \boldsymbol{\lambda}) + d \mathbb{E}_{\mathbb{P}} \left[\max_{t \in [T]} \{f_t(\mathbf{x}, \tilde{\mathbf{z}})\} \right]$$

- Motivation of using MGF
 - Determines non-negative random variables uniquely
 - Utilize independence: $\mathbb{E}_{\mathbb{P}} \left[e^{\sum_v \theta_v \tilde{z}_v} \right] = \prod_v \mathbb{E}_{\mathbb{P}} \left[e^{\theta_v \tilde{z}_v} \right]$

ECP approximation in uncapacitated case

In uncapacitated case, $\tilde{z} \sim \mathbb{P}$ are independent Poisson with rate λ .

► Bound the order statistic:

$$\begin{aligned} & \mathbb{E}_{\mathbb{P}} \left[\max_{t \in [T]} \{f_t(\mathbf{x}, \tilde{z})\} \right] \\ \leq & \inf_{\mu > 0} \mu \log \sum_{t \in [T]} \mathbb{E}_{\mathbb{P}} \left[\exp \left(\frac{f_t(\mathbf{x}, \tilde{z}) - f_t(\mathbf{x}, \lambda)}{\mu} \right) \right] + \max_{t \in [T]} \{f_t(\mathbf{x}, \lambda)\} \end{aligned}$$

ECP approximation in uncapacitated case

In uncapacitated case, $\tilde{z} \sim \mathbb{P}$ are independent Poisson with rate λ .

- Bound the order statistic:

$$\begin{aligned} & \mathbb{E}_{\mathbb{P}} \left[\max_{t \in [T]} \{f_t(\mathbf{x}, \tilde{z})\} \right] \\ & \leq \inf_{\mu > 0} \mu \log \sum_{t \in [T]} \mathbb{E}_{\mathbb{P}} \left[\exp \left(\frac{f_t(\mathbf{x}, \tilde{z}) - f_t(\mathbf{x}, \lambda)}{\mu} \right) \right] + \max_{t \in [T]} \{f_t(\mathbf{x}, \lambda)\} \end{aligned}$$

- Exploit stochastic independence:

$$\begin{aligned} & \mu \log \mathbb{E}_{\mathbb{P}} \left[\exp \left(\frac{f_t(\mathbf{x}, \tilde{z}) - f_t(\mathbf{x}, \lambda)}{\mu} \right) \right] \\ & = \sum_{v \in \mathcal{V}_t} \mu \log \mathbb{E}_{\mathbb{P}} \left[\exp \left(\frac{x_{v,t} \tilde{z}_v}{\mu} \right) \right] - f_t(\mathbf{x}, \lambda) \end{aligned}$$

ECP approximation in uncapacitated case

In uncapacitated case, $\tilde{z} \sim \mathbb{P}$ are independent Poisson with rate λ .

- Bound the order statistic:

$$\begin{aligned} & \mathbb{E}_{\mathbb{P}} \left[\max_{t \in [T]} \{f_t(\mathbf{x}, \tilde{z})\} \right] \\ & \leq \inf_{\mu > 0} \mu \log \sum_{t \in [T]} \mathbb{E}_{\mathbb{P}} \left[\exp \left(\frac{f_t(\mathbf{x}, \tilde{z}) - f_t(\mathbf{x}, \lambda)}{\mu} \right) \right] + \max_{t \in [T]} \{f_t(\mathbf{x}, \lambda)\} \end{aligned}$$

- Exploit stochastic independence:

$$\begin{aligned} & \mu \log \mathbb{E}_{\mathbb{P}} \left[\exp \left(\frac{f_t(\mathbf{x}, \tilde{z}) - f_t(\mathbf{x}, \lambda)}{\mu} \right) \right] \\ & = \sum_{v \in \mathcal{V}_t} \mu \log \mathbb{E}_{\mathbb{P}} \left[\exp \left(\frac{x_{v,t} \tilde{z}_v}{\mu} \right) \right] - f_t(\mathbf{x}, \lambda) \end{aligned}$$

- Exponential cone representable log-MGF of Poisson variables:

$$\mu \log \mathbb{E}_{\mathbb{P}} \left[\exp \left(\frac{x_{v,t} \tilde{z}_v}{\mu} \right) \right] = \lambda_v (\mu e^{x_{v,t}/\mu} - \mu) \text{ is } \mathcal{K}_{\text{exp}}\text{-representable}$$

ECP approximation in capacitated case

- ▶ Difficulty: No independent Poisson \Rightarrow No closed-form MGF

ECP approximation in capacitated case

- ▶ Difficulty: No independent Poisson \Rightarrow No closed-form MGF
- ▶ Upper bounds via DRO:

$$\mathbb{P} \in \mathcal{F} \Rightarrow \mathbb{E}_{\mathbb{P}} [c(\mathbf{x}, \tilde{\mathbf{z}})] \leq \sup_{\mathbb{P} \in \mathcal{F}} \mathbb{E}_{\mathbb{P}} [c(\mathbf{x}, \tilde{\mathbf{z}})]$$

ECP approximation in capacitated case

- ▶ Difficulty: No independent Poisson \Rightarrow No closed-form MGF
- ▶ Upper bounds via DRO:

$$\mathbb{P} \in \mathcal{F} \Rightarrow \mathbb{E}_{\mathbb{P}} [c(\mathbf{x}, \tilde{\mathbf{z}})] \leq \sup_{\mathbb{P} \in \mathcal{F}} \mathbb{E}_{\mathbb{P}} [c(\mathbf{x}, \tilde{\mathbf{z}})]$$

- ▶ Moments: truncated Poisson $\mathbb{P} \in \mathcal{F}^1$

$$\mathcal{F}^1 \triangleq \left\{ \mathbb{P} \mid \begin{array}{l} \tilde{\mathbf{z}} \sim \mathbb{P} \\ \log \mathbb{E}_{\mathbb{P}} \left[\exp \left(\boldsymbol{\theta}^\top \tilde{\mathbf{z}} \right) \right] \leq \sum_{v \in [V]} \lambda_v \left(e^{\theta_v} - 1 \right) \quad \forall \boldsymbol{\theta} \geq \mathbf{0} \end{array} \right\}$$

ECP approximation in capacitated case

- ▶ Difficulty: No independent Poisson \Rightarrow No closed-form MGF
- ▶ Upper bounds via DRO:

$$\mathbb{P} \in \mathcal{F} \Rightarrow \mathbb{E}_{\mathbb{P}} [c(\mathbf{x}, \tilde{\mathbf{z}})] \leq \sup_{\mathbb{P} \in \mathcal{F}} \mathbb{E}_{\mathbb{P}} [c(\mathbf{x}, \tilde{\mathbf{z}})]$$

- ▶ Moments: truncated Poisson $\mathbb{P} \in \mathcal{F}^1$

$$\mathcal{F}^1 \triangleq \left\{ \mathbb{P} \mid \begin{array}{l} \tilde{\mathbf{z}} \sim \mathbb{P} \\ \log \mathbb{E}_{\mathbb{P}} \left[\exp \left(\boldsymbol{\theta}^\top \tilde{\mathbf{z}} \right) \right] \leq \sum_{v \in [V]} \lambda_v \left(e^{\theta_v} - 1 \right) \quad \forall \boldsymbol{\theta} \geq \mathbf{0} \end{array} \right\}$$

- ▶ Support: $\mathbb{P} \in \mathcal{F}^2$

$$\mathcal{F}^2 \triangleq \left\{ \mathbb{P} \mid \begin{array}{l} \tilde{\mathbf{z}} \sim \mathbb{P} \\ \mathbb{E}_{\mathbb{P}} [\tilde{\mathbf{z}}] \leq \boldsymbol{\lambda} \\ \mathbb{P} [\tilde{\mathbf{z}} \in \mathcal{Z}] = 1 \end{array} \right\}$$

ECP approximation in capacitated case

- ▶ Difficulty: No independent Poisson \Rightarrow No closed-form MGF
- ▶ Upper bounds via DRO:

$$\mathbb{P} \in \mathcal{F} \Rightarrow \mathbb{E}_{\mathbb{P}} [c(\mathbf{x}, \tilde{\mathbf{z}})] \leq \sup_{\mathbb{P} \in \mathcal{F}} \mathbb{E}_{\mathbb{P}} [c(\mathbf{x}, \tilde{\mathbf{z}})]$$

- ▶ Moments: truncated Poisson $\mathbb{P} \in \mathcal{F}^1$

$$\mathcal{F}^1 \triangleq \left\{ \mathbb{P} \mid \begin{array}{l} \tilde{\mathbf{z}} \sim \mathbb{P} \\ \log \mathbb{E}_{\mathbb{P}} \left[\exp \left(\boldsymbol{\theta}^\top \tilde{\mathbf{z}} \right) \right] \leq \sum_{v \in [V]} \lambda_v \left(e^{\theta_v} - 1 \right) \quad \forall \boldsymbol{\theta} \geq \mathbf{0} \end{array} \right\}$$


- ▶ Support: $\mathbb{P} \in \mathcal{F}^2$

$$\mathcal{F}^2 \triangleq \left\{ \mathbb{P} \mid \begin{array}{l} \tilde{\mathbf{z}} \sim \mathbb{P} \\ \mathbb{E}_{\mathbb{P}} [\tilde{\mathbf{z}}] \leq \boldsymbol{\lambda} \\ \mathbb{P} [\tilde{\mathbf{z}} \in \mathcal{Z}] = 1 \end{array} \right\}$$

- ▶ ECP-C using $\mathcal{F} = \mathcal{F}_1 \cap \mathcal{F}_2$ via infimal convolution

ECP vs SAA with different sample size


- Expected cost under SAA at different sample size given $C = 30$




- SAA with 38,000 samples cannot be solved in 36 hours and is far from optimal upon termination
- ECP-C is near optimal: the optimality gap is at most $(805.00 - 773.67)/773.67 \times 100\% \approx 4.05\%$, even 0.71% when using the extrapolation results.

ECP vs DRO with mean-covariance information

Compare the upper bounds of $\mathbb{E}_{\mathbb{P}} \left[\max_{v \in [V]} \tilde{z}_v \right]$ where $\tilde{z}_v \sim \text{Poisson}(\lambda_v)$



(a) $\lambda_v = 7$



(b) $\lambda_v = 75$

Summary

- ▶ An ECP approach to a large-scale stochastic program
 - ▶ Bound the order statistic
 - ▶ Exploit stochastic independence
 - ▶ \mathcal{K}_{exp} -representable upper bounds of log-MGF
- ▶ Scalability, tractability, superior performance.

Outline

Introduction

The Key Methodology of The ECP Approach to EV Charging

Robust CARA Optimization

- ▶ Generalize the model of uncertainty and decision criteria
 - ▶ Risk neutral \rightarrow risk averse
 - ▶ No distribution ambiguity \rightarrow ambiguity averse
- ▶ Generalize the objective function

- ▶ Generalize the model of uncertainty and decision criteria
 - ▶ Risk neutral \rightarrow risk averse
 - ▶ No distribution ambiguity \rightarrow ambiguity averse
- ▶ Generalize the objective function
 - ▶ EV-SP is a very special two-stage stochastic linear program with fixed recourse:

$$\begin{aligned} f(\mathbf{x}, \mathbf{z}) = & \min_y y \\ \text{s.t. } & y \geq \sum_{s \in [T]} e_s \sum_{v \in \mathcal{V}_s} x_{v,s} z_v + d \sum_{v \in \mathcal{V}_t} x_{v,t} z_v, \quad \forall t \in [T] \end{aligned}$$

The dimension of recourse variable is one.

- ▶ Extend to two-stage and multi-stage linear optimization problems

Constant absolute risk aversion (CARA)

- ▶ Exponential utility: $u_E(v) \triangleq 1 - e^{-v/\kappa} \rightarrow$ risk aversion
- ▶ CARA: **risk tolerance level** $-u'_E/u''_E \equiv \kappa > 0$

Constant absolute risk aversion (CARA)

- ▶ Exponential utility: $u_E(v) \triangleq 1 - e^{-v/\kappa} \rightarrow$ risk aversion
- ▶ CARA: **risk tolerance level** $-u'_E/u''_E \equiv \kappa > 0$
- ▶ Assessment of κ :
 - ▶ Roughly indifferent to accepting or rejecting a gamble involving a
 - ▶ 50-50 chance of winning κ or losing $\kappa/2$
 - ▶ 75-25 chance of winning κ or losing κ

Constant absolute risk aversion (CARA)

- ▶ Exponential utility: $u_E(v) \triangleq 1 - e^{-v/\kappa} \rightarrow$ risk aversion
- ▶ CARA: **risk tolerance level** $-u'_E/u''_E \equiv \kappa > 0$
- ▶ Assessment of κ :
 - ▶ Roughly indifferent to accepting or rejecting a gamble involving a
 - ▶ 50-50 chance of winning κ or losing $\kappa/2$
 - ▶ 75-25 chance of winning κ or losing κ
- ▶ Tractability in economic analysis
- ▶ Popularity: about five times more commonly adopted than all other types of utility functions combined [CC95]

Robust optimization with CARA preference

A robust CARA optimization model:

$$\max_{\mathbf{x} \in \mathcal{X}} \inf_{\mathbb{P} \in \mathcal{F}} \mathbb{C}_{\mathbb{P}}^{\kappa} [f(\mathbf{x}, \tilde{\mathbf{z}})] \quad (1)$$

where the **CARA certainty equivalent**

$$\mathbb{C}_{\mathbb{P}}^{\kappa} [\tilde{v}] \triangleq u_E^{-1} (\mathbb{E}_{\mathbb{P}} [u_E(\tilde{v})]) = \begin{cases} \text{ess inf}_{\mathbb{P}} [\tilde{v}] & \text{if } \kappa = 0 \\ \mathbb{E}_{\mathbb{P}} [\tilde{v}] & \text{if } \kappa = \infty \\ -\kappa \log \mathbb{E}_{\mathbb{P}} \left[\exp \left(-\frac{\tilde{v}}{\kappa} \right) \right] & \text{if } \kappa \in (0, \infty) \end{cases}$$

$$\mathbb{C}_{\mathbb{P}}^{\kappa}[\tilde{v}] \triangleq u_E^{-1}(\mathbb{E}_{\mathbb{P}}[u_E(\tilde{v})]) = \begin{cases} \text{ess inf}_{\mathbb{P}}[\tilde{v}] & \text{if } \kappa = 0 \\ \mathbb{E}_{\mathbb{P}}[\tilde{v}] & \text{if } \kappa = \infty \\ -\kappa \log \mathbb{E}_{\mathbb{P}}\left[\exp\left(-\frac{\tilde{v}}{\kappa}\right)\right] & \text{if } \kappa \in (0, \infty) \end{cases}$$

Properties of $\mathbb{C}_{\mathbb{P}}^{\kappa}[\tilde{v}]$; Extend to $\mathbb{C}_{\mathcal{F}}^{\kappa}[\tilde{v}] \triangleq \inf_{\mathbb{P} \in \mathcal{F}} \mathbb{C}_{\mathbb{P}}^{\kappa}[\tilde{v}]$

1. $\lim_{\kappa \rightarrow \infty} \mathbb{C}_{\mathbb{P}}^{\kappa}[\tilde{v}] = \mathbb{E}_{\mathbb{P}}[\tilde{v}]$, $\lim_{\kappa \downarrow 0} \mathbb{C}_{\mathbb{P}}^{\kappa}[\tilde{v}] = \text{ess inf}_{\mathbb{P}}[\tilde{v}]$.
2. $\mathbb{C}_{\mathbb{P}}^{\kappa}[\tilde{v}]$ is increasing in $\kappa > 0$
3. $\mathbb{C}_{\mathbb{P}}^{\kappa}[\tilde{v}]$ is jointly concave in \tilde{v} and $\kappa > 0$.
4. $\mathbb{C}_{\mathbb{P}}^{\kappa}[\tilde{v} + \tilde{\nu}] = \mathbb{C}_{\mathbb{P}}^{\kappa}[\tilde{v}] + \mathbb{C}_{\mathbb{P}}^{\kappa}[\tilde{\nu}]$ if $\tilde{v}, \tilde{\nu}$ are **independent**.
5. **Super-additivity**: $\mathbb{C}_{\mathbb{P}}^{\kappa_1 + \kappa_2}[\tilde{v} + \tilde{\nu}] \geq \mathbb{C}_{\mathbb{P}}^{\kappa_1}[\tilde{v}] + \mathbb{C}_{\mathbb{P}}^{\kappa_2}[\tilde{\nu}]$ for any $\kappa_1, \kappa_2 \in \mathbb{R}_+$.

Independent factors with ambiguous marginals

- ▶ $\tilde{z} = (\tilde{z}_1, \tilde{z}_2, \dots, \tilde{z}_{I_z})$ has independent components
- ▶ $\tilde{z}_j \sim \mathbb{P}_j \in \mathcal{F}_j \subseteq \mathcal{P}_0([\underline{z}_j, \bar{z}_j])$ where $\underline{z}_j < \bar{z}_j$
- ▶ So $\tilde{z} \sim \mathbb{P} \in \mathcal{F} \triangleq \times_{j \in [I_z]} \mathcal{F}_j \subseteq \mathcal{P}_0(\mathcal{Z})$ where $\mathcal{Z} \triangleq [\underline{z}, \bar{z}]$

Independent factors with ambiguous marginals

- ▶ $\tilde{z} = (\tilde{z}_1, \tilde{z}_2, \dots, \tilde{z}_{I_z})$ has independent components
- ▶ $\tilde{z}_j \sim \mathbb{P}_j \in \mathcal{F}_j \subseteq \mathcal{P}_0([z_j, \bar{z}_j])$ where $z_j < \bar{z}_j$
- ▶ So $\tilde{z} \sim \mathbb{P} \in \mathcal{F} \triangleq \times_{j \in [I_z]} \mathcal{F}_j \subseteq \mathcal{P}_0(\mathcal{Z})$ where $\mathcal{Z} \triangleq [z, \bar{z}]$

For an affine function

$$f(\mathbf{x}, \mathbf{z}) = a^0(\mathbf{x}) + \sum_{j \in [I_z]} a^j(\mathbf{x}) z_j,$$

we have

$$\mathbb{C}_{\mathcal{F}}^{\kappa}[f(\mathbf{x}, \tilde{\mathbf{z}})] = a^0(\mathbf{x}) + \sum_{j \in [I_z]} \phi_j(\kappa, a^j(\mathbf{x}))$$

where the function $\phi_j : [0, \infty] \times \mathbb{R} \rightarrow \mathbb{R}$,

$$\phi_j(\kappa, \lambda) \triangleq \inf_{\mathbb{P}_j \in \mathcal{F}_j} \mathbb{C}_{\mathbb{P}}^{\kappa}[\lambda \tilde{z}_j].$$

Example 1

Let $\tilde{z}_j \sim \text{Unif}([0, 1])$, $a^j(\mathbf{x}) \equiv a_j < 0$ and $a^0(\mathbf{x}) \equiv a_0 > 0$, then

$$\begin{aligned}\mathbb{C}_{\mathbb{P}}^{\kappa}[f(\mathbf{x}, \tilde{\mathbf{z}})] &= a_0 - \sum_{j \in [I_z]} \kappa \log \int_0^1 \exp\left(-\frac{a_j z_j}{\kappa}\right) dz_j \\ &= a_0 - \sum_{j \in [I_z]} \kappa \log \left(\frac{\kappa - \kappa e^{-a_j/\kappa}}{a_j} \right).\end{aligned}$$

Example 1

Let $\tilde{z}_j \sim \text{Unif}([0, 1])$, $a^j(\mathbf{x}) \equiv a_j < 0$ and $a^0(\mathbf{x}) \equiv a_0 > 0$, then

$$\begin{aligned}\mathbb{C}_{\mathbb{P}}^{\kappa}[f(\mathbf{x}, \tilde{\mathbf{z}})] &= a_0 - \sum_{j \in [I_z]} \kappa \log \int_0^1 \exp\left(-\frac{a_j z_j}{\kappa}\right) dz_j \\ &= a_0 - \sum_{j \in [I_z]} \kappa \log \left(\frac{\kappa - \kappa e^{-a_j/\kappa}}{a_j} \right).\end{aligned}$$

In contrast, evaluating an expected concave piecewise linear utility such as

$$\mathbb{E}_{\mathbb{P}} \left[\min \left\{ a_0 + \sum_{j \in [I_z]} a_j \tilde{z}_j, 0 \right\} \right]$$

is known to be #P-hard.

Example 2

Consider a simple mean-support ambiguity set:

$$\mathcal{G} = \left\{ \mathbb{P} \in \mathcal{P}_0(\mathbb{R}^{I_z}) \left| \begin{array}{l} \tilde{z} \sim \mathbb{P} \\ \mathbb{E}_{\mathbb{P}}[\tilde{z}] = \boldsymbol{\mu} \\ \mathbb{P}[\tilde{z} \in [\underline{z}, \bar{z}]] = 1 \end{array} \right. \right\}.$$

Note that evaluating $\mathbb{C}_{\mathcal{G}}^{\kappa}[\mathbf{a}^{\top} \tilde{z}]$ requires

$$\begin{aligned} \sup_{\mathbb{P} \in \mathcal{G}} \mathbb{E}_{\mathbb{P}} \left[\exp \left(-\frac{\mathbf{a}^{\top} \tilde{z}}{\kappa} \right) \right] &= \inf_{\alpha, \boldsymbol{\beta}} \quad \alpha + \boldsymbol{\beta}^{\top} \boldsymbol{\mu} \\ \text{s.t.} \quad \alpha &\geq \sup_{\mathbf{z} \in [\underline{\mathbf{z}}, \bar{\mathbf{z}}]} \exp \left(\frac{-\mathbf{a}^{\top} \mathbf{z}}{\kappa} \right) - \boldsymbol{\beta}^{\top} \mathbf{z}, \end{aligned}$$

which involves a convex maximization problem.

Theorem 1

Let $g(\mathbf{x}, \kappa) = -\kappa \log \sum_{i \in [I]} p_i e^{-x_i/\kappa}$ with $\kappa > 0$ and $p_i > 0$ for all $i \in [I]$, then the closure of $\{(\mathbf{x}, \kappa, y) : y \leq g(\mathbf{x}, \kappa), \kappa > 0\}$ can be represented by

$$\left\{ (\mathbf{x}, \kappa, y) \mid \exists \mathbf{q} \in \mathbb{R}^I : \sum_{i \in [I]} p_i q_i \leq \kappa, (q_i, \kappa, y - x_i) \in \mathcal{K}_{\text{exp}} \quad \forall i \in [I] \right\}.$$

Theorem 1

Let $g(\mathbf{x}, \kappa) = -\kappa \log \sum_{i \in [I]} p_i e^{-x_i/\kappa}$ with $\kappa > 0$ and $p_i > 0$ for all $i \in [I]$, then the closure of $\{(\mathbf{x}, \kappa, y) : y \leq g(\mathbf{x}, \kappa), \kappa > 0\}$ can be represented by

$$\left\{ (\mathbf{x}, \kappa, y) \mid \exists \mathbf{q} \in \mathbb{R}^I : \sum_{i \in [I]} p_i q_i \leq \kappa, (q_i, \kappa, y - x_i) \in \mathcal{K}_{\text{exp}} \quad \forall i \in [I] \right\}.$$

Recall that for an affine function,

$$\mathbb{C}_{\mathcal{F}}^{\kappa} [f(\mathbf{x}, \tilde{\mathbf{z}})] = a^0(\mathbf{x}) + \sum_{j \in [I_z]} \phi_j(\kappa, a^j(\mathbf{x})).$$

ECP formulations

- $\phi(\kappa, \lambda) \triangleq \mathbb{C}_{\mathcal{G}}^{\kappa}[\lambda \tilde{z}]$ is \mathcal{K}_{exp} -representable for many ambiguity sets. [NS07]

Table 1 Equivalent representations of $\phi(\kappa, \lambda)$

Ambiguity set \mathcal{F}	$\phi(\kappa, \lambda)$
$\{\mathbb{P}[\tilde{z} \in [-1, 1]] = 1\}$	$- \lambda $
$\left\{ \begin{array}{l} \mathbb{P} \text{ is symmetric} \\ \mathbb{P}[\tilde{z} \in [-1, 1]] = 1 \end{array} \right\}$	$-\kappa \log \left(\frac{e^{\lambda/\kappa} + e^{-\lambda/\kappa}}{2} \right)$
$\left\{ \begin{array}{l} \mathbb{P} \text{ is unimodal w.r.t. } 0 \\ \mathbb{P}[\tilde{z} \in [-1, 1]] = 1 \end{array} \right\}$	$-\kappa \log \int_0^1 e^{s \lambda /\kappa} ds$
$\left\{ \begin{array}{l} \mathbb{P} \text{ is symmetric,} \\ \text{unimodal w.r.t. } 0 \\ \mathbb{P}[\tilde{z} \in [-1, 1]] = 1 \end{array} \right\}$	$\lambda - \kappa \log \int_0^1 e^{-2\lambda s/\kappa} ds$
$\left\{ \begin{array}{l} \mathbb{E}_{\mathbb{P}}[\tilde{z}] \in [\underline{\mu}, \overline{\mu}] \\ \mathbb{P}[\tilde{z} \in [-1, 1]] = 1 \end{array} \right\}$	$\min \left\{ \begin{array}{l} -\kappa \log \left(\frac{(1+\underline{\mu})e^{-\lambda/\kappa} + (1-\underline{\mu})e^{\lambda/\kappa}}{2} \right), \\ -\kappa \log \left(\frac{(1+\overline{\mu})e^{-\lambda/\kappa} + (1-\overline{\mu})e^{\lambda/\kappa}}{2} \right) \end{array} \right\}$
$\left\{ \begin{array}{l} \mathbb{E}_{\mathbb{P}}[\tilde{z}] = \mu \\ \mathbb{E}_{\mathbb{P}}[\tilde{z} - \mu] \leq \delta \\ \mathbb{P}[\tilde{z} \in [-1, 1]] = 1 \end{array} \right\}$	$-\kappa \log \left(\frac{\delta}{2(\mu+1)} e^{\lambda/\kappa} + \frac{\delta}{2(1-\mu)} e^{-\lambda/\kappa} + \left(1 - \frac{\delta}{2(\mu+1)} - \frac{\delta}{2(1-\mu)} \right) e^{-\mu\lambda/\kappa} \right)$
$\left\{ \begin{array}{l} \mathbb{E}_{\mathbb{P}}[\tilde{z}] = \mu \\ \mathbb{E}_{\mathbb{P}}[\tilde{z} ^2] \leq \sigma^2 \\ \mathbb{P}[\tilde{z} \in [-1, 1]] = 1 \end{array} \right\}$	$\min \left\{ \begin{array}{l} -\kappa \log \left(\frac{(1-\mu)^2 \exp\left(\frac{-(\mu-\sigma^2)\lambda}{(1-\mu)\kappa}\right) + (\sigma^2 - \mu^2) \exp(-\lambda/\kappa)}{1 - 2\mu + \sigma^2} \right), \\ -\kappa \log \left(\frac{(1+\mu)^2 \exp\left(\frac{-(\mu+\sigma^2)\lambda}{(1-\mu)\kappa}\right) + (\sigma^2 - \mu^2) \exp(\lambda/\kappa)}{1 + 2\mu + \sigma^2} \right) \end{array} \right\}$
$\left\{ \begin{array}{l} \mathbb{P} \text{ is symmetric,} \\ \mathbb{E}_{\mathbb{P}}[\tilde{z} ^2] \leq \sigma^2 \\ \mathbb{P}[\tilde{z} \in [-1, 1]] = 1 \end{array} \right\}$	$-\kappa \log \left(\frac{\sigma^2(e^{\lambda/\kappa} + e^{-\lambda/\kappa})}{2} + 1 - \sigma^2 \right)$

- ▶ $\phi(\kappa, \lambda) \triangleq \mathbb{C}_{\mathcal{G}}^{\kappa}[\lambda \tilde{z}]$ is \mathcal{K}_{exp} -representable for many ambiguity sets. [NS07]
- ▶ $\mathbb{C}_{\mathcal{F}}^{\kappa}[f(\mathbf{x}, \tilde{z})] = a^0(\mathbf{x}) + \sum_{j \in [I_z]} \phi_j(\kappa, a^j(\mathbf{x}))$ is \mathcal{K}_{exp} -representable if $a^0(\mathbf{x})$, $a^j(\mathbf{x})$ are affine.

Concave piecewise affine functions

- ▶ However, in practice the payoff functions $f(x, z)$ are usually nonlinear in z .

Concave piecewise affine functions

- ▶ However, in practice the payoff functions $f(\mathbf{x}, \mathbf{z})$ are usually nonlinear in \mathbf{z} .
- ▶ Consider a concave piecewise affine payoff function

$$f(\mathbf{x}, \mathbf{z}) = \min_{i \in \mathcal{I}} \left\{ a_i^0(\mathbf{x}) + \sum_{j \in [I_z]} a_i^j(\mathbf{x}) z_j \right\}.$$

- ▶ Recall evaluating $\mathbb{C}_{\mathbb{P}}^{\kappa} [f(\mathbf{x}, \tilde{\mathbf{z}})]$ under known distribution can be #P-hard.

Concave piecewise affine functions

- ▶ However, in practice the payoff functions $f(\mathbf{x}, \mathbf{z})$ are usually nonlinear in \mathbf{z} .
- ▶ Consider a concave piecewise affine payoff function

$$f(\mathbf{x}, \mathbf{z}) = \min_{i \in \mathcal{I}} \left\{ a_i^0(\mathbf{x}) + \sum_{j \in [I_z]} a_i^j(\mathbf{x}) z_j \right\}.$$

- ▶ Recall evaluating $\mathbb{C}_{\mathbb{P}}^{\kappa} [f(\mathbf{x}, \tilde{\mathbf{z}})]$ under known distribution can be #P-hard.
- ▶ Tractable approximations are needed.

A tractable lower bound

Let $\alpha_i = a_i^0(\mathbf{x})$, $\beta_i^j = a_i^j(\mathbf{x})$,

$$\begin{aligned} & \mathbb{C}_{\mathcal{F}}^{\kappa} \left[\min_{i \in \mathcal{I}} \left\{ \alpha_i + \beta_i^{\top} \tilde{\mathbf{z}} \right\} \right] \\ &= \mathbb{C}_{\mathcal{F}}^{\kappa} \left[\gamma^{\top} \tilde{\mathbf{z}} + \min_{i \in \mathcal{I}} \left\{ \alpha_i + (\beta_i - \gamma)^{\top} \tilde{\mathbf{z}} \right\} \right] \end{aligned}$$

A tractable lower bound

$$\begin{aligned} & \mathbb{C}_{\mathcal{F}}^{\kappa} \left[\min_{i \in \mathcal{I}} \left\{ \alpha_i + \beta_i^{\top} \tilde{\mathbf{z}} \right\} \right] \\ &= \mathbb{C}_{\mathcal{F}}^{\kappa} \left[\boldsymbol{\gamma}^{\top} \tilde{\mathbf{z}} + \min_{i \in \mathcal{I}} \left\{ \alpha_i + (\beta_i - \boldsymbol{\gamma})^{\top} \tilde{\mathbf{z}} \right\} \right] \\ &\geq \sup_{\kappa_0 + \kappa_1 = \kappa, \kappa \geq \mathbf{0}} \mathbb{C}_{\mathcal{F}}^{\kappa_0} \left[\boldsymbol{\gamma}^{\top} \tilde{\mathbf{z}} \right] + \mathbb{C}_{\mathcal{F}}^{\kappa_1} \left[\min_{i \in \mathcal{I}} \left\{ \alpha_i + (\beta_i - \boldsymbol{\gamma})^{\top} \tilde{\mathbf{z}} \right\} \right] \\ & \quad (\text{Super-additivity of } \mathbb{C}_{\mathcal{F}}^{\kappa} [\tilde{v}] \text{ w.r.t. } (\tilde{v}, \kappa) \text{ with } \kappa \geq 0) \end{aligned}$$

A tractable lower bound

$$\begin{aligned} & \mathbb{C}_{\mathcal{F}}^{\kappa} \left[\min_{i \in \mathcal{I}} \left\{ \alpha_i + \beta_i^{\top} \tilde{\mathbf{z}} \right\} \right] \\ &= \mathbb{C}_{\mathcal{F}}^{\kappa} \left[\gamma^{\top} \tilde{\mathbf{z}} + \min_{i \in \mathcal{I}} \left\{ \alpha_i + (\beta_i - \gamma)^{\top} \tilde{\mathbf{z}} \right\} \right] \\ &\geq \sup_{\kappa_0 + \kappa_1 = \kappa, \kappa \geq \mathbf{0}} \mathbb{C}_{\mathcal{F}}^{\kappa_0} \left[\gamma^{\top} \tilde{\mathbf{z}} \right] + \mathbb{C}_{\mathcal{F}}^{\kappa_1} \left[\min_{i \in \mathcal{I}} \left\{ \alpha_i + (\beta_i - \gamma)^{\top} \tilde{\mathbf{z}} \right\} \right] \\ &= \sup_{\kappa_0 + \kappa_1 = \kappa, \kappa \geq \mathbf{0}} \mathbb{C}_{\mathcal{F}}^{\kappa_0} \left[\gamma^{\top} \tilde{\mathbf{z}} \right] - \kappa_1 \log \sup_{\mathbb{P} \in \mathcal{F}} \mathbb{E}_{\mathbb{P}} \left[\exp \left(\frac{\max_{i \in \mathcal{I}} \left\{ -\alpha_i - (\beta_i - \gamma)^{\top} \tilde{\mathbf{z}} \right\}}{\kappa_1} \right) \right] \end{aligned}$$

A tractable lower bound

$$\begin{aligned}& \mathbb{C}_{\mathcal{F}}^{\kappa} \left[\min_{i \in \mathcal{I}} \left\{ \alpha_i + \beta_i^{\top} \tilde{\mathbf{z}} \right\} \right] \\&= \mathbb{C}_{\mathcal{F}}^{\kappa} \left[\gamma^{\top} \tilde{\mathbf{z}} + \min_{i \in \mathcal{I}} \{ \alpha_i + (\beta_i - \gamma)^{\top} \tilde{\mathbf{z}} \} \right] \\&\geq \sup_{\kappa_0 + \kappa_1 = \kappa, \kappa \geq \mathbf{0}} \mathbb{C}_{\mathcal{F}}^{\kappa_0} \left[\gamma^{\top} \tilde{\mathbf{z}} \right] + \mathbb{C}_{\mathcal{F}}^{\kappa_1} \left[\min_{i \in \mathcal{I}} \{ \alpha_i + (\beta_i - \gamma)^{\top} \tilde{\mathbf{z}} \} \right] \\&= \sup_{\kappa_0 + \kappa_1 = \kappa, \kappa \geq \mathbf{0}} \mathbb{C}_{\mathcal{F}}^{\kappa_0} \left[\gamma^{\top} \tilde{\mathbf{z}} \right] - \kappa_1 \log \sup_{\mathbb{P} \in \mathcal{F}} \mathbb{E}_{\mathbb{P}} \left[\exp \left(\frac{\max_{i \in \mathcal{I}} \{ -\alpha_i - (\beta_i - \gamma)^{\top} \tilde{\mathbf{z}} \}}{\kappa_1} \right) \right] \\&\geq \sup_{\kappa_0 + \kappa_1 = \kappa, \kappa \geq \mathbf{0}} \mathbb{C}_{\mathcal{F}}^{\kappa_0} \left[\gamma^{\top} \tilde{\mathbf{z}} \right] - \kappa_1 \log \sum_{i \in \mathcal{I}} \sup_{\mathbb{P} \in \mathcal{F}} \mathbb{E}_{\mathbb{P}} \left[\exp \left(\frac{-\alpha_i + (\gamma - \beta_i)^{\top} \tilde{\mathbf{z}}}{\kappa_1} \right) \right] \\&\quad \text{(Bound max by sum)}\end{aligned}$$

A tractable lower bound

$$\begin{aligned}
 & \mathbb{C}_{\mathcal{F}}^{\kappa} \left[\min_{i \in \mathcal{I}} \left\{ \alpha_i + \beta_i^{\top} \tilde{\mathbf{z}} \right\} \right] \\
 &= \mathbb{C}_{\mathcal{F}}^{\kappa} \left[\gamma^{\top} \tilde{\mathbf{z}} + \min_{i \in \mathcal{I}} \left\{ \alpha_i + (\beta_i - \gamma)^{\top} \tilde{\mathbf{z}} \right\} \right] \\
 &\geq \sup_{\kappa_0 + \kappa_1 = \kappa, \kappa \geq \mathbf{0}} \mathbb{C}_{\mathcal{F}}^{\kappa_0} \left[\gamma^{\top} \tilde{\mathbf{z}} \right] + \mathbb{C}_{\mathcal{F}}^{\kappa_1} \left[\min_{i \in \mathcal{I}} \left\{ \alpha_i + (\beta_i - \gamma)^{\top} \tilde{\mathbf{z}} \right\} \right] \\
 &= \sup_{\kappa_0 + \kappa_1 = \kappa, \kappa \geq \mathbf{0}} \mathbb{C}_{\mathcal{F}}^{\kappa_0} \left[\gamma^{\top} \tilde{\mathbf{z}} \right] - \kappa_1 \log \sup_{\mathbb{P} \in \mathcal{F}} \mathbb{E}_{\mathbb{P}} \left[\exp \left(\frac{\max_{i \in \mathcal{I}} \left\{ -\alpha_i - (\beta_i - \gamma)^{\top} \tilde{\mathbf{z}} \right\}}{\kappa_1} \right) \right] \\
 &\geq \sup_{\kappa_0 + \kappa_1 = \kappa, \kappa \geq \mathbf{0}} \mathbb{C}_{\mathcal{F}}^{\kappa_0} \left[\gamma^{\top} \tilde{\mathbf{z}} \right] - \kappa_1 \log \sum_{i \in \mathcal{I}} \sup_{\mathbb{P} \in \mathcal{F}} \mathbb{E}_{\mathbb{P}} \left[\exp \left(\frac{-\alpha_i + (\gamma - \beta_i)^{\top} \tilde{\mathbf{z}}}{\kappa_1} \right) \right] \\
 &= \sup_{\kappa_0 + \kappa_1 = \kappa, \kappa \geq \mathbf{0}} \mathbb{C}_{\mathcal{F}}^{\kappa_0} \left[\gamma^{\top} \tilde{\mathbf{z}} \right] - \kappa_1 \log \sum_{i \in \mathcal{I}} \exp \left(-\frac{\mathbb{C}_{\mathcal{F}}^{\kappa_1} \left[\alpha_i + (\beta_i - \gamma)^{\top} \tilde{\mathbf{z}} \right]}{\kappa_1} \right)
 \end{aligned}$$

All in all, $\mathbb{C}_{\mathcal{F}}^{\kappa}[f(\mathbf{x}, \tilde{\mathbf{z}})]$ has an ECP lower bound

$$\begin{aligned}\Lambda(\kappa, \mathbf{x}) &\triangleq \max r_0 + \rho \\ \text{s.t. } &\kappa_0 + \kappa_1 = \kappa \\ &\sum_{i \in \mathcal{I}} q_i \leq \kappa_1 \\ &(q_i, \kappa_1, \rho - r_i) \in \mathcal{K}_{\text{exp}} \quad \forall i \in \mathcal{I} \\ &\sum_{j \in [I_z]} \phi_j(\kappa_0, \gamma^j) \geq r_0 \\ &\alpha_i + \sum_{j \in [I_z]} \phi_j(\kappa_1, \beta_i^j - \gamma^j) \geq r_i \quad \forall i \in \mathcal{I} \\ &\boldsymbol{\gamma} \in \mathbb{R}^{I_z}, \boldsymbol{r} \in \mathbb{R}^{1+N}, \boldsymbol{\kappa} \in \mathbb{R}_+^2, \rho \in \mathbb{R}, \mathbf{q} \in \mathbb{R}^N\end{aligned}$$

if $a_i^0(\mathbf{x})$ and $a_i^j(\mathbf{x})$ are affine.

Theorem 2

For any $\mathbf{x} \in \mathcal{X}$, $\Lambda(\kappa, \mathbf{x})$ is nondecreasing in $\kappa \in [0, \infty]$ and satisfies $\Lambda(\kappa, \mathbf{x}) \geq \inf_{\mathbf{z} \in \mathcal{Z}} f(\mathbf{x}, \mathbf{z})$. Moreover, $\mathbb{C}_{\mathcal{F}}^{\kappa}[f(\mathbf{x}, \tilde{\mathbf{z}})] = \Lambda(\kappa, \mathbf{x})$ if there exists some $i^* \in \mathcal{I}$ such that

$$a_{i^*}^0(\mathbf{x}) + \sum_{j \in [I_z]} a_{i^*}^j(\mathbf{x}) z_j \leq a_i^0(\mathbf{x}) + \sum_{j \in [I_z]} a_i^j(\mathbf{x}) z_j \quad \forall \mathbf{z} \in \mathcal{Z}, i \in \mathcal{I}.$$


Cases of good approximation:

- ▶ Low risk tolerance κ
- ▶ Low coefficient of variations among the payoff components

Comparison with Monte-Carlo approximation

Approximate $\mathbb{C}_{\mathbb{P}}^{\kappa} \left[\min_{i \in [N]} \{ \mathbf{a}_i^{\top} \tilde{\mathbf{z}} \} \right]$ where $\tilde{\mathbf{z}}_j \sim \text{Unif}([0, 1])$


Figure: Comparison of our bound and Monte-Carlo approximation (10^6 samples) for $\kappa \geq 0.25$



Comparison with Monte-Carlo approximation

Approximate $\mathbb{C}_{\mathbb{P}}^{\kappa} \left[\min_{i \in [N]} \{ \mathbf{a}_i^{\top} \tilde{\mathbf{z}} \} \right]$ where $\tilde{\mathbf{z}}_j \sim \text{Unif}([0, 1])$

Figure: Ratio of Monte-Carlo approximation to our bound for $\kappa \leq 0.2$ at $N = 1$ where our bound is exact while **Monte-Carlo is upward biased**



Two-stage optimization

Consider the two-stage problem with fixed recourse

$$\begin{aligned} f(\mathbf{x}, \mathbf{z}) = & \max_{\mathbf{y}} \quad \mathbf{c}^\top \mathbf{y} \\ \text{s.t.} \quad & \mathbf{b}_i^\top \mathbf{y} \leq a_i^0(\mathbf{x}) + \mathbf{a}_i^\top(\mathbf{x})\mathbf{z} \quad \forall i \in \mathcal{I}, \\ & \mathbf{y} \in \mathbb{R}^{I_y} \end{aligned}$$

where \mathbf{y} is the recourse decision adaptive to uncertainty realization.

Two-stage optimization

Consider the two-stage problem with fixed recourse

$$\begin{aligned} f(\mathbf{x}, \mathbf{z}) = & \max_{\mathbf{y}} \quad \mathbf{c}^\top \mathbf{y} \\ \text{s.t.} \quad & \mathbf{b}_i^\top \mathbf{y} \leq a_i^0(\mathbf{x}) + \mathbf{a}_i^\top(\mathbf{x})\mathbf{z} \quad \forall i \in \mathcal{I}, \\ & \mathbf{y} \in \mathbb{R}^{I_y} \end{aligned}$$

where \mathbf{y} is the recourse decision adaptive to uncertainty realization.

- Assume no here-and-now variable in objective w.l.o.g.

Two-stage optimization

Consider the two-stage problem with fixed recourse

$$\begin{aligned} f(\mathbf{x}, \mathbf{z}) = & \max_{\mathbf{y}} \quad \mathbf{c}^\top \mathbf{y} \\ \text{s.t.} \quad & \mathbf{b}_i^\top \mathbf{y} \leq a_i^0(\mathbf{x}) + \mathbf{a}_i^\top(\mathbf{x}) \mathbf{z} \quad \forall i \in \mathcal{I}, \\ & \mathbf{y} \in \mathbb{R}^{I_y} \end{aligned}$$

where \mathbf{y} is the recourse decision adaptive to uncertainty realization.

- ▶ Assume no here-and-now variable in objective w.l.o.g.
- ▶ Assume $\mathbf{b}_i \neq \mathbf{0}$ w.l.o.g.

Two-stage optimization

Consider the two-stage problem with fixed recourse

$$\begin{aligned} f(\mathbf{x}, \mathbf{z}) = & \max_{\mathbf{y}} \quad \mathbf{c}^\top \mathbf{y} \\ \text{s.t.} \quad & \mathbf{b}_i^\top \mathbf{y} \leq a_i^0(\mathbf{x}) + \mathbf{a}_i^\top(\mathbf{x}) \mathbf{z} \quad \forall i \in \mathcal{I}, \\ & \mathbf{y} \in \mathbb{R}^{I_y} \end{aligned}$$

where \mathbf{y} is the recourse decision adaptive to uncertainty realization.

- ▶ Assume no here-and-now variable in objective w.l.o.g.
- ▶ Assume $\mathbf{b}_i \neq \mathbf{0}$ w.l.o.g.
- ▶ Assume inequality constraints only w.l.o.g.

Two-stage optimization

Consider the two-stage problem with fixed recourse

$$\begin{aligned} f(\mathbf{x}, \mathbf{z}) = & \max_{\mathbf{y}} \quad \mathbf{c}^\top \mathbf{y} \\ \text{s.t.} \quad & \mathbf{b}_i^\top \mathbf{y} \leq a_i^0(\mathbf{x}) + \mathbf{a}_i^\top(\mathbf{x})\mathbf{z} \quad \forall i \in \mathcal{I}, \\ & \mathbf{y} \in \mathbb{R}^{I_y} \end{aligned}$$

where \mathbf{y} is the recourse decision adaptive to uncertainty realization.

- ▶ Assume no here-and-now variable in objective w.l.o.g.
- ▶ Assume $\mathbf{b}_i \neq \mathbf{0}$ w.l.o.g.
- ▶ Assume inequality constraints only w.l.o.g.

Note: Generalize the concave piecewise affine payoff function where $I_y = 1$, $\mathbf{c} = 1$, and $\mathbf{b}_i = 1$, $i \in \mathcal{I}$.

Multi-deflected linear decision rule

► A multi-deflected linear decision rule (MLDR)

$$\hat{\mathbf{y}}(\mathbf{z}) \triangleq \bar{\mathbf{y}}(\mathbf{z}) + \sum_{\ell \in [m]} \mathbf{y}_*^\ell \left(\max_{i \in \mathcal{I}_\ell^o} \left\{ \frac{h_i(\mathbf{x}, \bar{\mathbf{y}}(\mathbf{z}), \mathbf{z})}{\|\mathbf{b}_i\|} \right\} \right)^+.$$

where

$$\begin{aligned} \bar{\mathbf{y}}(\mathbf{z}) &\triangleq \mathbf{y}^0 + \mathbf{Y}\mathbf{z} \\ h_i(\mathbf{x}, \mathbf{y}, \mathbf{z}) &\triangleq \mathbf{b}_i^\top \mathbf{y} - a_i^0(\mathbf{x}) - \mathbf{a}_i^\top(\mathbf{x})\mathbf{z} \quad \forall i \in \mathcal{I}. \end{aligned}$$

Multi-deflected linear decision rule

► A multi-deflected linear decision rule (MLDR)

$$\hat{\mathbf{y}}(\mathbf{z}) \triangleq \bar{\mathbf{y}}(\mathbf{z}) + \sum_{\ell \in [m]} \mathbf{y}_*^\ell \left(\max_{i \in \mathcal{I}_\ell^o} \left\{ \frac{h_i(\mathbf{x}, \bar{\mathbf{y}}(\mathbf{z}), \mathbf{z})}{\|\mathbf{b}_i\|} \right\} \right)^+.$$

where \mathbf{y}_*^ℓ and \mathcal{I}_ℓ^o are chosen as follows:

- Solve for each $i \in \mathcal{I}$,

$$\begin{aligned} \max_{\mathbf{y} \in \mathbb{R}^{I_y}} \quad & \mathbf{c}^\top \mathbf{y} \\ & \mathbf{b}_k^\top \mathbf{y} \leq 0 \quad \forall k \in \mathcal{I} \setminus \{i\} \\ & \mathbf{b}_i^\top \mathbf{y} = -\|\mathbf{b}_i\| \end{aligned}$$

and denote \mathbf{y}_*^i as its optimal solution if feasible.

- Partition feasible index set

$$\mathcal{I}^o = \bigcup_{\ell \in [m]} \mathcal{I}_\ell^o$$

such that $\mathbf{y}_*^{i_1} = \mathbf{y}_*^{i_2}$ if and only if i_1 and i_2 are in the same \mathcal{I}_ℓ^o .

- ▶ A **multi-deflected linear decision rule (MLDR)**

$$\hat{\mathbf{y}}(\mathbf{z}) \triangleq \bar{\mathbf{y}}(\mathbf{z}) + \sum_{\ell \in [m]} \mathbf{y}_*^\ell \left(\max_{i \in \mathcal{I}_\ell^o} \left\{ \frac{h_i(\mathbf{x}, \bar{\mathbf{y}}(\mathbf{z}), \mathbf{z})}{\|\mathbf{b}_i\|} \right\} \right)^+.$$

- ▶ Better than previous deflected LDR in [[CSSZ08](#)]

- ▶ A **multi-deflected linear decision rule (MLDR)**

$$\hat{\mathbf{y}}(\mathbf{z}) \triangleq \bar{\mathbf{y}}(\mathbf{z}) + \sum_{\ell \in [m]} \mathbf{y}_*^\ell \left(\max_{i \in \mathcal{I}_\ell^o} \left\{ \frac{h_i(\mathbf{x}, \bar{\mathbf{y}}(\mathbf{z}), \mathbf{z})}{\|\mathbf{b}_i\|} \right\} \right)^+.$$

- ▶ Better than previous deflected LDR in [CSSZ08]
- ▶ Can replicate optimal decision rule under complete recourse with $I_y = 1$

- ▶ A **multi-deflected linear decision rule (MLDR)**

$$\hat{\mathbf{y}}(\mathbf{z}) \triangleq \bar{\mathbf{y}}(\mathbf{z}) + \sum_{\ell \in [m]} \mathbf{y}_*^\ell \left(\max_{i \in \mathcal{I}_\ell^o} \left\{ \frac{h_i(\mathbf{x}, \bar{\mathbf{y}}(\mathbf{z}), \mathbf{z})}{\|\mathbf{b}_i\|} \right\} \right)^+.$$

- ▶ Better than previous deflected LDR in [CSSZ08]
- ▶ Can replicate optimal decision rule under complete recourse with $I_y = 1$
- ▶ Obtain ECP lower bound based on approximation for concave piecewise affine functions.

A multi-period consumption model

Given $\xi_t \triangleq (z_1, \dots, z_t)$,

$$\begin{aligned} \max_{\substack{\mathbf{x} \in \mathcal{X}, \\ \mathbf{y}_1, \dots, \mathbf{y}_T}} \quad & \mathbb{C}_{\mathcal{F}}^{\kappa, \theta} \left[\mathbf{c}_1^\top \mathbf{y}_1(\tilde{\xi}_1), \dots, \mathbf{c}_T^\top \mathbf{y}_T(\tilde{\xi}_T) \right] \\ \text{s.t.} \quad & \sum_{\tau \in [t]} \mathbf{b}_{t,i,\tau}^\top \mathbf{y}_\tau(\xi_\tau) \leq a_{t,i}^0(\mathbf{x}) + \mathbf{a}_{t,i}^\top(\mathbf{x}) \xi_t \quad \forall t \in [T], i \in \mathcal{I}_t, \mathbf{z} \in \mathcal{Z} \\ & \mathbf{y}_t \in \mathcal{R}^{I_{\xi_t}, I_{y_t}} \quad \forall t \in [T]. \end{aligned}$$

where the **multi-period ambiguity-averse CARA certainty equivalent**

$$\mathbb{C}_{\mathcal{F}}^{\kappa, \theta} [\tilde{\mathbf{v}}] \triangleq \begin{cases} \min_{t \in [T]: \theta_t > 0} \left\{ \inf_{\mathbb{P} \in \mathcal{F}} \text{ess inf}_{\mathbb{P}} [\tilde{v}_t] \right\} & \text{if } \kappa = 0 \\ \sum_{t \in [T]} \theta_t \inf_{\mathbb{P} \in \mathcal{F}} \mathbb{E}_{\mathbb{P}} [\tilde{v}_t] & \text{if } \kappa = \infty \\ -\kappa \log \left(\sum_{t \in [T]} \theta_t \sup_{\mathbb{P} \in \mathcal{F}} \mathbb{E}_{\mathbb{P}} \left[\exp \left(-\frac{\tilde{v}_t}{\kappa} \right) \right] \right) & \text{if } \kappa \in (0, \infty). \end{cases}$$

A multi-period consumption model

Given $\xi_t \triangleq (z_1, \dots, z_t)$,

$$\begin{aligned} \max_{\substack{\mathbf{x} \in \mathcal{X}, \\ \mathbf{y}_1, \dots, \mathbf{y}_T}} \quad & \mathbb{C}_{\mathcal{F}}^{\kappa, \theta} \left[\mathbf{c}_1^\top \mathbf{y}_1(\tilde{\xi}_1), \dots, \mathbf{c}_T^\top \mathbf{y}_T(\tilde{\xi}_T) \right] \\ \text{s.t.} \quad & \sum_{\tau \in [t]} \mathbf{b}_{t,i,\tau}^\top \mathbf{y}_\tau(\xi_\tau) \leq a_{t,i}^0(\mathbf{x}) + \mathbf{a}_{t,i}^\top(\mathbf{x}) \xi_t \quad \forall t \in [T], i \in \mathcal{I}_t, \mathbf{z} \in \mathcal{Z} \\ & \mathbf{y}_t \in \mathcal{R}^{I_{\xi_t}, I_{y_t}} \quad \forall t \in [T]. \end{aligned}$$

where the **multi-period ambiguity-averse CARA certainty equivalent**

$$\mathbb{C}_{\mathcal{F}}^{\kappa, \theta} [\tilde{\mathbf{v}}] \triangleq \begin{cases} \min_{t \in [T]: \theta_t > 0} \left\{ \inf_{\mathbb{P} \in \mathcal{F}} \text{ess inf}_{\mathbb{P}} [\tilde{v}_t] \right\} & \text{if } \kappa = 0 \\ \sum_{t \in [T]} \theta_t \inf_{\mathbb{P} \in \mathcal{F}} \mathbb{E}_{\mathbb{P}} [\tilde{v}_t] & \text{if } \kappa = \infty \\ -\kappa \log \left(\sum_{t \in [T]} \theta_t \sup_{\mathbb{P} \in \mathcal{F}} \mathbb{E}_{\mathbb{P}} \left[\exp \left(-\frac{\tilde{v}_t}{\kappa} \right) \right] \right) & \text{if } \kappa \in (0, \infty). \end{cases}$$

► Multi-period MLDR \implies Tractable ECP approximation

Multi-period inventory management

A consumption model [CSSLS07]:

- ▶ Initial inventory x_1 , wealth w_1
- ▶ In each period $t = 1, \dots, T$,

Multi-period inventory management

A consumption model [CSSLS07]:

- ▶ Initial inventory x_1 , wealth w_1
- ▶ In each period $t = 1, \dots, T$,
 - ▶ Order up to $y_t \geq x_t$ with ordering cost $c_t(y_t - x_t)$

Multi-period inventory management

A consumption model [CSSL07]:

- ▶ Initial inventory x_1 , wealth w_1
- ▶ In each period $t = 1, \dots, T$,
 - ▶ Order up to $y_t \geq x_t$ with ordering cost $c_t(y_t - x_t)$
 - ▶ Uncertain demand \tilde{z}_t realizes, obtain profit subtracting holding and backlogging cost $p_t \tilde{z}_t - h(y_t - \tilde{z}_t)^+ - b(\tilde{z}_t - y_t)^+$

Multi-period inventory management

A consumption model [CSSLS07]:

- ▶ Initial inventory x_1 , wealth w_1
- ▶ In each period $t = 1, \dots, T$,
 - ▶ Order up to $y_t \geq x_t$ with ordering cost $c_t(y_t - x_t)$
 - ▶ Uncertain demand \tilde{z}_t realizes, obtain profit subtracting holding and backlogging cost $p_t \tilde{z}_t - h(y_t - \tilde{z}_t)^+ - b(\tilde{z}_t - y_t)^+$
 - ▶ Determine consumption level f_t and obtain utility $1 - e^{-f_t/\kappa}$

Multi-period inventory management

A consumption model [CSSLS07]:

- ▶ Initial inventory x_1 , wealth w_1
- ▶ In each period $t = 1, \dots, T$,
 - ▶ Order up to $y_t \geq x_t$ with ordering cost $c_t(y_t - x_t)$
 - ▶ Uncertain demand \tilde{z}_t realizes, obtain profit subtracting holding and backlogging cost $p_t \tilde{z}_t - h(y_t - \tilde{z}_t)^+ - b(\tilde{z}_t - y_t)^+$
 - ▶ Determine consumption level f_t and obtain utility $1 - e^{-f_t/\kappa}$
 - ▶ Inventory update:

$$x_{t+1} = y_t - \tilde{z}_t$$

Multi-period inventory management

A consumption model [CSSLS07]:

- ▶ Initial inventory x_1 , wealth w_1
- ▶ In each period $t = 1, \dots, T$,
 - ▶ Order up to $y_t \geq x_t$ with ordering cost $c_t(y_t - x_t)$
 - ▶ Uncertain demand \tilde{z}_t realizes, obtain profit subtracting holding and backlogging cost $p_t \tilde{z}_t - h(y_t - \tilde{z}_t)^+ - b(\tilde{z}_t - y_t)^+$
 - ▶ Determine consumption level f_t and obtain utility $1 - e^{-f_t/\kappa}$
 - ▶ Inventory update:

$$x_{t+1} = y_t - \tilde{z}_t$$

- ▶ Wealth update:

$$w_{t+1} = (w_t + q_t - f_t)(1 + \beta)$$

where the income

$$q_t = p_t \tilde{z}_t - h(y_t - \tilde{z}_t)^+ - b(\tilde{z}_t - y_t)^+ - c_t(y_t - \tilde{z}_t)$$

- ▶ Assume $w_{T+1} = 0$.

Multi-period inventory management

$$\begin{aligned} \max_{\mathbf{x}, \mathbf{y}, \mathbf{f}, \mathbf{w}, \mathbf{q}} \quad & \mathbb{E}_{\mathbb{P}} \left[\sum_{t \in [T]} \theta_t (1 - e^{-f_t(\tilde{\xi}_t)/\kappa}) \right] \\ \text{s.t.} \quad & f_t(\tilde{\xi}_t) = w_t(\tilde{\xi}_{t-1}) - \frac{w_{t+1}(\tilde{\xi}_t)}{1 + \beta} + q_t(\tilde{\xi}_t) \quad \forall t \in [T] \\ & q_t(\tilde{\xi}_t) \leq p_t \tilde{z}_t - h(y_t(\tilde{\xi}_{t-1}) - \tilde{z}_t) - c_t(y_t(\tilde{\xi}_{t-1}) - x_t(\tilde{\xi}_{t-1})) \quad \forall t \in [T] \\ & q_t(\tilde{\xi}_t) \leq p_t \tilde{z}_t - b(\tilde{z}_t - y_t(\tilde{\xi}_{t-1})) - c_t(y_t(\tilde{\xi}_{t-1}) - x_t(\tilde{\xi}_{t-1})) \quad \forall t \in [T] \\ & y_t(\tilde{\xi}_{t-1}) \geq x_t(\tilde{\xi}_{t-1}) \quad \forall t \in [T] \\ & x_{t+1}(\tilde{\xi}_t) = y_t(\tilde{\xi}_{t-1}) - \tilde{z}_t \quad \forall t \in [T-1] \\ & w_{T+1}(\tilde{\xi}_T) = 0 \end{aligned}$$

- Approach 1: dynamic programming (DP)
- Approach 2: Fourier-Motzkin elimination + multi-period MLDR

- ▶ Similar parameter setting as in [CSSLS07]:
 $h = 6, b = 3, c_t = 1, p_t = 8, \beta = 0.1$ for all $t \in [T]$

Numerical experiments

- ▶ Similar parameter setting as in [CSSLS07]:
 $h = 6, b = 3, c_t = 1, p_t = 8, \beta = 0.1$ for all $t \in [T]$
- ▶ Risk tolerance $\kappa \in \{0.25, 1, 4, 16, 64, 256\}$

Numerical experiments

- ▶ Similar parameter setting as in [CSSLS07]:
 $h = 6, b = 3, c_t = 1, p_t = 8, \beta = 0.1$ for all $t \in [T]$
- ▶ Risk tolerance $\kappa \in \{0.25, 1, 4, 16, 64, 256\}$
- ▶ Demand data: $S \in \{20, 200\}$ i.i.d. samples from uniform distribution over $\{0, 1, 2, \dots, 20\}$.

Numerical experiments


- ▶ Similar parameter setting as in [CSSLS07]:
 $h = 6, b = 3, c_t = 1, p_t = 8, \beta = 0.1$ for all $t \in [T]$
- ▶ Risk tolerance $\kappa \in \{0.25, 1, 4, 16, 64, 256\}$
- ▶ Demand data: $S \in \{20, 200\}$ i.i.d. samples from uniform distribution over $\{0, 1, 2, \dots, 20\}$.
- ▶ Solving the problem by (i) DP; (ii) MLDR under empirical distribution of S samples, then evaluate the solution (policy) over 10,000 samples generated from the same distribution

Numerical experiments

- ▶ Similar parameter setting as in [CSSLS07]:
 $h = 6, b = 3, c_t = 1, p_t = 8, \beta = 0.1$ for all $t \in [T]$
- ▶ Risk tolerance $\kappa \in \{0.25, 1, 4, 16, 64, 256\}$
- ▶ Demand data: $S \in \{20, 200\}$ i.i.d. samples from uniform distribution over $\{0, 1, 2, \dots, 20\}$.
- ▶ Solving the problem by (i) DP; (ii) MLDR under empirical distribution of S samples, then evaluate the solution (policy) over 10,000 samples generated from the same distribution
- ▶ The results are averaged over 50 random instances.

Numerical experiments

Figure: Multi-period CARA certainty equivalent under different risk tolerance parameters



Summary

- ▶ A robust decision model with CARA preference

$$\max_{\mathbf{x} \in \mathcal{X}} \mathbb{C}_{\mathcal{F}}^{\kappa} [f(\mathbf{x}, \tilde{\mathbf{z}})]$$

- ▶ Tractable ECP approximations for a hierarchy of payoff functions:
 - ▶ Affine perturbations
 - ▶ Concave piecewise affine perturbations
 - ▶ Two-stage optimization with fixed recourse
- ▶ Extend to a multi-period consumption model
- ▶ Robust performance in data-driven setting when risk tolerance is low.

- ▶ Other entropy related decision criteria such as entropic value-at-risk, entropic risk measure.
- ▶ Faster computation for large-scale ECP (with integer constraints)
- ▶ Handle correlated uncertain factors in robust CARA optimization
- ▶ Stronger probability bound of functions of (partially) independent random variables

Acknowledgements



Dr. Melvyn Sim



Dr. Kim-Chuan Toh

Acknowledgements



Dr. Zhenyu Hu



Dr. Napat Rujeerapaiboon

Acknowledgements

- ▶ Prof. Teo Chung-Piaw

Acknowledgements

- ▶ Prof. Teo Chung-Piaw
- ▶ IORA & DAO students

Acknowledgements

- ▶ Prof. Teo Chung-Piaw
- ▶ IORA & DAO students
- ▶ IORA faculty members

Acknowledgements

- ▶ Prof. Teo Chung-Piaw
- ▶ IORA & DAO students
- ▶ IORA faculty members
- ▶ IORA staff

Acknowledgements

- ▶ Prof. Teo Chung-Piaw
- ▶ IORA & DAO students
- ▶ IORA faculty members
- ▶ IORA staff
- ▶ Research collaborators

Acknowledgements

- ▶ Prof. Teo Chung-Piaw
- ▶ IORA & DAO students
- ▶ IORA faculty members
- ▶ IORA staff
- ▶ Research collaborators
- ▶ Sun Qinghe

Acknowledgements

- ▶ Prof. Teo Chung-Piaw
- ▶ IORA & DAO students
- ▶ IORA faculty members
- ▶ IORA staff
- ▶ Research collaborators
- ▶ Sun Qinghe
- ▶ My family

References I

-  James L Corner and Patricia D Corner, *Characteristics of decisions in decision analysis practice*, Journal of the Operational Research Society **46** (1995), no. 3, 304–314.
-  Xin Chen, Melvyn Sim, David Simchi-Levi, and Peng Sun, *Risk aversion in inventory management*, Operations Research **55** (2007), no. 5, 828–842.
-  Xin Chen, Melvyn Sim, Peng Sun, and Jiawei Zhang, *A linear decision-based approximation approach to stochastic programming*, Operations Research **56** (2008), no. 2, 344–357.
-  Itzhak Gilboa and David Schmeidler, *Maxmin expected utility with non-unique prior*, Journal of Mathematical Economics **18** (1989), no. 2, 141–153.
-  Arkadi Nemirovski and Alexander Shapiro, *Convex approximations of chance constrained programs*, SIAM Journal on Optimization **17** (2007), no. 4, 969–996.

References II



John Von Neumann and Oskar Morgenstern, *Theory of games and economic behavior (commemorative edition)*, Princeton university press, 1947.