

Conditional Robust Optimization

Saturday Seminar

Li Chen

May 2022

Outline

Conditional stochastic optimization

Robust optimization to conditional stochastic optimization

Stochastic optimization: given uncertain parameters $\tilde{\mathbf{v}} \sim \mathbb{P}$, find a decision $\mathbf{x} \in \mathcal{X}$ to minimize the expected cost:

$$\min_{\mathbf{x} \in \mathcal{X}} \mathbb{E}_{\mathbb{P}} [g(\mathbf{x}, \tilde{\mathbf{v}})]$$

Stochastic optimization: given uncertain parameters $\tilde{v} \sim \mathbb{P}$, find a decision $x \in \mathcal{X}$ to minimize the expected cost:

$$\min_{x \in \mathcal{X}} \mathbb{E}_{\mathbb{P}} [g(x, \tilde{v})]$$

Example 1 (Newsvendor)

A newsvendor decides to order x units in face with stochastic demand \tilde{v} to maximize the expected profit:

$$\max_{x \geq 0} \mathbb{E}_{\mathbb{P}} [p \min\{\tilde{v}, x\}] - cx$$

where c is the unit ordering cost, p is the unit revenue.

Stochastic optimization: given uncertain parameters $\tilde{\mathbf{v}} \sim \mathbb{P}$, find a decision $\mathbf{x} \in \mathcal{X}$ to minimize the expected cost:

$$\min_{\mathbf{x} \in \mathcal{X}} \mathbb{E}_{\mathbb{P}} [g(\mathbf{x}, \tilde{\mathbf{v}})]$$

In practice, the distribution \mathbb{P} is estimated from historical data $\{\hat{\mathbf{v}}_k\}_{k \in [N]}$, for example, the empirical distribution $\hat{\mathbb{P}} = \frac{1}{N} \sum_{k \in [N]} \delta_{\hat{\mathbf{v}}_k}$.

Motivation

Stochastic optimization: given uncertain parameters $\tilde{\mathbf{v}} \sim \mathbb{P}$, find a decision $\mathbf{x} \in \mathcal{X}$ to minimize the expected cost:

$$\min_{\mathbf{x} \in \mathcal{X}} \mathbb{E}_{\mathbb{P}} [g(\mathbf{x}, \tilde{\mathbf{v}})]$$

In practice, the distribution \mathbb{P} is estimated from historical data $\{\hat{\mathbf{v}}_k\}_{k \in [N]}$, for example, the empirical distribution $\hat{\mathbb{P}} = \frac{1}{N} \sum_{k \in [N]} \delta_{\hat{\mathbf{v}}_k}$.

However, we might have auxiliary data $\{\hat{\mathbf{u}}_k\}_{k \in [N]}$ that can help predict $\tilde{\mathbf{v}}$.

Conditional stochastic optimization: replacing distribution by conditional distribution and solve

$$\min_{\mathbf{x} \in \mathcal{X}} \mathbb{E}_{\mathbb{P}} [g(\mathbf{x}, \tilde{\mathbf{v}}) \mid \tilde{\mathbf{u}} = \mathbf{u}]$$

is motivated to utilize these data for better decision-making.

Conditional stochastic optimization (CSO) framework

- ▶ $\tilde{\mathbf{u}}$ is observable before decision-making, it is called feature/ covariates/ side information/ explanatory variables/ contexts
- ▶ $\tilde{\mathbf{v}}$: the uncertain parameters of interest
- ▶ Assume $(\tilde{\mathbf{u}}, \tilde{\mathbf{v}}) \sim \mathbb{P}^*$, then ideally, we would like to solve

$$\min_{\mathbf{x} \in \mathcal{X}} \mathbb{E}_{\mathbb{P}^*} [g(\mathbf{x}, \tilde{\mathbf{v}}) \mid \tilde{\mathbf{u}} = \mathbf{u}] \quad (1)$$

for each given $\mathbf{u} \in \mathcal{U}$.

- ▶ Equivalently, we write the CSO problem as

$$\begin{aligned} \min_{\mathbf{x}(\cdot)} \quad & \mathbb{E}_{\mathbb{P}^*} [g(\mathbf{x}(\tilde{\mathbf{u}}), \tilde{\mathbf{v}})] \\ \text{s.t.} \quad & \mathbf{x}(\mathbf{u}) \in \mathcal{X} \quad \forall \mathbf{u} \in \mathcal{U} \end{aligned} \quad (2)$$

where $\mathbf{x}(\cdot)$ is a mapping from feature to decision.

Example 1 (Regression)

$$g(\mathbf{x}, \mathbf{v}) = \|\mathbf{x} - \mathbf{v}\|_2^2$$

Example 2 (Joint production and procurement problem)

- ▶ Feature-demand historical data $\{(\hat{\mathbf{u}}_k, \hat{\mathbf{v}}_k)\}_{k \in [N]}$
- ▶ Production and procurement decision

$$\mathcal{X} = \{(\mathbf{x}, \mathbf{r}) \mid \mathbf{A}\mathbf{x} \leq \mathbf{r}, \mathbf{x} \geq \mathbf{0}, \mathbf{r} \geq \mathbf{0}\},$$

- ▶ Cost function:

$$g(\mathbf{x}, \mathbf{r}, \mathbf{v}) = \mathbf{c}^\top \mathbf{r} - \sum_{i \in [n_x]} p_i \min\{x_i, v_i\}$$

Comparison to supervised learning

Recall the equivalent form of CSO:

$$\begin{array}{ll} \min_{\boldsymbol{x}(\cdot)} & \mathbb{E}_{\mathbb{P}^*} [g(\boldsymbol{x}(\tilde{\boldsymbol{u}}), \tilde{\boldsymbol{v}})] \\ \text{s.t.} & \boldsymbol{x}(\boldsymbol{u}) \in \mathcal{X} \quad \forall \boldsymbol{u} \in \mathcal{U} \end{array}$$

- ▶ Mathematically, CSO can be regarded as a generalized supervised learning problem with more complex objective function and **constraints**.
- ▶ From modeling perspective, since supervised learning focus on **prediction** while CSO focus on **decision-making**, the latter requires more interpretability on the policy $\boldsymbol{x}(\cdot)$.

Comparison to supervised learning

Recall the equivalent form of CSO:

$$\begin{array}{ll} \min_{\mathbf{x}(\cdot)} & \mathbb{E}_{\mathbb{P}^*} [g(\mathbf{x}(\tilde{\mathbf{u}}), \tilde{\mathbf{v}})] \\ \text{s.t.} & \mathbf{x}(\mathbf{u}) \in \mathcal{X} \quad \forall \mathbf{u} \in \mathcal{U} \end{array}$$

- ▶ Mathematically, CSO can be regarded as a generalized supervised learning problem with more complex objective function and **constraints**.
- ▶ From modeling perspective, since supervised learning focus on **prediction** while CSO focus on **decision-making**, the latter requires more interpretability on the policy $\mathbf{x}(\cdot)$.

In business analytics, “prescriptive analytics” refers to CSO while “predictive analytics” refers to supervised learning (prediction).

Solution frameworks [BK20]

In practice, \mathbb{P}^* is unknown: we only have data $\{(\hat{\mathbf{u}}_k, \hat{\mathbf{v}}_k)\}_{k \in [N]}$ of feature-outcome pairs.

- Use density estimation to approximate conditional distribution

$$\begin{aligned}\mathbb{P}^* [\tilde{\mathbf{v}} \mid \tilde{\mathbf{u}} = \mathbf{u}] &\longrightarrow \sum_{k \in [N]} w_k(\mathbf{u}) \delta_{\hat{\mathbf{v}}_k} \\ \min_{\mathbf{x} \in \mathcal{X}} \mathbb{E}_{\mathbb{P}^*} [g(\mathbf{x}, \tilde{\mathbf{v}}) \mid \tilde{\mathbf{u}} = \mathbf{u}] &\longrightarrow \min_{\mathbf{x} \in \mathcal{X}} \sum_{k \in [N]} w_k(\mathbf{u}) g(\mathbf{x}, \hat{\mathbf{v}}_k)\end{aligned}$$

- Pros: fix \mathbf{u} , CSO is the same as traditional stochastic optimization
- Cons: density estimation is difficult for high-dim or small N cases
- usually called "predict-then-optimize" approach

Solution frameworks [BK20]

In practice, \mathbb{P}^* is unknown: we only have data $\{(\hat{\mathbf{u}}_k, \hat{\mathbf{v}}_k)\}_{k \in [N]}$ of feature-outcome pairs.

- Use density estimation to approximate conditional distribution

$$\begin{aligned}\mathbb{P}^*[\tilde{\mathbf{v}} \mid \tilde{\mathbf{u}} = \mathbf{u}] &\longrightarrow \sum_{k \in [N]} w_k(\mathbf{u}) \delta_{\hat{\mathbf{v}}_k} \\ \min_{\mathbf{x} \in \mathcal{X}} \mathbb{E}_{\mathbb{P}^*}[g(\mathbf{x}, \tilde{\mathbf{v}}) \mid \tilde{\mathbf{u}} = \mathbf{u}] &\longrightarrow \min_{\mathbf{x} \in \mathcal{X}} \sum_{k \in [N]} w_k(\mathbf{u}) g(\mathbf{x}, \hat{\mathbf{v}}_k)\end{aligned}$$

- Pros: fix \mathbf{u} , CSO is the same as traditional stochastic optimization
- Cons: density estimation is difficult for high-dim or small N cases
- usually called "predict-then-optimize" approach
- Approximate joint distribution \mathbb{P}^* by $\hat{\mathbb{P}}$ and optimize over a chosen policy set \mathcal{A} :

$$\begin{aligned}\min_{\mathbf{x}(\cdot) \in \mathcal{A}} \quad & \mathbb{E}_{\hat{\mathbb{P}}}[g(\mathbf{x}(\tilde{\mathbf{u}}), \tilde{\mathbf{v}})] \\ \text{s.t.} \quad & \mathbf{x}(\mathbf{u}) \in \mathcal{X} \quad \forall \mathbf{u} \in \mathcal{U}\end{aligned}$$

- Pros: directly output a policy; fully utilize available data
- Cons: choice of \mathcal{A} is difficult; more complicated optimization
- Rmk: same as Empirical Risk Minimization (ERM)

CSO via density estimation

Given $\mathbf{u} \in \mathcal{U}$, solve

$$\min_{\mathbf{x} \in \mathcal{X}} \sum_{k \in [N]} w_k(\mathbf{u}) g(\mathbf{x}, \hat{\mathbf{v}}_k)$$

with conditional distribution estimation via

- ▶ Nadaraya-Watson Kernel density estimation (NW-KDE) [[HPB10](#)]

$$w_k(\mathbf{u}) = \frac{K_h(\mathbf{u} - \hat{\mathbf{u}}_k)}{\sum_{j \in [N]} K_h(\mathbf{u} - \hat{\mathbf{u}}_j)}$$

where $K(\cdot)$ is a Kernel function and h is the width hyperparameter.

- ▶ Local learning methods such as k-NN
- ▶ CART
- ▶ Random Forest
- ▶ Parametric statistical models + residual distribution

The choice of the policy set needs to balance the trade-off of tractability, interpretability, and optimality.

- ▶ Affine policy set [BR19]

$$\boldsymbol{x}(\boldsymbol{u}) = \boldsymbol{x}^0 + \boldsymbol{X}\boldsymbol{u}$$

- ▶ Decision tree policy set [BDM19]

$$\boldsymbol{x}(\boldsymbol{u}) = \sum_{\ell \in [L]} \boldsymbol{x}^{\ell} \mathbb{I}[\boldsymbol{u} \in \mathcal{U}_{\ell}]$$

requires solving mixed-integer optimization.

- ▶ Reproducing Kernel Hilbert Space (RKHS) [BK22, NP22]

$$\boldsymbol{x}(\boldsymbol{u}) = \sum_{k \in [N]} K(\boldsymbol{u}, \boldsymbol{u}_k) \boldsymbol{x}^k$$

the constraint $\boldsymbol{x}(\boldsymbol{u}) \in \mathcal{X}$ is difficult to enforce and often ignored.

- ▶ more can be explored...

Outline

Conditional stochastic optimization

Robust optimization to conditional stochastic optimization

Motivation for robust optimization

For machine learning or data-driven stochastic optimization, the goal is to obtain a predictor/ decision with good **out-of-sample** performance.

- ▶ Although many generalization bounds exist for ERM, it requires large N

Motivation for robust optimization

For machine learning or data-driven stochastic optimization, the goal is to obtain a predictor/ decision with good **out-of-sample** performance.

- ▶ Although many generalization bounds exist for ERM, it requires large N
- ▶ Distributionally robust optimization (DRO) accounts distribution shift (discrepancy of training set and test set)

$$\min_{\mathbf{x} \in \mathcal{X}} \mathbb{E}_{\hat{\mathbb{P}}} [g(\mathbf{x}, \tilde{\mathbf{v}})] \implies \min_{\mathbf{x} \in \mathcal{X}} \sup_{\mathbb{P}: d(\mathbb{P}, \hat{\mathbb{P}}) \leq \Gamma} \mathbb{E}_{\mathbb{P}} [g(\mathbf{x}, \tilde{\mathbf{v}})]$$

- ▶ focus on Type-I Wasserstein distance [EK18]

$$d(\mathbb{P}_1, \mathbb{P}_2) = \inf_{\mathbb{Q} \in \mathcal{P}_0(\mathcal{V} \times \mathcal{V})} \{ \mathbb{E}_{\mathbb{Q}} [\|\tilde{\mathbf{v}}_1 - \tilde{\mathbf{v}}_2\|] : (\tilde{\mathbf{v}}_1, \tilde{\mathbf{v}}_2) \sim \mathbb{Q} \text{ with marginal } \mathbb{P}_1, \mathbb{P}_2 \}$$

Motivation for robust optimization

For machine learning or data-driven stochastic optimization, the goal is to obtain a predictor/ decision with good **out-of-sample** performance.

- ▶ Although many generalization bounds exist for ERM, it requires large N
- ▶ Distributionally robust optimization (DRO) accounts distribution shift (discrepancy of training set and test set)

$$\min_{\mathbf{x} \in \mathcal{X}} \mathbb{E}_{\hat{\mathbb{P}}} [g(\mathbf{x}, \tilde{\mathbf{v}})] \implies \min_{\mathbf{x} \in \mathcal{X}} \sup_{\mathbb{P}: d(\mathbb{P}, \hat{\mathbb{P}}) \leq \Gamma} \mathbb{E}_{\mathbb{P}} [g(\mathbf{x}, \tilde{\mathbf{v}})]$$

- ▶ focus on Type-I Wasserstein distance [EK18]

$$d(\mathbb{P}_1, \mathbb{P}_2) = \inf_{\mathbb{Q} \in \mathcal{P}_0(\mathcal{V} \times \mathcal{V})} \{ \mathbb{E}_{\mathbb{Q}} [\|\tilde{\mathbf{v}}_1 - \tilde{\mathbf{v}}_2\|] : (\tilde{\mathbf{v}}_1, \tilde{\mathbf{v}}_2) \sim \mathbb{Q} \text{ with marginal } \mathbb{P}_1, \mathbb{P}_2 \}$$

- ▶ Regularization via robustification [GCK17, SAKE19]

Example 3 (Linear regression)

If $\tilde{\mathbf{u}}$ and $\tilde{\mathbf{v}}$ has no support constraints, then

$$\min_{\mathbf{x} \in \mathbb{R}^{n_u}} \sup_{\mathbb{P}: d(\mathbb{P}, \hat{\mathbb{P}}) \leq \Gamma} \mathbb{E}_{\mathbb{P}} [\|\mathbf{x}^\top \tilde{\mathbf{u}} - \tilde{\mathbf{v}}\|] \iff \min_{\mathbf{x} \in \mathbb{R}^{n_u}} \mathbb{E}_{\hat{\mathbb{P}}} [\|\mathbf{x}^\top \tilde{\mathbf{u}} - \tilde{\mathbf{v}}\|] + \Gamma \|\mathbf{x}; -1\|_*$$

Conditional robust optimization (CRO)

- Robust optimization for “predict-then-optimize” model
[SWHH21, EPM21, KBL20, BVP21]

$$\min_{\mathbf{x} \in \mathcal{X}} \sup_{\mathbb{P}: d(\mathbb{P}, \hat{\mathbb{P}}(\mathbf{u})) \leq \Gamma} \mathbb{E}_{\mathbb{P}} [g(\mathbf{x}, \tilde{\mathbf{v}})]$$

where $\hat{\mathbb{P}}(\mathbf{u}) \triangleq \sum_{k \in [N]} w_k(\mathbf{u}) \delta_{\hat{\mathbf{v}}_k}$

Conditional robust optimization (CRO)

- Robust optimization for “predict-then-optimize” model
[SWHH21, EPM21, KBL20, BVP21]

$$\min_{\mathbf{x} \in \mathcal{X}} \sup_{\mathbb{P}: d(\mathbb{P}, \hat{\mathbb{P}}(\mathbf{u})) \leq \Gamma} \mathbb{E}_{\mathbb{P}} [g(\mathbf{x}, \tilde{\mathbf{v}})]$$

where $\hat{\mathbb{P}}(\mathbf{u}) \triangleq \sum_{k \in [N]} w_k(\mathbf{u}) \delta_{\hat{\mathbf{v}}_k}$

[SWHH21] adopts a variance regularization and establish its connection to DRO.

[EPM21] add additional layer of robustness on the reference distribution $\hat{\mathbb{P}}(\mathbf{u})$

[KBL20] use Wasserstein DRO for the residual distribution with a parametric model

[BVP21] analyze statistical performance of DRO models on bootstrapping sample sets.

Conditional robust optimization (CRO)

- ▶ Robust optimization for “predict-then-optimize” model
[SWHH21, EPM21, KBL20, BVP21]

$$\min_{\mathbf{x} \in \mathcal{X}} \sup_{\mathbb{P}: d(\mathbb{P}, \hat{\mathbb{P}}(\mathbf{u})) \leq \Gamma} \mathbb{E}_{\mathbb{P}} [g(\mathbf{x}, \tilde{\mathbf{v}})]$$

where $\hat{\mathbb{P}}(\mathbf{u}) \triangleq \sum_{k \in [N]} w_k(\mathbf{u}) \delta_{\hat{\mathbf{v}}_k}$

- ▶ The computation is the same as Wasserstein DRO
- ▶ Most focus on statistical consistency

Conditional robust optimization (CRO)

- ▶ Robust optimization for “predict-then-optimize” model [[SWHH21](#), [EPM21](#), [KBL20](#), [BVP21](#)]

$$\min_{\mathbf{x} \in \mathcal{X}} \sup_{\mathbb{P}: d(\mathbb{P}, \hat{\mathbb{P}}(\mathbf{u})) \leq \Gamma} \mathbb{E}_{\mathbb{P}} [g(\mathbf{x}, \tilde{\mathbf{v}})]$$

where $\hat{\mathbb{P}}(\mathbf{u}) \triangleq \sum_{k \in [N]} w_k(\mathbf{u}) \delta_{\hat{\mathbf{v}}_k}$

- ▶ The computation is the same as Wasserstein DRO
 - ▶ Most focus on statistical consistency
- ▶ Robust optimization for “ERM” model [[CSZZ22](#)]
 - ▶ lack of study

The robust ERM model

Consider solving a DRO model over a chosen policy set \mathcal{A} :

$$\begin{aligned} \min_{\mathbf{x}(\cdot) \in \mathcal{A}} \quad & \sup_{\mathbb{P} \in \mathcal{F}(\Gamma)} \mathbb{E}_{\mathbb{P}} [g(\mathbf{x}(\tilde{\mathbf{u}}), \tilde{\mathbf{v}})] \\ \text{s.t.} \quad & \mathbf{x}(\mathbf{u}) \in \mathcal{X}, \forall \mathbf{u} \in \mathcal{U} \end{aligned} \tag{RERM}$$

where the type-I Wasserstein ambiguity set is

$$\mathcal{F}(\Gamma) \triangleq \left\{ \mathbb{P} \in \mathcal{P}_0(\mathcal{Z}) \mid \begin{array}{l} (\tilde{\mathbf{u}}, \tilde{\mathbf{v}}) \sim \mathbb{P} \\ \Delta(\mathbb{P}, \hat{\mathbb{P}}) \leq \Gamma \end{array} \right\},$$

$\mathcal{Z} = \mathcal{U} \times \mathcal{V}$ and

$$\Delta(\mathbb{P}, \hat{\mathbb{P}}) \triangleq \inf_{\mathbb{Q} \in \mathcal{P}_0(\mathcal{Z}^2)} \left\{ \mathbb{E}_{\mathbb{Q}} [\|\tilde{\mathbf{u}} - \tilde{\mathbf{u}}_1\| + \|\tilde{\mathbf{v}} - \tilde{\mathbf{v}}_1\|] \mid \begin{array}{l} (\tilde{\mathbf{u}}, \tilde{\mathbf{v}}, \tilde{\mathbf{u}}_1, \tilde{\mathbf{v}}_1) \sim \mathbb{Q}, \\ (\tilde{\mathbf{u}}, \tilde{\mathbf{v}}) \sim \mathbb{P}, (\tilde{\mathbf{u}}_1, \tilde{\mathbf{v}}_1) \sim \hat{\mathbb{P}} \end{array} \right\}.$$

The robust ERM model

Consider solving a DRO model over a chosen policy set \mathcal{A} :

$$\begin{aligned} \min_{\mathbf{x}(\cdot) \in \mathcal{A}} \quad & \sup_{\mathbb{P} \in \mathcal{F}(\Gamma)} \mathbb{E}_{\mathbb{P}} [g(\mathbf{x}(\tilde{\mathbf{u}}), \tilde{\mathbf{v}})] \\ \text{s.t.} \quad & \mathbf{x}(\mathbf{u}) \in \mathcal{X}, \forall \mathbf{u} \in \mathcal{U} \end{aligned} \tag{RERM}$$

where the type-I Wasserstein ambiguity set is

$$\mathcal{F}(\Gamma) \triangleq \left\{ \mathbb{P} \in \mathcal{P}_0(\mathcal{Z}) \mid \begin{array}{l} (\tilde{\mathbf{u}}, \tilde{\mathbf{v}}) \sim \mathbb{P} \\ \Delta(\mathbb{P}, \hat{\mathbb{P}}) \leq \Gamma \end{array} \right\},$$

$\mathcal{Z} = \mathcal{U} \times \mathcal{V}$ and

$$\Delta(\mathbb{P}, \hat{\mathbb{P}}) \triangleq \inf_{\mathbb{Q} \in \mathcal{P}_0(\mathcal{Z}^2)} \left\{ \mathbb{E}_{\mathbb{Q}} [\|\tilde{\mathbf{u}} - \tilde{\mathbf{u}}_1\| + \|\tilde{\mathbf{v}} - \tilde{\mathbf{v}}_1\|] \mid \begin{array}{l} (\tilde{\mathbf{u}}, \tilde{\mathbf{v}}, \tilde{\mathbf{u}}_1, \tilde{\mathbf{v}}_1) \sim \mathbb{Q}, \\ (\tilde{\mathbf{u}}, \tilde{\mathbf{v}}) \sim \mathbb{P}, (\tilde{\mathbf{u}}_1, \tilde{\mathbf{v}}_1) \sim \hat{\mathbb{P}} \end{array} \right\}.$$

► We focus on linear optimization with recourse:

$$\begin{aligned} g(\mathbf{x}, \mathbf{v}) = \quad & \min \quad \mathbf{d}^\top \mathbf{y} \\ \text{s.t.} \quad & \mathbf{F}(\mathbf{v})\mathbf{x} + \mathbf{B}\mathbf{y} \geq \mathbf{f}(\mathbf{v}) \\ & \mathbf{y} \in \mathbb{R}^{n_y}, \end{aligned}$$

The robust ERM model

Consider solving a DRO model over a chosen policy set \mathcal{A} :

$$\begin{aligned} \min_{\mathbf{x}(\cdot) \in \mathcal{A}} \quad & \sup_{\mathbb{P} \in \mathcal{F}(\Gamma)} \mathbb{E}_{\mathbb{P}} [g(\mathbf{x}(\tilde{\mathbf{u}}), \tilde{\mathbf{v}})] \\ \text{s.t.} \quad & \mathbf{x}(\mathbf{u}) \in \mathcal{X}, \forall \mathbf{u} \in \mathcal{U} \end{aligned} \tag{RERM}$$

where the type-I Wasserstein ambiguity set is

$$\mathcal{F}(\Gamma) \triangleq \left\{ \mathbb{P} \in \mathcal{P}_0(\mathcal{Z}) \mid \begin{array}{l} (\tilde{\mathbf{u}}, \tilde{\mathbf{v}}) \sim \mathbb{P} \\ \Delta(\mathbb{P}, \hat{\mathbb{P}}) \leq \Gamma \end{array} \right\},$$

$\mathcal{Z} = \mathcal{U} \times \mathcal{V}$ and

$$\Delta(\mathbb{P}, \hat{\mathbb{P}}) \triangleq \inf_{\mathbb{Q} \in \mathcal{P}_0(\mathcal{Z}^2)} \left\{ \mathbb{E}_{\mathbb{Q}} [\|\tilde{\mathbf{u}} - \tilde{\mathbf{u}}_1\| + \|\tilde{\mathbf{v}} - \tilde{\mathbf{v}}_1\|] \mid \begin{array}{l} (\tilde{\mathbf{u}}, \tilde{\mathbf{v}}, \tilde{\mathbf{u}}_1, \tilde{\mathbf{v}}_1) \sim \mathbb{Q}, \\ (\tilde{\mathbf{u}}, \tilde{\mathbf{v}}) \sim \mathbb{P}, (\tilde{\mathbf{u}}_1, \tilde{\mathbf{v}}_1) \sim \hat{\mathbb{P}} \end{array} \right\}.$$

- We focus on linear optimization with recourse:

$$\begin{aligned} g(\mathbf{x}, \mathbf{v}) = \quad & \min \quad \mathbf{d}^\top \mathbf{y} \\ \text{s.t.} \quad & \mathbf{F}(\mathbf{v})\mathbf{x} + \mathbf{B}\mathbf{y} \geq \mathbf{f}(\mathbf{v}) \\ & \mathbf{y} \in \mathbb{R}^{n_y}, \end{aligned}$$

- Polyhedral feasibility set

$$\mathcal{X} = \{\mathbf{x} \in \mathbb{R}^{n_x} : \mathbf{A}\mathbf{x} \leq \mathbf{b}\}$$

Adaptive DRO perspective

- Technically, we can regard Problem (RERM) as a three-stage robust optimization problem:

$$\mathbf{w} \rightarrow \tilde{\mathbf{u}} \rightarrow \mathbf{x}(\tilde{\mathbf{u}}) \rightarrow \tilde{\mathbf{v}} \rightarrow \mathbf{y}(\tilde{\mathbf{u}}, \tilde{\mathbf{v}})$$

where \mathbf{w} are auxiliary decisions that are independent of uncertainty.

- Standard duality approach leads to the DRO reformulation [EK18]

$$\begin{aligned} \min \quad & \kappa\Gamma + \frac{1}{N} \sum_{k \in [N]} t_k \\ \text{s.t.} \quad & \sup_{(\mathbf{u}, \mathbf{v}) \in \mathcal{Z}} \{g(\mathbf{x}(\mathbf{u}), \mathbf{v}) - \kappa(\|\mathbf{v} - \hat{\mathbf{v}}_k\| + \|\mathbf{u} - \hat{\mathbf{u}}_k\|)\} \leq t_k \quad \forall k \in [N] \\ & \mathbf{x}(\mathbf{u}) \in \mathcal{X} \quad \forall \mathbf{u} \in \mathcal{U} \\ & \mathbf{x} \in \mathcal{A}, \kappa \geq 0. \end{aligned} \tag{3}$$

Scenario-wise lifted equivalence

Recall that

$$g(\mathbf{x}, \mathbf{v}) = \min_{\substack{\mathbf{d}^\top \mathbf{y} \\ \text{s.t. } \mathbf{F}(\mathbf{v})\mathbf{x} + \mathbf{B}\mathbf{y} \geq \mathbf{f}(\mathbf{v}) \\ \mathbf{y} \in \mathbb{R}^{n_y},}}$$

then for each $k \in [N]$:

$$\begin{aligned} & g(\mathbf{x}(\mathbf{u}), \mathbf{v}) - \kappa(\|\mathbf{v} - \hat{\mathbf{v}}_k\| + \|\mathbf{u} - \hat{\mathbf{u}}_k\|) \leq t_k \\ \iff & \mathbf{d}^\top \mathbf{y}_k(\mathbf{u}, \mathbf{v}) - \kappa(\|\mathbf{v} - \hat{\mathbf{v}}_k\| + \|\mathbf{u} - \hat{\mathbf{u}}_k\|) \leq t_k \\ & \mathbf{F}(\mathbf{v})\mathbf{x}(\mathbf{u}) + \mathbf{B}\mathbf{y}_k(\mathbf{u}, \mathbf{v}) \geq \mathbf{f}(\mathbf{v}) \quad \forall (\mathbf{u}, \mathbf{v}) \in \mathcal{Z} \\ \iff & \mathbf{d}^\top \mathbf{y}_k(\mathbf{u}, \mathbf{v}, \sigma, \nu) - \kappa(\sigma + \nu) \leq t_k \quad \forall (\mathbf{u}, \mathbf{v}, \sigma, \nu) \in \bar{\mathcal{Z}}_k \\ & \mathbf{F}(\mathbf{v})\mathbf{x}(\mathbf{u}) + \mathbf{B}\mathbf{y}_k(\mathbf{u}, \mathbf{v}, \sigma, \nu) \geq \mathbf{f}(\mathbf{v}) \quad \forall (\mathbf{u}, \mathbf{v}, \sigma, \nu) \in \bar{\mathcal{Z}}_k \end{aligned}$$

where the lifted uncertainty set is

$$\bar{\mathcal{Z}}_k \triangleq \{(\mathbf{u}, \mathbf{v}, \sigma, \nu) \in \mathcal{U} \times \mathcal{V} \times \mathbb{R} \times \mathbb{R} \mid \sigma \geq \|\mathbf{u} - \hat{\mathbf{u}}_k\|, \nu \geq \|\mathbf{v} - \hat{\mathbf{v}}_k\|\}$$

Last equivalence: given $\mathbf{y}_k(\mathbf{u}, \mathbf{v})$, construct $\bar{\mathbf{y}}_k(\mathbf{u}, \mathbf{v}, \sigma, \nu) = \mathbf{y}_k(\mathbf{u}, \mathbf{v})$;
given $\bar{\mathbf{y}}_k(\mathbf{u}, \mathbf{v}, \sigma, \nu)$, construct $\mathbf{y}_k(\mathbf{u}, \mathbf{v}) = \bar{\mathbf{y}}_k(\mathbf{u}, \mathbf{v}, \|\mathbf{u} - \hat{\mathbf{u}}_k\|, \|\mathbf{v} - \hat{\mathbf{v}}_k\|)$.

Tractable approximations

The difficult constraints are

$$\begin{aligned} \mathbf{d}^\top \mathbf{y}_k(\mathbf{u}, \mathbf{v}, \sigma, \nu) - \kappa(\sigma + \nu) &\leq t_k & \forall (\mathbf{u}, \mathbf{v}, \sigma, \nu) \in \bar{\mathcal{Z}}_k \\ \mathbf{F}(\mathbf{v})\mathbf{x}(\mathbf{u}) + \mathbf{B}\mathbf{y}_k(\mathbf{u}, \mathbf{v}, \sigma, \nu) &\geq \mathbf{f}(\mathbf{v}) & \forall (\mathbf{u}, \mathbf{v}, \sigma, \nu) \in \bar{\mathcal{Z}}_k \end{aligned}$$

Note that even $\mathbf{x}(\cdot)$ is affine, we have bilinear optimization to solve.

► First approximation: Bi-affine approx of $\mathbf{y}_k(\cdot)$: Let

$$\begin{aligned} \mathbf{y}_k(\mathbf{u}, \mathbf{v}, \sigma, \nu) = & \mathbf{y}_k^0 + \sum_{i \in [n_u]} \mathbf{y}_k^i u_i + \sum_{j \in [n_v]} \mathbf{y}_k^{n_u+j} v_j + \mathbf{y}_k^{n_u+n_v+1} \sigma \\ & + \mathbf{y}_k^{n_u+n_v+2} \nu + \sum_{i \in [n_u]} \sum_{j \in [n_v]} \mathbf{y}_k^{ij} u_i v_j \end{aligned}$$

► It remains to approximate bilinear constraints of the form:

$$\mathbf{q}_1^\top \mathbf{u} + \mathbf{q}_2^\top \mathbf{v} + \mathbf{v}^\top \mathbf{Q}_3 \mathbf{u} \leq q_4 \sigma + q_5 \nu + q_6, \quad \forall (\mathbf{u}, \mathbf{v}, \sigma, \nu) \in \bar{\mathcal{Z}}_k \quad (4)$$

Note q_4 and q_5 are necessarily non-negative.

Bilinear optimization via adaptive robust optimization

Assume \mathcal{U} and \mathcal{V} are non-empty bounded polytope, for simplicity consider

$$\mathcal{U} = \{\mathbf{u} \in \mathbb{R}^{n_u} \mid \underline{\mathbf{u}} \leq \mathbf{u} \leq \bar{\mathbf{u}}\}, \quad \mathcal{V} = \{\mathbf{v} \in \mathbb{R}^{n_v} \mid \underline{\mathbf{v}} \leq \mathbf{v} \leq \bar{\mathbf{v}}\},$$

Bilinear optimization via adaptive robust optimization

Assume \mathcal{U} and \mathcal{V} are non-empty bounded polytope, for simplicity consider

$$\mathcal{U} = \{\mathbf{u} \in \mathbb{R}^{n_u} \mid \underline{\mathbf{u}} \leq \mathbf{u} \leq \bar{\mathbf{u}}\}, \quad \mathcal{V} = \{\mathbf{v} \in \mathbb{R}^{n_v} \mid \underline{\mathbf{v}} \leq \mathbf{v} \leq \bar{\mathbf{v}}\},$$

Then we dualize \mathbf{v} to obtain the following equivalence

$$\begin{aligned} & \sup_{(\mathbf{u}, \mathbf{v}, \sigma, \nu) \in \bar{\mathcal{Z}}_k} \{ \mathbf{q}_1^\top \mathbf{u} + \mathbf{q}_2^\top \mathbf{v} + \mathbf{v}^\top \mathbf{Q}_3 \mathbf{u} - q_4 \sigma - q_5 \nu \} \\ = & \sup_{\mathbf{u} \in \mathcal{U}, \mathbf{v} \in \mathcal{V}} \{ \mathbf{q}_1^\top \mathbf{u} + \mathbf{q}_2^\top \mathbf{v} + \mathbf{v}^\top \mathbf{Q}_3 \mathbf{u} - q_4 \|\mathbf{u} - \hat{\mathbf{u}}_k\| - q_5 \|\mathbf{v} - \hat{\mathbf{v}}_k\| \} \\ = & \sup_{\mathbf{u} \in \mathcal{U}} \inf_{\bar{\boldsymbol{\rho}} \geq \mathbf{0}, \underline{\boldsymbol{\rho}} \geq \mathbf{0}} \{ \mathbf{q}_1^\top \mathbf{u} + \bar{\boldsymbol{\rho}}^\top \bar{\mathbf{v}} - \underline{\boldsymbol{\rho}}^\top \underline{\mathbf{v}} + (\mathbf{q}_2 + \mathbf{Q}_3 \mathbf{u} + \underline{\boldsymbol{\rho}} - \bar{\boldsymbol{\rho}})^\top \hat{\mathbf{v}}_k \\ & - q_4 \|\mathbf{u} - \hat{\mathbf{u}}_k\| : \|\mathbf{q}_2 + \mathbf{Q}_3 \mathbf{u} - \bar{\boldsymbol{\rho}} + \underline{\boldsymbol{\rho}}\|_* \leq q_5 \} . \end{aligned}$$

Bilinear optimization via adaptive robust optimization

Assume \mathcal{U} and \mathcal{V} are non-empty bounded polytope, for simplicity consider

$$\mathcal{U} = \{\mathbf{u} \in \mathbb{R}^{n_u} \mid \underline{\mathbf{u}} \leq \mathbf{u} \leq \bar{\mathbf{u}}\}, \quad \mathcal{V} = \{\mathbf{v} \in \mathbb{R}^{n_v} \mid \underline{\mathbf{v}} \leq \mathbf{v} \leq \bar{\mathbf{v}}\},$$

Then we dualize \mathbf{v} to obtain the following equivalence

$$\begin{aligned} & \sup_{(\mathbf{u}, \mathbf{v}, \sigma, \nu) \in \bar{\mathcal{Z}}_k} \{ \mathbf{q}_1^\top \mathbf{u} + \mathbf{q}_2^\top \mathbf{v} + \mathbf{v}^\top \mathbf{Q}_3 \mathbf{u} - q_4 \sigma - q_5 \nu \} \\ = & \sup_{\mathbf{u} \in \mathcal{U}, \mathbf{v} \in \mathcal{V}} \{ \mathbf{q}_1^\top \mathbf{u} + \mathbf{q}_2^\top \mathbf{v} + \mathbf{v}^\top \mathbf{Q}_3 \mathbf{u} - q_4 \|\mathbf{u} - \hat{\mathbf{u}}_k\| - q_5 \|\mathbf{v} - \hat{\mathbf{v}}_k\| \} \\ = & \sup_{\mathbf{u} \in \mathcal{U}} \inf_{\bar{\boldsymbol{\rho}} \geq \mathbf{0}, \underline{\boldsymbol{\rho}} \geq \mathbf{0}} \{ \mathbf{q}_1^\top \mathbf{u} + \bar{\boldsymbol{\rho}}^\top \bar{\mathbf{v}} - \underline{\boldsymbol{\rho}}^\top \underline{\mathbf{v}} + (\mathbf{q}_2 + \mathbf{Q}_3 \mathbf{u} + \underline{\boldsymbol{\rho}} - \bar{\boldsymbol{\rho}})^\top \hat{\mathbf{v}}_k \\ & - q_4 \|\mathbf{u} - \hat{\mathbf{u}}_k\| : \|\mathbf{q}_2 + \mathbf{Q}_3 \mathbf{u} - \bar{\boldsymbol{\rho}} + \underline{\boldsymbol{\rho}}\|_* \leq q_5 \} . \end{aligned}$$

We regard it as an adaptive robust optimization problem with uncertainty \mathbf{u} and recourse decision $\bar{\boldsymbol{\rho}}, \underline{\boldsymbol{\rho}}$.

Bilinear optimization via adaptive robust optimization

Assume \mathcal{U} and \mathcal{V} are non-empty bounded polytope, for simplicity consider

$$\mathcal{U} = \{\mathbf{u} \in \mathbb{R}^{n_u} \mid \underline{\mathbf{u}} \leq \mathbf{u} \leq \bar{\mathbf{u}}\}, \quad \mathcal{V} = \{\mathbf{v} \in \mathbb{R}^{n_v} \mid \underline{\mathbf{v}} \leq \mathbf{v} \leq \bar{\mathbf{v}}\},$$

We regard it as an adaptive robust optimization problem with uncertainty \mathbf{u} and recourse decision $\bar{\boldsymbol{\rho}}, \underline{\boldsymbol{\rho}}$.

$$\inf_{\bar{\boldsymbol{\rho}}(\mathbf{u}) \geq \mathbf{0}, \underline{\boldsymbol{\rho}}(\mathbf{u}) \geq \mathbf{0}} \sup_{\mathbf{u} \in \mathcal{U}} \left\{ \mathbf{q}_1^\top \mathbf{u} + \bar{\boldsymbol{\rho}}(\mathbf{u})^\top \bar{\mathbf{v}} - \underline{\boldsymbol{\rho}}(\mathbf{u})^\top \underline{\mathbf{v}} - q_4 \|\mathbf{u} - \hat{\mathbf{u}}_k\| \right. \\ \left. + (\mathbf{q}_2 + \mathbf{Q}_3 \mathbf{u} + \underline{\boldsymbol{\rho}}(\mathbf{u}) - \bar{\boldsymbol{\rho}}(\mathbf{u}))^\top \hat{\mathbf{v}}_k : \|\mathbf{q}_2 + \mathbf{Q}_3 \mathbf{u} - \bar{\boldsymbol{\rho}}(\mathbf{u}) + \underline{\boldsymbol{\rho}}(\mathbf{u})\|_* \leq q_5 \right\}$$

Bilinear optimization via adaptive robust optimization

Assume \mathcal{U} and \mathcal{V} are non-empty bounded polytope, for simplicity consider

$$\mathcal{U} = \{\mathbf{u} \in \mathbb{R}^{n_u} \mid \underline{\mathbf{u}} \leq \mathbf{u} \leq \bar{\mathbf{u}}\}, \quad \mathcal{V} = \{\mathbf{v} \in \mathbb{R}^{n_v} \mid \underline{\mathbf{v}} \leq \mathbf{v} \leq \bar{\mathbf{v}}\},$$

We regard it as an adaptive robust optimization problem with uncertainty \mathbf{u} and recourse decision $\bar{\boldsymbol{\rho}}, \underline{\boldsymbol{\rho}}$.

$$\inf_{\bar{\boldsymbol{\rho}}(\mathbf{u}) \geq \mathbf{0}, \underline{\boldsymbol{\rho}}(\mathbf{u}) \geq \mathbf{0}} \sup_{\mathbf{u} \in \mathcal{U}} \left\{ \mathbf{q}_1^\top \mathbf{u} + \bar{\boldsymbol{\rho}}(\mathbf{u})^\top \bar{\mathbf{v}} - \underline{\boldsymbol{\rho}}(\mathbf{u})^\top \underline{\mathbf{v}} - q_4 \|\mathbf{u} - \hat{\mathbf{u}}_k\| \right. \\ \left. + (q_2 + \mathbf{Q}_3 \mathbf{u} + \underline{\boldsymbol{\rho}}(\mathbf{u}) - \bar{\boldsymbol{\rho}}(\mathbf{u}))^\top \hat{\mathbf{v}}_k : \|\mathbf{q}_2 + \mathbf{Q}_3 \mathbf{u} - \bar{\boldsymbol{\rho}}(\mathbf{u}) + \underline{\boldsymbol{\rho}}(\mathbf{u})\|_* \leq q_5 \right\}$$

We apply affine recourse approx: $\bar{\boldsymbol{\rho}}(\mathbf{u}) = \bar{\mathbf{p}} + \bar{\mathbf{P}}\mathbf{u}$, $\underline{\boldsymbol{\rho}}(\mathbf{u}) = \underline{\mathbf{p}} + \underline{\mathbf{P}}\mathbf{u}$.

$$\sup_{\mathbf{u} \in \mathcal{U}} \left\{ \mathbf{q}_1^\top \mathbf{u} + (\bar{\mathbf{p}} + \bar{\mathbf{P}}\mathbf{u})^\top \bar{\mathbf{v}} - (\underline{\mathbf{p}} + \underline{\mathbf{P}}\mathbf{u})^\top \underline{\mathbf{v}} \right. \\ \left. + (q_2 + \mathbf{Q}_3 \mathbf{u} + \underline{\mathbf{p}} + \underline{\mathbf{P}}\mathbf{u} - \bar{\mathbf{p}} - \bar{\mathbf{P}}\mathbf{u})^\top \hat{\mathbf{v}}_k - q_4 \|\mathbf{u} - \hat{\mathbf{u}}_k\| \right\} \leq q_6 \\ \sup_{\mathbf{u} \in \mathcal{U}} \left\{ \|\mathbf{q}_2 + \mathbf{Q}_3 \mathbf{u} - \bar{\mathbf{p}} - \bar{\mathbf{P}}\mathbf{u} + \underline{\mathbf{p}} + \underline{\mathbf{P}}\mathbf{u}\|_* \right\} \leq q_5 \\ \inf_{\mathbf{u} \in \mathcal{U}} \left\{ \mathbf{e}_k^\top (\bar{\mathbf{p}} + \bar{\mathbf{P}}\mathbf{u}) \right\} \geq 0 \quad \forall i \in [n_v] \\ \inf_{\mathbf{u} \in \mathcal{U}} \left\{ \mathbf{e}_k^\top (\underline{\mathbf{p}} + \underline{\mathbf{P}}\mathbf{u}) \right\} \geq 0 \quad \forall i \in [n_v]$$

Maximize dual norm over a polytope

Recall \mathcal{U} is a bounded polytope. However, norm maximization over polytope is difficult in general. Indeed,

$$\max_{\mathbf{x}} \{ \mathbf{x}^\top \mathbf{\Lambda} \mathbf{x} : \|\mathbf{x}\|_\infty \leq 1 \}$$

is NP-hard for general $\mathbf{\Lambda} \succeq \mathbf{0}$ [Hås01].

Maximize dual norm over a polytope

Recall \mathcal{U} is a bounded polytope. However, norm maximization over polytope is difficult in general. Indeed,

$$\max_{\mathbf{x}} \{ \mathbf{x}^\top \mathbf{\Lambda} \mathbf{x} : \|\mathbf{x}\|_\infty \leq 1 \}$$

is NP-hard for general $\mathbf{\Lambda} \succeq \mathbf{0}$ [Hås01].

$$\begin{aligned} & \sup_{\mathbf{u} \in \mathcal{U}} \{ \|\mathbf{q}_2 + \mathbf{Q}_3 \mathbf{u} - \bar{\mathbf{p}} - \bar{\mathbf{P}} \mathbf{u} + \underline{\mathbf{p}} + \underline{\mathbf{P}} \mathbf{u}\|_* \} \leq q_5 \\ \iff & \sup_{\mathbf{u} \in \mathcal{U}, \|\boldsymbol{\gamma}\| \leq 1} \{ \boldsymbol{\gamma}^\top (\mathbf{q}_2 + \mathbf{Q}_3 \mathbf{u} - \bar{\mathbf{p}} - \bar{\mathbf{P}} \mathbf{u} + \underline{\mathbf{p}} + \underline{\mathbf{P}} \mathbf{u}) \} \leq q_5 \\ \iff & \sup_{\|\boldsymbol{\gamma}\| \leq 1} \inf_{\bar{\boldsymbol{\theta}} \geq \mathbf{0}, \boldsymbol{\theta} \geq \mathbf{0}} \{ \boldsymbol{\gamma}^\top (\mathbf{q}_2 - \bar{\mathbf{p}} + \underline{\mathbf{p}}) + \bar{\mathbf{u}}^\top \bar{\boldsymbol{\theta}} - \underline{\mathbf{u}}^\top \boldsymbol{\theta} : \\ & (\mathbf{Q}_3 - \bar{\mathbf{P}} + \underline{\mathbf{P}})^\top \boldsymbol{\gamma} = \bar{\boldsymbol{\theta}} - \boldsymbol{\theta} \} \leq q_5 \end{aligned}$$

Maximize dual norm over a polytope

Recall \mathcal{U} is a bounded polytope. However, norm maximization over polytope is difficult in general. Indeed,

$$\max_{\mathbf{x}} \{ \mathbf{x}^\top \mathbf{\Lambda} \mathbf{x} : \|\mathbf{x}\|_\infty \leq 1 \}$$

is NP-hard for general $\mathbf{\Lambda} \succeq \mathbf{0}$ [Hås01].

$$\begin{aligned} & \sup_{\mathbf{u} \in \mathcal{U}} \{ \|\mathbf{q}_2 + \mathbf{Q}_3 \mathbf{u} - \bar{\mathbf{p}} - \bar{\mathbf{P}} \mathbf{u} + \underline{\mathbf{p}} + \underline{\mathbf{P}} \mathbf{u}\|_* \} \leq q_5 \\ \iff & \sup_{\mathbf{u} \in \mathcal{U}, \|\boldsymbol{\gamma}\| \leq 1} \{ \boldsymbol{\gamma}^\top (\mathbf{q}_2 + \mathbf{Q}_3 \mathbf{u} - \bar{\mathbf{p}} - \bar{\mathbf{P}} \mathbf{u} + \underline{\mathbf{p}} + \underline{\mathbf{P}} \mathbf{u}) \} \leq q_5 \\ \iff & \sup_{\|\boldsymbol{\gamma}\| \leq 1} \inf_{\bar{\boldsymbol{\theta}} \geq \mathbf{0}, \underline{\boldsymbol{\theta}} \geq \mathbf{0}} \{ \boldsymbol{\gamma}^\top (\mathbf{q}_2 - \bar{\mathbf{p}} + \underline{\mathbf{p}}) + \bar{\mathbf{u}}^\top \bar{\boldsymbol{\theta}} - \underline{\mathbf{u}}^\top \underline{\boldsymbol{\theta}} : \\ & (\mathbf{Q}_3 - \bar{\mathbf{P}} + \underline{\mathbf{P}})^\top \boldsymbol{\gamma} = \bar{\boldsymbol{\theta}} - \underline{\boldsymbol{\theta}} \} \leq q_5 \end{aligned}$$

Again, we can apply affine recourse approx of $\bar{\boldsymbol{\theta}}, \underline{\boldsymbol{\theta}}$ with uncertainty $\boldsymbol{\gamma}$:

$$\bar{\boldsymbol{\theta}}(\boldsymbol{\gamma}) = \bar{\boldsymbol{\theta}} + \bar{\boldsymbol{\Theta}}^\top \boldsymbol{\gamma}, \quad \underline{\boldsymbol{\theta}}(\boldsymbol{\gamma}) = \underline{\boldsymbol{\theta}} + \underline{\boldsymbol{\Theta}}^\top \boldsymbol{\gamma} - (\mathbf{Q}_3 - \bar{\mathbf{P}} + \underline{\mathbf{P}})^\top \boldsymbol{\gamma}$$

Maximize dual norm over a polytope

Recall \mathcal{U} is a bounded polytope. However, norm maximization over polytope is difficult in general. Indeed,

$$\max_{\mathbf{x}} \{ \mathbf{x}^\top \mathbf{\Lambda} \mathbf{x} : \|\mathbf{x}\|_\infty \leq 1 \}$$

is NP-hard for general $\mathbf{\Lambda} \succeq \mathbf{0}$ [Hås01].

$$\begin{aligned} & \sup_{\mathbf{u} \in \mathcal{U}} \{ \|\mathbf{q}_2 + \mathbf{Q}_3 \mathbf{u} - \bar{\mathbf{p}} - \bar{\mathbf{P}} \mathbf{u} + \underline{\mathbf{p}} + \underline{\mathbf{P}} \mathbf{u}\|_* \} \leq q_5 \\ \iff & \sup_{\mathbf{u} \in \mathcal{U}, \|\boldsymbol{\gamma}\| \leq 1} \{ \boldsymbol{\gamma}^\top (\mathbf{q}_2 + \mathbf{Q}_3 \mathbf{u} - \bar{\mathbf{p}} - \bar{\mathbf{P}} \mathbf{u} + \underline{\mathbf{p}} + \underline{\mathbf{P}} \mathbf{u}) \} \leq q_5 \\ \iff & \sup_{\|\boldsymbol{\gamma}\| \leq 1} \inf_{\bar{\boldsymbol{\theta}} \geq \mathbf{0}, \underline{\boldsymbol{\theta}} \geq \mathbf{0}} \{ \boldsymbol{\gamma}^\top (\mathbf{q}_2 - \bar{\mathbf{p}} + \underline{\mathbf{p}}) + \bar{\mathbf{u}}^\top \bar{\boldsymbol{\theta}} - \underline{\mathbf{u}}^\top \underline{\boldsymbol{\theta}} : \\ & (\mathbf{Q}_3 - \bar{\mathbf{P}} + \underline{\mathbf{P}})^\top \boldsymbol{\gamma} = \bar{\boldsymbol{\theta}} - \underline{\boldsymbol{\theta}} \} \leq q_5 \end{aligned}$$

Again, we can apply affine recourse approx of $\bar{\boldsymbol{\theta}}, \underline{\boldsymbol{\theta}}$ with uncertainty $\boldsymbol{\gamma}$:

$$\bar{\boldsymbol{\theta}}(\boldsymbol{\gamma}) = \boldsymbol{\theta} + \boldsymbol{\Theta}^\top \boldsymbol{\gamma}, \quad \underline{\boldsymbol{\theta}}(\boldsymbol{\gamma}) = \boldsymbol{\theta} + \boldsymbol{\Theta}^\top \boldsymbol{\gamma} - (\mathbf{Q}_3 - \bar{\mathbf{P}} + \underline{\mathbf{P}})^\top \boldsymbol{\gamma}$$

Rmk: For 1-norm $\|\cdot\|_1$ with dual norm $\|\cdot\|_\infty$, one can enumerate $2n_\gamma$ vertices of 1-norm unit ball.

Summary of deriving safe approximations

1. Start from Problem **RERM** (DRO model)
2. Obtain equivalent robust optimization problem (3) for each $k \in [N]$ by standard duality approach
3. For each $k \in [N]$, we adopt a lifted bi-affine approx of $\mathbf{y}(\mathbf{u}, \mathbf{v}, \sigma, \nu)$
4. Approximate the robust bilinear constraints via adaptive robust optimization (repeatedly use LP duality + affine recourse approx)







Summary of deriving safe approximations

1. Start from Problem [RERM](#) (DRO model)
2. Obtain equivalent robust optimization problem (3) for each $k \in [N]$ by standard duality approach
3. For each $k \in [N]$, we adopt a lifted bi-affine approx of $\mathbf{y}(\mathbf{u}, \mathbf{v}, \sigma, \nu)$
4. Approximate the robust bilinear constraints via adaptive robust optimization (repeatedly use LP duality + affine recourse approx)
 - ▶ Close relation of adaptive robust opt and bilinear opt, some connection of affine recourse approx and reformulation-linearization-technique (RLT) is shown [[ZMdM⁺22](#)]
 - ▶ Generalize the approx from bilinear opt to multi-linear opt





Conclusion

- ▶ Conditional stochastic optimization (CSO) unifies supervised learning and data-driven stochastic optimization
- ▶ Most literature focus on statistical properties
- ▶ Robustifying CSO brings additional challenges for optimization
- ▶ We study a RERM model and derive its tractable convex approximations
 - ▶ we use a tree-based piecewise affine policy in [CSZZ22]
 - ▶ more numerical study is needed
 - ▶ more research on modeling and optimization: u and v are not disjoint, non-linear decision rule with robust opt, endogenous uncertainty...

References I

-  Dimitris Bertsimas, Jack Dunn, and Nishanth Mundru, *Optimal prescriptive trees*, INFORMS Journal on Optimization **1** (2019), no. 2, 164–183.
-  D. Bertsimas and N. Kallus, *From predictive to prescriptive analytics*, Management Science **66** (2020), no. 3, 1025–1044.
-  Dimitris Bertsimas and Nihal Koduri, *Data-driven optimization: A reproducing kernel Hilbert space approach*, Operations Research **70** (2022), no. 1, 454–471.
-  Gah-Yi Ban and Cynthia Rudin, *The big data newsvendor: Practical insights from machine learning*, Operations Research **67** (2019), no. 1, 90–108.
-  Dimitris Bertsimas and Bart Van Parys, *Bootstrap robust prescriptive analytics*, Mathematical Programming (2021), 1–40.
-  Li Chen, Melvyn Sim, Xun Zhang, and Minglong Zhou, *Robust explainable prescriptive analytics*, Available at SSRN 4106222 (2022).

References II

-  Peyman Mohajerin Esfahani and Daniel Kuhn, *Data-driven distributionally robust optimization using the Wasserstein metric: Performance guarantees and tractable reformulations*, Mathematical Programming **171** (2018), no. 1-2, 115–166.
-  Adrián Esteban-Pérez and Juan M Morales, *Distributionally robust stochastic programs with side information based on trimmings*, Mathematical Programming (2021), 1–37.
-  R. Gao, X. Chen, and A.J Kleywegt, *Wasserstein distributional robustness and regularization in statistical learning*, Available at arXiv:1712.06050, 2017.
-  Johan Håstad, *Some optimal inapproximability results*, Journal of the ACM (JACM) **48** (2001), no. 4, 798–859.
-  Lauren Hannah, Warren Powell, and David Blei, *Nonparametric density estimation for stochastic optimization with an observable state variable*, Advances in Neural Information Processing Systems **23** (2010).

References III

-  Rohit Kannan, Güzin Bayraksan, and James R Luedtke, *Residuals-based distributionally robust optimization with covariate information*, arXiv preprint arXiv:2012.01088 (2020).
-  Pascal M Notz and Richard Pibernik, *Prescriptive analytics for flexible capacity management*, Management Science **68** (2022), no. 3, 1756–1775.
-  Soroosh Shafieezadeh-Abadeh, Daniel Kuhn, and Peyman Mohajerin Esfahani, *Regularization via mass transportation*, Journal of Machine Learning Research **20** (2019), no. 103, 1–68.
-  Prateek R Srivastava, Yijie Wang, Grani A Hanasusanto, and Chin Pang Ho, *On data-driven prescriptive analytics with side information: A regularized nadaraya-watson approach*, arXiv preprint arXiv:2110.04855 (2021).

References IV



Jianzhe Zhen, Ahmadreza Marandi, Danique de Moor, Dick den Hertog, and Lieven Vandenbergh, *Disjoint bilinear programming: a two-stage robust optimization perspective*, INFORMS Journal on Computing (2022).