

Particle Physics 3: Supersymmetry and Grand Unification

Lecture 7: Superspace and the Construction of Supersymmetric Dynamics

Leonard Susskind

Original lecture by Leonard Susskind

Transcript-derived notes curated by [LazyingArt LLC](#) with [Video2Book](#)

Chapter 1

Superspace and the Construction of Supersymmetric Dynamics

Overview. We rebuild supersymmetry from the smallest system in which it can act: one bosonic oscillator and one fermionic oscillator of equal energy. The contrast between their second- and first-order equations leads to the supercharges and their algebra. We then reinterpret that algebra as geometry on a space with the ordinary coordinate (t) and the Grassmann coordinates $\theta, \bar{\theta}$. Superfields package bosons, fermions, and an auxiliary field into a single object; Berezin integration extracts an invariant ordinary Lagrangian; covariant derivatives make symmetry-compatible constraints possible; and the chiral constraint provides the first important example. The lecture deliberately exposes the sign and ordering difficulties of this machinery, then closes by explaining why the symmetry itself is more fundamental than a catalogue of Feynman rules.

1.1 Return to the One-Mode System

Let us begin again with fermions and bosons at rest. We suppress every spatial coordinate and every momentum label. A quantum is either present or absent in a given mode, and creation and annihilation operators are enough to describe it. Write $a^\dagger, a$ for a bosonic mode and $c^\dagger, c$ for a fermionic mode. They obey

$$[a, a^\dagger] = 1, \quad \{c, c^\dagger\} = 1, \quad c^2 = (c^\dagger)^2 = 0. \quad (1.1)$$

If the two quanta have the same energy m , then, apart from an irrelevant vacuum constant,

$$H = m (a^\dagger a + c^\dagger c). \quad (1.2)$$

The number operator $a^\dagger a$ may take any nonnegative integer value, while $c^\dagger c$ can only be zero or one. This is the entire kinematics of the toy model.

The simplification is strategic. We want to understand the operation that removes a boson and inserts a fermion, or does the reverse, without changing the energy. Before constructing that operation, however, we translate the same system from operator language into Lagrangian language.


1.2 The Bosonic Oscillator

For the boson, introduce a real coordinate $\phi(t)$. With no spatial variation, a free scalar field has become an ordinary harmonic oscillator. The first expression written on the board was

$$\frac{1}{2}\dot{\phi}^2 + \frac{1}{2}m^2\phi^2.$$

That expression is kinetic plus potential energy, not the Lagrangian. The sign is corrected immediately in the lecture:

$$L_B = \frac{1}{2}\dot{\phi}^2 - \frac{1}{2}m^2\phi^2. \quad (1.3)$$



Lecture frame: 00:03:25.431.

Figure 1.1: The one-mode Hamiltonian above the bosonic Lagrangian. The frame preserves the initial plus sign before the spoken correction to $T - V$.

Question from the audience. What is wrong with the sign in the displayed oscillator Lagrangian? ^a

Response. The energy is $T + V$, but the Lagrangian is $T - V$. The potential term must therefore carry a minus sign. The correction changes the equation from exponential behavior to the intended harmonic oscillation.

^aLecture timestamp: 00:04:00.

Indeed, the Euler–Lagrange equation gives

$$\frac{d}{dt} \frac{\partial L_B}{\partial \dot{\phi}} = \frac{\partial L_B}{\partial \phi}, \quad (1.4)$$

$$\ddot{\phi} = -m^2\phi, \quad (1.5)$$

or

$$\ddot{\phi} + m^2\phi = 0. \quad (1.6)$$

Its frequency is m , not m^2 . Quantizing this oscillator produces the same ladder of states described by $a^\dagger$ and a in Eq. (1.2).

1.3 Why a Fermionic Action Is First Order

The fermionic Lagrangian looks simpler because it contains only one time derivative. The reason is already visible in a deliberately naive trial. For an ordinary commuting variable,

$$L_{\text{trial}} = \phi \dot{\phi} = \frac{1}{2} \frac{d}{dt} \phi^2. \quad (1.7)$$

It is a total derivative. Its Euler–Lagrange equation says only $\dot{\phi} = \dot{\phi}$, so every differentiable path is allowed. Removing a total derivative from the action changes endpoint data but not the bulk equation of motion.

Now rename the variable ψ and declare it Grassmann odd. Both ψ and $\dot{\psi}$ anticommute. With a fixed choice of left variation, reordering the odd factors introduces a minus sign, so the same formal trial gives

$$\frac{d}{dt} \frac{\partial L_{\text{trial}}}{\partial \dot{\psi}} = \dot{\psi}, \quad \frac{\partial L_{\text{trial}}}{\partial \psi} = -\dot{\psi}. \quad (1.8)$$

The equation is therefore

$$\dot{\psi} = -\dot{\psi} \implies \dot{\psi} = 0. \quad (1.9)$$

The expression that was a tautology for a commuting coordinate has acquired dynamical content for an anticommuting one. It has too much content for an oscillator, since it permits only a constant, but this is precisely the clue we need.

Question from the audience. Is the solution simply that ψ is constant? ^a

Response. Yes. Equation (1.9) is a genuine first-order differential equation, and its solution is an arbitrary Grassmann-valued constant. A mass term will turn that static solution into an oscillating one.

^aLecture timestamp: 00:13:57.

The obvious attempt, $m\psi^2$, fails because $\psi^2 = 0$. We need two real odd variables ψ_1, ψ_2 , or equivalently one complex variable and its conjugate. The lecture works through the two-real-variable form on the board, repeatedly checking signs and factors of two. A conventionally normalized complex form is

$$L_F = \frac{i}{2} (\bar{\psi} \dot{\psi} - \dot{\bar{\psi}} \psi) - m \bar{\psi} \psi. \quad (1.10)$$

Up to a total derivative, this is $i\bar{\psi} \dot{\psi} - m\bar{\psi} \psi$. Varying $\bar{\psi}$ gives

$$i\dot{\psi} = m\psi, \quad \psi(t) = \psi(0)e^{-imt}. \quad (1.11)$$

The conjugate equation has the opposite phase. Thus the fermionic mode oscillates with energy m , just as the bosonic mode does, although its equation is first order rather than second order.

¹

¹Editorial clarification: Equation (1.10) fixes one standard ordering and normalization for the exploratory two-real-variable calculation. The lecture itself inserts the factor of i , adjusts a factor of two, and leaves one real-component sign unsettled. These changes do not alter its central conclusion: fermion kinetics are first order, the mass appears linearly, and a complex fermion supplies a nonvanishing bilinear.

This is the zero-dimensional shadow of a familiar relativistic contrast. The Klein–Gordon equation is second order in time, while the Dirac equation is first order. With space suppressed, the boson is one harmonic oscillator and the fermion is one complex Grassmann oscillator.

Question from the audience. Does the order in which ψ , $\dot{\psi}$, and their conjugates are written matter? ^a

Response. The intermediate formulas do depend on ordering because odd factors anticommute. The physical theory does not depend on an arbitrary writing preference, provided one uses a single convention for left or right variation, conjugation, and integration throughout. Reversing an odd pair changes a sign that must be changed everywhere else consistently.

^aLecture timestamp: 00:18:58.

Combining Eqs. (1.3) and (1.10) gives a Lagrangian description of the same matched pair represented by Eq. (1.2). We are now ready to identify the symmetry between them.

1.4 Supercharges and Their Algebra

An operator that removes a fermion and inserts a boson is

$$Q^\dagger = \sqrt{2m} a^\dagger c, \quad (1.12)$$

while its conjugate removes a boson and inserts a fermion,

$$Q = \sqrt{2m} a c^\dagger. \quad (1.13)$$

The normalization is chosen so that the algebra takes its standard form. Using $cc^\dagger = 1 - c^\dagger c$, $aa^\dagger = a^\dagger a + 1$, and the fact that bosonic and fermionic operators commute with one another,

$$\begin{aligned} \{Q, Q^\dagger\} &= 2m (ac^\dagger a^\dagger c + a^\dagger cac^\dagger) \\ &= 2m [(a^\dagger a + 1)c^\dagger c + a^\dagger a(1 - c^\dagger c)] \\ &= 2m (a^\dagger a + c^\dagger c) = 2H. \end{aligned} \quad (1.14)$$

The bosonic operators may be moved through the fermionic ones. The same nilpotence that limits fermion occupation gives

$$Q^2 = (Q^\dagger)^2 = 0. \quad (1.15)$$

Finally,

$$[H, Q] = [H, Q^\dagger] = 0. \quad (1.16)$$

Each supercharge changes the boson number and fermion number in opposite directions, so the two changes in energy cancel when the frequencies are equal.

Question from the audience. What is $[H, Q]$? Is the answer proportional to Q ? ^a

Response. No. The answer is zero. A lone creation or annihilation operator would acquire its energy under commutation with H , but the bosonic and fermionic contributions in Q cancel. That cancellation is exactly the equal-energy condition.

^aLecture timestamp: 00:25:10.

Question from the audience. Was that commutator not already written on the board? ^a

Response. It was. The rediscovery is useful because it forces us to interpret rather than merely copy the relation: commuting with the Hamiltonian means that the transformation is a symmetry and that its generator is conserved.


^aLecture timestamp: 00:25:45.

The adjective *charge* reflects this conservation law. We should not, however, treat Q as an ordinary simultaneously measurable number attached to a state. It is Grassmann odd, nilpotent, and maps a bosonic state into a fermionic one. The useful statement is that it is a conserved generator acting within degenerate supermultiplets. ²

1.5 Generators as Derivatives

The algebra is compact, but by itself it does not tell us how to build interacting theories. The next step is geometric. Familiar symmetry generators already act as differential operators. In the convention of the lecture,

$$H = i \frac{\partial}{\partial t}, \quad P = -i \frac{\partial}{\partial x}. \quad (1.17)$$



Lecture frame: 00:28:12.720.

Figure 1.2: The Hamiltonian represented as the generator of time translations.

Angular momentum is similarly a coordinate times a derivative. If supercharges are odd generators, it is natural to ask whether they can be represented by derivatives with respect to odd coordinates. We therefore enlarge the time axis to the superspace

$$(t, \theta, \bar{\theta}), \quad \theta^2 = \bar{\theta}^2 = 0, \quad \theta \bar{\theta} = -\bar{\theta} \theta. \quad (1.18)$$

Fields on this space depend on an ordinary time and a conjugate pair of Grassmann coordinates.

²Editorial clarification: This wording refines the classroom analogy with labeling states by energy or electric charge. Individual non-Hermitian odd supercharges are not ordinary observables with numerical quantum labels. Their conserved action and the representation of the full superalgebra provide the correct classification.

The lecture deliberately exposes how treacherous the signs are. A sign from complex conjugation, a sign from moving an odd parameter, and a sign from choosing left rather than right derivatives are different operations. Rather than preserve mutually incompatible board stages, we now fix one convention: all odd derivatives are left derivatives, and

$$q = \partial_\theta + i\bar{\theta}\partial_t, \quad \bar{q} = \partial_{\bar{\theta}} + i\theta\partial_t. \quad (1.19)$$

Then

$$q^2 = \bar{q}^2 = 0, \quad \{q, \bar{q}\} = 2i\partial_t = 2H. \quad (1.20)$$

The factor of two comes from two cross terms: $\{\partial_\theta, \theta\} = 1$ and $\{\partial_{\bar{\theta}}, \bar{\theta}\} = 1$. If the generators contained only ∂_θ and $\partial_{\bar{\theta}}$, their anticommutator would vanish and could not reproduce the Hamiltonian.

Question from the audience. Should the two time-derivative terms in q and $\bar{q}$ carry the same sign? ^a

Response. They must be chosen consistently so that the mixed anticommutator is $2H$. Either an overall plus convention or its consistently conjugated alternative can be used. The calculation, not visual intuition about the symbol i , fixes the relative signs.

^aLecture timestamp: 00:31:01.

The room repeatedly checks the signs, and the difficulty becomes a joke: the signs appear to change during the commute to work. The joke is apt. Anticommuting a Grassmann parameter through a coordinate can change a sign even when no physical convention has changed. Equation (1.19) is the fixed reference for everything that follows.

1.5.1 Supersymmetry as a coordinate shift

Let $\xi, \bar{\xi}$ be infinitesimal odd parameters. Acting on a superfield with

$$\delta_\xi = \xi q + \bar{\xi} \bar{q} \quad (1.21)$$

is equivalent, in the convention above, to the infinitesimal coordinate change

$$\theta \longrightarrow \theta + \xi, \quad (1.22)$$

$$\bar{\theta} \longrightarrow \bar{\theta} + \bar{\xi}, \quad (1.23)$$

$$t \longrightarrow t + i(\xi\bar{\theta} + \bar{\xi}\theta). \quad (1.24)$$

The change in a scalar superfield is therefore

$$\delta_\xi \Phi = (\xi\partial_\theta + \bar{\xi}\partial_{\bar{\theta}} + i\xi\bar{\theta}\partial_t + i\bar{\xi}\theta\partial_t) \Phi = (\xi q + \bar{\xi} \bar{q}) \Phi. \quad (1.25)$$

This is the geometric content of supersymmetry. It is a translation in odd directions tied to a translation in ordinary time. Two odd shifts close into a time translation because of Eq. (1.20).

Question from the audience. Could the disputed minus sign represent a motion backward in time? ^a

Response. A different sign convention can reverse the displayed infinitesimal time shift, but no separate physical backward-time process is being introduced. What matters is that the coordinate transformation and the differential generators obey the same algebra.

^aLecture timestamp: 00:58:13.

Question from the audience. Does this extended time have a physical realization, or is it only formal bookkeeping? ^a

Response. It is bookkeeping in one useful sense: it packages the algebra into geometry. But the packaging becomes physically valuable when interactions are introduced. Superspace gives a systematic calculus for constructing Lagrangians that preserve the boson–fermion symmetry at every step, just as tensor calculus helps us construct Lorentz scalars.

^aLecture timestamp: 00:58:23.

1.6 Superfields and the Auxiliary Component

Because there are only two odd coordinates, the expansion of an even superfield terminates:

$$\Phi(t, \theta, \bar{\theta}) = \phi(t) + \bar{\theta}\psi(t) + \bar{\psi}(t)\theta + D(t)\bar{\theta}\theta. \quad (1.26)$$

The coefficients of even monomials, ϕ and D , are bosonic; the coefficients of odd monomials, ψ and $\bar{\psi}$, are fermionic. No θ^2 , $\bar{\theta}^2$, or higher term survives. We leave the time dependence unexpanded because an ordinary Taylor series in t would not terminate and would serve no purpose here.

Question from the audience. What is the component $D(t)$ in the top term? ^a

Response. At this stage it is simply the final bosonic coefficient in the finite expansion. In the component action it normally has no kinetic term and is eliminated by an algebraic equation of motion, so it is called an auxiliary field. The lecture playfully suggests “D for dummy,” which captures its nondynamical role without being the historical definition.

^aLecture timestamp: 00:41:46.

The superfield is one object, while $\phi, \psi, \bar{\psi}, D$ are its components. A supersymmetry transformation mixes them. Thus a boson and a fermion are related in much the same structural sense that the x - and y -components of a vector are related by a rotation, even though Grassmann directions cannot be pictured as ordinary axes.

1.7 The Superaction and the Top Component

To obtain equations of motion we need an action. In ordinary field theory one forms a scalar Lagrangian and integrates it over spacetime. Here the analogous expression is

$$S = \int dt d\theta d\bar{\theta} \Lambda(\Phi, \text{superspace derivatives of } \Phi), \quad (1.27)$$

where Λ is an even superfield. Berezin integration is algebraic. We choose

$$\int d\theta d\bar{\theta} \bar{\theta}\theta = 1. \quad (1.28)$$

Therefore, if

$$\Lambda = A + \bar{\theta}B + \bar{B}\theta + C\bar{\theta}\theta, \quad (1.29)$$

then

$$\int d\theta d\bar{\theta} \Lambda = C. \quad (1.30)$$


The odd integral extracts the top component, leaving an ordinary Lagrangian to be integrated over time.

Why is the result invariant? A supersymmetry variation translates the integration variables. The variation of the top component is an ordinary total derivative in time, while odd total derivatives vanish under Berezin integration. With suitable endpoint conditions, the action is unchanged. This is the superspace analogue of building a Lorentz scalar before integrating it over spacetime.

1.8 Multiplying Superfields

The simplest way to make a new superfield from an old one is to multiply it by itself. Starting from Eq. (1.26), every term containing θ^2 or $\bar{\theta}^2$ vanishes. Keeping the order $\bar{\theta}\theta$ and writing barred component fields to the left gives

$$\begin{aligned}\Phi^2 &= \phi^2 + 2\phi\bar{\theta}\psi + 2\phi\bar{\psi}\theta \\ &\quad + (2\phi D - 2\bar{\psi}\psi)\bar{\theta}\theta.\end{aligned}\tag{1.31}$$



Lecture frame: 01:07:57.844.

Figure 1.3: The component-by-component superfield product and the collection of its $\bar{\theta}\theta$ terms.

The two terms ϕD and $D\phi$ add because both fields are even. The two fermion cross terms also add after each is reordered, but their common sign is negative:

$$(\bar{\theta}\psi)(\bar{\psi}\theta) + (\bar{\psi}\theta)(\bar{\theta}\psi) = -2\bar{\psi}\psi\bar{\theta}\theta.\tag{1.32}$$

Integrating the square over the odd coordinates yields

$$\int d\theta d\bar{\theta} \Phi^2 = 2\phi D - 2\bar{\psi}\psi. \quad (1.33)$$

This term is supersymmetric by construction, but it is not yet a kinetic Lagrangian. It contains no time derivatives.

The board then proposes Φ^3 , Φ^4 , and Φ^n as exercises. There is a compact rule. For any ordinary function W ,

$$W(\Phi)|_{\bar{\theta}\theta} = W'(\phi)D - W''(\phi)\bar{\psi}\psi. \quad (1.34)$$

For $W(\phi) = \phi^2$, this reproduces Eq. (1.31); for $W(\phi) = \phi^n$, it gives

$$\Phi^n|_{\bar{\theta}\theta} = n\phi^{n-1}D - n(n-1)\phi^{n-2}\bar{\psi}\psi. \quad (1.35)$$

3

At this point the lecturer asks whether the room is exhausted. The audience says no; he answers that he is. That exchange is more than comic relief. Powers of superfields are easy because the expansion terminates, but derivatives and constraints are where the sign bookkeeping becomes genuinely demanding.

1.9 Which Derivatives Respect Supersymmetry?

There is one correction to make before following the next board argument. The ordinary time derivative is harmless:

$$[\partial_t, q] = [\partial_t, \bar{q}] = 0. \quad (1.36)$$

Consequently $\partial_t\Phi$ transforms as a superfield whenever Φ does. The subtle objects are the bare odd derivatives $\partial_\theta\Phi$ and $\partial_{\bar{\theta}}\Phi$, because they do not anticommute with both supercharges. We need modified odd derivatives for constraints and for manifestly supersymmetric component constructions.

4

1.9.1 The “magic field” analogy

The lecture motivates the criterion with ordinary rotations. Let $f(x, y)$ satisfy

$$\partial_y f = 0. \quad (1.37)$$

Call such a function a “magic field.” It depends on x but not on y . Is the class of magic fields preserved by rotations? For

$$L_z = -i(x\partial_y - y\partial_x), \quad (1.38)$$

we find

$$[\partial_y, L_z] = i\partial_x \neq 0. \quad (1.39)$$

³Editorial clarification: The lecture assigns the n -th power as an exercise and states the expected kinds of terms. Equation (1.34) supplies the completed rule in the fixed ordering convention.

⁴Editorial clarification: The spoken claim that time differentiation fails to produce another superfield is not correct in the convention used here. Since ∂_t commutes with both supercharges, it is covariant. The subsequent construction of modified odd derivatives remains necessary and is the actual mathematical point of the lecture.

Therefore $L_z f$ generally acquires y -dependence even when f initially obeys Eq. (1.37). The constraint is not rotationally invariant.

The general lesson is simple. If a field obeys $C\Phi = 0$ and a symmetry is generated by G , the transformed field remains constrained when the graded commutator of C and G vanishes. For two odd operators, the graded commutator is an anticommutator.

Question from the audience. When is a constraint on a field consistent with a symmetry? ^a

Response. When the constraint operator graded-commutes with the symmetry generators. Then $C(G\Phi)$ can be moved past G and reduced to $G(C\Phi) = 0$. The rotational example fails because ∂_y does not commute with the rotation generator.

^aLecture timestamp: 01:17:56.

1.9.2 Covariant odd derivatives

With the supercharges fixed by Eq. (1.19), define

$$\mathcal{D} = \partial_\theta - i\bar{\theta}\partial_t, \quad \bar{\mathcal{D}} = \partial_{\bar{\theta}} - i\theta\partial_t. \quad (1.40)$$

The only change is the sign of the time-derivative term. Direct calculation gives

$$\{\mathcal{D}, q\} = \{\mathcal{D}, \bar{q}\} = \{\bar{\mathcal{D}}, q\} = \{\bar{\mathcal{D}}, \bar{q}\} = 0, \quad (1.41)$$

$$\{\mathcal{D}, \bar{\mathcal{D}}\} = -2i\partial_t. \quad (1.42)$$

The two potentially nonzero cross terms cancel. This is the odd analogue of a covariant derivative: it differentiates while preserving the transformation law.

The lecture uses D both for the auxiliary component and for this derivative. We reserve $D(t)$ for the component and use calligraphic $\mathcal{D}$ for the operator.

1.10 The Chiral Constraint

We can now impose a nontrivial constraint that survives supersymmetry:

$$\bar{\mathcal{D}}\Phi = 0. \quad (1.43)$$

Such a field is called chiral. If it is chiral before a supersymmetry transformation, then

$$\bar{\mathcal{D}}(q\Phi) = -q(\bar{\mathcal{D}}\Phi) = 0, \quad \bar{\mathcal{D}}(\bar{q}\Phi) = -\bar{q}(\bar{\mathcal{D}}\Phi) = 0. \quad (1.44)$$

Thus the entire transformed superfield remains inside the constrained class. Unlike the magic-field condition, chirality is compatible with the symmetry.

To solve the constraint, define a shifted time coordinate

$$\tau = t + i\bar{\theta}\theta. \quad (1.45)$$

Because we use left derivatives,

$$\partial_{\bar{\theta}}\tau = i\theta, \quad (1.46)$$

and hence

$$\bar{D}F(\tau, \theta) = (\partial_{\bar{\theta}} - i\theta\partial_t)F(\tau, \theta) = i\theta\partial_\tau F - i\theta\partial_\tau F = 0. \quad (1.47)$$

The general local chiral superfield therefore depends on $\bar{\theta}$ only through τ :

$$\Phi_{\text{ch}}(\tau, \theta) = \phi(\tau) + \bar{\psi}(\tau)\theta. \quad (1.48)$$

Question from the audience. Is dependence on τ and θ the only form that satisfies the chiral constraint?^a

Response. Locally, yes. The differential equation $\bar{D}\Phi = 0$ says precisely that the independent $\bar{\theta}$ -dependence can be absorbed into the shifted coordinate $\tau = t + i\bar{\theta}\theta$. The remaining finite expansion is a function of τ plus one term proportional to θ .

^aLecture timestamp: 01:32:35.

Re-expanding in the original time coordinate,

$$\phi(\tau) = \phi(t) + i\bar{\theta}\theta\dot{\phi}(t), \quad (1.49)$$

$$\bar{\psi}(\tau)\theta = \bar{\psi}(t)\theta, \quad (1.50)$$

because the correction to the second line contains θ^2 . Thus

$$\Phi_{\text{ch}}(t, \theta, \bar{\theta}) = \phi(t) + \bar{\psi}(t)\theta + i\bar{\theta}\theta\dot{\phi}(t). \quad (1.51)$$

Question from the audience. Should the Taylor expansion of $\phi(t + i\bar{\theta}\theta)$ contain a second-order term?^a

Response. No. The first correction is $i\bar{\theta}\theta\dot{\phi}$, and every higher correction contains $(\bar{\theta}\theta)^2 = 0$. The audience catches this immediately; nilpotence has already supplied the first two terms and terminated the series.

^aLecture timestamp: 01:33:51.

This is the economy of chirality. The unconstrained superfield in Eq. (1.26) has two bosonic and two fermionic component slots. The chiral condition removes the independent auxiliary slot in this minimal one-dimensional example and ties the top component to $\dot{\phi}$.

1.11 The Calculation That Properly Stops

The natural next move is to combine the chiral field with its conjugate and integrate over superspace. The conjugate expansion is started, and an audience member asks whether the order $\theta\bar{\theta}$ must reverse under conjugation. At that point the signs no longer look trustworthy. Rather than bluff, the calculation stops.

Question from the audience. Can we have the punchline even if the conjugate-field multiplication is deferred?^a

Response. Yes. With the signs and conjugations handled consistently, a superspace expression built from a chiral field and its conjugate produces the desired ordinary bosonic and fermionic kinetic terms. The important result for this lecture is the construction principle, not an unreliable final line copied from a tired calculation.

^aLecture timestamp: 01:37:30.

The stopping point should remain visible. The formalism is unforgiving: one misplaced odd factor changes a sign, and the same convention must survive conjugation, differentiation, multiplication, and integration. The correct response to losing that chain is to repair it before proceeding. In a standard convention the familiar kinetic action is built from a conjugate product, but its detailed component reduction belongs to the next clean calculation, not to a reconstruction of an unfinished board line.

1.12 Why the Symmetry Is the Punchline

A supersymmetric Lagrangian does more than place a boson and a fermion side by side. It constrains their masses and interactions so that they form a representation of the superalgebra. In an exactly supersymmetric vacuum, members of a supermultiplet have equal masses. Their loop contributions are correlated, and bosonic and fermionic ultraviolet effects cancel in precisely the combinations enforced by the symmetry.

This is why the construction machinery matters. One could list Feynman vertices and verify cancellations diagram by diagram, but the cancellations would then look accidental. Supersymmetry explains them in the same way Lorentz symmetry explains why scattering amplitudes transform consistently in every inertial frame.

5

Question from the audience. How is “chiral” spelled? ^a

Response. C-H-I-R-A-L. The word labels the constraint $\bar{D}\Phi = 0$; it is not the ordinary geometric handedness of an object in three-dimensional space, although the terminology descends from the left- and right-handed structure of relativistic spinors.

^aLecture timestamp: 01:39:48.

Question from the audience. Could one bypass the Lagrangian and go directly to supersymmetric Feynman rules? ^a

Response. One can certainly state the rules, but then the relations among couplings and the cancellations they produce look like unexplained coincidences. The deeper content is the generalized spacetime symmetry. The Lagrangian or an equivalent invariant formulation makes that reason visible and generates the rules systematically.

^aLecture timestamp: 01:40:00.

Suppose every perturbative order happened to give Lorentz-invariant amplitudes although no Lorentz transformations had been mentioned. After enough calculations, we would demand an explanation. The answer would be the spacetime symmetry that organizes all orders at once. The role of superspace is analogous: shifts in $\theta, \bar{\theta}$ mix bosonic and fermionic components, and their algebra closes on translations in spacetime.

⁵Editorial clarification: Exact supersymmetry does not imply that every individual self-energy correction vanishes. It relates renormalizations and removes particular ultraviolet sensitivities; nonrenormalization statements depend on the quantity, dimension, and amount of supersymmetry. The lecture’s broad cancellation claim is retained in this qualified form.

Question from the audience. Should ordinary spatial coordinates also be expanded in a finite power series, as the Grassmann coordinates are? ^a

Response. No. A power series in an ordinary coordinate does not terminate and is not usually the natural decomposition. A Fourier transform is more useful because it resolves a field into waves and momenta. Expansion in θ is special precisely because nilpotence makes it finite.

^aLecture timestamp: 01:44:55.

Question from the audience. If a new particle appeared at the LHC, how could we tell that it was a superpartner rather than merely an unknown particle? ^a

Response. Supersymmetry predicts correlated quantum numbers and interactions, not just one extra state. Spins differ by one half, gauge charges are related, and many cubic and quartic couplings are governed by fewer independent parameters than in a generic theory. Cross sections, branching fractions, and the rest of the spectrum would have to satisfy those relations.

^aLecture timestamp: 01:45:42.

One unfamiliar resonance would not by itself prove supersymmetry. A network of coupling and spectrum relations would. That is the experimental counterpart of the superspace construction: apparently independent component interactions descend from one invariant object.

Question from the audience. Observed particles and their hypothetical superpartners do not have equal masses. How does that change the theory? ^a

Response. It changes it enormously. Supersymmetry must be broken, usually through spontaneous breaking communicated to the observed sector. The breaking splits multiplet masses while retaining enough of the structure to control interactions and, if the scale is low enough, ultraviolet corrections. Working out that mechanism is a substantial theory of its own, especially when gravity is included.

^aLecture timestamp: 01:47:01.

1.13 Simple Principle, Difficult Calculus

The closing assessment is unusually candid. The starting idea sounds simple: bosons and fermions are related by a symmetry. Yet implementation rapidly becomes a discipline of indices, conjugations, Grassmann orderings, and signs. Even after constructing the symmetry, realistic particle physics requires breaking it; coupling it to gravity leads toward supergravity and adds another layer of structure.

This difficulty is not evidence that the idea is empty. Nature is under no obligation to make a simple principle easy to calculate with. Specialists learn the subject by doing the details repeatedly until the sign rules become second nature. The frustration comes from the distance between the compact algebra

$$\{Q, Q^\dagger\} = 2H \tag{1.52}$$

and the amount of machinery required to realize it in an interacting theory.

Question from the audience. Does another layer of abstraction eventually simplify the formalism, or does the complication keep accumulating? ^a

Response. No general simplification has removed the detailed work. The underlying rules may be short, but using them reliably requires repeated control of the conventions. That is why a full treatment normally occupies an entire technical course rather than a fraction of a broad particle-physics course.

^aLecture timestamp: 01:49:33.

Question from the audience. Do the transformations tell us what fermions and bosons really are, or why their ordinary behavior is different? ^a

Response. They tell us that the two are components of one superfield and images of one another under the supercharges. Acting with Q maps a bosonic component to a fermionic one, while the next transformation produces derivatives and auxiliary components. This is analogous to a rotation mixing vector components, but it cannot be translated into an equally literal picture: the appropriate language is the Grassmann geometry itself.

^aLecture timestamp: 01:51:42.

That answer returns us to the central idea. A supersymmetry transformation is a small motion in superspace. It mixes $\phi, \psi, \bar{\psi}, D$ just as an ordinary rotation mixes components, and it does so while closing on a translation in time. The picture is abstract, but it is not arbitrary: every term in the superfield, every sign in the covariant derivatives, and every relation among component interactions exists to realize the algebra.

1.14 What We Have Built

We began with the matched oscillator Hamiltonian and recovered its two different Lagrangian descriptions. The boson obeys a second-order oscillator equation; the fermion requires a first-order Grassmann action and a complex pair of odd components. The supercharges exchange the two modes, square to zero individually, anticommute to the Hamiltonian, and commute with it.

Representing those charges as differential operators led to superspace. A finite superfield expansion assembled bosonic, fermionic, and auxiliary components; Berezin integration selected the top component of a super-Lagrangian; and multiplication generated invariant interaction candidates. The demand for symmetry-compatible odd differentiation then led to $\mathcal{D}, \bar{\mathcal{D}}$, the chiral constraint, and the shifted coordinate $\tau = t + i\bar{\theta}\theta$.

The unfinished conjugate-product calculation is part of the lesson rather than a defect to hide. Supersymmetry begins with a simple physical demand, but its calculus is sensitive to every ordering choice. Once those choices are controlled, the reward is powerful: masses and couplings are organized by symmetry, quantum corrections satisfy nonaccidental relations, and bosons and fermions become components of one geometric object.