

Chris Henson
Project Euler Questions 137/140

This short write-up will walk through some of the mathematical derivations that I used to solve Project Euler questions 137/140.

First, let us recall the definition of the Fibonacci numbers. We start with:

$$F_0 = 0, \quad F_1 = 1$$

and define further numbers by the relation:

$$F_n = F_{n-1} + F_{n-2}$$

This gives us the first few terms:

$$0, 1, 1, 2, 3, 5, 8, 13, 21, 34, 55, 89, 144, \dots$$

(Please note that Project Euler omits the first index, but this will not change our calculations in any meaningful way, and I will try to be as explicit as possible about this.)

The natural generalization is to consider a different starting point. I will write

$$G(a, b, n) = G(a, b, n - 1) + G(a, b, n - 2)$$

given the starting point:

$$G(a, b, 0) = a, \quad G(a, b, 1) = b$$

and when we are working with a set a and b I will simply write G_n . You can see that $G(0, 1, n)$ gives the Fibonacci numbers themselves.

These two Project Euler questions are concerned with the power series with coefficients given by the above sequences. That is, we are interested in the value of the summations:

$$\sum_{k=1}^{\infty} G(a, b, n)x^k = G(a, b, 1)x + G(a, b, 2)x^2 + G(a, b, 3)x^3 + \dots$$

Again, if we take $a = 0$ and $b = 1$, we will have the Fibonacci numbers:

$$\begin{aligned} \sum_{k=1}^{\infty} G(0, 1, n)x^k &= F_1x + F_2x^2 + F_3x^3 + \dots \\ &= x + x^2 + 2x^3 + 3x^4 + 5x^5 \dots \end{aligned}$$

Interestingly, it is not too difficult to derive a closed form solution to these series. Let us fix $G_0 = a$ and $G_1 = b$. Then the power series is:

$$\sum_{k=0}^{\infty} G_k x^k = G_0 + G_1 x + G_2 x^2 + \dots$$

Please note that I have included the initial term G_0 that Project Euler omits. We can correct for this at the end by simply subtracting this term.

First, I will pull out the first two given terms from the series:

$$\sum_{k=0}^{\infty} G_k x^k = G_0 + G_1 x + \sum_{k=2}^{\infty} G_k x^k$$

Now I can Replace G_k with the relation that defines it, and split apart the summation further:

$$\begin{aligned} \sum_{k=0}^{\infty} G_k x^k &= G_0 + G_1 x + \sum_{k=2}^{\infty} (G_{k-1} + G_{k-2}) x^k \\ &= G_0 + G_1 x + \sum_{k=2}^{\infty} G_{k-1} x^k + \sum_{k=2}^{\infty} G_{k-2} x^k \end{aligned}$$

Next, I notice that multiplying my original series by x and x^2 gives:

$$\begin{aligned} x \sum_{k=0}^{\infty} G_k x^k &= G_0 x + G_1 x^2 + G_2 x^3 + \dots \\ x^2 \sum_{k=0}^{\infty} G_k x^k &= G_0 x^2 + G_1 x^3 + G_2 x^4 + \dots \end{aligned}$$

Substituting these into the above equation I have:

$$\sum_{k=0}^{\infty} G_k x^k = G_0 + G_1 x + x \sum_{k=0}^{\infty} G_k x^k - G_0 x + x^2 \sum_{k=0}^{\infty} G_k x^k$$

Denoting the original summation as $s(x)$ this is:

$$s(x) = G_0 + G_1x + xs(x) - G_0x + x^2s(x)$$

and solving gives:

$$s(x) = \frac{G_0 - G_0x + G_1x}{1 - x - x^2}$$

Remember that this includes our first term. So we have¹:

$$\sum_{k=0}^{\infty} G_k x^k = G_0 + G_1x + G_2x^2 + \dots = \frac{G_0 - G_0x + G_1x}{1 - x - x^2}$$

For the Fibonacci series we set $G_0 = 0$ and $G_1 = 1$ giving:

$$\sum_{k=0}^{\infty} F_k x^k = F_0 + F_1x + F_2x^2 + \dots = \frac{x}{1 - x - x^2}$$

1

Note Project Euler omits the first term, considering instead the series:

$$\sum_{k=1}^{\infty} G_k x^k = G_1x + G_2x^2 + \dots = \frac{G_0 - G_0x + G_1x}{1 - x - x^2} - G_0$$

The next question under consideration is which rational values of x give an integer value for our power series? I will write:

$$s(x) = \sum_{k=1}^{\infty} G_k x^k = G_0 + G_1 x + G_2 x^2$$

Above we had the equation:

$$s(x) = G_0 + G_1 x + x s(x) - G_0 x + x^2 s(x)$$

which solving for x using the quadratic formula² gives us:

$$x = \frac{-(G_1 - G_0 + s) + \sqrt{(G_1 - G_0 + s)^2 - 4s(G_0 - s)}}{2s}$$

Remember that we included the term G_0 . If we would like to correct for this, we can make the transformation $s \rightarrow S - G_0$ where:

$$\begin{aligned} S(x) &= \sum_{k=1}^{\infty} G_k x^k = \\ &= \sum_{k=0}^{\infty} G_k x^k - G_0 \\ &= s(x) - G_0 \end{aligned}$$

which gives instead:

$$x = \frac{-(G_1 + S) + \sqrt{(G_1 + S)^2 - 4(S + G_0)(-S)}}{2(S + G_0)}$$

²We actually have two choices for roots, but one is outside the radius of convergence for the power series

Using the second equation and taking $G_0 = 0$ and $G_1 = 1$ to give the Fibonacci power series, we have have the following values:

$$\begin{aligned} S(\sqrt{2} - 1) &= 1 \\ S\left(\frac{1}{2}\right) &= 2 \\ S\left(\frac{\sqrt{13} - 2}{3}\right) &= 3 \\ S\left(\frac{\sqrt{89} - 5}{8}\right) &= 4 \\ &\vdots \end{aligned}$$

We can see for instance that $S = 2$ has the desired property that x is rational. What conditions must be met for this? First, look at our equation for x we we substitue the particular values for G_0 and G_1 :

$$x = \frac{-(1 + S) + \sqrt{(1 + S)^2 - (2S)^2}}{2S}$$

This will be rational exactly when the discriminant $(1 + S)^2 - (2S)^2$ is a perfect square ³

³The following arguement follows from this article : [Link](#)

Since the discriminant is itself a sum of two squares, we can consider them as Pythagorean triples so that for integers m, n :

$$S + 1 = m^2 - n^2, \quad 2S = 2mn$$

This gives the equation:

$$m^2 - n^2 = mn + 1$$

which can be written as

$$(2m - n)^2 = 4 + 5n^2$$

We need $4 + 5n^2$ to be a perfect square to ensure a rational x . Unexpectedly, using $n = F_{2j}$ (where j is a positive integer) ensures this is satisfied:

$$\begin{aligned} 4 + 5n^2 &= 4 + 5F_{2j}^2 = 4 + 5 \left(\frac{(\varphi^{2j} - \psi^{2j})^2}{5} \right) \\ &= 4 + \varphi^{4j} - 2(\varphi\psi)^{2j} + \psi^{4j} \\ &= \varphi^{4j} + 2 + \psi^{4j} \\ &= \varphi^{4j} + 2(\varphi\psi)^{2j} + \psi^{4j} \\ &= (\varphi^{2j} + \psi^{2j})^2 \end{aligned}$$

Furthermore, substituting into the equation $(2m - n)^2 = 4 + 5n^2$ above gives:

$$\begin{aligned} m &= \frac{1}{2}(\varphi^{2j} + \psi^{2j} + F_{2j}) \\ &= \frac{1}{2} \left(\varphi^{2j} + \psi^{2j} + \frac{\varphi^{2j} - \psi^{2j}}{\sqrt{5}} \right) \\ &= \frac{\varphi\varphi^{2j} - \psi\psi^{2j}}{\sqrt{5}} = \frac{\varphi^{2j+1} - \psi^{2j+1}}{\sqrt{5}} \\ &= F_{2j+1} \end{aligned}$$

Finally, substituting this back into the original series gives:

$$\sum_{k=1}^{\infty} F_k (F_{2j}/F_{2j+1})^k = F_{2j} F_{2j+1}, \quad j \geq 1$$

What an unexpected result! That we could consider successive coefficients in identifying these rational points is certainly not intuitive. We see that these numbers also grow very quickly:

$$F_{2j} F_{2j+1}, n \in \mathbb{Z}^+ = \{2, 15, 104, 714, 4895, 33552, 229970, 1576239, 10803704, 74049690, \\ 507544127, 3478759200, 23843770274, 163427632719, 1120149658760, \dots\}$$

The last number in this list is the solution to problem 137. Problem 140 considers the initialization $G_0 = 3$ and $G_1 = 1$, which give a more complicated sort of equation that determines the rational points. Still, they can be written entirely in terms of the Fibonacci series.